

# Freshwater treatment of amoebic gill disease and sea-lice in seawater salmon production: considerations of water chemistry and fish welfare



**Main Office**

Gaustadalléen 21  
 NO-0349 Oslo, Norway  
 Phone (47) 22 18 51 00  
 Telefax (47) 22 18 52 00  
 Internet: www.niva.no

**NIVA Region South**

Jon Lilletuns vei 3  
 NO-4879 Grimstad, Norway  
 Phone (47) 22 18 51 00  
 Telefax (47) 37 04 45 13

**NIVA Region East**

Sandvikaveien 59  
 NO-2312 Ottestad, Norway  
 Phone (47) 22 18 51 00  
 Telefax (47) 62 57 66 53

**NIVA Region West**

Thormøhlens gate 53 D  
 NO-5006 Bergen Norway  
 Phone (47) 22 18 51 00  
 Telefax (47) 55 31 22 14

**NIVA Region Mid-Norway**

Høgskoleringen 9  
 NO-7034 Trondheim  
 Phone (47) 22 18 51 00  
 Telefax (47) 73 54 63 87

Title <b>Freshwater treatment of amoebic gill disease                  and sea-lice in seawater salmon production:                  considerations of water chemistry and fish                  welfare.</b>	Report No.. <b>6632-2014</b>	Date 19.02.2014
	Project No. 13392	Pages      Price 38
Author(s) Powell M.D. and Kristensen, T.	Topic group Fish health	Distribution Open
	Geographical area	Printed NIVA

Client(s) Fiskeri og Havbruksnæringens Forskningsfond	Client ref.
--	-------------

Abstract

Amoebic gill disease (AGD) and sea lice are two of the most significant disease issues facing the Norwegian Atlantic salmon aquaculture industry. Although both diseases respond to various extents, to freshwater treatment, the chemistry, interactions and efficacy of treatment can be variable. These variations can have significant impacts upon the success and failure of treatment and costs to the production cycle. Although it is known that soft freshwater is most effective in bathing of Atlantic salmon with AGD and that most of the freshwaters in Norway fall into the soft category, the low alkalinity and buffering capacity of such water s may impact on the pH and metal toxicity of the water source in use. Similarly dissolved organic carbon can be beneficial in treatment, although sequestration of metal ions can be reversed as the water pH drops due to high densities of fish and accumulations of carbon dioxide. Alternative treatments such as the use of oxidative disinfectants such as hydrogen peroxide used for AGD and sea lice control may have potential although the interactions in seawater with organic loads and dissolved organic carbon are unclear. Similarly the use of oxidative disinfectants in freshwater will depend upon the water chemistry and interactions with treatment chemicals, fish and water organic content. It is recommended that best practice model is required to understand the interactions of water chemistry, fish loading and treatment chemicals in the efficacy of treatments for AGD and sea lice under Norwegian conditions.

4 keywords, Norwegian

1. Amøbisk Gjelle Sykdom
2. Ferskvann
3. Lakselus
4. Vannkjemi og vannkvalitet

4 keywords, English

1. Amoebic Gill Disease
2. Freshwater
3. Sea Lice
4. Water chemistry and quality

Project Manager

Research Manager

ISBN 978-82-577-6367-1

**Freshwater treatment of amoebic gill disease and  
sea-lice in seawater salmon production:  
considerations of water chemistry and fish welfare.**

## Preface

Through dialogue with representatives from the aquaculture industry and FHL, the need for a review on current knowledge about freshwater treatment against AGD and sea-lice was identified. NIVAs current expertise on water quality in aquaculture, and especially the hands-on experience with AGD treatments was therefore collected and supplemented with information from the scientific literature in this report.

We hope this document will help in improving the efficiency of such treatments, and thank FHF for the assignment.

Bodø, 19 February 2014



Torstein Kristensen

# Contents

<b>Sammendrag</b>	<b>6</b>
<b>1. Introduction</b>	<b>7</b>
<b>2. Norwegian freshwater quality</b>	<b>8</b>
2.1 Chemistry of natural water sources	8
2.2 Metabolite accumulation: effects on water chemistry	11
<b>3. Amoebic Gill disease</b>	<b>13</b>
3.1 General description	13
3.2 Approaches to control of AGD – practical limitations	15
3.3 Freshwater treatment of AGD	16
3.4 Oxidative disinfectants for AGD control in seawater	18
<b>4. Sea lice</b>	<b>20</b>
4.1 General description	20
4.2 Approaches to control of sea lice – practical limitations	21
4.3 Freshwater treatment of sea lice	21
4.4 Oxidative disinfectants for control of sea lice in seawater	21
<b>5. Implications for fish welfare</b>	<b>23</b>
<b>6. Conclusions</b>	<b>23</b>
<b>7. Recommendations</b>	<b>24</b>

---

## Summary

Amoebic gill disease (AGD) and sea lice are two of the most significant disease issues facing the Norwegian Atlantic salmon aquaculture industry. Although both diseases respond to various extents, to freshwater treatment, the chemistry, interactions and efficacy of treatment can be variable. These variations can have significant impacts upon the success and failure of treatment and costs to the production cycle. Although it is known that soft freshwater is most effective in bathing of Atlantic salmon with AGD and that most of the freshwaters in Norway fall into the soft category, the low alkalinity and buffering capacity of such waters may impact on the pH and metal toxicity of the water source in use. Similarly dissolved organic carbon can be beneficial in treatment, although sequestration of metal ions can be reversed as the water pH drops due to high densities of fish and accumulations of carbon dioxide. Alternative treatments such as the use of oxidative disinfectants such as hydrogen peroxide used for AGD and sea lice control may have potential although the interactions in seawater with organic loads and dissolved organic carbon are unclear. Similarly the use of oxidative disinfectants in freshwater will depend upon the water chemistry and interactions with treatment chemicals, fish and water organic content. It is recommended that best practice model is required to understand the interactions of water chemistry, fish loading and treatment chemicals in the efficacy of treatments for AGD and sea lice under Norwegian conditions.

## Sammendrag

Amøbisk gjellesykdom (AGD) og lakselus er to av de mest alvorlige sykdomsproblemene for norsk lakseoppdrett. Selv om begge sykdommene responderer på ferskvannsbehandling, vil varierende vannkjemien og ulike interaksjoner påvirke effekten av behandlingen. Disse faktorene kan ha vesentlige innvirkning på behandlingskostnadene og være avgjørende om behandlingen blir en suksess eller fiasko. Det er kjent at bløtt ferskvann er mest effektiv i bading av atlantisk laks med AGD. De fleste vannkildene i Norge havner i denne kategorien, hvor lav alkalitet og lav bufferkapasiteten er styrende for pH-verdien og metall toksisiteten i den aktuelle vannkilden. Mengden oppløst organisk karbon i ferskvannet kan og påvirke behandlingseffektiviteten. Det har vært vist at noe organisk karbon kan være fordelaktig, selv om partikkelbundet metallioner kan frigjøres ved endret pH og potensielt akkumuleres på gjellene. Alternative behandlinger som for eksempel bruk av oksidative desinfeksjonsmidler (bla. hydrogenperoksid) brukes mot AGD og lakselus kan ha et potensial selv om flere interaksjoner i sjøvann (organisk belastning) er uklare. Tilsvarende anvendelse av oksidative desinfeksjonsmidler i ferskvann vil være avhengig av vannkjemien og interaksjoner med behandling kjemikalier, fisk og organisk innhold i vannet. Det anbefales at det benyttes en beste praksis modell for økt behandlingseffektivitet. Samtidig er det avgjørende å forstå samspillet mellom vannkjemien, lasting og lossing av fisk, og bruk av eventuelle kjemikalier mot AGD og lakselus under norske forhold.

### List of abbreviations

AGD	Amoebic gill disease
CO <sub>2</sub>	Carbon dioxide
TAN	Total ammonia nitrogen
SW	Sea water
FW	Fresh water
TOC	Total organic carbon
DOC	Dissolved organic carbon
pK <sub>a</sub>	Dissociation constant of acid-base reaction
NIVA	Norwegian Institute for Water Research
PCR	Polymerase chain reaction



# 1. Introduction

The use of short-term freshwater treatments of Atlantic salmon during the marine phase of the production cycle has increased drastically in Norway. The increased occurrence of Amoebic Gill Disease (AGD) and infections with resistant/multiresistant strains of sea-lice has caused this development. If this treatment strategy is to be developed and applied in the industry, a number of issues concerning water quality on fish welfare and treatment efficiency needs to be addressed, and knowledge-gaps identified. Performing such treatments as part of the production requires at the very least some rule-of-thumb, and in the longer run more stringent guidelines in order to adhere to legislation and internal company guidelines is required. The Norwegian legislation (Lov om Dyrevelferd, LOV-2009-06-19-97, § 8) states:

*«Dyreholder skal påse at driftsformer, metoder, utstyr og tekniske løsninger som brukes til dyr, er egnet til å ivareta hensynet til dyrenes velferd»*

*«Den som markedsfører eller omsetter nye driftsformer, metoder, utstyr og tekniske løsninger til bruk på dyr eller i dyrehold, skal påse at disse er utprøvd og funnet egnet ut fra hensynet til dyrevelferd»*

Subjecting seawater (SW, hyperosmotic) adapted teleost fish to a procedure combining abrupt transfer to a hypoosmotic freshwater (FW) environment at high fish densities, crowding and handling is a procedure likely to cause a degree of stress in the fish. Maintaining fish in these conditions also cause metabolite accumulation (CO<sub>2</sub>/TAN) in the water with subsequent water quality changes that may further aggravate this stress. Thus, a fundamental knowledge about the effects of FW treatment on stress and physiology alone, and combined with water quality changes is needed to ensure fish welfare and optimal treatment effect. Treatment efficacy may also be influenced by the quality and chemistry of freshwater used, as well as metabolite accumulation associated with the various chemical processes. This document aims at summarizing current knowledge on the subject, and providing preliminary guidelines to ensure treatment efficacy and animal welfare during FW treatments.

## 2. Norwegian freshwater quality

### 2.1 Chemistry of natural water sources

Norwegian surface waters are characterized by being soft, i.e. having a low bicarbonate buffering capacity and low  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  content (Henriksen et al. 1989; Skjelkvåle et al. 2007, Kristensen et al. 2009) (Fig. 1). High precipitation rates and low evaporation due to temperate climate, combined with acidic and weathering-resistant bedrock give rise to this chemical composition of the surface waters. Water pH is therefore also naturally low in many sites (Fig 1), with additional reduction caused by acidification in the southern and south-western regions (Skjelkvåle et al. 2005). Two major concerns arises from a low buffering capacity and/or pH, namely a strong further pH decrease when  $\text{CO}_2$  accumulates in the water and an increased gill permeability caused by low Ca saturation of ion channels in the gills (Evans et al. 2005). The first may cause mobilization of metal ions (if present) (Fivelstad et al. 2003a), while the latter results in increased susceptibility to metals (Leivestad et al. 1980). Low pH increases the efflux of Na and Cl across the gill surface due to an osmotic gradient of about  $350 \text{ mOsm L}^{-1}$  between the fish and the freshwater environment (Fig. 2). This problem is exacerbated by  $\text{H}^+$  ions competing for gill binding sites with  $\text{Ca}^{2+}$  (Pagenkopf 1983; Wilson 2012). Additionally, metals such as Al may be mobilized to gill reactive forms.




Figure 1. Range ( $\pm$  min/max) of  $\text{Ca}^{2+}$  and  $\text{Na}^+$  concentrations in Norwegian fresh waters. Adapted from Kristensen et al. (2009).




Figure 2. Range ( $\pm$  min/max) of water pH and alkalinity of Norwegian freshwaters. Adapted from Kristensen et al. (2009).

Total organic carbon (TOC) levels are, in general, relatively high in Norwegian water sources with a high degree of variability (Fig. 3). Fulvic acids in TOC of humic origin contribute to the low water pH in Norway (Lydersen et al., 2002), and also contribute to transport of associated metals. Metals bound to humic substances are generally less bioavailable than low molecular weight metal species (also denoted free metal ions), and elevated TOC may thus serve to protect fish from harmful effects of metals provided remobilization is not enhanced by decreased pH (Rosseland and Staurnes 1994; Andren et al. 2006), and/or increased ionic strength (Bjerknes et al. 2003; Teien et al. 2006a) in the rearing water.




Figure 3 range ( $\pm 1$  SD) of total organic carbon (TOC) and aluminium concentration in Norwegian fresh water.

Water chemistry is important in the survival of Atlantic salmon smolts when stressed by other pathogens such as sea lice with the main focus studies to date being acidification of freshwater and its associated implication with the mobility of toxic transitional metal ion species such as  $Al^{3+}$  (Finstad et al. 2007, 2012) (Fig. 4). In particular the episodic and fluctuating effects of acidified freshwater enhances the stress effects and reduced survival of post-smolts infected with sealice (Finstad et al. 2012). Not all freshwater can be deemed suitable or optimal for the treatment of Atlantic salmon is a parasite control regime.




Figure 4. Gill bound aluminium concentrations ( $\pm 1$  SD) with respect to salinity in Atlantic salmon. Adapted from Kroglund et al. (2011)

## 2.2 Metabolite accumulation: effects on water chemistry

Oxygen levels in treatment water must be maintained by addition, and it is vital to maintain levels above 80% saturation. In the following discussion on metabolites, adequate oxygenation is assumed.

Carbon dioxide ( $\text{CO}_2$ ) is generated as the end product of aerobic metabolism in a theoretical molar ratio of 1-0,7 to consumed oxygen. In practice, and in an aquaculture setting, about 1.1 g  $\text{CO}_2$  is produced for each mg  $\text{O}_2$  consumed (Fivelstad and Binde 1994). The solubility of  $\text{CO}_2$  in water and body fluids is very high due to reaction with water and generation of  $\text{HCO}_3^-$  and  $\text{H}^+$  (the bicarbonate system). Reactions of the bicarbonate system is described (simplified) below (Equation 1)



The amount of  $\text{CO}_2$  dissolved in water or blood is through the  $\text{H}^+$  generation a strong determinant of pH. While solubility is high (30 times more soluble than  $\text{O}_2$ ), the gas tension of  $\text{CO}_{2(g)}$  is low in equilibrium conditions due to low atmospheric partial pressure ( $\sim 0,04\%$ , 0,04 kPa). However, the relative amount of excreted  $\text{CO}_2$  that is converted to  $\text{HCO}_3^-$  in water is strongly dependent on pH. The pKa of the first equilibrium-reaction of the bicarbonate system (Equation 1) is about 6.4. This means that a balance between  $\text{CO}_{2(g)}$ , which is the primary concern, and  $\text{HCO}_3^-$  vary

substantially in the low range of pH and buffering capacity values observed. In closed aquaculture transport/treatment systems this has to be accounted for when determining safe biomass/treatment durations.

Gaseous CO<sub>2</sub> accumulation over time in closed treatment systems causes pH depression through H<sup>+</sup>/HCO<sub>3</sub><sup>-</sup> generation and elevated CO<sub>2(g)</sub> tension, termed hypercapnia. External hypercapnia forces blood CO<sub>2(g)</sub> and HCO<sub>3</sub><sup>-</sup> levels to increase in order to maintain excretion by diffusion across the gills (Wood and Jackson 1980; Perry and Gilmore 2006). Where proliferative gill disease occurs, the accumulation of CO<sub>2(g)</sub> in the blood is already increased due to diffusion limited CO<sub>2</sub> excretion (see Powell and Perry 1999, Powell 2006; 2007), thus under conditions of hypercapnia, the resulting respiratory acidosis (drop in blood pH due to accumulations of CO<sub>2(g)</sub>) is even further enhanced. The rate of CO<sub>2</sub> accumulation is dependent on water volume to biomass ratio, and the metabolic rate of the fish. Metabolic rate depends on temperature, fish size and the state (stress, active swimming) of the fish. For practical purposes, a maximal metabolic rate at a given temperature and fish size should be assumed in bath treatments to provide a safety factor when calculating biomass loading. Equations given in Thorarensen and Farrell (2011) are recommended used for this purpose. In order to calculate CO<sub>2</sub> accumulation over time in closed transport systems (FW and SW), a model based on empirical data has been established (NIVA). The resulting need for water exchange and/or CO<sub>2</sub> aeration can thus be calculated based on this (preliminary) model.

Atlantic salmon is ammonotelic, i.e. excreting the bulk of nitrogenous waste from deamination of proteins as ammonia (NH<sub>3(g)</sub>). A pH dependent equilibrium exists between NH<sub>3</sub> and ionized ammonium NH<sub>4</sub><sup>+</sup> with a pKa of about 9,2 (Emerson et al. 1975). NH<sub>3</sub> is equilibrated between body compartments while NH<sub>4</sub><sup>+</sup> is distributed according to pH (Randall and Wright 1995). The gills are the main site of ammonium and ammonia excretion (Evans 2005; Terjesen 2008). Ammonia is excreted through passive diffusion, and ammonium by NH<sub>4</sub><sup>+</sup>/Na<sup>+</sup> exchange (Randall and Wright 1995). The two forms are collectively measured in water as total ammonia nitrogen (TAN) and NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup> ratio calculated as a function of pH and temperature (Emerson et al.

1975). Unionized  $\text{NH}_3$  is regarded as the main toxic form of the two, and toxicity is therefore highly pH dependent (Thorarensen and Farrell 2011). The primary toxic effect is regarded to be disruption of oxidative metabolism and draining of energy stores in the brain. Acute responses include disruption of enzyme activity, reduced swimming capacity, increased ventilation rate and osmoregulatory disturbances, while chronic exposure reduces growth and disease resistance (reviewed by Thorarensen and Farrell 2011). Recommended safe levels for salmonids range from 12 to 30  $\mu\text{g NH}_3\text{-N L}^{-1}$  (Thorarensen and Farrell 2011). In closed transport or treatment systems, pH depression will help detoxify ammonia. The main concern arises if  $\text{CO}_2$  aeration is applied to a degree where ammonia build-up may cause problems, and if sudden shifts to a higher pH occurs, i.e. if seawater is added to the treatment water. Equations in Thorarensen and Farrell (2011) and/or the NIVA transport model may provide the tools to calculate TAN accumulation. However, pH changes must be also taken into account to determine risk for toxic effects.

### 3. Amoebic Gill disease

#### 3.1 General description

Amoebic Gill Disease (AGD) of marine Atlantic salmon is caused by the amoeba *Paramoeba perurans*. *Paramoeba perurans* appears to be a facultative parasite of fish, having been identified to infect a number of different marine and euryhaline species including rainbow trout (*Oncorhynchus mykiss*), brown trout (*Salmo trutta*), chinook salmon (*O. Tsawyschta*), turbot (*Psetta psetta*), Ballan wrasse (*Labrus bergylta*), sharpnose sea bream (*Diplodus puntazzo*), seabass (*Dicentrarchus labrax*), ayu (*Plecoglossus altivelis*), blue warehou (*Seriolella brama*) (Nowak et al. 2002, Crosbie et al. 2010, Karlsbakk et al. 2013). It is speculated to have a relatively simple lifecycle, although this has yet to be confirmed. The impact of this pathogen has been seen in Tasmania, Australia since the late 1980s where cost estimates of 10-15% of the value of production being attributed to its control. However, the agent *Paramoeba perurans* and indeed AGD has been recognized and diagnosed in many other countries besides Australia including Chile (Bustos et al.

2011); Ireland, Japan, New Zealand, Norway, USA, Scotland and Spain (Young et al 2007; 2008; Steinum et al 2008; Nylund et al 2008; Nowak et al. 2002).

The pathology of AGD specifically (Powell et al. 2008) and related infectious and non-infectious gill disorders have been widely reviewed in the past (Mitchell and Rodger 2011; Rodger et al. 2010). Specific to AGD is the interaction of the gill epithelium with *Paramoeba perurans* whereby attachment of the parasite, results in acute cellular necrosis (Powell et al. 2008) and filamental epithelial cell hyperplasia giving rise to a compensatory plaque of tissue infiltrated with inflammatory immune cells and specifically eosinophils (Lovy et al 2006) that essentially prevents further damage to the gill. The filamental hyperplasia, reduces the functional gill surface area and the associated accumulation in mucus production causes inhibition of carbon dioxide excretion across the gill leading to a persistent respiratory acidosis (Powell et al. 2000). However, respiratory disturbances are only part of the pathology, an acute cardiovascular compromise occurs whereby systemic hypertension develops causing to circulatory collapse and finally death – particularly in fish susceptible to stress (eg. triploid fish, Powell et al. 2008).

This means that assessment of the success of treatments for AGD can either focus upon:

1. The presence or absence of the parasite
2. The presence or absence of gill lesions
3. Fish survival, mortality or other indicators of performance such as appetite or growth

In reality all three approaches make for a wholistic assessment of treatment success. In recent years tools have been developed to assess the presence or absence of *Paramoeba perurans* using real-time PCR of gill swabs or tissue samples (Young et al 2008). This test has since been commercialized by a number of diagnostic companies including Pharmaq Analytiq AS and Patogen AS in Norway. Although exquisitely sensitive; good correlations between actual amoeba numbers, pathology caused and treatment success have not been established.



### 3.2 Approaches to control of AGD – practical limitations

In commercial farm situations the number of fish to be treated, size of the fish to be treated, and location of net pens pose significant logistical limitations. The experience from sealice treatments have given Norwegian farmers extensive experience in handling cage-based bath treatments although the time and expense and frequency for treatment of large numbers of Atlantic salmon impose significant constraints upon the frequency, speed and success for treating an entire AGD affected site. Important with any form of bathing treatments, cages of Atlantic salmon need to be confined either by tarpaulin (Fig. 5), cage skirt or else transferred to a vessel such as



Figure 5. Process of commercial freshwater bathing for AGD control in Tasmania (Clockwise) Preparation of a tarpaulin liner into a cage prior to treatment, transfer of fish by pump into freshwater bath held within a tarpaulin, active oxygenation of the water within the bath, removal of the tarpaulin allowing the fish to fall into the waiting cage below.

a well boat for treatment. This imposes a handling effect, resulting in acute stress on the fish with the implications to water quality (see discussion above) and fish welfare (see discussion below). Some treatments are likely to be relatively less stressful than other, although even a cage skirt (open at the bottom but enclosing the water at the surface of the cage) can have significant effects on water circulation, oxygenation and stress effects on the fish (Stein et al. 2012). Indeed other studies have suggested that the oxygenation of any treatment is likely to be a constraint to bath treating Atlantic salmon under commercial conditions (Treasurer et al. 2000). Current treatment practices are to either treat at cage-side or else to remove the cage (or fish) from the main production site to a designated treatment site. In Australia, a country with extensive experience of this practice, cages are towed to a treatment barge prior to treatment. This handling of fish means that feeding is stopped 24h prior to scheduled treatment, and fish are not fed again until after treatment and replacement of the cage onto the production site. In some cases this may represent 48h with the associated loss of growth. The other implication of this treatment strategy is that typically one cage at a time is treated and then the clean cage is replaced alongside a potentially heavily infected cage. The risk of cross contamination and a re-infection is therefore magnified. Indeed the epidemiology and infection risk associated with the movement of cages, treatment and use of fallowing have been explored (Douglas-Helders et al. 2004) and although clear benefits were seen by placing newly bathed fish onto virgin and partly fallowed sites, the occurrence of AGD was delayed and not prevented – subsequently this practice has been all but abandoned in Australia, citing the lack of non-AGD affected sites or available sites to allow fallowing to occur.

### **3.3 Freshwater treatment of AGD**

In Australia, the treatment of choice for the control of AGD is freshwater bathing. This treatment was quickly identified as the primary control option in the early 1990s. Subsequently, the efficacy of such treatments was characterized (Parsons et al. 2001a and b) and a demonstrated positive physiological effect on the fish shown (Powell et al. 2001). The effects of treatment on re-infection showed that although gill lesions (hyperplastic patches) were removed, amoebic re-colonisation of the gill

occurred quickly with amoeba numbers equalling pre-treatment levels in as few as 10 days with the most aggressive infections at the height of the Tasmanian summer (Clark et al. 2003).

The process of freshwater bathing involves the filling of a large plastic tarpaulin (approximately 1 ML) with water piped to the bathing site. Under the tarpaulin is a clean net cage. Fish to be treated are transferred (typically by air-lift pump) to the freshwater filled tarpaulin and maintained for 3-4 h with additional oxygenation. Oxygen levels are targeted at 120-150% air saturation. Following the bathing period the tarpaulin is removed by winch and the fish fall into the awaiting cage. The process takes approximately 1 working day to complete.

Amoebae isolated from the gills of AGD affected salmon (specific diagnostics for *Paramoeba perurans* were not available at the time) were used in a number of short-term, in vitro bioassays that allowed determination of some of the key chemical characteristics that favoured survival or killed gill amoebae. Isolated gill amoebae showed resilience to freshwater treatment where the Na<sup>+</sup> concentration was increased, however, the effects of increased concentrations of Mg<sup>2+</sup> and Ca<sup>2+</sup> ions significantly promoted survival over 3 hours of exposure. Even when Na<sup>+</sup> concentrations were relatively high (250 mg L<sup>-1</sup>), low Ca<sup>2+</sup> concentrations (10 mg L<sup>-1</sup>) were as effective as unionized freshwater after 3 hours of exposure (Powell and Clark 2003). This suggested that soft freshwater sources were more favourable for the control of AGD in bathing compared to hard water sources. This was confirmed under field conditions where artificially softened freshwater (ion exchange with Na<sup>+</sup>) (Roberts and Powell 2003a) and the selection of soft freshwater sources produced increased efficacy of bathing and a 113 degree day (13% increase in the inter-bath interval) delay in subsequent bathing (Powell et al. 2005). Other approaches of removing Ca<sup>2+</sup> from freshwater and seawater were also examined using the ionic chelator Calgon T<sup>TM</sup> although at effective concentrations (6 mg L<sup>-1</sup> per mg L<sup>-1</sup> of hardness) to soften hardwater (225 mg L<sup>-1</sup> CaCO<sub>3</sub> equivalents) the resultant discharge of phosphate and cost would be prohibitive under commercial conditions (Powell et al. 2005).

Dissolved organic carbon (in the form of humic and tannic acid) has been shown to enhance the efficacy of freshwater at killing Paramoebae (Green et al. 2005). In combination with different concentrations of  $\text{Ca}^{2+}$ , a combination of soft water with high concentrations of organic acids resulted in the best conditions for killing amoebae in freshwater both in vitro as well as in an experimental freshwater bath (Green et al. 2005). The effects of DOC also resulted in the decrease in the number of hyperplastic gill lesions following the bath. The mechanism by which this effect acts is unclear but it is possible that the organic acids (tannic and humic) resulted in chelation of divalent cations so enhancing the efficacy of the freshwater treatment. Alternatively, the organic acid load may have had a direct toxic effect on the amoebae (Green et al. 2005).

The effects of combining freshwater with oxidative disinfectants have been investigated with some limited success. It was found that chloramine-T and hydrogen peroxide both enhanced the efficacy of freshwater baths, although the benefits were small (Powell and Clark 2002). However, the variability in treatment success may have been as a result of different water qualities. Combined treatments offer the advantage of ensuring that amoebae are killed once removed from the gills of affected fish so reducing the chance of re-infection once the bath is ended.

### **3.4 Oxidative disinfectants for AGD control in seawater**

Alternatives to freshwater bathing have included the investigation of a number of oxidative disinfectants. In general most oxidative disinfectants work through the release of reactive oxygen or chlorine species thus destabilizing or permeating cell membranes. Two main oxidative disinfectants have been the focus of significant investigation for the control of AGD: Chlorine-based Chloramine-T (*n*-sodium-*n*-chloroparatoluenesulphonamide) and oxygen-based hydrogen peroxide ( $\text{H}_2\text{O}_2$ ). Chloramine-T, hydrogen peroxide and chlorine dioxide were all toxic in vitro to isolated gill amoebae from AGD-affected salmon primarily *Neoparamoeba pemaquidensis* (Howard and Carson 1993; Powell and Clark 2002; 2003) and specifically hydrogen peroxide with *Paramoeba perurans* (Adams et al. 2012). In

preliminary tests with fish suggested that AGD was reduced when added to enhance the efficacy of freshwater baths (Powell and Clark 2002).

In vitro and medium scale treatment investigations with chloramine-T demonstrated that chloramine-T was acutely toxic to isolated gill amoebae (Powell and Clark 2003). In small scale field studies, exposure of AGD affected Atlantic salmon to chloramine-T at 10 mg L<sup>-1</sup> for 1 hour in seawater were moderately successful with significant reductions in gill amoeba numbers (Harris et al. 2004; Harris et al 2005).

Furthermore, treatment of Chinook salmon (*Onchorynchus tsawhytcha*) smolts was also successful at removing gill amoebae with minimal adverse effects, although AGD in this species is not thought to be a significant health issue (Powell personal observations). However, the toxicity of chloramine-T to Atlantic salmon smolt is enhanced in seawater and further enhanced when oxygen levels are increased to 200% air saturation (Powell and Harris 2004). Moreover, attempts to upscale treatment to commercial operations proved challenging with mixed results and have not been developed further (Powell personal observations).

Hydrogen peroxide was tested in salt water laboratory bath treatments of AGD affected Atlantic salmon smolts at 10 and 15°C with post-treatment survival dependent upon exposure duration. Toxicity increased with exposure time at both temperatures at concentrations exceeding 1000 mg L<sup>-1</sup>. However total gill amoeba counts were not significantly reduced compared to untreated controls and the variability between fish was high (Powell et al. 2005). More recently this approach has been re-examined suggestion that at 12°C hydrogen peroxide (1250 mg L<sup>-1</sup>) for 15 min reduced the number of gill lesions (Adams et al. 2012). However, as with the previous study (Powell et al. 2003) the results were variable with slightly longer durations of exposure resulted in a highly variable efficacy (Adams et al. 2012). Oxidative disinfectants (such as chloramine-T and hydrogen peroxide) have a number of well documented patho-physiological effects on healthy and damaged gill tissue of salmonids in both fresh (Powell and Perry 1995; 1997a; 1997b; 1997c) and marine conditions (Powell and Harris 2004). In general, the effects of acute oxidative disinfectant exposure include an acute congestion of the gill filament and central

venous sinus, most typically as a result of an increased vascular pressure caused by elevated cardiac output or intra-branchial pressure increased due to adrenergic responses induced by the release of adrenalin and noradrenalin. The gill lamellar epithelium is often either crenated or denuded often associated with epithelial cell necrosis. The consequences of this are in freshwater, a net influx of water (and efflux of plasma electrolytes), increased vascular volume, haemolysis and eventual cardiovascular collapse due to increase vascular viscosity and ultimately haemostasis. Under marine conditions, haemo-concentration occurs with an apparent efflux of water, and potential influx of  $\text{Na}^+$  and  $\text{Cl}^-$ , an associated hypernatriuremia and ultimately electrolytic imbalance resulting in death. Sublethal effects of oxidative disinfectants result in permeabilisation of epithelial cell membranes and trans-cellular efflux (freshwater) or potentially influx (seawater) of  $\text{Na}^+$  and  $\text{Cl}^-$  ions. This process results in acid-base disturbances (Powell and Perry 1997d;1998). These ionic disturbances are often manifest in hypertrophy of chloride and mitochondrial rich cells in the gill (Powell and Harris 2004).

## 4. Sea lice

### 4.1 General description

The issue of sea lice infestation of farmed Atlantic salmon (primarily by *Lepeophtheirus salmonis* in the Northern hemisphere and *Caligus rogercresseyi* in the Southern hemisphere) has been a constant challenge for commercial farm production of the species for many decades. The primary issue is that the salmon is infected by a free swimming phototactic nauplius stage that moults and attaches onto a host as the attached chalimus stage, after successive moults, the motile copepodite stages and adult stages move over the epithelial surface of the salmon grazing upon the skin and mucus. The subsequent result is acute erosive lesions leading to osmoregulatory distress. Although much has been discussed regarding the immune responses to infestation, control measures still rely mostly on chemical de lousing and disinfection to control the level of infestation of fish on commercial farms.

#### 4.2 Approaches to control of sea lice – practical limitations

Sealice control under commercial farm conditions in some countries is highly regulated. Typically the number of gravid female lice is the primary treatment trigger and the issue of surveillance and accurate enumeration has recently been in focus (Heuch et al. 2011; Revie et al. 2007). Of particular importance, as with bathing fish for AGD treatments, cages of Atlantic salmon need to be confined either by tarpaulin, cage skirt or else transferred to a vessel such as a well boat for treatment. The consequent handling effects result in acute stress on the fish (see discussion below) with implications for water quality (see discussion above). Some treatments are likely to be relatively less stressful than other, although even a cage skirt (open at the bottom but enclosing the water at the surface of the cage) can have significant effects on water circulation, oxygenation and stress effects on the fish (Stein et al. 2012). Indeed other studies have suggested that the oxygenation of any treatment is likely to be a constraint to bath treating Atlantic salmon under commercial conditions (Treasurer et al. 2000).

#### 4.3 Freshwater treatment of sea lice

Exposure of sea louse infested Atlantic salmon to salinities below 29 ppt results in the gradual loss of sea lice. *Lepeophtheirus salmonis* is rapidly killed in freshwater (Connors et al. 2008), and the survival of free-swimming copepodids was severely compromised at salinities below 29 ppt, irrespective of their attachment to a host (Bricknell et al. 2006). However, earlier work by Stone et al. indicated that short duration (3 h) bath exposure was likely ineffective treatment for controlling sea lice (Stone et al. 2002). Little work has been undertaken with respect to commercial treatments for sea lice with freshwater where the water chemistry and its implications have been examined. However, it should suffice that similar constraints of water quality and chemistry as those discussed above should remain valid.

#### 4.4 Oxidative disinfectants for control of sea lice in seawater

Among a number of chemical treatments for sealice (for example see reviews by Torrissen et al. 2013; BurrIDGE et al. 2010; Robertson et al. 2009), hydrogen peroxide has been identified as a potential disinfectant (Toovey et al. 2000; Treasurer and

Grant 1997; Johnson et al. 1993). Primarily, hydrogen peroxide is believed to act by either killing the copepod directly through oxidation of cell membranes, or else causing it to detach from the skin surface (Torrissen et al. 2013). Although having a relatively narrow window of safety, treatments up to 1500 mg L<sup>-1</sup> for up to 20 min (depending upon temperature) are reported. The challenge remains to introduce, distribute and mix sufficient quantities of chemical into a bath treatment and eliminate the residual peroxide following the treatment, to prevent overdosing the salmon. Thus it is generally considered more effective against the motile stages of the parasite rather than the attached chalimus stages. However, recently the success of using hydrogen peroxide has been questioned with evidence of resistance developing in sea louse populations (Treasurer et al. 2000) and limited success of this treatment option with other species besides *Lepiophtheirus salmonis* such as against *Caligus rogercresseyi* in Chile (Bravo et al. 2010).

Oxidative disinfectants (such as chloramine-T and hydrogen peroxide) have a number of well documented patho-physiological effects on healthy and damaged gill tissue of salmonids in both fresh (Powell and Perry 1995; 1997a;1997b;1997c) and marine conditions (Keimer and Black 1997; Powell and Harris 2004;). In general, the effects of acute oxidative disinfectant exposure include an acute congestion of the gill filament and central venous sinus, most typically as a result of an increased vascular pressure caused by elevated cardiac output or intra-branchial pressure increased due to adrenergic responses induced by the release of adrenalin and noradrenalin. The gill lamellar epithelium is often either crenated or denuded often associated with epithelial cell necrosis. The consequences of this are, a net influx of water (and efflux of plasma electrolytes), in freshwater, increased vascular volume, haemolysis and eventual cardiovascular collapse due to increased vascular viscosity and ultimately haemostasis. Under marine conditions, haemo-concentration occurs with an apparent efflux of water, and potential influx of Na<sup>+</sup> and Cl<sup>-</sup>, an associated hypernatremia and ultimately electrolytic imbalance resulting in death. Sublethal effects of oxidative disinfectants result in permeabilisation of epithelial cell membranes and trans-cellular efflux (freshwater) or potentially influx (seawater) of Na<sup>+</sup> and Cl<sup>-</sup> ions. This process results in acid-base disturbances (Powell and Perry 1997d;1998).



These ionic disturbances are often manifesting in hypertrophy of chloride and mitochondrial rich cells in the gill (Powell and Harris 2004).

## 5. Implications for fish welfare

Atlantic salmon are a euryhaline species and post-smoltification exhibit a high capacity for ion and osmoregulation. The transfer of AGD affected post-smolt salmon into freshwater results in an acute net efflux of  $\text{Na}^+$  and  $\text{Cl}^-$  ions that peaks at 2 h post-transfer and is quickly reduced by 3 h exposure (Roberts and Powell 2003b). The net titrateable alkalinity flux was significant for both AGD affected and non-affected salmon with AGD affected fish having a larger net titrateable alkalinity efflux (Roberts and Powell 2003b). Thus, ionic disturbances are relatively minor. Indeed freshwater bathing enhances the break-up of the hyperplastic gill lesions (Roberts and Powell 2003a) and physiological disturbances caused by AGD including respiratory acidosis are reduced (Powell et al. 2001). Additionally, hyperoxic freshwater bathing also reduced systemic hypertension in AGD-affected Atlantic salmon (Powell et al. 2002).

## 6. Conclusions

It is apparent from the existing experience and data presented; freshwater treatments of Atlantic salmon smolts affected by AGD are effective and generally pose low risk to overall fish health with a widely accepted large margin of safety, despite some short term physiological effects. However, there are a number of potential risks and large gaps in the knowledgebase surrounding bath treatments for this disease. Despite the work carried out so far as presented in this review (summarized in a number of reports (Powell and Clark 2002, Powell et al. 2005; Powell et al. 2007), the water characteristics of Norwegian freshwater sources, nor the scale of Norwegian production has been taken into account in describing a best practice for the treatment of AGD. In particular the interactions between DOC and POC and both divalent cations (e.g.  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ ) and transition metal ions such as  $\text{Cu}^{2+}$  and  $\text{Al}^{3+}$  have not been investigated with respect to the effects on the amoebae, efficacy of treatment

and lesion resolution post-treatment and physiological effects in the fish.

Additionally, interactions between the water chemistry, effects of organic load, fish biomass, temperature, salinity, oxygenation status and treatment chemicals such as hydrogen peroxide are poorly understood. In this respect it is imperative that these factors be investigated with respect to each other so leading to improvements in bath efficacy, treatment safety, and fish welfare.

## 7. Recommendations

1. Water sources for freshwater bathing target optimal water chemistry and quality (summarized in Table 1). These parameters should include low  $\text{Ca}^{2+}$  and  $\text{Na}^{2+}$  content (characteristically soft waters) with a moderate pH optimally approximating 6.5. The water source would benefit from a high dissolved organic carbon content although it is recognized that the interactions of metal ions, divalent base metal ions (e.g.  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ ) are unknown at pH values characteristic of bathing operations. Water sources with high alkalinity should be targeted although these often are buffered mostly by  $\text{CaCO}_3$  and thus the risk is increases in the  $\text{Ca}^{2+}$  concentration. Alternatives may include the use of other buffering/neutralizing agents added to the bath water (e.g.  $\text{NaOH}$ ,  $\text{Na}_2\text{CO}_3$ ,  $\text{NaHCO}_3$ ) although these have not been tested on a commercial scale. The use of carbonate based buffers would require the use of an active degassing process to facilitate the stripping of  $\text{CO}_2$  produced in the buffering process.
2. It is recommended that all water sources used for commercial bathing of Atlantic salmon should be analysed as close to the time of use (for bathing) for water chemistry characteristics. Many of the historical analyses may be changed over time depending upon catchment use, seasonal and annual fluctuations in rainfall etc.
3. Aversive action with fresh water treatments are rarely required based upon the experiences in Tasmania although large drops in pH and oxygen saturation would be clear indicators of failed handling of fish during the treatment. It

should also be noted that the more severe the AGD score, the greater the risk of respiratory compromise in the fish and resultant treatment-related mortality.

4. The continued investigation of water quality and chemistry parameters and their interactions in freshwater treatment baths to optimize treatment efficacy and identify constraints for the control of AGD.
5. Use of existing information and the further investigation of water chemistry interactions in treatment efficacy to further develop an “industry best practice” for AGD treatments for the control of gill health on Norwegian sea farms.
6. The incorporation of farm site and environmental data into best practice AGD treatment including preceding environmental characteristics (e.g. algal blooms, jellyfish, freshwater runoff, DOC, temperature, salinity etc).
7. The up-skilling and further training of farm personnel in the diagnosis of AGD, scoring of gill lesions, handling of fish and monitoring of water quality and fish welfare during treatment baths.
8. The development of novel, alternatives to freshwater bathing (e.g. In-feed treatments) that could be used in conjunction with topical treatment of the sustainable control of AGD.

**Summary of critical research needs and priorities for strengthening existing data and protocols for effective treatment.**

- a. **Research Priority Action 1:** These investigations should include the interactions of metal ions and organic carbon load within the baths with regard to efficacy of bathing.
- b. **Research Priority Action 2:** The interactions of hydrogen peroxide with metal ions, organic carbon, oxygenation level, salinity etc and efficacy and toxicity to *Paramoeba perurans* and Atlantic salmon in both freshwater and seawater.
- c. **Research Priority Action 3:** Inclusion of environmental data into an updated “industry best practice” and the incorporation of information from currently funded research projects from different agencies (e.g. FHF, NFR etc) regarding the causes of gill disease and their interactions with AGD.
- d. **Research Priority Action 4:** Freshwater bathing or the use of hydrogen peroxide treatments poses extensive logistical challenges and high risk (in the case of H<sub>2</sub>O<sub>2</sub> treatments). Alternative treatment and therapeutic strategies need further investigation. The aim of identifying additives to bath treatments that could improve efficacy (or combined treatments of freshwater and oxidative disinfectants) should be prioritized along with the further development of in feed treatments and treatment regimes.

Table 1. Recommended values and limits for water chemistry of freshwater baths for the treatment of AGD affected Atlantic salmon with gill scores < 3<sup>1</sup> and the percentile of Norwegian waters with the appropriate characteristics. Blank boxes mean no action can be expected to be taken.

Parameters	Prior to bathing	%ile of waters	During bathing	Aversive action options
Conductivity	< 500 $\mu\text{S cm}^{-1}$	0-100	< 1000 $\mu\text{S cm}^{-1}$	Add low conductivity water to treatment Continue treatment monitoring closely
pH	6.0-6.7	19-80	6.0-6.8	Increase buffer capacity if possible Continue treatment monitoring closely Terminate treatment
ORP <sup>2</sup>	FW <sup>3</sup> 40-100 mV SW <sup>3</sup> 140-170 mV		< 350 mV <sup>4</sup>	Terminate treatment immediately
TOC/DOC	< 3 mg L <sup>-1</sup>	0-70	If possible sample for later analysis	
Ca <sup>2+</sup> Concentration	< 10 mg L <sup>-1</sup>	0-100	If possible sample for later analysis	
Na <sup>2+</sup> Concentration	< 10 mg L <sup>-1</sup>	0-90		
O <sub>2</sub> saturation	90-110%		90-110%	Increase oxygen input and solubilization
CO <sub>2</sub> concentration	< 5 mg L <sub>-1</sub>		< 25 mg L <sup>-1</sup>	Actively degass using compressed air Continue treatment monitoring closely Terminate treatment
Water characteristics	Freshwater < 5ppt salinity		Freshwater < 5ppt salinity	

<sup>1</sup> The gill score is likely to have a direct effect upon the resultant mortality associated with a treatment bath. With a higher gill score, the risks of respiratory compromise and cardiovascular collapse is increased.

<sup>2</sup> Recommended monitoring continuously when using H<sub>2</sub>O<sub>2</sub> treatments as an indicator of oxidative toxicity. This can be used in combination with Total Residual Oxidation measurements using spectrophotometric analysis.

<sup>3</sup> Values will vary between different water sources and decision s and actions should be dependent upon and relative to baseline values

<sup>4</sup> Critical value based upon data from Harris et al. (2004).

### **Acknowledgements**

Parts of this report were presented at the FHF AGD seminar Scandic Airport Hotel Bergen, 16 January 2014. Thanks to Bjørn-Olav Rosseland for his advice and contributions in the preparation of this report and Ole-Kristian Hess-Erga for his comments and suggestions. Cover photo: Fresh water bathing for AGD in Atlantic salmon in Tasmania, courtesy of J. Harris.

## References

- Adams, M.B., Crosbie, P.B.B. and Nowak, B.F. (2012). Preliminary success using hydrogen peroxide to treat Atlantic salmon, *Salmo salar* L., affected with experimentally induced amoebic gill disease (AGD). *Journal of Fish Diseases* 35: 839-848.
- Andren, C.M., Kroglund, F. and Teien, H.C. (2006) Controlled exposure of brown trout to humic water limed to different pH and inorganic aluminium concentrations, *Verh. International Reviews in Limnology* 29: 1548–1552.
- Bravo, S., Treasurer, J., Sepulveda, M and Lagos, C. (2010). Effectiveness of hydrogen peroxide in the control of *Caligus rogercresseyi* in Chile and implications for sea louse management. *Aquaculture* 303: 22-27.
- Bricknell, I.R., Dalesman, S.J., O’Shea, B., Pert, C.C. and Luntz, A.J.M. (2006). Effect of environmental salinity on sea lice *Lepeptheirus salmonis* settlement success. *Diseases of Aquatic Organisms* 71: 201-212.
- Bjerknes, V., Fyllingen, I., Holtet, L., Teien, H-C., Rosseland, B.O. and Kroglund, F. (2003). Aluminium in acidic river water causes mortality of farmed Atlantic Salmon (*Salmo salar* L.) Norwegian fjords. *Marine Chemistry* 83: 169–174
- Burridge, L., Weis, J.S., Cabello, F., Pizarro, J. and Bostick, K. (2010). Chemical use in salmon aquaculture: a review of current practices and possible environmental effects. *Aquaculture* 306: 7-23.
- Bustos, P.A., Young, N.D., Rozas, M.A., Bohle, H.M., Ildefonso, R.S., Morrison, R.N. and Nowak, B.F. (2011). Amoebic gill disease (AGD) in Atlantic salmon (*Salmo salar*) farmed in Chile. *Aquaculture* 310: 281-288.

Clark, G., Powell, M. and Nowak, B. (2003). Effect of commercial freshwater bathing on reinfection of Atlantic salmon, *Salmo salar* with amoebic gill disease. *Aquaculture* 219: 135-142.

Crosbie, P.B.B., Ogawa, K., Nakano, D. and Nowak, B.F. (2010). Amoebic gill disease in hatchery-reared ayu, *Plecoglossus altivelis* (Temmick & Schegel), in Japan is caused by *Neoparamoeba perurans*. *Journal of Fish Diseases* 33: 455-458.

Connors, B.M., Juarez-Colunga, E and Dill, L.M. (2008). Effects of varying salinities on *Lepeoptheirus salmonis* survival on juvenile pink and chum salmon. *Journal of Fish Biology* 72: 1825-1830.

Douglas-Helders, G.M., Weir, I.J., O'Brien, D.P., Carson, J. and Nowak, B.F. (2004). Effects of husbandry on prevalence of amoebic gill disease and performance of reared Atlantic salmon (*Salmo salar* L.). *Aquaculture* 241: 21-30.

Emerson, K., Russo, R., Lund, R. and Thurson, R. (1975). Aqueous ammonia equilibrium calculations: effects of pH and temperature. *Journal of the Fisheries Research Board of Canada* 32: 2379–2383.

Evans, D.H., Piermarini, P.M., and Choe, K.P. (2005). The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiological Reviews* 85: 97-177.

Finstad, B., Kroglund, F., Strand, R., Stefansson, S.O., Bjørn, P.A., Rosseland, B.O., Nilsen, T.O. and Salbu, B. (2007). Salmon lice or suboptimal water quality – reasons for reduced post-smolt survival? *Aquaculture* 273: 374-383.

Finstad, B., Kroglund, F., Bjørn, P.A., Nilsen, R., Pettersen, K., Rosseland, B.O., Teien, H-C., Nilsen, T.O., Stefansson, S.O., Salbu, B., Fiske, P and Ebbesson, L.O.E. (2012). Salmon lice- induced mortality of Atlantic salmon postsmolts



experiencing episodic acidification and recovery in freshwater. *Aquaculture* 362-363: 193-199.

Fivelstad, S. and Binde, M. (1994). Effects of reduced waterflow (increased loading) in soft water on Atlantic salmon smolts (*Salmo salar* L.) while maintaining oxygen at constant level by oxygenation of the inlet water. *Aquaculture Engineering*. 13: 211–238.

Fivelstad, S., Waagbø, R., Zeitz, S.F., Hosfeld, A.C.D., Olsen, A.B. and Stefansson, S. (2003). A major water quality problem in smolt farms: combined effects of carbon dioxide, reduced pH and aluminium on Atlantic salmon (*Salmo salar* L.) smolts: physiology and growth. *Aquaculture* 215:339–357.

Green, T., Powell, M.D., Harris, J.O. and Butler, R. (2005). Effects of dissolved organic carbon (DOC) and hardness in freshwater used to treat amoebic gill disease (AGD). *Aquaculture Research* 36: 398-404.

Harris, J.O., Powell, M.D., Attard, M.G. and DeHayr, L. (2005). Clinical assessment of chloramine-T as a treatment for the control of gill amoebae in farmed Tasmanian Atlantic salmon, *Salmo salar* L. *Aquaculture Research* 36: 776-784.

Harris, J.O., Powell, M.D., Attard, M. G. and Green, T. (2004). Efficacy of chloramine-T as a treatment for amoebic gill disease (AGD) in marine Atlantic salmon (*Salmo salar* L.). *Aquaculture Research* 35: 1448-1456.

Heuch, P., Gettinby, G., Revie, C.W. (2011). Counting sea lice on Atlantic salmon farms – empirical and theoretical observations. *Aquaculture* 320: 149-153.

Henriksen, A., Lien, L., Rosseland, B.O., Traaen, T., Sevaldrud, I. (1989). Lake acidification in Norway - present and predicted fish status. *Ambio* 18: 314-321.

Howard, T. and Carson J. (1993). Progress report: are there alternatives to freshwater treatment of AGD? *In: Seeking and solving: Saltas R&D Review seminar pp 81-87, Hobart, Tasmania, Australia.*

Johnson, S.C., Constible, J.M. and Richard, J. (1993). Laboratory investigations on the efficacy of hydrogen peroxide against the salmon louse *Lepeoptheirus salmonis* and its toxicological and histopathological effects on Atlantic salmon *Salmo salar* and chinook salmon *Oncorhynchus tshawytscha*. *Diseases of Aquatic Organisms 17: 197-204.*

Karlsbakk., E., Olsen, A.B., Einen, A-C, B., Mo, T.A., Fiksdal, I.U., Aase, H., Kalgraff, C., Skår, S-Å. and Hansen, H. (2013) Amoebic gill disease due to *Paramoeba perurans* in ballan wrasse (*Labrus bergylta*). *Aquaculture 412-413: 41-44.*

Keimer, M.C.B. and Black, K.D. (1997). The effects of hydrogen peroxide on the gill tissues of Atlantic salmon, *Salmo salar* L. *Aquaculture 153: 181-189.*

Kristensen, T., Åtland, Å., Rosten, T., Urke, H.A. and Rosseland, B.O. (2009). Important influent-water quality parameters at freshwater production sites in two salmon producing countries. *Aquacultural Engineering 41: 53-59.*

Leivestad, H., Muniz, I.P. and Rosseland, B.O. (1980). Acid stress in trout from a dilute mountain stream. Pp 336-337 in: Tollan, A. & Drabløs, D. (eds.) *Ecological impact of acid precipitation. - SNSF-project FA 105/80.*

Kroglund, F., Teien, H-C., Guttrup, J. and Johansen, Å. (2011). Påvirkes laksesmolt av aluminium i brakkvann?. *Storelva I Holt, Aust-Agdeer, 2010.*

Lovy, J., Becker, J.A., Speare, D.J., Wadowska, D.W., Wright, G.M. and Powell M.D. (2007). Ultrastructural examination of the host cellular response in the gills of Atlantic salmon, *Salmo salar*, with Amoebic Gill Disease. *Veterinary Pathology 44: 663-671.*

Lydersen, E., Lofgren, S. and Arnesen, R.T. (2002). Metals in Scandinavian surface waters: effects of acidification, liming and potential reacidification. *Critical Reviews in Environmental Science and Technology*. 32: 73–295.

Mitchell, S.O. and Rodger, H.D. (2011). A review of infectious gill disease in marine salmonid fish. *Journal of Fish Diseases* 34: 411-432.

Nowak, B.F., Carson, J., Powell, M.D. and Dykova, I. (2002). Amoebic gill disease in the marine environment. *Bulletin of the European Association of Fish Pathologists* 22: 144-148.

Nylund, A., Watanabe, K., Nylund, S., Karlsen, M., Saether, P., A., Arnesen, C.E. and Karlsbakk, E. (2008). Morphogenesis of salmonid gill poxvirus associated with proliferative gill disease in farmed Atlantic salmon (*Salmo salar*) in Norway. *Archives of Virology* 153: 1299-1309.

Pagenkopf, G.K. (1983). Gill Surface Interaction Model for Trace-Metal Toxicity to Fishes: Role of complexation, pH and Water Hardness. *Environmental Science and Technology* 17: 342–347.

Parsons, H., Nowak, B., Fisk, D. and Powell, M. (2001). Effectiveness of freshwater bathing as a treatment against Amoebic Gill Disease in Atlantic salmon. *Aquaculture* 195: 205-210.

Parsons, H.J., Nowak, B.F, Powell, M.D. and Dix, T. (2001). Changes in water quality during freshwater bathing of Atlantic salmon, *Salmo salar*, in Tasmania. *Bulletin of the European Association of Fish Pathologists* 21: 71-76.

Perry, S.F. and Gilmour, K.M. (2006). Acid-base balance and CO<sub>2</sub> excretion in fish: unanswered questions and emerging models. *Respiratory Physiology and Neurobiology* 154:199–215.

Powell, M.D. and Perry, S.F. (1996). Respiratory and acid-base disturbances in rainbow trout (*Oncorhynchus mykiss*) blood during exposure to chloramine-T, paratoluenesulphonamide and hypochlorite. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 701-708.

Powell, M.D. and Perry, S.F. (1997). Respiratory and acid-base disturbances in rainbow trout (*Oncorhynchus mykiss*) blood during exposure to chloramine-T under hypoxia and hyperoxia. *Journal of Fish Biology* 50: 418-428.

Powell, M.D. and Perry, S.F. (1997). Respiratory and acid-base pathophysiology of hydrogen peroxide in rainbow trout (*Oncorhynchus mykiss* Walbaum). *Aquatic Toxicology* 27: 99-112.

Powell, M.D. and Perry, S.F. (1997). Branchial ionic flux responses in rainbow trout to chloramine-T after acclimation to different levels of water hardness. *Journal of Aquatic Animal Health* 9: 196-202.

Powell, M.D. and Perry, S.F. (1998). Acid-base and ionic fluxes in rainbow trout (*Oncorhynchus mykiss*) during exposure to chloramine-T. *Aquatic Toxicology* 43: 13-24.

Powell, M.D. and Perry, S.F. (1999). Cardio-respiratory effects of chloramine-T exposure in rainbow trout. *Experimental Biology Online* 4: 5.

Powell, M.D., Parsons, H.J, Nowak, B.F. (2001). Physiological effects of freshwater bathing of Atlantic salmon (*Salmo salar*) as a treatment for amoebic gill disease. *Aquaculture* 199: 259-266.

Powell, M.D. and Clark, G.A. (2002). Atlantic Salmon Aquaculture Subprogram: effective treatments for the control of AGD. *Fisheries Research and Development Corporation final report project 2000/266 ISBN 0 85901 986 85p.*

Powell, M.D. and Clark, G.A. (2003). *In vitro* survival and the effect of effects of water chemistry and oxidative chemical treatments on isolated gill amoebae from AGD-affected Atlantic salmon. *Aquaculture* 220: 135-144.

Powell, M.D. and Harris, J.O. (2004). Influence of oxygen on the toxicity of chloramine-T to Atlantic salmon (*Salmo salar* L.) smolts in fresh and salt water. *Journal of Aquatic Animal Health* 16: 83-92.

Powell, M.D., Attard, M., Harris, J.O., Roberts, S.D., and Leef, M.J. (2005). Atlantic Salmon Aquaculture Subprogram: Why fish die – treatment and pathophysiology of Amoebic Gill Disease. *Fisheries Research and Development Corporation final report project 2001/205 ISBN 1 86295 259 0 115p.*

Powell, M.D. (2006). Physiological and clinical pathology. In: “*Systemic Pathology of Fish 2nd edition*” (ed. H.W. Ferguson) pp. 336-356. *Scotian Press, London, UK.*

Powell, M.D. (2007). Respiration in infectious and non-infectious gill diseases. In: “*Fish Respiration and Environment*” (eds. M.N. Fernandes , F.T. Rantin, M.L. Glass & B.G. Kapoor) pp. 317-339. *Science Publishers, Inc. Enfield (NH), USA.*

Powell, M.D., Becker, J.A., Ransome, J., Florent, R.L, and Jones, M.J. (2007). Atlantic Salmon Aquaculture Subprogram: Commercial AGD and salmon health. *Fisheries Research and Development Corporation final report project 2004/213 ISBN 978 1 86295 378 9 85p.*

Powell, M.D., Leef, M.L., Roberts, S.D. and Jones, M.A. (2008) Neoparamoebic gill infections: host response and physiology in salmonids. *Journal of Fish Biology* 73: 2161-2183.

Randall DJ, Wright PA (1995) Circulation and gas transfer. In: Groot G, Margolis L, Clarke WC (Eds.), *Physiological Ecology of Pacific Salmon*. *UBC Press, Vancouver, pp. 441-458*

Revie, C.W., Hollinger, E., Gettinby, G., Lees, F and Heuch, P.A. (2007). Clustering of parasites within cages on Scottish and Norwegian salmon farms: alternative sampling strategies illustrated using simulation. *Preventative Veterinary Medicine* 81: 135-147.

Roberts, S.D. and Powell, M.D. (2003). Reduced total hardness of freshwater enhances the efficacy of bathing as a treatment for amoebic gill disease in Atlantic salmon (*Salmo salar* L.). *Journal of Fish Diseases* 26: 591-599.

Roberts, S.D. and Powell, M.D. (2003). Comparative ionic flux and mucous cell histochemistry: effects of salinity and disease status in the euryhaline Atlantic salmon (*Salmo salar* L.). *Comparative Biochemistry and Physiology Part A* 134: 525-537.

Robertson, P.K.J., Black, K.D., Adams, M., Willis, K., Buchan, F., Orr, H., Lawton, L and McCullagh, C. (2009). A new generation of biocides for control of crustacean in fish farms. *Journal of Photochemistry and Photobiology B: biology* 95: 58-63.

Rodger, H.D., Henry, L. and Mitchell, S.O. (2010). Non-infectious gill disorders of marine salmonid fish. *Reviews in Fish Biology and Fisheries* DOI: 10.1007/s 11 160-010-9182-6.

Rosseland, B.O. and Staurnes, M. (1994). Physiological mechanisms for toxic effects and resistance to acidic water: an ecophysiological and ecotoxicological approach. In: Steinberg CEW, Wright RW (Eds.), *Acidification of Freshwater Ecosystems: Implications for the Future*. John Wiley & Sons, Ltd., pp. 227–246.

Skjelkvåle, B.L., Stoddard, J.L., Jeffries, D.S., Tørseth, K., Høgåsen, T., Bowman, J., Mannio, J., Monteith, D.T., Mosello, R., Rogora, M., Rzychon, D., Vesely, J., Wieting, J., Wilander, A. and Worsztynowicz, A. (2005). Regional scale evidence for improvements in surface water chemistry 1990-2001. *Environmental Pollution* 137:165-176.

Stein, L.H., Nilsson, J., Hevrøy, E.M., Oppedal, F., Kristiansen, T.S., Lien, A.M. and Folkedal, O. (2012). Skirt around a salmon cage to reduce infestation of salmon lice resulted in low oxygen levels. *Aquaculture Engineering* 51: 21-25.

Steinum, T., Kvellestad, A., Rønneberg, L.B., Nilsen, H., Asheim, A., Fjell, K., Nygård, S.M.R., Olsen, A.B. and Dale, O.B. (2008). First cases of amoebic gill disease (AGD) in Norwegian seawater farmed Atlantic salmon, *Salmo salar* L., and phylogeny of the causative agent using 18S cDNA sequences. *Journal of Fish Diseases* 31: 205-213.

Stone, J., Boyd, S., Sommerville, C. and Rae, G.H. (2002). An evaluation of freshwater bath treatments for the control of sea lice, *Lepeophtheirus salmonis* (Krøyer), infections in Atlantic salmon, *Salmo salar* L. *Journal of Fish diseases* 25: 371-373.

Teien, H-C., Garmo, Ø.A., Åtland, Å. and Salbu, B. (2008). Transformation of iron species in mixing zones and accumulation on fish gills. *Environmental Science and Technology* 42: 1780-1786.

Terjesen, B.F. (2008). Nitrogen excretion. In: *Fish Larval Physiology* (ed. by R.N. Finn & B. Kapoor), pp. 263–302. Science Publishers, New York.

Thorarensen, H., and Farrell, A.P. (2011). The biological requirements for post-smolt Atlantic salmon in closed-containment systems. *Aquaculture* 02/2011; DOI:10.1016/j.aquaculture.2010.11.043

Toovey, J.P.G., and Lyndon, A.R. (2000). The effects of hydrogen peroxide, dichlorvos and cypermethrin on subsequent fecundity of sea lice, *Lepeophtheirus salmonis* under fish farm conditions. *Bulletin of the European Association of Fish Pathologists* 20: 224-228.

Torrissen, O., Jones, S., Asche, F., Guttormsen, A., Skilbrei, O.T., Nilsen, F., Horsberg, T.E. and Jackson, D. (2013). Salmon lice – impact on wild salmonids and salmon aquaculture. *Journal of Fish Diseases* 36: 171-194.

Treasurer, J.W. and Grant, A. (1997). The efficacy of hydrogen peroxide for the treatment of farmed Atlantic salmon, *Salmo salar* L. infested with sea lice (Copepoda: Caligidae). *Aquaculture* 148: 265-275.

Treasurer, J., Grant, A and Davis, P.J. (2000). Physical constraints of bath treatments of Atlantic salmon (*Salmo salar*) with a sea lice burden (Copepoda: Caligidae). *Contributions to Zoology* 69: 129-136.

Wilson, R.W. (2012). Aluminum. In: Wood CM, Farrell AP, Brauner CJ (Eds). Homeostasis and Toxicology of Non-Essential Metals. *Fish Physiology Vol. 31B*. 68-104.

Wood, C.M. and Jackson, E.B. (1980). Blood acid–base regulation during environmental hyperoxia in the rainbow trout (*Salmo gairdneri*). *Respiration Physiology* 42:351–372

Young, N.D., Crosbie, P.B., Adams, M.B., Nowak, B.F. and Morrison, R.N. (2007). *Neoparamoeba perurans* n. sp., an agent of amoebic gill disease of Atlantic salmon (*Salmo salar*). *International Journal of Parasitology* 37: 1469-1481.

Young, N.D., Dykova, I., Nowak, B.F. and Morrison, R.N. (2008). Development of a diagnostic PCR to detect *Neoparamoeba perurans*, agent of amoebic gill disease. *Journal of Fish Diseases* 29: 1-11.

Young, N.D., Dykova, I., Nowak, B.F. and Morrison, R.N. (2008). *Neoparamoeba perurans* is a cosmopolitan aetiological agent of amoebic gill disease. *Diseases of Aquatic Organisms* 78: 217-223.



NIVA: Norway's leading centre of competence in aquatic environments

NIVA provides government, business and the public with a basis for preferred water management through its contracted research, reports and development work. A characteristic of NIVA is its broad scope of professional disciplines and extensive contact network in Norway and abroad. Our solid professionalism, interdisciplinary working methods and holistic approach are key elements that make us an excellent advisor for government and society.



Norwegian Institute for Water Research

Gaustadalléen 21 • NO-0349 Oslo, Norway  
Telephone: +47 22 18 51 00 • Fax: 22 18 52 00  
[www.niva.no](http://www.niva.no) • [post@niva.no](mailto:post@niva.no)