


Acid Rain Research

REPORT 15/1988

Natural Organic
Acids:

**Their Role
in Freshwater
Acidification
and Aluminium
Speciation**



NIVA – REPORT

Norwegian Institute for Water Research  NIVA

Main Office
P.O.Box 33, Blindern
N-0313 Oslo 3
Norway
Phone (47 2) 23 52 80
Telefax (47 2) 39 41 29

Regional Office, Sørlandet
Grooseveien 36
N-4890 Grimstad
Norway
Phone (47 41) 43 033
Telefax (47 41) 42 709

Regional Office, Østlandet
Rute 866
N-2312 Ottestad
Norway
Phone (47 65) 76 752

Regional Office, Vestlandet
Brevikven 5
N-5035 Bergen - Sandviken
Norway
Phone (47 5) 95 17 00
Telefax (47 5) 25 78 90

Report No.:
Sub-No.: 2108
Serial No.:
Limited distribution:

Report Title: Natural organic acids: Their role in freshwater-acidification and aluminium speciation	Date: 11 March 1988
Author (s): Egil Gjessing Magne Grande Eirin Røgeberg	Project No 0-85244
	Topic group: Acid precipitation
	Geographical area: Norway
	Number of pages (incl. app.)

Contractor: Surface Water Acidification Program (SWAP)	Contractors ref. (or NTFN-No)
---	-------------------------------

Abstract: Samples containing humic substances, from an acidified area have been compared with corresponding samples from a less acidified area. The studies are based on an equilibrium-dialyzation, using artificial membranes. The waters are thoroughly characterized before and after dialyzation. Results may suggest that the sulphur reacts with the humus, to form a "sulfonic humus" having a lower pK-value. Toxicity tests, using fish, suggest that dialyzed samples from acidified areas are more toxic than samples from less acidified areas.

4 keywords, Norwegian

1. Humus
2. Forurning
3. Toksisitet
4. Dialyse


4 keywords, English

1. Humic Substances
2. Acidification
3. Toxicity
4. Dialyzation

Project leader


Egil Gjessing

For the Administration


Svein Stene-Johansen

ISBN - 82-577-1387-2

0-85244

NATURAL ORGANIC ACIDS: THEIR ROLE IN FRESHWATER ACIDIFICATION AND
ALUMINIUM SPECIATION

Oslo, 20. april 1988

Project leader: Egil T. Gjessing

CONTENTS

<u>Seksjon</u>	<u>Side</u>
1. INTRODUCTION	1
2. METHODS AND EXPERIMENTAL APPROACH	2
2.1 Strong and weak acids	2
2.1.1 Dialyzing technique	4
2.2 "Burning off"	4
2.3 Toxicity tests with fish	4
2.4 Samples	5
3. RESULTS	6
3.1 UV - H ₂ O ₂ -treatment	6
3.2 Dialyzation	6
4. DISCUSSION	9
4.1 UV/H ₂ O ₂	9
4.2 Dialyzation technique	10
4.3 Acidified surface waters	11
4.3.1 "Egil"-area	11
4.3.2 "Kim"-area	12
4.4 "Egil"-area/"Kim"-area/Hellerudmyra	13
4.4.1 H ⁺ -strong and weak acids	13
4.4.2 Aluminum	14
4.4.3 Sulphate/chloride	14
4.5 Artificially acidified Hellerudmyr water	14
4.6 Toxicity tests	16
5. CONCLUSION	16
5.1 Summary	17
6. References	19
7. 9 tables	20

1. INTRODUCTION

Research on freshwater acidification and loss of fisheries have centered on clear waters. The role of natural organic acids and in particular the interaction between organic acids and the strong acids in soils and surface water is poorly understood. Weak acids in natural coloured waters are mainly weak organic acids and aluminum compounds. Other weak acids such as silica and ammonia usually play only a minor role, except at very low aluminum and organic matter concentrations.

The natural organic matter in water originates from the soil and is a result of chemical and microbiological decomposition of terrestrial plants. This organic matter is acidic and consists of slightly negatively charged macromolecules which are retained in the soil by adsorption to mineral surfaces (Kodama and Schnitzer 1968, 1969; Gjessing and Gjerdahl 1975; Gjessing 1976). The ability to be retained in the soil depends on pH and the nature of the mineral surfaces (Gjessing 1976). This suggests that acid precipitation will effect the release of the soil organic matter into surface water both quantitatively and qualitatively (Gjessing 1976).

The natural organic matter present in water, the aquatic humus, will also change with changes in water chemistry such as pH. A change in pH effects the colour, the molecular size, complexation, the degree of solubility in organic solvents and ion - fixation properties (Gjessing 1976). The bioavailability of inorganic and organic aluminum species, associated with humus, is thus changed.

The purpose of the work presented below is:

1. to introduce and test a modified method for determination of weak acids in water and to use this method to characterize the content of weak acids in coloured water from acidified and less acidified areas.
2. based on the assumption that the toxic effect of species in water is related to the ability to penetrate cell walls, studies on the dialyzability of different types of water, using artificial membranes, are performed.

2. METHODS AND EXPERIMENTAL APPROACH

2.1 Strong and weak acids

Normally, the determinations of strong and weak acids are based on Gran's method (Gran, 1952). The method includes potentiometric pH titration and transformation of titration data to a straight line, so called "Gran plot". Several variations of the Gran procedure have been reported, partly due to an overestimate of total acidity due to CO_2 contamination of the titrant, and partly due to an overestimate of strong acidity at the expense of an underestimate of the weak acidity due to dissociated weak acids present in the low pH region of the titration curve. From a critical survey of the literature the modified Gran procedure described by Molvørsmyr and Lund (1983) seemed to be the best choice of method in order to avoid the most serious errors.

The titrations are carried out by coulometric generation of hydroxide ions, thus avoiding carbonate contamination of the titrant. A pH titration-range between 3.6 and 10.3 was assumed adequate for a majority of the fresh waters. With minor modifications this method was adapted for routine analysis of strong and weak acids (for details see Røgeberg, 1987).




Figure 1.

System used for dialyzing humus water. The dialyzing tube have a diameter of 16 mm and a theoretical "cut off" of about 12 000 MW. The purpose of the UV-radiation is to keep the system as sterile as possible.

2.1.1 Dialyzing technique

The equipment used for dialyzing the different water samples is illustrated on figure 1. The samples are dialyzed in distilled water. As can be seen at the upper right on the figure, the humus water is recirculated by a pump through several meters of a dialyzing tube (up to 18 meters in length) contained in a tank with a volume of approximately 30 liters. The dialyzing tube is surrounded by distilled water which is circulated in the tank and renewed several times during the run. The conductivity of this "washing water" never exceeded 0.4 mS/m, and before ending the run the conductivity of this water was below 0.15 mS/m. During the dialyzation the conductivity of the recirculated humus sample was measured. Some typical patterns of the reduction of conductivity of the water sample as a function of dialyzation time, are given in figure 2.

2.2 "Burning off"

The organic matter in the water samples is mineralized, using UV and H_2O_2 .

Ultraviolet (UV) radiation of water, to which is added minute amounts of hydrogen peroxide (H_2O_2), will mineralize the organic matter nearly completely.

Some of the samples referred to below have been UV/ H_2O_2 - treated and analyzed on relevant inorganic compounds.

The main purpose of this treatment is to see whether this is a possible approach to distinguish between inorganic and organic acids in water.

2.3 Toxicity tests with fish

Using Atlantic salmon (*Salmo Salar* L.), about one half of the samples are tested regarding fish toxicity (see table 7).

In these fish/water studies the toxicity of water is compared before and after membrane-dialyzation.

The screening test involves one liter of water, in which the survival of two small fishes (Atlantic Salmon about one gram each) is studied during a 96 hour period.

2.4 Samples

Hellerudmyra is a highly coloured water from a marsh area outside Oslo. This water source has been extensively used in our humus research during the last twenty years, and is apparently not affected by acid precipitation to any great extent. This water has also been the base for the Nordic Reference Humus and for the "NIVA-concentrate" (see table 3). The water used in the present experiment is the same water as that used for the isolation of the "Nordic" and the "NIVA-concentrate".

Artificial acidification is performed by adding either concentrated HCl or concentrated H_2SO_4 . The Hellerudmyr-water is acidified to below pH 2 and stored for 24 hours before dialyzation.

"Egil" and "Kim" are samples from the RAIN-project-area at Risdalsheia, an acidified region in southern Norway.

"Kim" is water from the outlet of a small "roofed" catchment where all the acid is removed from the collected precipitation, before it is sprinkled back into the catchment.

"Egil" is the water from the reference-area, where the collected acid-precipitation is sprinkled back to the catchment without any treatment.

Artificial salt-solution is prepared from the following inorganic salts (regarding chemical composition of the solution see table 8): $NaCl$, Na_2SO_4 , $Ca(NO_3)_2 \cdot 4H_2O$, $MgSO_4 \cdot 7H_2O$, K_2SO_4 , H_2SO_4 .

3. RESULTS

3.1 UV - H₂O₂-treatment

Seasonal samples from Hellerudmyra are compared. These four samples are also compared with one sample from "Egil". The results are summarized in table 1.

In general this treatment mineralizes essentially all the DOC (97%).

Comparing the pH in the water-samples from Hellerudmyra, with the pH of the sample from the Egil-catchment, it appears that essentially all the organic acid is removed from the Hellerudmyr samples whereas UV/H₂O₂-treated "Egil" retains a considerable concentration of H⁺-ions.

With regard to aluminum this oxidation treatment removes most of the reactive aluminum (RAL) from Hellerudmyr-water, however little from the Egil-sample. All the "non labile" aluminum (ILAL), which consists of organically bound complexes, is removed, which agrees with theory.

It should be emphasized that both the concentration of sulphate and in particular the concentration of nitrate are increased after oxidation with UV/H₂O₂. This increase might partly be due to organic nitrogen and sulphate connected to the humus molecule, but also due to a "complexation" of nitrate and sulphate to the humus molecule.

3.2 Dialyzation

The "membrane performance" is illustrated on figure 2. The dialyzation pattern on humus water (Hellerudmyra) is compared to salt and mineral acid solution.

According to the membrane specification (SPECTRATOR membrane tubing, SPECTRUM Medical Industries Inc.) the dialyzing tube operates in the molecular weight range of 12 000.




Figure 2

The change of conductivity of sample with time of dialyzation. The results suggest that SO_4^- -ions generally are "held back" by the membrane to a greater extent than the other inorganic ions.

As can be seen from figure 2 the discrimination of molecules is not only a matter of molecular size; most probably there is also an electrical charge effect, or a combination of size and charge. The results given in figure 2 indicate that sulphate ions are more resistant to penetrating the membrane than other anions, such as chloride. This again suggests that this artificial membrane surface, as natural membrane surfaces, has a net negative charge.

The chemical data of Humus water from Hellerudmyra, before and after dialyzation, are given in table 2 and table 3. Table 2 represents samples from four seasons and table 3 is based on water, sampled in July 1986. The results given in table 3 are partly "natural" Hellerudmyr-water, partly artificially acidified (HCl or H_2SO_4) and partly reference humus (Nordic fulvic acid and NIVA concentrate (before and after dialyzation)).

Comparing column 6, 7, 8 and 9 in table 3, which is the same water dialyzed (and analyzed) in four different runs, there is generally good agreement in pH (H^+), conductivity, aluminum and weak acid, regarding degree of retainment. The discrepancy regarding the other components is partly, probably mostly, due to analytical errors.

The middle section of table 3 represents acidified water from Hellerudmyra (using HCl or H_2SO_4), before and after dialyzation. As it appears from fig. 2 the dialyzing patterns are very different, comparing HCl and H_2SO_4 . In addition to Cl and SO_4 it is in particular the pH of the dialysates that are significantly different; it should be emphasized that there is lack of agreement between the concentration of H^+ of the dialyzed " H_2SO_4 -sample" and conductivity.

The right section of table 3 represents reference humus, the "Nordic fulvic acid reference humus" and "NIVA-concentrate". The former is isolated according to internationally accepted procedures, which involves adsorption of the organic matter to a synthetic adsorbant (XAD-8) and the "NIVA-concentrate", which is organic matter isolated from the same water as the "Nordic reference humus", however, by evaporation of membrane filtered water (0.45 μm) at reduced pressure and 30 $^{\circ}C$.

Comparing the dialysates of the two "Reference humics," the lower content of weak acids of the "Nordic reference fulvic acid" should be emphasized. It is also remarkable that both dialysates, as also the dialysates of the "normal" Hellerudmyr-water, contain a significant amount of strong acids.

In table 4, 5 and 6 are the results of membrane filtered and dialyzed samples from the "Egil-area" (reference area of the RAIN-product at Risdalsheia) and the "Kim-area" (the catchment at Risdalsheia where the acid precipitation is removed). These are samples taken during summer and fall 1986. In general the results suggest relatively large differences between the samples. As the duplicate runs (column 16 and 17 in Table 4, and column 31 and 32 in Table 6) show essentially the same results, indicating reasonably good reproducibility, these differences are probably due to changes with time in the chemical composition; implying that there are episodic changes.

Neither "Egil nor Kim" (with a few exceptions) have any strong acids in the dialysate, in contrast to the sample from Hellerudmyra. As the biological "activity" of these waters is an essential part of the acidification problem, the toxicity of several of these waters have been tested, using fish. The results in table 7 indicate the time of survival. The dotted line indicates: 1 fish still living.

4. DISCUSSION

4.1 UV/H₂O₂

The oxidation process involved in this treatment, removes essentially all the organic matter.

Comparing the four seasonal samples from Hellerudmyra (table 1), the pH increases to above 5.8 (from less than 4.5). This suggests that organic acid dominates the acidic properties of these waters.

With regard to aluminum the results in table 1 show that the "non labile aluminum" (ILAL) is "removable", and suggests that most of the aluminum is hydrolyzed.

Comparing the anions chloride, sulphate and nitrate, before and after oxidation, it appears from table 1 that there is a general increase after UV/H₂O₂-treatment. For the summer-humus containing more than 22 mg of carbon pr. liter the increase in sulphate concentration is in the range of 15 µeq/l.

This might be organic sulphur oxidized to SO₄, SO₄ "complexed" with humus or sulphate liberated from a "sulphonic humus ester". With regard to the NO₃ increase this is most probably due to oxidation of organically bound nitrogen.

The one sample from the "Egil-area" (ref. catchment at Rindalsheia), gives oxidation products which are similar to the Hellerudmyr-samples except from the pH. The pH of the oxidized sample from this acidified area is 4.8, which is more than a pH unit lower than in Hellerudmyr-water, indicating 15 µeq/l of mineral acids.

4.2 Dialyzation technique

The size of the pores of the dialyzing membrane is, according to the specification, equivalent to a molecular weight in the range of 12 000. Theoretically inorganic salt solutions should therefore easily penetrate through.

As shown in figure 2 this is not the case. It appears from the figure that CaCl₂ and HCl penetrate relatively easily, whereas Na₂SO₄ and the mixture of salts (see table 8) are resistant to penetrating the membrane.

Comparing the chemical composition of dialyzed inorganic salt-solution with that of natural water (Hellerudmyra) the results are relatively similar.

It should be emphasized that there are several uncertainties connected with the analytical results, as there is an ionic imbalance, particularly for the dialyzed inorganic salt solution.

Nevertheless, the performance of the dialyzing tube was unexpected regarding inorganic salts; in particular the penetration of sulphate ions, appeared to be slow. This suggests that the membrane surface has a negative charge, resulting in a rejection of anions. As SO_4^- -ions in aqueous solutions are relatively large, they are rejected to a greater extent than for instance the smaller chloride ions.

However, in spite of these unexpected properties, it is important to emphasize that cell membranes as such also generally have a negatively charged surface, so that the dialyzing experiments are valuable in interpreting the biological implications of surface waters containing humic substances.

4.3 Acidified surface waters

4.3.1 "Egil"-area

Samples from the "Egil-area" are representative for humus water from acidified waters. It appears indirectly from table 4 and 5 that in spite of pronounced differences in H^+ -concentration (50-160 $\mu\text{eq}/\text{l}$) in the membrane filtered waters, the resulting H^+ -concentration in the dialyzed water is relatively constant (10-20 $\mu\text{eq H}^+/\text{l}$). Sulphate in the sample-fraction that passes the dialyzing membrane also varies considerably (11-86 %, or 4-104 $\mu\text{eq SO}_4/\text{l}$). The SO_4^- -concentration in the dialyzed sample is also relatively constant (28-62 $\mu\text{eq SO}_4/\text{l}$, with exception of one value). However, as indicated earlier, there are generally remarkably high concentrations of sulphate in the dialyzed waters.

With regard to the penetration of SO_4^- through the dialyzing membrane, a comparison of the results of the synthetic solution (see fig.2 and table 8) and the "Egil-water sample" (for instance No.23-24) reveals some interesting considerations:

Of the SO_4 in the synthetic water, only 23% (1.4 mg SO_4 /l) has passed the dialyzing membrane at "steady state", whereas in the natural sample more than 60% (4.5mg SO_4 /l) has passed the membrane under the same experimental conditions. This may imply that SO_4 is "helped" to the membrane when natural water is dialyzed, possibly together with the organic matter.

The aluminum results are very complex and confusing. Generally 50% of the total aluminium passes the dialyzing membrane, whereas roughly 1/4 to 1/2 of the labile form penetrates the membrane.

The lack of "persistent" aluminum results, reveals the complexity of the aluminum chemistry: During the dialyzation the chemistry will gradually change, and the pattern of these changes will depend on the chemistry of the original water. This implies that the concentration gradients of the different aluminum species during the dialyzation will differ and consequently the amounts that are penetratable through the dialyzing membrane depend on numerous conditional factors.

Essentially all the strong acids in the "Egil"-samples apparently penetrate through the membrane, suggesting that all mineral acids are removed from the dialyzates. With regard to the weak acids the degree of penetration through the membrane varies considerably. In spite of the fact that the concentration of weak acids in the original samples is relatively constant, the degree of retainment by the membrane differs significantly. Any attempts to correlate these differences with other parameters have so far failed.

4.3.2 "Kim"-area

These samples are from the deacidified area at Risdalsheia, the area where the acids in the precipitation are removed.

PH in the dialyzed samples from the "Kim"-area is not significantly different from that of the "Egil"-area (in the range of 4.6 - 5.0).

The sulphate in the original samples from the "Kim"-area does not vary to any great extent (with one exception, within 2.0 - 2.7 mg SO₄/l), in contrast to the samples from the "Egil"-area. Only 1/4 the sulphate penetrates the membrane. Presently it is not known whether this resistance to dialysis is due to low concentration gradient, charge phenomena or humus complexation.

4.4 "Egil"-area/"Kim"-area/Hellerudmyra

Chemical composition before and after dialyzation.

4.4.1 H⁺-strong and weak acids

In table 9 the results of the three different sample-types are summarized.

More H⁺ and almost all strong acids (SA) are removed from the naturally acidified water-samples compared to water from Hellerudmyra where only 50% SA is removed. With regard to the total content of strong acid, the samples from the acidified areas do contain about twice as much as water from the less acidified area (Hellerudmyra). Comparing the content of weak acids in these two sample-sets, it appears from table 9 that there are insignificantly less weak acids in the water from Hellerudmyra. However, by relating the content of weak acids to that of organic matter (DOC) the results indicate that "Egil"-water contains two times more and "Kim"-water about 1 1/2 times more than Hellerudmyra. The same ratios are apparently valid for weak acids in dialyzed samples.

4.4.2 Aluminum

Significantly less aluminum passes through the dialyzing membrane in water from the "Kim"-area compared to the sample from the "Egil"-area.

With regard to RAL (Reactive aluminum) the results indicate a relationship between percent of membrane penetrable aluminum and pH in the original water. In water from Hellerudmyra no LAL (Labile aluminum) apparently passes the membrane, whereas in acidified water ("Kim" and "Egil") significant amounts of LAL penetrate the dialyzing membrane.

4.4.3 Sulphate/chloride

Of the major anions, nitrate and chloride behave differently with regard to penetration through the membrane compared to the divalent sulphate. However, there is no significant difference between the three sample sets ("Hellerud", "Egil" and "Kim") in this respect. Generally most of the univalent anions (Cl and NO_3), passes the dialyzing membrane, whereas more than 1/2 of the sulphate is retained.

As indicated earlier, this difference in behaviour might be due to charge phenomena, molecular size or due to an association between sulphate and the humus molecule.

4.5 Artificially acidified Hellerudmyr water

The results from dialyzation of acidified humus water given in table 3 and in figure 2, clearly show that the addition of mineral acids to humus water does change the dialyzation behaviour dramatically and also the nature of the final dialyzation product:

Comparing HCl acidification (for instance, sample no 56, with dialyzed natural sample no 54), it appears from table 3 that all the components, including DOC and SO_4 , pass the membrane to a significantly greater extent. There are several possible explanations for these important findings:

1. The molecular size of humus is reduced; the difference in DOC retainment between natural and HCl acidified water is, however, not remarkable (18.4 versus 16.1). This indicates that the transport of ions through the membrane is probably not by means of the organic matter.
2. The acidification results in a release of humus complexed cations; particularly Mg, ILAL and Fe apparently pass the membrane more easily.
3. The negative charge of the membrane-surface might be reduced (so also the charge of the humus molecule).

The conductivity of the HCl-sample (figure 2 "Hellerudmyra 55/56") apparently levels off at the same value as the natural sample, however, at a longer dialyzing time.

Comparing the dialyzing pattern of the H_2SO_4 -acidified humus water, with that of HCl-acidified humus-water (figure 2), "Steady state" is apparently not reached after 6 days!

With regard to the composition of the final dialyzation product of this H_2SO_4 treated water sample, it appears from table 3 (sample no 58), that except for sulphate and DOC the results are not remarkably different from that of the HCl-treated sample (sample no.54). The relatively high concentration of SO_4 is probably due to, as indicated earlier, charge effects on the membrane, as the concentration of SO_4 in the dialysate is relatively high even at a late stage of the dialyzation run when the pH is relatively high.

4.6 Toxicity tests

As expected "Egil and Kim"-water (from Risdalsheia) are generally more toxic to yearlings of salmon than Hellerudmyr water. It can also be seen from table 7 that the waters are less toxic after dialyzation; which also should be expected. These experiments indicate, however, some results of significant importance: Dialysates of the most acidic waters (sample 33, 31, 29, 27, 23, 19 and 11) are apparantly more toxic than dialysates from less acidic samples. It should also be pointed out that in 5 of these 7 apparantly toxic "dialysates" there has also been found a significant amount of strong acids.

5. CONCLUSION

- * UV/H₂O₂ - treatment of natural water removes essentially all the organic matter. The composition of the "residual", such as pH, show differences between water from acidified and less acidified areas. This may indicate differences in the mineral acid/organic acids ratio.
- * Membrane dialyzation, based solely on concentration gradient (sample/distilled water, without pressure) gives satisfactorily reproducible results and is apparently a more useful technique than most other dialyzation methods for estimating passage of compounds into organisms.
- * Acid and acidified samples penetrate the membrane used more slowly than less acidified samples or deacidified samples.
- * All the strong acids (SA), and most of the labile aluminum (LAL) pass the membrane from "natural" acidified waters. In contrast, only 50 per cent SA and essentially no LAL penetrate the membrane from the less acidified samples.

- * There is no significant correlation between weak acids (WA) and DOC, either within sample types nor all samples considered.
- * The sample from the acidified area apparently contains more weak acids (WA) per mg carbon (min, 20 $\mu\text{eq}/\text{mgC}$) than samples from the less acidified area (13 $\mu\text{eq}/\text{mgC}$).
- * As the sulphate concentration per mg carbon in dialyzed samples are significantly higher in the acid waters compared to the less acidified, the results may suggest that the sulphur reacts with the humus, to form a "sulfonic humus" having a lower pK-value.
- * Toxicity tests, using fish, suggest that dialyzed samples from acidified areas are more toxic than samples from less acidified areas.

5.1 Summary

Samples, containing humic substances, from an acidified area have been compared with corresponding samples from a less acidified area. Samples from a deacidified area (RAIN-project) and artificially acidified humus samples (laboratory acidified) are also included.

The studies are based on an equilibrium-dialyztion, using artificial membranes. The waters are thoroughly characterized before and after dialyztion. In the studies are also included toxicity tests of dialyzed and undialyzed water.

A potential method (using UV/H₂O₂ mineralization) for estimating the content of mineral acids in waters is described. The description of a modified method for the determination of weak and strong acids is also included.

The results suggest that the artificial dialyzation tubes have a negative surface-charge in the actual pH region, which hinders the penetration of anions, such as SO_4 . However, as these properties are typical also for natural membranes, the technique may be valuable for the interpretation of biological effects.

Equilibrium dialyzation of three types of samples: acidified, non-acidified and deacidified humus water, show that even with considerable variance in the pH of the original samples, the pH of the dialyzed "product" are remarkably constant (pH 4.7 to 5.0).

The content of strong acid (SA) is, as expected, much higher in the water from the acidified area. In contrast to the non-acidified waters, the dialyzation removes almost all the SA from these waters.

In the dialyzed water of the non-acidified sample, there are apparently insignificantly less weak acids (WA). However, when related to DOC, the WA-content is one half compared to the acidified water.

Labile aluminum (LAL, the potentially most toxic Al-fraction to fish) does not penetrate the membrane from non-acidified waters, whereas in acidified waters, a significant fraction of LAL penetrates through the membrane.

Sulphate dialyzes differently compared to the other major anions, indicating a chemical association with the humic substances.

The fish toxicity of dialyzed acidified humus water is apparently higher than the corresponding non-acidified waters. The chemical parameters used to describe differences in the chemical composition of the various water samples, are apparently not sufficient to explain why this may be so.

6. References

Gjessing, E.T. and Gjerdahl, T.: (1975) "Electromobility of aquatic humus. Fractionation by the use of the isoelectric focusing technique". pp. 43-51 (1975). In: Povoledo, D. and Golterman, H.L. (ed.). "Humic substances - Their structure and function in the biosphere". Proceedings of an international meeting held at Nieuwersluis, the Netherlands May 29.-31., 1972. Pudoc, Wageningen 1975.

Gjessing, E.T.: (1976) "Physical and chemical characteristics of aquatic humus". Ann Arbor Science Publishers Inc. pp. 1-120. Ann Arbor, Mich.

Gran, G.: (1952) "Determination of equivalence point in potentiometric titrations". Part II. Analyst 17, 661-671.

Kodama, H. and Schnitzer, M.: (1968) "Effects of interlayer cations on the adsorption of a soil humic compound by montmorillonite", Soil.Sci. 106(1), 73-74.

Kodama, H. and Schnitzer, M.: (1969) "Thermal analysis of a fulvic acid-montmorillonite complex", Proceedings of the International Clay Conference, Tokyo, (Jerusalem: Israel University Press, 1969), pp. 765-774.

Molværsmyr, K. and Lund, W.: (1983) "Acids and bases in fresh-water. Interpretations of results from Gran plots". Water Res. 17, 303-307.

Røgeberg, E.: (1987) "A coulometric Gran titration method for the determination of strong and weak acids in freshwater". Acid Rain Res. Report 12/1987, (Norwegian Inst. Water Research, Oslo), 28 pp.

Table 1. Membrane filtered (MF: 0.45 µm) and Mineralized (UV/H₂O₂) samples from Hellerudmyra and "Egil area".

	HELLERUDMYRA (less acidified area)														"Egil" - catchment acidified area					
	Winter				Spring				Summer				Fall				Fall			
	MF conc µeq /l		UV/H ₂ O ₂ conc µeq /l		MF conc µeq /l		UV/H ₂ O ₂ conc µeq /l		MF conc µeq /l		UV/H ₂ O ₂ conc µeq /l		MF conc µeq /l		UV/H ₂ O ₂ conc µeq /l		MF conc µeq /l		UV/H ₂ O ₂ conc µeq /l	
pH	4.53	29	6.20	0	4.51	31	5.84	1	4.35	44	6.79	0	4.37	43	5.91	1	4.00	100	4.82	15
mS/m COND	2.77	-	1.82	-	2.15	-	1.34	-	2.88	-	1.56	-	3.28	-	2.04	-	7.19	-	4.38	-
µg/l Fe	300	-	190	-	260	-	180	-	713	-	530	-	306	-	195	-	240	-	220	-
µg/l Mn	21	-	15	-	15	-	15	-	13	-	12	-	17	-	12	-	12	-	18	-
mg Pt/Color	105	-	10	-	107	-	8	-	242	-	16	-	133	-	7	-	75	-	2	-
mg/l Ca	1.13	56	1.16	58	0.78	40	0.79	39	1.01	50	1.00	50	1.30	65	1.30	65	0.46	23	0.49	25
mg/l Mg	0.25	21	0.24	20	0.16	13	0.16	13	0.18	15	0.18	15	0.27	22	0.26	21	0.36	30	0.39	32
mg/l K	0.07	2	0.09	2	0.20	5	0.4	5	0.03	1	0.03	1	0.07	2	0.10	3	0.39	10	0.39	10
mg/l Na	1.01	44	1.05	46	0.64	28	0.63	27	0.75	33	0.72	31	0.98	43	1.02	44	2.62	114	2.69	117
µg/l Al	298	-	215	-	185	-	159	-	380	-	381	-	319	-	200	-	590	-	450	-
µg/l RAL	235	-	25	-	155	-	0**	-	304	-	46	-	268	-	19	-	500	-	390	-
µg/l ILAL	183	-	0**	-	114	-	0**	-	271	-	0**	-	204	-	0**	-	218	-	0**	-
µg/l LAL	52	6	25	3	41	5	0**	-	35	4	46	5	64	7	19	2	282	31	390	43
mg/l Cl	1.3	37	1.5	42	1.5	42	1.6	45	0.7	20	0.9	25	1.3	37	1.3	37	2.6	73	2.7	75
mg/l SO ₄	3.5	73	3.7	77	3.5	73	3.4	71	1.5	31	2.2	46	4.6	96	4.5	94	4.4	92	4.8	100
µg N/l NO ₃	15	1	93	7	9	1	82	6	7	1	54	4	5	0	91	7	620	44	685	49
mg C/l DOC	11.8	-	0.3	-	9.4	-	0.2	-	22.9	-	0.5	-	13.9	-	0.5	-	9.4	-	0.6	-
I/cm UV-abs	0.49	-	0.00	-	0.44	-	0.00	-	1.09	-	0.00	-	.615	-	0.00	-	0.39	-	0.00	-
ACID																				
Σ cations	-	158	-	129	-	122	-	85	-	147	-	102	-	182	-	136	-	308	-	242
Σ anions	-	111	-	126	-	116	-	122	-	52	-	75	-	133	-	138	-	209	-	224
Deficit	-	47	-	3	-	6	-	-37	-	95	-	27	-	49	-	-2	-	99	-	18

**> <10=0

Table 2. Membrane filtered (MF: 0.45 µm) and Dialyzed (MW 12.000) samples from Hellerudmyra and "Egil area".

Column No	1				2				3				4				5							
	HELLERUDMYRA (less acidified area)																"Egil" - catchment acidified area							
	Winter				Spring				Summer				Fall				Fall							
	MF*)		Dial		MF*)		Dial		MF*)		Dial		MF*)		Dial		MF*)		Dial		MF*)		Dial	
	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l	conc	µeq /l
pH	4.53	29	4.91	12	4.51	31	4.87	14	4.35	44	4.67	21	4.37	43	4.86	14	4.00	100	4.78	14				
mS/m COND	2.77	-	0.94	-	2.15	-	0.94	-	2.88	-	1.43	-	3.28	-	1.10	-	7.19	-	1.11	-				
µg/l Fe	300	-	260	-	260	-	310	-	713	-	630	-	306	-	290	-	240	-	178	-				
µg/l Mn	21	-	10	-	15	-	9	-	13	-	9	-	17	-	8	-	12	-	4	-				
mgPt/Color	105	-	104	-	107	-	120	-	242	-	219	-	133	-	124	-	75	-	40	-				
mg/l Ca	1.13	56	0.50	25	0.78	40	0.46	23	1.01	50	0.68	34	1.30	65	0.57	28	0.46	23	0.19	10				
mg/l Mg	0.25	21	0.09	7	0.16	13	0.08	7	0.18	15	0.12	9	0.27	22	0.11	9	0.36	30	0.11	9				
mg/l K	0.07	2	0.02	1	0.20	5	0.05	1	0.03	1	0.01	0	0.07	2	0.04	1	0.39	10	0.05	1				
mg/l Na	1.01	44	0.13	6	0.64	28	0.13	6	0.75	33	0.13	6	0.98	43	0.15	7	2.62	114	0.26	11				
µg/l Al	298	-	240	-	185	-	205	-	380	-	347	-	319	-	258	-	590	-	263	-				
µg/l RAL	235	-	178	-	155	-	146	-	304	-	283	-	268	-	197	-	500	-	219	-				
µg/l ILAL	183	-	112	-	114	-	89	-	271	-	224	-	204	-	131	-	218	-	156	-				
µg/l LAL	52	6	66	7	41	5	57	6	35	4	59	7	64	7	66	7	282	31	63	7				
mg/l Cl	1.3	37	0.2	6	1.5	42	0.1	3	0.7	20	0.1	3	1.3	37	0.0	0	2.6	73	0.2	6				
mg/l SO ₄	3.5	73	1.1	23	3.5	73	1.1	23	1.5	31	0.5	10	4.6	96	1.1	23	4.4	92	1.7	35				
µg N/l NO ₃	15	1	3	0	9	1	3	0	7	1	8	1	5	0	7	1	620	44	5	0				
mg C/l DOC	11.8	-	7.3	-	9.4	-	6.3	-	22.9	-	15.3	-	13.9	-	10.5	-	9.4	-	4.6	-				
I/cm UV-abs	0.49	-	0.34	-	0.44	-	0.33	-	1.09	-	0.85	-	0.62	-	0.43	-	0.39	-	0.23	-				
ACID																								
Σ cations	-	158	-	58	-	122	-	57	-	147	-	77	-	182	-	66	-	308	-	52				
Σ anions	-	111	-	29	-	116	-	26	-	52	-	11	-	133	-	24	-	209	-	41				
Deficit		47	-	29	-	6	-	31	-	95	-	66	-	49	-	42	-	99	-	11				

*) Membrane filter (0.45 µm)

Table 3. Membrane filtered (MF:0.45µm) and dialyzed (MW 12,000) samples from "Hellerudmyra", "Hellerudmyra" artificially acidified, "NIVA-CONC" and Nordic Fulvic acid", (sampled July -86) Z denotes: passage through dialyzing membrane.

Column No	"HELLERUDMYRA"										"HELLERUDMYRA" ARTIFICIALLY ACIDIFIED										"REFERENCE HUMUS"																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
	7					8					9					10					11					12					13		14																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441

Table 4. Membrane filtered (MF: 0.45µm) and Dialyzed (MW 12.000) samples from "Egji-ara" ("natural acidified"). Z denotes: passage through dialyzing membrane.

Sample No	15			16			17			18			19			20			21			22			23			24			25							
	1	2		3	4		5	6		7	8		9	10		11	12		13	14		15	16		17	18		19	20		21	22		23	24		25	26
	Duplicate																																					
pH	4.1	4.7	75	4.0	4.7	80	4.0	4.8	85	4.0	5.0	90	4.3	4.8	68	4.3	4.7	60	4.0	4.9	87	3.9	4.9	50	3.8	4.8	90	4.1	4.8	79	3.8	4.9	92					
ms/m Cond	3.64	1.52	58	6.70	1.65	75	6.75	1.73	74	6.93	1.12	84	3.14	1.45	54	3.54	1.59	55	6.23	1.70	73	7.60	1.63	79	8.81	1.77	80	4.80	1.44	70	8.10	1.31	84					
µg/l Fe	350	300	14	330	220	33	330	240	27	320	110	66	120	110	8	430	410	5	210	200	5	170	120	29	170	130	24	130	120	7	170	100	41					
µg/l Mn	20	20	0	<20	20	-	20	<20	0	20	<20	-	<20	<20	-	20	<20	-	40	30	25	30	20	33	20	<20	-	<20	<20	-	<20	<20	-					
mgPt/Color	160	160	0	110	86	12	110	110	0	110	80	27	90	83	8	180	180	0	110	90	18	70	55	21	64	55	14	87	86	1	60	50	17					
mg/l Ca	1.0	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1	<1	-	<1.0	<1.0	-					
mg/l Mg	0.15	0.20	-	0.30	0.20	33	0.30	0.20	33	0.30	0.10	66	0.10	<0.1	-	0.20	0.15	25	0.40	0.25	38	0.40	0.20	50	0.50	0.30	40	0.20	0.15	25	0.45	0.15	67					
mg/l K	0.15	0.10	33	0.20	0.10	50	0.20	0.10	50	0.85	0.10	88	0.20	0.10	50	0.70	0.20	71	0.25	<0.1	-	0.55	0.10	82	0.60	0.10	83	0.50	<0.1	>80	0.55	0.10	82					
mg/l Na	1.1	0.4	64	2.0	0.6	70	2.60	0.7	73	3.0	0.2	93	1.2	1.1	7	1.8	0.8	56	2.4	0.5	79	3.3	0.7	79	2.9	0.6	79	2.3	0.7	70	2.9	0.5	83					
µg/l Al	560	440	21	670	360	46	650	380	41	640	390	39	240	180	67	580	580	0	550	330	40	600	280	53	600	400	33	300	310	0	600	310	48					
µg/l ILAL	370	340	8	320	110	66	310	120	61	290	120	59	60	30	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
µg/l LAL	190	100	47	350	250	29	340	260	24	350	270	23	180	150	17	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
mg/l Cl	2.0	0.6	70	3.0	0.2	93	3.0	0.3	90	5.0	0.4	92	1.7	0.4	76	2.6	0.7	73	1.7	0.4	76	4.6	0.2	96	5.3	0.2	96	4.6	0.4	91	7.0	0.2	97					
mg/l SO ₄	2.6	2.0	23	6.6	2.2	67	6.8	2.9	57	5.4	1.4	74	1.8	1.6	11	3.0	2.0	33	7.0	2.5	64	8.0	3.0	63	8.0	4.0	50	2.5	2.0	-	5.0	2.0	60					
µg N/l NO ₃	130	<50	-	220	<50	-	220	<50	-	530	<50	>91	110	<50	-	250	<50	>80	920	<50	>95	670	<50	>93	700	<50	>93	180	<50	>72	440	<50	>89					
mg C/l DOC	17.7	16.3	-	13.8	9.3 ¹	-	13.8	9.8 ¹	-	13.7	8.6 ¹	-	9.3 ¹	8.8 ¹	-	-	-	-	-	-	-	9.2 ¹	6.4 ¹	-	8.6 ¹	6.3 ¹	-	9.7 ¹	8.8 ¹	-	7.8 ¹	5.5 ¹	-					
I/cm UV-abs	.785	.728	7	.613	.415	32	.613	.437	29	.610	.385	37	.417	.389	7	-	-	-	-	-	-	.412	.284	31	.378	.278	26	.428	.388	9	.347	.249	28					
FTU	0.2	1.3	-	0.25	0.75	-	0.25	0.55	-	0.35	0.75	-	0.20	0.45	-	0.30	0.60	-	0.25	0.70	-	0.25	0.50	-	0.20	0.50	-	0.30	0.45	-	0.25	0.40	-					
µgN/l Ammon	110	5	95	260	80	69	240	140	42	460	100	78	170	65	-	220	120	45	500	240	52	500	280	44	460	210	54	310	250	19	230	190	17					
µmol/l Strong acid	37	-15	100	79	-12	100	85	-18	100	58	-30	100	3	-14	100	29	-21	100	105	142	87	105	27	74	135	3	98	72	21	71	141	-7	100					
µmol/l Weak acid	249	199	20	274	171	38	263	187	29	292	166	43	193	149	23	275	253	8	244	182	24	272	144	47	250	186	26	189	162	13	223	171	23					

¹) TOT.AL - ILAL

²) Estimated from UV-abs.

ILAL is determined manually and both Al and ILAL is determined by atomic abs.

Table 5. Membrane filtered (MF: 0.45 µm) and Dialyzed (MW 12.000) samples from "Egil-area" ("natural acidified"). % denotes; per cent passage through dialyzing membrane.

Column No	26					27					28				
Sample No	41		42			47		48			49		50		
	MF		DIAL		%	MF		DIAL		%	MF		DIAL		%
	conc	µeq /l	conc	µeq /l		conc	µeq /l	conc	µeq /l		conc	µeq /l	conc	µeq /l	
pH	4.05	90	5.13	8	91	4.06	87	4.95	11	87	4.03	93	5.03	10	89
mS/m COND	8.05	-	1.29	-	84	7.41	-	1.36	-	82	8.57	-	1.30	-	85
µg/l Fe	148	-	66	-	55	166	-	126	-	32	116	-	74	-	36
µg/l Mn	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
mgPt/Color	56	-	47	-	16	72	-	61	-	15	53	-	38	-	28
mg/l Ca	0.52	26	0.47	23	10	0.39	19	0.28	14	28	0.46	23	0.33	16	28
mg/l Mg	0.42	35	0.18	15	57	0.23	27	0.19	16	42	0.44	36	0.22	19	50
mg/l K	0.50	13	0.10	3	80	0.46	12	0.09	2	80	0.52	13	0.08	2	85
mg/l Na	2.83	123	0.30	13	89	2.65	115	0.35	15	87	3.30	144	0.33	14	90
µg/l RAL	430	-	170	-	60	385	-	210	-	45	440	-	194	-	56
µg/l ILAL	94	-	99	-	0	118	-	125	-	0	91	-	94	-	0
µg/l LAL *)	336	37	71	8	79	267	30	85	9	68	349	39	100	11	71
mg/l Cl	4.8	135	<0.1	2	99	5.7	161	<0.1	2	99	6.8	192	<0.1	2	99
mg/l SO ₄	8.2	129	2.5	52	60	4.5	94	2.3	48	49	5.6	117	2.7	56	52
µg N/l NO ₃	930	66	6	0	99	575	41	4	0	99	725	52	4	0	99
mg C/l DOC	6.4	-	5.5	-	14	7.7	-	6.1	-	21	5.9	-	3.8	-	36
I/cm UV-abs	.284	-	.221	-	22	.331	-	.275	-	18	.224	-	.173	-	29
Σ cations	-	324	-	70	78	-	290	-	67	77	-	348	-	72	79
Σ anions	-	330	-	54	84	-	296	-	59	80	-	361	-	58	84
Deficit	-	-6	-	16	-	-	-	-	0	-	-	-	-	14	-

*) RAL - ILAL

Table 6. Membrane filtered (MF: 0.45 µm) and dialyzed (MW 12,000) samples from "Kim-area" ("artificially deacidified"). Z denotes passage through dialyzing membrane.

Column No	29			30			31			32			33			34										
	Sample No	7	8	MF Dial pass	Z MF Dial pass	MF conc µeq/l	Dial conc µeq/l	Z MF Dial pass	MF conc µeq/l	Dial conc µeq/l	Z MF Dial pass	MF conc µeq/l	Dial conc µeq/l	Z MF Dial pass	MF conc µeq/l	Dial conc µeq/l	Z MF Dial pass									
pH	4.3	4.6	50	3.9	4.9	90	4.03	93	5.01	10	89	4.03	93	4.96	11	88	4.17	68	4.77	17	75	4.19	65	4.90	13	80
mS/m COND	3.57	1.62	55	6.78	1.04	85	7.89	-	1.47	-	81	7.89	-	1.32	-	83	4.99	-	1.59	-	68	5.03	-	1.30	-	34
µg/l Fe	340	300	12	200	160	20	182	-	104	-	44	182	-	104	-	44	310	-	2.60	-	16	270	-	210	-	22
µg/l Mn	< 20	< 20	-	20	< 20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
mgPt/Color	160	150	6	93	94	0	60	-	55	-	8	60	-	52	-	13	141	-	144	-	0	99	-	95	-	4
mg/l Ca	< 1	< 1	< 1	< 1	< 1	-	0.54	27	0.52	26	4	0.54	27	0.42	21	22	0.22	11	0.29	14	0	0.26	13	0.25	12	4
mg/l Mg	0.15	0.10	33	0.35	0.15	57	0.41	34	0.18	15	56	0.41	34	0.21	17	49	0.20	16	0.20	16	0	0.22	18	0.18	15	18
mg/l K	0.25	0.10	60	0.50	0.10	80	0.38	10	0.11	3	71	0.38	10	0.07	2	82	0.28	7	0.12	3	57	0.25	6	0.08	2	68
mg/l Na	2.1	0.9	57	2.5	0.5	80	2.6	113	0.37	16	86	2.6	113	0.28	12	89	1.77	77	0.50	22	72	1.91	83	0.43	19	77
µg/l RAL	-	-	-	-	-	-	370	-	140	-	62	370	-	189	-	49	415	-	355	-	14	385	-	313	-	19
µg/l ILAL	140	130	7	-	-	-	85	-	88	-	0	85	-	102	-	0	197	-	224	-	0	194	-	194	-	0
µg/l LAL *	-	-	-	-	-	-	285	32	52	6	82	285	32	87	10	69	218	24	131	15	40	191	21	119	13	38
mg/l Cl	3.8	0.6	84	8.0	0.4	95	5.8	164	0.1	3	99	5.8	164	<0.1	3	99	3.2	90	<0.1	3	99	3.8	107	<0.1	3	99
mg/l SO ₄	2.2	2.0	10	2.0	1.5	25	4.5	94	30	62	33	4.5	94	3.2	67	29	2.5	54	1.9	40	24	2.7	56	1.9	40	30
µg N/l NO ₃	< 50	< 50	-	60	< 50	-	650	46	5	0	99	650	46	5	0	99	117	8	4	0	95	178	13	4	0	97
mg C/l DOC	18.7	16.7	-	-	9.7	-	6.7	-	5.3	-	21	6.7	-	5.3	-	21	13.3	-	11.8	-	11	10.8	-	8.9	-	18
I/cm UV-abs	.861	.779	10	-	.445	-	.290	-	.247	-	15	.290	-	.232	-	20	.624	-	.569	-	9	.474	-	.420	-	11
FTU	0.30	0.60	-	0.30	0.45	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
µgN/l Ammonia	30	15	-	55	130	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
µmol/l Strong acid	29	-	35	119	13	89	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
µmol/l Weak acid	283	253	10	210	189	10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
I cations	-	-	-	-	-	-	309	-	76	-	304	-	74	-	74	-	203	-	87	-	87	-	206	-	74	-
I anions	-	-	-	-	-	-	304	-	71	-	304	-	70	-	70	-	152	-	42	-	42	-	163	-	43	-
Deficit	-	-	-	-	-	-	5	-	5	-	5	-	4	-	4	-	51	-	45	-	45	-	43	-	31	-

*) RAL - ILAL 1) Estimated from UV-abs.

Table (7) Toxicity test of membranefiltered and dialyzed water

SAMPLE	pH	ms/l	µg/l		mg/l		µgN/l	UV-abs	µmol/l		FTU	xx)	REMARKS
			TOT AL	ILAL	SO ₄	NO ₃			Strong acid	Weak acid			
1	4.2	3.64	560	370	2.6	130	0.785	37	249	0.20	Em		
2	4.7	1.52	440	340	2.0	<50	0.728	-15	199	1.3	Ed		†
3	4.0	6.70	670	350	6.6	220	0.613	79	274	0.25	Em		†
4	4.7	1.65	360	250	1.1	<50	0.415	-12	171	0.75	Ed		
5	4.0	6.93	640	350	5.4	530	0.601	58	292	0.35	Em		††
6	5.0	1.12	390	270	1.7	<50	0.385	-30	166	0.75	Ed		
7	4.3	3.57	640	500	140	<50	0.861	29	283	0.30	Km		†
8	4.6	1.62	610	480	130	2.0	0.779	-35	253	0.60	Kd		
9	4.3	3.14	240	180	60	1.0	0.417	3	193	0.20	Em		
10	4.8	1.45	180	150	30	1.6	0.385	-14	149	0.45	Ed		†
11	4.0	6.75	650	340	6.8	220	0.613	85	263	0.25	Em		†
12	4.8	1.73	380	260	120	2.9	0.437	-18	187	0.55	Ed		†
13	4.3	3.54	580	-	3.0	250	-	29	275	0.30	Em		
14	4.7	1.59	580	-	2.0	<50	-	-21	253	0.60	Ed		
15	4.3	3.00	530	-	90	3.0	<50	-	23	283	0.25	Hm	
16	4.9	1.40	500	-	2.5	<50	-	18	211	0.80	Hd		
17	4.3	3.00	530	-	3.0	<50	-	50	281	0.25	Hm		
18	4.8	1.66	500	-	3.0	<50	-	20	233	0.45	Hd		
19	1.8	-	550	-	4.0	<50	-	-	-	0.50	-		HCl
20	4.2	2.40	280	-	2.0	<50	-	100	170	1.1	Hd		††
21	4.3	3.27	550	-	40	3.0	<50	-	36	272	0.55	Hm	†
22	4.7	1.76	480	-	2.5	<50	-	19	243	1.1	Hd		†
23	4.0	6.23	550	-	7.0	920	-	105	244	0.25	Em		†
24	4.9	1.70	330	-	2.5	<50	0.437	14	182	0.70	Ed		†
25	4.2	2.11	20	0	1.0	<50	0.487	56	143	0.30	Hm		†
26	4.7	1.08	140	-	1.0	<50	-	28	154	0.50	Hd		
27	3.9	7.60	600	-	8.0	670	0.412	105	272	0.25	Em		†
28	4.9	1.63	280	-	3.0	<50	0.284	27	144	0.50	Ed		†
29	3.8	8.10	600	-	5.0	440	0.347	140	223	0.25	Em		†
30	4.9	1.31	310	-	2.0	<50	0.249	-7	171	0.40	Ed		†
31	3.8	8.81	600	-	8.0	700	0.378	135	250	0.20	Em		†
32	4.8	1.77	400	-	4.0	<50	0.278	3	186	0.50	Ed		†
33	3.9	6.78	500	-	2.0	60	-	119	210	0.30	Km		†
34	4.9	1.04	430	-	1.5	<50	0.445	13	189	0.45	Kd		†
35	4.1	4.80	300	-	2.5	180	0.428	72	189	0.30	Em		†
36	4.8	1.44	310	-	2.0	<50	0.388	21	162	0.45	Ed		†

xx) H = Hellerudmyra
 E = "Egill-area"
 K = "Kim-area"
 m = membrane filtered
 d = dialyzed

Hours of exposure
 ★ Fish jumped out
 † 1 fish dead

Table 8. Composition of a salt solution and sample from Hellerudmyra (July -86) before and after dialyzation.

Sample	61		62		59		60	
	Salt solution				Hellerudmyra			
	conc	MF µeq/l	conc	DIAL µeq/l	conc	MF µeq/l	conc	DIAL µeq/l
pH	4.22	60	4.89	13	4.39	41	4.91	12
COND mS/m	4.14	-	1.75	-	3.03	-	1.76	-
Colour mg Pt/l	-	-	-	-	188	-	176	-
DOC mg C/l	-	-	-	-	19.6	-	12.5	-
Fe µg Fe/l	-	-	-	-	570	-	480	-
Ca mg Ca/l	0.27	13	0.07	3	1.15	57	0.98	49
Mg mg Mg/l	0.32	26	0.25	21	0.24	20	0.24	20
RAL µg Al/l	-	-	-	-	274	-	237	-
ILAL - " -	-	-	-	-	190	-	196	-
LAL - " -	-	-	-	-	84	0	41	5
K mg K/l	0.29	7	0.09	2	0.04	1	0.07	2
Na mg NA/l	1.88	82	0.51	22	1.09	47	0.65	28
So ₄ mg So ⁴ /l	5.9	123	4.5	94	1.9	40	1.5	31
NO ₃ µg N/l	44	3	2	-	2	-	5	-
Cl mg Cl/l	2.4	68	<0.1	2	1.7	48	0.8	23
UV abs. I/cm	.00	-	.00	-	.91	-	0.87	-
Σ cations		188		61		175		116
Σ anions		194		96		88		54

Table 9. Mean values from all tables (except for ionbalance, see foot notes).
 % denotes; passing through membrane.

	HELLERUDMYRA					EGIL					KIM					EGIL, Samples corresponding with KIM				
	MF		DIAL		%	MF		DIAL		%	MF		DIAL		%	MF		DIAL		%
	conc	¹⁾ $\mu\text{eq/l}$	conc	¹⁾ $\mu\text{eq/l}$		conc	³⁾ $\mu\text{eq/l}$	conc	³⁾ $\mu\text{eq/l}$		conc	⁵⁾ $\mu\text{eq/l}$	conc	⁵⁾ $\mu\text{eq/l}$		conc	⁶⁾ $\mu\text{eq/l}$	conc	⁶⁾ $\mu\text{eq/l}$	
pH	4.38	37	4.84	14	67	4.03	93	4.83	14	85	4.08	80	4.82	15	82	3.98	90	4.92	12	87
mS/m Cond	2.92	-	1.33	-	54	6.50	-	1.46	-	78	6.03	-	1.39	-	77	7.35	-	1.41	-	81
$\mu\text{g/l Fe}$	549	-	528	-	4	227	-	167	-	26	247	-	190	-	23	148	-	101	-	32
$\mu\text{g/l Mn}$	40	-	23	-	-	<20	-	<20	-	-	<20	-	<20	-	-	<20	-	<20	-	-
mgPt/Color	175	-	170	-	3	94	-	81	-	14	102	-	98	-	4	66	-	56	-	15
mg/l Ca	1.09 ¹	54	0.63 ¹	31	42	0.46 ³	23	0.26 ³	13	43	0.39 ⁵	20	0.37 ⁵	18	51	0.47 ⁶	24	0.39 ⁶	19	21
mg/l Mg	0.26	18	0.14	9	46	0.39 ³	32	0.18 ³	15	53	0.31 ⁵	25	0.19 ⁵	16	39	0.40 ⁶	33	0.19 ⁶	16	52
mg/l K	0.08 ¹	2	0.02 ¹	0	75	0.47 ³	12	0.08 ³	2	83	0.32 ⁵	8	0.10 ⁵	3	69	0.50 ⁶	13	0.09 ⁶	2	82
mg/l Na	1.03	39	0.33	9	70	2.46	124	0.54	13	78	2.30	100	0.50	17	78	2.63	126	0.53	14	89
$\mu\text{g/l RAL}$	266 ¹	-	249 ¹	-	6	439 ³	-	198 ³	-	55	385 ⁵	-	249 ⁵	-	35	421 ⁶	-	186 ⁶	-	56
$\mu\text{g/l ILAL}$	211 ¹	-	176 ¹	-	17	130 ³	-	119 ³	-	9	140 ⁵	-	152 ⁵	-	0	99 ⁶	-	104 ⁶	-	0
$\mu\text{g/l LAL}$	55 ¹	6	73 ¹	8	0	309 ³	34	79 ³	9	74	245 ⁵	27	97 ⁵	11	60	322 ⁶	36	82 ⁶	9	75
mg/l Cl	1.48	33	0.38	4	74	4.03	140	0.3	3	93	5.07	131	0.1	<3	98	5.2	156	0.1	<3	98
mg/l SO ₄	3.01	63	1.63	33	48	5.16	108	2.32	48	55	3.07	74	2.25	52	27	5.2	117	2.5	52	52
$\mu\text{g N/l NO}_3$	9	0	3	0	67	482	51	6	0	99	284	28	4	0	99	580	56	4	0	99
mg C/l DOC	15.9 ¹	-	11.6 ¹	-	27	6.6 ³	-	5.0 ³	-	24	9.4 ⁵	-	7.8 ⁵	-	17	6.5 ⁶	-	5.2 ⁶	-	20
I/cm UV-abs	.720	-	.550	-	24	.450	-	.342	-	24	.508	-	.449	-	12	.333	-	.264	-	21
$\mu\text{mol/l Strong acid}$	37 ²		19 ²		(49)	77 ⁴		5 ⁴		94	74		11		100					
$\mu\text{mol/l Weak}$	282 ²		222 ²		(21)	248 ⁴		179 ⁴		28	247		221		10					
Σ cations		156	-	72	-		318		63			260		78			322		69	
Σ anions		96	-	25	-		299		51			233		54			329		52	
Deficit		60		47			19		12			27		24			-7		17	

1) Column No 1, 2, 3, 4 & 9
 2) column 6 & 8

3) column 5, 26, 27 & 28
 4) " 15 - 25

5) column 31, 32, 33 & 34
 6) " 26(x2), 27 & 28

Acid Rain Research Reports

- 1/1982** Henriksen, A. 1982. Changes in base cation concentrations due to freshwater acidification. 50pp. Out of print.
- 2/1982** Henriksen, A. and Andersen, S. 1982. Forsuringssituasjonen i Oslomarkas vann. 45 pp. Out of print.
- 3/1982** Henriksen, A. 1982. Preacidification pH-values in Norwegian rivers and lakes. 24pp. Out of print.
- 4/1983** Wright, R.F. 1983. Predicting acidification of North American lakes. 165 pp.
- 5/1983** *Schoen, R., Wright, R.F. and Krieter, M.* 1983. Regional survey of freshwater acidification in West Germany (FRG). 15 pp.
- 6/1984** Wright, R.F. 1984. Changes in the chemistry of Lake Hovvatn, Norway, following liming and reacidification. 68 pp.
- 7/1985** Wright, R.F. 1985. RAIN project. Annual report for 1984. 39 pp.
- 8/1985** *Lotse, E and Otabbong, E.* 1985. Physiochemical properties of soils at Risdalsheia and Sogndal: RAIN project. 48 pp.
- 9/1986** Wright, R.F. and Gjessing, E. 1986. RAIN project. Annual report for 1985. 33 pp.
- 10/1986** Wright, R.F., Gjessing, E., Semb, A. and Sletaune, B. 1986. RAIN project. Data report 1983–85. 62 pp.
- 11/1986** Henriksen, A., Røgeberg, E., Andersen, S., and Veidel, A. 1986. MOBILLAB–NIVA, a complete station for monitoring water quality. 44 pp.
- 12/1987** Røgeberg, E. 1987. A coulometric Gran titration method for the determination of strong and weak acids in freshwater. 28 pp.
- 13/1987** Wright, R.F. 1987. RAIN project. Annual report for 1986. 90 pp.
- 14/1988** Hauhs, M. 1988. Water and ion movement through a mini-catchment at Risdalsheia, Norway (RAIN–project). 74 pp.