

Fagrådet

for vann- og avløpsteknisk
samarbeid i indre Oslofjord



Statlig program for
forurensningsovervåkning

Rapport nr. 621/95


Overvåkning av forurensnings- situasjonen i indre Oslofjord

Effekter av forurensning og
dypvannsutsiftning på
faunaen langs bunnen av
Oslofjorden basert på materiale
samlet siden 1952

(Effects of pollution and
deepwater exchange on the
fauna along the bottom of
Oslofjorden, Norway, based on
material collected since 1952)

Del II

Referanser, tabeller, figurer og
english summary



NIVA - RAPPORT

Norsk institutt for vannforskning



NIVA

Prosjektnr.: O-71094	Undernr.:
Løpenr.: 3 3 2 4 b	Begr. distrib.: Fri

Hovedkontor Postboks 173, Kjelsås 0411 Oslo Telefon (47) 22 18 51 00 Telefax (47) 22 18 52 00	Sørlandsavdelingen Televeien 1 4890 Grimstad Telefon (47) 37 04 30 33 Telefax (47) 37 04 45 13	Østlandsavdelingen Rute 866 2312 Ottestad Telefon (47) 62 57 64 00 Telefax (47) 62 57 66 53	Vestlandsavdelingen Thormøhlensgt 55 5008 Bergen Telefon (47) 55 32 56 40 Telefax (47) 55 32 88 33	Akvaplan-NIVA A/S Søndre Tollbugate 3 9000 Tromsø Telefon (47) 77 68 52 80 Telefax (47) 77 68 05 09
--	---	--	---	--

Rapportens tittel: Overvåking av forurensnings situasjonen i indre Oslofjord. Effekter av forurensning og dypvannsutsiftning på faunaen langs bunnen av Oslofjorden basert på materiale samlet siden 1952. (Effects of pollution and deepwater exchange on the fauna along the bottom of Oslofjorden, Norway, based on material collected since 1952). Overvåkingsrapport nr. 621 /95). TA nr.1257 /1995	Dato: 1.10.1995 Trykket: NIVA 1995
Forfatter(e): Fredrik Beyer, UiO Jane Indrehus, UiO	Faggruppe: Mø Geografisk område: Oslo, Akershus, Buskerud Antall sider: 143+153 Opplag:

Oppdragsgiver: Fagrådet for vann-og avløpsteknisk samarbeid i indre Oslofjord	Oppdragsg. ref.: A. Rosendahl
--	----------------------------------


Ekstrakt: Til innsamling av dyr like over bunnen (hyperbenthos) er det i ca. 40 år benyttet en lukkbar, finmasket (0.5mm) hov montert i en slede. Litt av det øverste, løse bunnsedimentet og mange slags dyr som er knyttet til dette, havner også i håven. Redskapet har en distansemåler. Prøver tatt på forskjellige lokaliteter viser markante forskjeller, og det samme kan være tilfelle for prøver tatt til forskjellig tid på samme sted. Materialet viser en kraftig forringelse av faunaen innover fjorden, som hovedsakelig kan tilskrives oksygenmangel. Denne har hatt akkumulert effekt på sediment og fauna. Utslipp av kloakk fra VEAS hadde lokalt uheldig innflytelse på faunaen i 1982-85. Med tilsynelatende utgangspunkt i Gråøyrenna i 1984 har 3 rekearter senere vist bemerkelsesverdig øket antall. Overføringen av kloakk fra Bunnefjordområdet har ikke gitt tilstrekkelig avlastning av dette. Yngel av kommersiell reke samt en mindre rekeart ble funnet i stort antall i Gråøyrennen fra 1984 og ble noe senere også funnet lengre innover. Muddereke ble etter mange år først funnet rikelig i 1992 i Gråøyrenna, men synes ikke å akseptere forholdene lengre inne. Bunnsledeprøver viser meget større utslag av miljøforskjeller og - forandringer enn tilsvarende prøver tatt med grabb eller skrape.
--


4 emneord, norske

1. Forurensningsovervåking
2. Oslofjorden
3. Hyperbenthos-benthos
4. Dypvannsutsiftning

4 emneord, engelske

1. Pollution monitoring
2. Oslofjorden
3. Hyperbenthos-benthos
4. Deepwater exchange

Prosjektleder

Jan Magnusson

For administrasjonen

Torgeir Bakke

SBN 82-577-2857-8

EFFEKTER AV FORURENSNING OG DYPVANNSSUTSKIFTNING PÅ
FAUNAEN LANGS BUNNEN AV OSLOFJORDEN
BASERT PÅ MATERIALE SAMLET SIDEN 1952

(EFFECTS OF POLLUTION AND DEEPWATER EXCHANGE ON THE FAUNA ALONG THE
BOTTOM OF OSLOFJORDEN, NORWAY, BASED ON MATERIAL COLLECTED SINCE 1952)

av

FREDRIK BEYER *

og

JANE INDREHUS *

DEL II

REFERANSER, TABELLER, FIGURER, ENGLISH SUMMARY

* Adresse: Universitetet i Oslo, Biologisk institutt, Avdeling for marin zoologi og marin kjemi, P.B. 1064, Blindern, 0316 Oslo 3.

INNHOOLD

	Side
REFERANSER	3
FANGSTSLISTER. Individider/100 m ³ "filtrert" av identifiserte arter og andre taksonomiske enheter.	
Elle. E53 og E59 betyr Elle 1953 og Elle 1959, etc.	18
Gråøyrenna. G53 betyr Gråøyrenna 1953, etc.	21
Svartedypet. SD81 betyr Svartedypet 1981, etc.	24
Vesthullet. VH81 betyr Vesthullet 1981, etc.	27
Steilene. ST73 betyr Steilene 1973, etc.	30
Svartskog. SS85 betyr Svartskog 1985, etc.	33
 AUTORLISTE	 36
 TEKNISKE DATA	 40
Elle	40
Gråøyrenna	40
Langårrenna	41
Sprodypet	41
Ytre Stødyp	42
Blåmagen	42
Svartedypet	42
Vesthullet	43
Steilene	43
Gåsøyrenna	45
Lysakerfjorden	45
Helviktangen	46
Kirkevikbanken	46
Svartskog	47
 FIGURLISTE	 48
 FIGURER	 52
 ENGLISH SUMMARY	 139
Introduction	139
Topography, hydrography and biochemical processes	139
Character of the bottom	140
Methods	140
Material	141
The fauna variations with time and location	142
Indicator species	144
Causes of the observed fauna variations	146
Comparisons with other investigations	147
Recommendations	148
 LIST OF TEXT TABLES	 149
 LIST OF FIGURES	 150

REFERANSER

I tillegg til de referanser som er angitt i teksten omfatter denne listen også arbeider som er benyttet ved identifiseringen av dyrene.

- Agassiz, A. 1888. Three cruises of the United States Coast and Geodetic Survey steamer "Blake" in the Gulf of Mexico, in the Caribbean Sea, and along the Atlantic coast of the United States, from 1877 to 1880, vol. I. *Bulletin of the Museum of Comparative Zoölogy at Harvard College, in Cambridge*. Vol. XIV: i-xxii, 1-314.
- Allen, J. A. 1960. On the biology of *Crangon allmani* Kinahan in Northumberland waters. *Journal of the Marine Biological Association of the United Kingdom* 39: 481-508.
- Andrén, L., P. E. Lindgren, K. G. Nyholm & I. Olsson 1968. II.7 Marinbiologi. Pp. 127-155 in: *Göteborgs Vattenvårdsanläggningar. Undersökningar för havsutsläpp år 1966-1967. Huvudrapport*. Göteborgs Stads Vatten- och Avloppsverk, Göteborg. 210 pp.
- Arntz, W. E. 1980. Predation by demersal fish and its impact on the dynamics of macrobenthos. Pp. 121-149 in: K. R. Tenore & B. C. Coull (eds) *Marine benthic dynamics*. University of South Carolina Press. 451 pp.
- Aschan, M. M. & A. M. Skullerud 1990. Effects of changes in sewage pollution on soft-bottom macrofauna communities in the Inner Oslofjord, Norway. *Sarsia* 75: 169-190.
- Baalsrud, K. (red.) 1968. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Samlerapport*. 63 pp. Norsk Institutt for Vannforskning, Oslo.
- Baalsrud, K. 1987. *Oslofjorden vårt nærmiljø*. Reclamo/Ryen. Oslo. 85 pp.
- Bakke, J. E. 1980. Drøbacterskelens betydning for zooplankton i Indre Oslofjord, samt registrering av enkelte lite undersøkte arter i Oslofjorden. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 197 pp.
- Barnard, J. L. 1962. South Atlantic abyssal amphipods collected by R. V. Vema. Pp. 1-78 in: *Abyssal Crustacea* ed. by J. L. Barnard, R. J. Menzies & M. C. Bacescu. Columbia University Press. New York and London. 223 pp.
- Berner, R. 1972. Sulfate reduction, pyrite formation, and the oceanic sulfur budget. Pp. 347-361 in: D. Dyrssen & D. Jagner (eds) *The changing chemistry of the oceans. Proceedings of the Twentieth Nobel Symposium held 16-20th August, 1971 at Aspenäsgråden, Lerum and Chalmers University of Technology, Göteborg, Sweden*. Almqvist & Wiksell, Stockholm. 365 pp.
- Beyer, F. 1956. Kappløpet mellom forråtnelse og fornyelse av indre Oslofjord. *Teknisk Ukeblad* 1956: 1045-1053.
- Beyer, F. 1958. A new, bottom-living trachymedusa from the Oslo Fjord. Description of the species, and a general discussion of the life conditions and fauna of the fjord deeps. *Nytt magasin for Zoologi* 6: 121-143.
- Beyer, F. 1967 a. Bunn sedimenter og bunnfauna i indre og midtre Oslofjord i 1938 og 1962 - 1966. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 6: 1-119 + Fig. 1-33*. Norsk Institutt for Vannforskning, Oslo.
- Beyer, F. 1967 b. Representativiteten av undersøkelsesperioden 1962-1966. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 18: 1-11 + Fig. 1-4*. Norsk Institutt for Vannforskning, Oslo.
- Beyer, F. 1968. Zooplankton, zoobenthos, and bottom sediments in the Oslofjord. *Helgoländer wissenschaftliche Meeresuntersuchungen* 17: 496-509.

- Beyer, F. 1971. Om vannutvekslingen i Oslofjorden og dens betydning for faunanen. (With English summary.) *Universitetet i Oslo. Institutt for marin biologi avd. A & C. Rapport nr. 3*:1-25.
- Beyer, F. 1992. *Meganyctiphanes norvegica* (M. SARS) (Euphausiacea) a voracious predator on *Calanus*, other copepods, and ctenophores, in Oslofjorden, southern Norway. *Sarsia* 77: 189-206.
- Beyer, F., Å. Dybwad og J. Versvik 1967. Zooplankton. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 5*: 1-66 + Fig. 1-24. Norsk Institutt for Vannforskning, Oslo.
- Beyer, F. og E. Føyn 1951. Surstoffmangel i Oslofjorden. En kritisk situasjon for fjordens dyrebestand. *Naturen nr. 10*: 289-306.
- Beyer, F. og J. Versvik 1968. Undersøkelse av virvelløse dyr langs med bunnen av indre Oslofjord i 1962-1966. MS.
- Boeck, A. 1871. Crustacea amphipoda borealia et arctica. *Forhandlinger, Videnskabselskabet i Christiania 1870*: 83-280.
- Boesch, D. F. & R. Rosenberg 1981. Response to stress in marine benthic communities. Pp. 179-200 in: G. W. Barrett & R. Rosenberg (eds). *Stress effects on natural ecosystems*. John Wiley & Sons, Chichester. 305 pp.
- Borchert, H. 1965. Formation of marine sedimentary ores. Pp. 159-204 in: J. P. Riley & G. Skirrow (eds). *Chemical oceanography. vol. 2*. Academic Press, London. 508 pp.
- Braarud, T. 1945 a. A phytoplankton survey of the polluted waters of Inner Oslo Fjord. *Hvalrådets Skrifter* 28: 1-142.
- Braarud, T. 1945 b. Forurensning og selvrensning av sjøvann. Undersøkelser i Oslofjorden. *Naturen* 1945: 212-235.
- Braarud, T. & I. Nygaard 1967. Fytoplankton. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 4*: 1-171. Norsk Institutt for Vannforskning, Oslo.
- Braarud, T. & J. T. Ruud 1937. The hydrographic conditions and aeration of the Oslo Fjord 1933-1934. *Hvalrådets Skrifter* 15: 1-56.
- Braaten, B. R. 1968. En biologisk helårsundersøkelse (1966-1967) av et forurenset basseng i indre Oslofjord. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 94 pp.
- Brand, T. von 1946. *Anaerobiosis in invertebrates*. No. 4 of the "biodynamica Monographs" ed. by B. J. Luyet. Biodynamica, Missouri. 328 pp.
- Brattegard, T. 1978. *Bestemmelsestabell for Mysidacea kjent fra norske farvann*. Institutt for Marinbiologi, Universitetet i Bergen. 21 pp.
- Brattegard, T. og T. Høisæter 1973. *Rapport. Supplerende undersøkelser av Fensfjordens dype bløtbunners dyreliv*. Biologisk Stasjon, Espegrend, Blomsterdalen. 71 pp.
- Breck, W. G. 1974. Redox levels in the sea. Pp. 153-179 in: E. D. Goldberg (ed.) *The sea. Vol. 5. Marine chemistry*. John Wiley & Sons, N. Y. 895 pp.
- Broch, H. 1936. Die Bedeutung der Drøbak-Schwelle für die Bodenfauna der Garnelen-Felder. *Aohandlinger utgitt av Det Norske Videnskaps-Akademi i Oslo. I. Matem.-Naturvid. Klasse 1935. No5*: 1-32.
- Calman, W. T. 1899. On the British Pandalidae. *Annual Magazine of Natural History III, Series 7*: 27-39.
- Christensen, J. M., S. Larsen, & B. O. Nyström 1978. *Muslinger*. Norsk utgave ved M. E. Christiansen. J. W. Cappelens Forlag A.S. 125 pp.

- Christiansen, M. E. 1972. *Bestemmelsestabell over Crustacea Decapoda. Tifotkreps*. Universitetsforlaget. Oslo - Bergen - Tromsø. 71 pp.
- Christie, H. 1980. Methods for ecological monitoring: Biological interactions in a rocky subtidal community. *Helgoländer Meeresuntersuchungen*, 33: 473-483.
- Colt, J. E. & D. A. Armstrong 1981. Nitrogen toxicity to crustaceans, fish and molluscs. Pp. 34-47 in: L. J. Allen & E. C. Kinney (eds) *Proceedings of the Bio-Engineering Symposium for Fish Culture. Fish Culture Section*. American Fisheries Society, Northeast Society of Conservation Engineers. Bethesda, Maryland.
- Cook, R. H. & C. M. Boyd 1965. The avoidance by *Gammarus oceanicus* SEGERSTRÅLE (Amphipoda, Crustacea) of anoxic regions. *Canadian Journal of Zoology* 43: 971-975.
- Costello, M. J. & P. Read 1994. Toxicity of sewage sludge to marine organisms: A review. *Marine Environmental Research* 37: 23-46.
- Dahl, F. E. 1973 a. Toktoversikt. Hydrografisk tokt 04. Oslofjorden 23. - 26.7.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-3 + Fig. 1-3 + 11 tabeller.*
- Dahl, F. E. 1973 b. Toktoversikt. Hydrografisk tokt 05. Oslofjorden 6. - 8.8.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-4 + Fig. 1-4 + 11 tabeller.*
- Dahl, F. E. 1973 c. Toktoversikt. Hydrografisk tokt 06. Oslofjorden 20. - 21.8.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-3 + Fig. 1-4 + 16 tabeller.*
- Dahl, F. E. 1973 d. Toktoversikt. Hydrografisk tokt 07. Oslofjorden 3. - 4.9.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-3 + Fig. 1-3 + 18 tabeller.*
- Dahl, F. E. 1973 e. Toktoversikt. Hydrografisk tokt 08. Oslofjorden 17. - 18.9.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-4 + Fig. 1-5 + 12 tabeller.*
- Dahl, F. E. 1973 f. Toktoversikt. Hydrografisk tokt 09. Oslofjorden 1. - 2.10.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-14 + 18 tabeller.*
- Dahl, F. E. 1973 g. Toktoversikt. Hydrografisk tokt 10. Oslofjorden 15. - 17.10.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-8 + 12 tabeller.*
- Dahl, F. E. 1973 h. Toktoversikt. Hydrografisk tokt 11. Oslofjorden 5. - 6.11.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-10 + 12 tabeller.*
- Dahl, F. E. 1973 i. Toktoversikt. Hydrografisk tokt 12. Oslofjorden 14. - 15.11.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-12 + 24 tabeller.*
- Dahl, F. E. 1974 a. Toktoversikt. Hydrografisk tokt 13. Oslofjorden 26. - 27.11.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-17 + 34 tabeller.*
- Dahl, F. E. 1974 b. Toktoversikt. Hydrografisk tokt 14. Oslofjorden 10. - 12.12.1973. *Vassdrags- og Havnelaboratoriet. Notat: 1-8 + 17 tabeller.*
- Day, J. H. 1978. The effect of plants and animals on the chemistry of estuarine sediments. Pp. 216-221 in: *Biogeochemistry of estuarine sediments. Proceedings of a Unesco/SCOR workshop held in Melreux, Belgium 29 November to 3 December 1976*. UNESCO, Paris. 293 pp.
- Dejours, P. 1975. *Principles of comparative respiratory physiology*. North-Holland Publishing Company, Amsterdam. 253 pp.
- Demel, K. & Z. Mulicki 1959. Studia ilosciowe nad fauna przydenna południowego Bałtyku. (Quantitative studies on the near-bottom Southern Baltic fauna.) *Prace Morskiego Instytutu Rybackiego w Gdyni. Czesc oOceanograficzno - ichtiologiczna. (Reports of the Sea Fisheries Institute in Gdynia. Oceanographical and ichthyological part.) Tom 10/A: 19-33.*

- Doetsch, R. N. & T. M. Cook 1973. *Introduction to bacteria and their ecobiology*. Medical and Technical Publishing Co. Ltd. Lancaster. 371 pp.
- Dons, C. 1915. Nord-Norges decapoder. *Tromsø Museums Aarshefter* 37 1914: 15-153 + Pl. I-II.
- Einarsson, H. 1945. Euphausiacea I. Northern Atlantic species. *Dana-Report* 27: 1-191.
- Ekman, S. 1933. Über einen neuen Bodengreifer für marin-zoologische Zwecke. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 28: 312-329.
- Ekman, S. 1947. Über die Festigkeit der marinen Sedimente als Faktor der Tierverbreitung, ein Beitrag zur Associationsanalyse. *Zoologiska Bidrag från Uppsala*. XXV: 1-20.
- Ekman, S. 1953. *Zoogeography of the sea*. Sidgwick and Jackson Limited, London. 417 pp.
- Elofson, O. 1941. Zur Kenntnis der marinen Ostracoden Schwedens mit besonderer Berücksichtigung des Skageraks. *Zoologiska bidrag från Uppsala* XIX: 215-534.
- Enequist, P. 1950. Studies on the soft-bottom amphipods of the Skagerak. *Zoologiska bidrag från Uppsala* XXVIII: 297-492 + Chart 6.
- Enoksson, V., F. Sörensson & W. Granéli 1990. Nitrogen transformations in the Kattegat. *Ambio*. Vol. XIX(3): 159-166.
- Fage, L. 1951. Cumacés. *Faune de France* 54: 1-136. Paris.
- Farran, G. P. 1905. Report on the Copepoda of the Atlantic slope off counties Mayo and Galway. *Department of Agriculture and Technical Instruction for Ireland. Report on the Sea and Inland Fisheries of Ireland for 1902 and 1903. Part II. Scientific Investigations*: 23-52 + Pl. 3-13.
- Fauchald, K. 1974. Spaeodoridae (Polychaeta: Errantia) from world-wide areas. *Journal of Natural History*, vol. 8: 257-289.
- Fauvel, P. 1923. Polychètes errantes. *Faune de France* 5: 1-488. Paris.
- Fauvel, P. 1927. Polychètes sédentaires. *Faune de France* 16: 1-494. Paris.
- Foreningen til fremme av fiskeriet i Oslofjorden innenfor Drøbak. 1946. *Beretning* 1945: 1-7.
- Foreningen til fremme av fiskeriet i Oslofjorden innenfor Drøbak. 1947. *Beretning* 1946: 1-7.
- Foreningen til fremme av fiskeriet i Oslofjorden innenfor Drøbak. 1952. *Beretning* 1951: 1-9.
- Foreningen til fremme av fiskeriet i Oslofjorden innenfor Drøbak. 1953. *Beretning* 1952: 1-7.
- Forsman, B. 1940. Untersuchungen über die Cumaceen des Skageraks. *Zoologiska bidrag från Uppsala* XVIII: 1-161.
- Fosshagen, A. 1979. Dyreplankton i Ryfylkefjordene 1973 - 1975. *Rådgivende utvalg for fjordundersøkelser. Ryfylkeprosjektet. Rapp. Nr. 2*: 1-100 + Tab. 1-52.
- Føyn, E. 1958. Sprangskikt, oksygenminima og sperreflater for fisk. *Fauna*. Årg. 11 (4): 121-131.
- Fritzvold, J. 1981. Vertikale vandringer hos hyperbenthos sett i relasjon til dyp og årstid. *Hovedfagsoppgave. Universitetet i Oslo, Biologisk institutt, avd. marin zoologi og marin kjemi*. 137 pp.
- Frobisher, M., R. D. Hinsdill, K. T. Crabtree & C. R. Goodheart 1974. *Fundamentals of microbiology*. W. B. Saunders Comp., Philadelphia. 850 pp.
- Gade, H. G. 1963. Some hydrographic observations of the Inner Oslofjord during 1959. *Hvalrådets Skrifter* 46: 1-62.

- Gade, H. G. 1967. Hydrografi. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 2: 1-104 + Fig. 1-68.*
- Gade, H. G. 1968. Horizontal and vertical exchanges and diffusion in the water masses of the Oslo Fjord. *Helgoländer wissenschaftliche Meeresuntersuchungen. 17: 462-475.*
- Gade, H. G. 1970. Hydrographic investigations in the Oslofjord, a study of water circulation and exchange processes. Vol I, II & III. *Geophysical Institute. Div. A. Physical Oceanography. Report 24: 1-193 & 1-92.*
- Gade, H. G. & A. Edwards 1980. Deep water renewal in fjords. Pp. 453-489 in: *Fjord oceanography*, ed. by H. J. Freeland, D. M. Farmer & C. D. Levings. Plenum Publishing Corp., New York. 715 pp.
- Giesbrecht, W. 1892. Systematik und Faunistik der pelagischen Copepoden des Golfes von Neapel und der angrenzenden Meeres-Abschnitte. *Fauna und Flora des Golfes von Neapel XIX. Monographie: i-ix, 1-831 + Tafel 1-54 .Berlin.*
- Gislén, T. 1929. Epibioses of the Gullmar Fjord I. A study in marine sociology. *Kristinebergs Zoologiska Station 1877 - 1927, 3: 1-123.* K. Svenska Akademien. Uppsala.
- Gislén, T. 1930. Epibioses of the Gullmar Fjord II. *Kristinebergs Zoologiska Station 1877 - 1927, 4: 1-380.* K. Svenska Akademien. Uppsala.
- Gjermundsen, B. 1974. Faunaen langsmed bunnen i de dypeste områder av Oslofjorden - En sammenlikning av prøver fra 1960, 1962-63 og 1971-72 fanga med Beyers bunnsleder. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi. 154 pp.*
- Gotto, R. V. 1964. The Anomoclausidae: a new family of cyclopoid copepods, with a description of *Anomoclausia indrehusae* gen. nov., sp. nov. *Crustaceana 6 (3): 220-228.*
- Gran, H. H. & T. Gaarder 1918. Über den Einfluss der atmosphärischen Veränderungen Nordeuropas auf die hydrographischen Verhältnisse des Kristianiafjords bei Dröbak im März 1916. *Conseil permanent international pour l'Exploration de la Mer. Publications de Circonstance 71: 1-29.*
- Grassle, J. F., H. L. Sanders, R. R. Hessler, G. T. Rowe & T. McLellan 1975. Pattern and zonation: a study of the bathyal megafauna using research submersible Alvin. *Deep Sea Research 22: 457-481.*
- Grassle, J. P. & J. F. Grassle 1976. Sibling species in the marine pollution indicator *Capitella* (Polychaeta). *Science, N. Y. 192: 567-569.*
- Gray, J. S. 1982. Effects of pollutants on marine ecosystems. *Netherlands Journal of Sea Research 16: 424-443.*
- Gunkel, W. 1970. Light - Bacteria, fungi and blue-green algae. Pp. 103-124 in: *Marine ecology. A comprehensive, integrated treatise on life in oceans and coastal waters.* Ed. by O. Kinne. Vol. I. *Environmental factors. Part 1.* Wiley-Interscience, London.
- Gunkel, W. 1972. Organic substances - Bacteria, fungi and blue-green algae. Pp. 1533-1549 in: *Marine ecology. A comprehensive, integrated treatise on life in oceans and coastal waters.* Ed. by O. Kinne. Vol. I. *Environmental factors. Part 3.* Wiley-Interscience, London.
- Hansen, H. J. 1916. Crustacea Malacostraca III. The order Isopoda. *The Danish Ingolf-Expedition III (5): 1-264 + Pl. I-XVI.*
- Hartmann-Schröder, G. 1971. Annelida, Borstenwürmer, Polychaeta. In: M. Dahl & F. Peus (eds) *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise. 58. Teil. 594 pp.*
- Hernroth, L. & F. Gröndahl 1985. On the biology of *Aurelia aurita* (L.) 3. Predation by *Coryphella verrucosa* (Gastropoda, Opisthobranchia), a major factor regulating the development of *Aurelia* populations in the Gullmar Fjord, western Sweden. *Ophelia, 24 (1): 37-45.*

- Hesthagen, I. H. 1970. On the near-bottom plankton and benthic invertebrate fauna of the Josephine Seamount and the Great Meteor Seamount. "*Meteor*" *Forschungs-Ergebnisse. Reihe D*, 8: 61-70.
- Hesthagen, I. H. 1971. On the biology of the bottom-dwelling trachymedusa *Tesserogastria musculosa* Beyer. *Norwegian Journal of Zoology* 19 (1): 1-19.
- Hesthagen, I. H. & B. Gjermundsen 1978. The replicability of sampling the hyperbenthic region by means of Beyer's 50 cm epibenthic closing net. *Meeresforschung* 26 (1977/78): 1-10.
- Hjort, J. 1900. The deep sea fauna in the fiords that have been investigated. Ch. IV, pp. 75-95 in: Fishing experiments in Norwegian fiords by J. Hjort & K. Dahl. *Report on Norwegian Fishery and Marine Investigations I* (1): 1-215.
- Hjort, J. & K. Dahl 1900. Fishing experiments in Norwegian fiords. *Report on Norwegian Fishery and Marine Investigations I* (1): 1-215.
- Hjort, J. & H. H. Gran 1900. Hydrographic-biological investigations of the Skagerrak and the Christiania Fjord. *Report on Norwegian Fishery and Marine Investigations I* (2): 1-41 + Tab. 1-7.
- Hjort, J. & J. T. Ruud 1938. Rekefisket som naturhistorie og samfundssak. *Fiskeridirektoratets Skrifter, Serie Havundersøkelser V* (4): 1-158.
- Hoffmann, H. 1926. I. Opisthobranchia. *Die Tierwelt der Nord- und Ostsee IX.c*: 1-52.
- Höglund, H. 1947. Foraminifera in the Gullmar Fjord and the Slagerak. *Zoologiska Bidrag från Uppsala*, XXVI: 1-328 + 32 plates.
- Holmlund, J. 1963. En undersøkelse over forplantnings- og vekstforholdene hos *Pontophilus norvegicus* i Oslofjorden. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 54 pp.
- Holthuis, L. B. 1946. Note on the genus *Pandalina* (Crustacea Decapoda), with the description of a new species from European waters. *Rijksmuseum van natuurlijke historie te Leiden. Zoologische mededeelingen XXVI*: 281-286.
- Holthuis, L. B. 1947. The Decapoda of the Siboga Expedition. Part IX. The Hippolytidae and Rhynchocinetidae collected by the Siboga and Snellius expeditions, with remarks on other species. *The Siboga Expedition. Monographs* 39: 1-100.
- Holthuis, L. B. 1955. The recent genera of the caridean and stenopodidean shrimps (class Crustacea, order Decapoda, supersection natantia) with keys for their determination. *Rijksmuseum van natuurlijke historie te Leiden. Zoologische verhandelingen* 26: 1-157.
- Horne, R. A. 1969. *Marine chemistry. The structure of water and the chemistry of the hydrosphere*. Wiley-Interscience, New York. 568 pp.
- Hovde, H. R. 1972. Forandringer i planktonfaunaen i indre Oslofjord de senere decennier. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 307 pp.
- Hult, J. 1936. On some species and genera of Parasellidae. *Arkiv för Zoologi. Band 29 A* (6): 1-14.
- Hult, J. 1936. On some species and genera of Parasellidae. II. *Arkiv för Zoologi. Band 29 B* (5): 1-6.
- Hult, J. 1937. Marina isopoder från svenska västkusten. *Göteborgs Kungl. Vetenskaps- och Vitterhets-Samhälles Handlingar. Femte följden. Ser. B. Band 5* (3): 1-48.
- Hult, J. 1942. On the soft-bottom isopods of the Skager Rak. *Zoologiska Bidrag från Uppsala XXI*: 1-234.
- Indrehus, J. 1967. A quantitative and qualitative study of the bottom natant fauna found at two stations in the Oslofjord, one inside and the other outside the Drøbak sill at the fjord entrance. *MS*, 256 pp.

- Jaeckel, S. 1952. Zur Verbreitung und Lebensweise der Opisthobranchier in der Nordsee. *Kieler Meeresforschungen*, 8: 249-259.
- Jakobsen, T. 1971. On the biology of *Sagitta elegans* Verrill and *Sagitta setosa* J. Müller in Inner Oslofjord. *Norwegian Journal of Zoology* 19 (2): 201-225.
- Jensen, A. S. og R. Spärck 1934. Bløddyr II. Saltvandmuslinger. *Danmarks Fauna* 40: 1-208. København.
- Johannessen, O. M. 1965. Strømundersøkelser i Drøbaksundet mai - juni 1963. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport i serie hydrografi (nr. 19)*: 1-53 + Fig. 1-23 + Tab. I-VIII.
- Johannessen, T. & Skau, E. 1970. Fisken kveles i Oslofjorden. *Verdens Gang. Lørdag 21. november 1970*: 9.
- Kaartvedt, S. 1982. Hyperfauna i Fanafjordens dypbasseng: Døgnvandringsmønstre, døgn- og sesongvariasjoner i hyperfaunaens sammensetning. *Hovedfagsoppgave i marinbiologi. Universitetet i Bergen*. 122 pp.
- Kaplan, I. R. & S. C. Rittenberg 1963. Basin sedimentation and diagenesis. Pp 583-619 in: M. N. Hill (ed.) *The sea. Ideas and observations on progress in the study of the seas. Vol. 3. The earth beneath the sea. History*. Interscience Publishers. New York.
- Kemp, S. 1910. The Decapoda Natantia of the coasts of Ireland. *Department of Agriculture and Technical Instruction for Ireland. Fisheries Branch. Scientific Investigations, 1908. No. I*: 1-190 + Pl. I-XXIII.
- Kilenyi, T. I. 1969. *The problems of ostracod ecology in the Thames estuary. The taxonomy, morphology and ecology of recent Ostracoda*. Oliver and Boyd. Edinburgh. 532 pp.
- Kinne, O. 1970. Temperature - Invertebrates. Pp. 407-514 in: *Marine ecology. A comprehensive, integrated treatise on life in oceans and coastal waters*. Ed. O. Kinne. Vol. I. Environmental factors. Part 1. Wiley-Interscience, London. 681 pp.
- Kirkerud, L. A., P. O. Martinsen, C. G. Christophersen & Ø. Bjerk 1975. Nitrite toxicity in a polluted marine environment. An experimental study. *Repr. from Annual Report 1974. Appendix II. Norwegian National IBP Committee, Oslo*. 65 pp.
- Klavestad, N. 1967. Undersøkelse av benthos-algevegetasjonen i indre Oslofjord i 1962 - 1965. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 9*. 163 pp.
- Konieczny, R. M. 1994. Miljøgiftundersøkelser i Indre Oslofjord. Delrapport 4. Miljøgifter i sedimenter. *NIVA-rapport, 0-92131, l. nr. 3094*. 134 pp.
- Künitzer, A. 1990. A comparison of the *Amphiura filiformis*-associations north-east of the Dogger Bank and of the German Bight. *Netherlands Journal of Sea Research* 25 (1/2): 199-208.
- Lang, K. 1948. *Monographie der Harpacticiden*. Lund. Reprinted 1975, Vol. 1-2. Otto Koeltz Science Publishers, Koenigstein. 1683 pp.
- Lauritsen, O. 1983. Foraminiferfaunaen i Slemmestadområdet, Indre Oslofjord. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 151 pp. + Pl. 1-6.
- Lie, U. 1967. The natural history of the Hardangerfjord. 8. Quantity and composition of the zooplankton, september 1955 - september 1956. *Sarsia* 30: 49-74.
- Lindley, J. A. 1986. Dormant eggs of calanoid copepods in sea-bed sediments of the English Channel and southern North Sea. *Journal of Plankton Research* 8 (2): 399-400.
- Linke, O. 1939. Die Biota des Jadebusenwattes. *Helgoländer wissenschaftliche Meeresuntersuchungen* 1 (3): 201-348.

- Løyning, P. 1922. Nudibranchfaunaen i Drøbaksundet. I. Fam. Aeolidae. *Skrifter utgitt av Videnskapsselskapet i Kristiania 1922. I. Matematisk-Naturvidensk. Klasse. No.6:* 1-103 + Pl. I-IV.
- Løyning, P. 1927. Nudibranchs from Bergen, collected in the neighbourhood of the Biological Station at Herdla. *Nyt Magazin for Naturvidenskaberne. Bind 65:* 243-264.
- McLeese, D. W. 1956. Effects of temperature, salinity and oxygen on the survival of the American lobster. *Journal of the Fisheries Research Board of Canada, 13:* 247-272.
- MacLeod, P. & I. Valiela 1975. The effect of density and mutual interference by a predator: A laboratory study of predation by the nudibranch *Coryphella rufibranchialis* on the hydroid *Tubularia larynx*. *Hydrobiologia vol. 47, 3-4:* 339-346.
- Magnusson, J., T. Bokn & G. Larsen 1991. Overvåking av forurensningssituasjonen i indre Oslofjord 1989 - 90. *NIVA-rapport O-7160 91, l. nr. 2581.* 52 pp.
- Magnusson, J., T. Bokn, F. Moy, A. Pedersen & G. Larsen 1992. Overvåking av forurensningssituasjonen i Indre Oslofjord 1991. *NIVA-rapport O-716091, l. nr.2722.* 116 pp.
- Magnusson, J. & T. Johnsen 1993. Overvåking av forurensningssituasjonen i indre Oslofjord 1992. *NIVA-rapport O-710 92, l. nr. 2890.* 39 pp.
- Mare, M. 1942. A study of a marine benthic community with special reference to the micro-organisms. *Journal of the Marine Biological Association of the United Kingdom XXV:* 517-554.
- Marshall, S. M., A. G. Nicholls & A. P. Orr 1935. On the biology of *Calanus finmarchicus*. VI. Oxygen consumption in relation to the phytoplankton. *Journal of the Marine Biological Association of the United Kingdom XX:* 1-28.
- Marshall, S. M. & A. P. Orr 1952. On the biology of *Calanus finmarchicus*. VII. Factors affecting egg production. *Journal of the Marine Biological Association of the United Kingdom XXX:* 527-548.
- Matthews, J. B. L. 1964. On the biology of some bottom-living copepods (Aetideidae and Phaennidae) from western Norway. *Sarsia 16:* 1-46.
- Mattson, S. 1981. The food of *Galeus melastomus*, *Gadiculus argenteus thori*, *Trisopterus esmarkii*, *Rhinonemus cimbrius*, and *Glyptocephalus cynoglossus* (Pisces) caught during the day with shrimp trawl in a West-Norwegian fjord. *Sarsia 66 (2):* 109-127.
- Menzies, R. J. 1962. The isopods of abyssal depths in the Atlantic Ocean. Pp. 79-206 in: *Abyssal Crustacea* ed. by J. L. Barnard, R. J. Menzies & M. C. Bacescu. Columbia University Press. New York and London. 223 pp.
- Mielck, W. 1935. Arbeitsweisen und Ergebnisse über die Eier und Larven von Fischen. 1. Teil, pp. 3-53 + Tafel 1 in: *Fischbrut- und Plankton-Untersuchungen auf dem Reichsforschungsdampfer "Poseidon" in der Ostsee, Mai - Juni 1931 von W. Mielck und C. Künne. Wissenschaftliche Meeresuntersuchungen. Neue Folge. Abteilung Helgoland 19 (7):* 1-120 + Taf. 1-8.
- Mill, P. J. 1972. *Respiration in the Invertebrates*. MacMillan St. Martin's Press, London. 212 pp.
- Millar, R. H. 1966. Tunicata, Ascidiacea. *Marine invertebrates of Scandinavia. 1.* Universitetsforlaget, Oslo. 123 pp.
- Miller, M. C. 1961. Distribution and food of the nudibranchiate Mollusca of the south of the Isle of Man. *Journal of animal Ecology 30:* 95-116.
- Mirza, F. B. & J. S. Gray 1981. The fauna of benthic sediments from the organically enriched Oslofjord, Norway. *Journal of experimental marine Biology and Ecology 54:* 181-207.
- Molander, A. 1928. Animal communities on soft bottom areas in the Gullmar fjord. *Kristinebergs Zoologiska Station 1877 - 1927, 2:* 1-90. K. Svenska Akademien. Uppsala.

- Moore, H. B. 1931. Muds of the Clyde Sea area. III. Chemical and physical conditions; rate and nature of sedimentation; and fauna. *Journal of the Marine Biological Association of the United Kingdom* XVII: 325-358.
- Morse, M. P. 1969. On the feeding of the nudibranch, *Coryphella verrucosa rufibranchialis*, with a discussion of its taxonomy. *Nautilus*, 83: 37-40.
- Mortensen, T. 1924. Pighude (Echinodermer). *Danmarks Fauna* 27: 1-274.
- Mothershead, R. F. II & R. C. Hale 1992. Influence of ecdysis on the accumulation of polycyclic aromatic hydrocarbons in field exposed blue crabs (*Callinectes Sapidus*). *Marine Environmental Research* 33: 145-156.
- Mulicki, Z. 1959. Warunki srodowiskowe wieloszeta Harmothoe sarsi Kinberg w Baltyku. (The polichaet Harmothoe sarsi Kinberg and its environments in the Baltic Sea.) *Prace Morskiego Instytutu Rybackiego w Gdyni. Czesc oOceanograficzno - ichtiologiczna. (Reports of the Sea Fisheries Institute in Gdynia. Oceanographical and ichthyological part.) Tom 10/A: 163-174.*
- Munthe-Kaas, H. 1968. Surface pollution and light extinction in the Oslofjord. *Helgoländer wissenschaftliche Meeresuntersuchungen* 17: 476-488.
- Norges Geografiske Oppmåling. Den Geodetiske Avdeling. *Vannstandsobservasjoner. Stasjon Oslo. 1951.*
- Nuglisch, K. 1985. Foraminiferen - marine Mikroorganismen. *Die Neue Brehm-Bücherei* 560:1-174.
- Odhner, N. H. 1939. Opisthobranchiate Mollusca from the western and northern coasts of Norway. *Det Kgl. Norske Videnskabers Selskabs Skrifter* 1939 NR 1: 1-92.
- Olsgard, F. 1993. Do toxic algal blooms affect subtidal soft-bottom communities? *Marine Ecology Progress Series. Vol. 102: 269-286.*
- Olsgard, F. 1994. Miljøgiftundersøkelser i Indre Oslofjord. Delrapport 7. Bløtbunnsfauna i Oslo havneområde. *NIVA-rapport, 0-92131, l. nr. 3104. 25 pp.*
- Olsgard, F. 1995. Overvåking av forurensningssituasjonen i Indre Oslofjord. Undersøkelser av bløtbunnsfauna 1993. *Statlig program for forurensningsovervåkning. Rapport 622/95. TA-1258/1995: 1-106.*
- Østvedt, O.-J. 1955. Zooplankton investigations from Weater Ship M in the Norwegian Sea, 1948-49. *Hvalrådets Skrifter. Scientific Results of Marine Biological Research. 40: 1-93.*
- Paasche, E. & S. R. Erga 1988. Phosphorus and nitrogen limitation of phytoplankton in the Inner Oslofjord (Norway). *Sarsia* 73: 229-243.
- Pearson, T. H., J. S. Gray & P. J. Johannessen 1983. Objective selection of sensitive species indicative of pollution-induced change in benthic communities. 2. Data analyses. *Marine Ecology - Progress Series* 12: 237-255.
- Pearson, T. H. & R. Rosenberg 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography and Marine Biology. Annual Revue. 16: 229-311.*
- Pelczar, M. J., R. D. Reid & E. C. S. Chan 1977. *Microbiology*. Fourth ed. McGraw-Hill Book Comp., N. Y. 952 pp.
- Petersen, C. G. J. 1913. Havets Bonitering II. Om Havbundens Dyresamfund og om disses Betydning for den marine Zoogeografi. *Beretning til Landbrugsministeriet fra den danske biologiske Station XXI: 1-42 + Pl. 1-6 + Kort Nr. I-III.*
- Petersen, C. G. J. 1915. Om Havbundens Dyresamfund i Skagerak, Kristianiafjord og de danske Farvande. *Beretning til Landbrugsministeriet fra den danske biologiske Station XXIII: 3-26 + 6 tabeller og et kart.*

- Petersen, C. G. J. 1918. Havbunden og Fiskenes Ernæring. En Oversigt over Arbejderne vedrørende vore Farvandes Bonitering i 1883-1917. *Beretning til Landbrugsministeriet fra den danske biologiske Station XXV*: 1-57 + Tavle I-X + kart.
- Pettibone, M. H. 1963. Marine polychaete worms of the New England Region. 1. Families Aphroditidae through Trochochaetidae. *United States National Museum Bulletin 227. Part 1*: 1-356.
- Rachor, E. 1990. Changes in sublittoral zoobenthos in the German Bight with regard to eutrophication. *Netherlands Journal of Sea Research 25 (1/2)*: 209-214.
- Rasmussen, B. 1953. On the geographical variation in growth and sexual development of the deep sea prawn. *Fiskeridirektoratets Skrifter. Serie Havundersøkelser. Vol. X (3)*: 1-160.
- Redfield, A. C., B. H. Ketchum & F. A. Richards 1963. The influence of organisms on the composition of seawater. Pp. 26-77 in: M. N. Hill (ed.). *The sea. Ideas and observations on progress in the study of the seas. Vol. 2. The composition of sea-water. Comparative and descriptive oceanography*. Interscience Publishers. New York. 554 pp.
- Rees, C. B. 1949. Continuous plankton records: The distribution of *Calanus finmarchicus* (GUNN.) and its two forms in the North Sea, 1938-39. *Hull Bulletins of Marine Ecology No. 14, vol. II*: 215-275.
- Reish, D. J. 1988. The use of toxicity testing in marine environmental research. Pp. 231-245 in: D. F. Soule & G. S. Kleppel (eds). *Marine organisms as indicators*. Springer-Verlag, New York. 342 pp.
- Remane, A. 1933. Verteilung und Organisation der benthonischen Mikrofauna der Kieler Bucht. *Wissenschaftliche Meeresuntersuchungen. Abteilung Kiel, N. F. 21*: 161-221.
- Remane, A. 1940. Einführung in die zoologische Ökologie der Nord- und Ostsee. *Die Tierwelt der Nord- und Ostsee Ia*: 1-238.
- Reppe, B. 1986. Hydroacoustic assessment and ecology of krill stocks in the Inner Oslofjord. *Hovedfagsoppgave. Universitetet i Oslo, Biologisk institutt, avd. marin zoologi og marin kjemi*. 88 pp.
- Rheinheimer, G. 1972. Dissolved gases - Bacteria, fungi and blue-green algae. Pp. 1459-1469 in: *Marine ecology. A comprehensive, integrated treatise on life in oceans and coastal waters*. Ed. by O. Kinne. Vol. I. *Environmental factors. Part 3*. Wiley-Interscience, London.
- Rhoads, D. C. & L. F. Boyer 1982. The effects of marine benthos on physical properties of sediments: A successional perspective. Pp. 3-52 in: P. McCall & M. J. S. Tevesz (eds) *Animal - Sediment relations*. Plenum Press, N. Y. 336 pp.
- Richards, F. A. 1965. Anoxic basins and fjords. Pp. 611-645 in: J. P. Riley & G. Skirrow (eds) *Chemical oceanography. Vol. I*. Academic Press, London. 712 pp.
- Risdal, D. 1963. Foraminiferfaunaen i en del sedimentkjerner fra indre Oslofjord. *Norges Geologiske Undersøkelse Nr. 224*: 1-90.
- Rittenberg, S. C., K. O. Emery & W. L. Orr 1955. Regeneration of nutrients in sediments of marine basins. *Deep-Sea Research. Vol. 3*: 23-45.
- Rose, M. 1933. Copépodes pélagiques. *Faune de France 26*: 1-374. Paris.
- Rosenberg, R. 1976. Benthic faunal dynamics during succession following pollution abatement in a Swedish estuary. *Oikos 27*: 414-427.
- Rosenberg, R., J. S. Gray, A. B. Josefson & T. H. Pearson 1987. Petersen's benthic stations revisited. II. Is the Oslofjord and eastern Skagerrak enriched? *Journal of Experimental Marine Biology and Ecology. Vol. 105*: 219-251.
- Russell, F. S. 1925. The vertical distribution of marine macroplankton. An observation on diurnal changes. *Journal of the Marine Biological Association of the United Kingdom XIII*: 769-809.

- Russell, F. S. 1928 a. The vertical distribution of marine macroplankton . VI. Further observations on diurnal changes. *Journal of the Marine Biological Association of the United Kingdom* XV: 81-103.
- Russell, F. S. 1928 b. A net for catching plankton near the bottom. *Journal of the Marine Biological Association of the United Kingdom* XV: 105-108.
- Russell, F. S. 1953. *The Medusae of the British Isles*. Cambridge University Press. 530 pp. + Pl. I-XXXV.
- Ruud, J. T. 1968. Introduction to the studies of pollution in the Oslofjord. *Helgoländer wissenschaftliche Meeresuntersuchungen*. 17: 455-461.
- Ruud, J. T. & J. Versvik 1966. Fisket i Oslofjorden. *Oslofjorden og dens forurensningsproblemer. I. Undersøkelsen 1962 - 1965. Delrapport nr. 3: 1-38 + Fig. 1-3*. Norsk Institutt for Vannforskning, Oslo.
- Salle, A. J. 1973. *Fundamental principles of bacteriology*. Seventh edition. McGraw-Hill Book Company, N. Y. 1094 pp.
- Sanders, H. L. 1968. Marine benthic diversity: a comparative study. *American Naturalist* 102: 243-282.
- Sars, G. O. 1864. Om en anomal Gruppe af Isopoder. *Forhandlinger, Videnskabselskabet i Christiania* 1863: 205-221.
- Sars, G. O. 1865. Om den aberrante Krebsdyrgruppe Cumacea, og dens nordiske Arter. *Forhandlinger, Videnskabselskabet i Christiania* 1864: 128-208.
- Sars, G. O. 1866. Oversigt af Norges marine Ostracoder. *Forhandlinger, Videnskabs-Selskabet i Christiania* 1865: 1-130.
- Sars, G. O. 1868. Beretning om en i Sommeren 1865 foretagen zoologisk Reise ved Kysterne af Christianias og Christiansands Stifter. *Nyt Magazin for Naturvidenskaberne* XV: 84-128.
- Sars, G. O. 1869. *Undersøgelser over Christianiafjordens Dyboandsfauna, anstillede paa en i Sommeren 1868 foretagen zoologisk Reise*. Johan Dahl, Christiania. 58 pp.
- Sars, G. O. 1870 a. Nye Dybvandscrustaceer fra Lofoten. *Forhandlinger, Videnskabs-Selskabet i Christiania* 1869: 147-174.
- Sars, G. O. 1870 b. *Carcinologiske Bidrag til Norges Fauna. I. Monographi over de ved Norges Kyster forekommende Mysider. Første Hefte: 1-64 + Tabula I-V*. Christiania.
- Sars, G. O. 1872. *Carcinologiske Bidrag til Norges Fauna. I. Monographi over de ved Norges Kyster forekommende Mysider. Andet Hefte: 1-34 + Tabula VI-VIII*. Christiania.
- Sars, G. O. 1878. *Bidrag til Kundskaben om Norges arktiske Fauna. I. Mollusca regionis arcticæ Novægiæ*. Christiania. 466 pp. + Kart + Tab. 1-34 + Tab. I-XVIII.
- Sars, G. O. 1879. *Carcinologiske Bidrag til Norges Fauna. I. Monographi over de ved Norges Kyster forekommende Mysider. Tredje Hefte: 1-131 + Tabula IX-XLII*. Christiania.
- Sars, G. O. 1883. Oversigt af Norges Crustaceer med foreløbige Bemærkninger over de nye eller mindre bekjendte Arter. *Forhandlinger, Videnskabselskabet i Christiania* 1882: 1-124.
- Sars, G. O. 1895 a. *An account of the Crustacea of Norway. I. Amphipoda (Text): 1-711*. Bergen Museum, Bergen.
- Sars, G. O. 1895 b. *An account of the Crustacea of Norway. I. Amphipoda (Plates): Pp. I-XIII + Pl. 1-240 + Suppl. Pl. I-VI*. Bergen Museum, Bergen.
- Sars, G. O. 1899. *An account of the Crustacea of Norway. II. Isopoda: 1-270 + Pl. 1-100*. Bergen Musuem, Bergen.

- Sars, G. O. 1900. *An account of the Crustacea of Norway. III. Cumacea*: 1-115 + Pl. I-LXXII. Bergen Museum, Bergen.
- Sars, G. O. 1903. *An account of the Crustacea of Norway. IV. Copepoda Calanoida*: 1-171 + Pl. I-CII + Suppl. Pl. I-VI. Bergen Museum, Bergen.
- Sars, G. O. 1911 a. *An account of the Crustacea of Norway. V. Copepoda Harpacticoida (Text)*: 1-449. Bergen Museum, Bergen.
- Sars, G. O. 1911 b. *An account of the Crustacea of Norway. V. Copepoda Harpacticoida (Plates)*: I-CCXXX + Supplm. Pl. 1-54. Bergen Museum, Bergen.
- Sars, G. O. 1918. *An account of the Crustacea of Norway. VI. Copepoda Cyclopoida*: 1-225 + Pl. I-CXVIII. Bergen Museum, Bergen.
- Sars, G. O. 1921 a. *An account of the Crustacea of Norway. VII. Copepoda. Supplement*: 1-121 + Pl. I-LXXVI. Bergen Museum, Bergen.
- Sars, G. O. 1921 b. *An account of the Crustacea of Norway. VIII. Copepoda Monstrilloida & Notodelphyoida*: 1-91 + Pl. I-XXXVII. Bergen Musuem, Bergen.
- Sars, G. O. 1928. *An account of the Crustacea of Norway. IX. Ostracoda*: 1-277 + Pl. I-CXIX. Bergen Museum, Bergen.
- Sars, M. 1829. *Bidrag til Söedyrenes Naturhistorie. 1*. Bergen. 59 pp.
- Sars, M. 1866. Om arktiske Dyreformer i Christianiafjorden. *Forhandlinger, Videnskabs-Selskabet i Christiania 1865*: 196-200.
- Sars, M. 1868. Bidrag til Kundskab om Christianiafjordens Fauna. *Nyt Magazin for Naturvidenskaberne XV*: 241-344.
- Sars, M. 1869. Fortsatte Bemærkninger over det dyriske Livs Udbredning i Havets Dybder. *Forhandlinger, Videnskabselskabet i Christiania 1868*: 246-275.
- Schram, T. A. 1970. Studies on the meroplankton in the Inner Oslofjord. II. Regional differences and seasonal changes in the specific distribution of larvae. *Nytt magasin for Zoologi 18 (1)*: 1-21.
- Schulz, S. 1969. Rückgang des Benthos in der Lübecker Bucht - Ein Vergleich mit den Verhältnissen im Bornholmbecken. *Limnologica (Berlin) 7 (1)*: 23-25.
- Simensen, T. og C.-H. Knudsen (red.) 1970. Oslofjorden og dens forurensningsproblemer. II. Utredning av tekniske løsninger. 111 pp. Norsk Institutt for Vannforskning, Oslo.
- Sivertsen, E. & L. B. Holthuis 1956. Crustacea Decapoda (The Penaeidea and Stenopodidea excepted). *Reports on Scientific Results of the "Michael Sars" North Atlantic Deep-Sea Expedition 1910. Vol. V*.
- Sjöstedt, L. G. 1936. Untersuchungen aus dem Öresund XXII. Beitrag zur Hydrographie des Sundes. *Acta Universitatis Lundensis, N. F. Avd. 2, 31 (12)*: 1-61.
- Skogsberg, T. 1920. Studies on marine ostracods. I. *Zoologiska bidrag från Uppsala 1 (Supplement)*: 1-784.
- Skullerud, A. M. 1985. Klassifikasjon av bunndyrsamfunn i Vestfjorden, indre Oslofjord. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi*. 111 pp. + tabeller.
- Smedstad, O. M. 1972. On the biology of *Aglantha digitale rosea* (Forbes) [Coelenterata: Trachymedusae] in the Inner Oslofjord. *Nytt magasin for Zoologi 20 (2)*: 111-135.
- Smith Jr, K. L., R. S. Kaufmann & W. W. Wakefield 1993. Mobile megafaunal activity monitored with a time-lapse camera in the abyssal North Pacific. *Deep Sea Research. I. Oceanographic research papers. Vol. 40 (11/12)*: 2307-2324.

- Sømme, J. D. 1934. Animal plankton of the Norwegian coast waters and the open sea. I. Production of *Calanus finmarchicus* (Gunner) and *Calanus hyperboreus* (Krøyer) in the Lofoten area. *Fiskeridirektoratets Skrifter. Serie Havundersøkelser. IV (9): 1-163.*
- Stålesen, G. 1964. En sammenlikning mellom bunnfaunaen på reke-feltene i Oslofjorden like innenfor og utenfor Drøbakterskelen i 1933/34 og 1962. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi. 50 pp.*
- Stålesen, O. 1963. Fisken på rekefeltene i Oslofjorden. En undersøkelse av de forandringer som har funnet sted de siste 30 år. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi. 47 pp. + tabell.*
- Stebbing, T. R. R. 1906. Amphipoda I Gammaridea. *Das Tierreich. 21. Lieferung. 806 pp.*
- Stene, R. O. 1968. Utbredelsen av *Metridium senile* og endel andre vanlige arter fra hardbunnsfaunaen i Indre Oslofjord. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi. 87 pp.*
- Stephensen, K. 1910. Storkrebs I. Skjoldkrebs. *Danmarks Fauna 9: 1-193. København.*
- Stephensen, K. 1923. Crustacea Malacostraca. V. (Amphipoda. I.) *The Danish Ingolf-Expedition III (8): 1-100 + Tab.*
- Stephensen, K. 1925. Crustacea Malacostraca. VI. (Amphipoda II.) *The Danish Ingolf-Expedition III (9): 101-178 + Tab.*
- Stephensen, K. 1928. Storkrebs II. Ringkrebs 1. Tanglopper (Amfipoder). *Danmarks Fauana 32: 1-399.*
- Stephensen, K. 1931. Crustacea Malacostraca. VII. (Apmphipoda. III) *The Danish Ingolf-Expedition III (11): 179-290 + Tab.*
- Stephensen, K. 1935. The Amphipoda of N. Norway and Spitsbergen with adjacent waters. *Tromsø Museums Skrifter vol. III. Part I: 1-140 + 2 maps.*
- Stephensen, K. 1944. Crustacea Malacostraca. VIII. (Amphipoda. IV.) *The Danish Ingolf-Expedition III (13): 1-51 + Tab.*
- Stephensen, K. 1948. Storkrebs IV. Ringkrebs 3. Tanglus (marine isopoder) og Tanaider. *Danmarks Fauna 53: 1-187. København.*
- Steven, G. A. 1930. Bottom fauna and the food of fishes. *Journal of the Marine Biological Association of the United Kingdom XVI: 677-700.*
- Strömngren, T. 1974. Zooplankton and hydrography in Trondheimsfjorden on the west coast of Norway. *Det Kgl. Norske Videnskabers Selskab. Museet. Miscellana 17: 1-35.*
- Sundene, O. 1953. The algal vegetation of Oslofjord. *Skrifter utgitt av Det Norske Videnskaps-Akademi i Oslo. I. Mat.-Naturv. Klasse. 1953. No. 2: 1-245.*
- Sverdrup, H. U., M. W. Johnson & R. H. Fleming 1942. *The oceans. Their physics, chemistry, and general biology.* Prentice-Hall, Inc., New York. 1087 pp.
- Swennen, C. 1961. Data on distribution, reproduction and ecology of the nudibranchiate molluscs occurring in the Netherlands. *Netherlands Journal of Sea Research 1, 1/2: 191-240.*
- Sylvester-Bradley, P. C. 1950. The identity of the ostracod *Philomedes brenda* (Baird). *Annual Magazine of Natural History (12) 3: 777-778.*
- Tardent, P. 1979. *Meeresbiologie. Eine Einführung. 116 Abbildungen, 26 Tabellen.* Georg Thieme Verlag Stuttgart. 381 pp.

- Thamdrup, H. M. 1935. Beiträge zur Ökologie der Wattenfauna auf experimenteller Grundlage. *Meddelelser fra Kommissionen for Danmarks Fiskeri- og Havundersøgelser. Serie Fiskeri. Bind X(2):* 1-135.
- Theede, H. 1973. Comparative studies on the influence of oxygen deficiency and hydrogen sulphide on marine bottom invertebrates. *Netherlands Journal of Sea Research* 7: 244-252.
- Thompson, T. E. 1960. Defensive adaptations in opisthobranchs. *Journal of the Marine Biological Association of the United Kingdom* 39: 123-134.
- Thompson, T. E. & G. H. Brown, 1984. *Biology of Opisthobranch Molluscs. Vol. II.* The Ray Society, London. 229 pp.
- Thorson, G. 1946. Reproduction and larval development of Danish marine bottom invertebrates, with special reference to the planktonic larvae in the Sound (Øresund). *Meddelelser fra Kommissionen for Danmarks Fiskeri- og Havundersøgelser. Serie Plankton. Bind 4 (1):* 1-523.
- Thorson, G. 1957. Sampling the benthos. Pp. 61-73 in: J. W. Hedgpeth (ed.) *Treatise on marine ecology and paleoecology. The Geological Society of America Memoir* 67.1297 pp.
- Thorson, G. 1968. Infaunaen, den jævne havbunds dyresamfund. Pp. 82-166 in: A. Nørrevang & T. J. Meyer (eds) *Danmarks natur. Bind 3. Havet.* Politikens Forlag. 488 pp.
- Todd, C. D. 1981. The ecology of nudibranch molluscs. Pp. 141-234 in: M. Barnes (ed.) *Oceanography and Marine Biology. An annual review. Vol. 19.* Aberdeen University Press. 655 pp.
- Tranter, D. J. & P. E. Smith 1968. Filtration performance. Pp. 27-56 in: *Zooplankton sampling*, published by UNESCO, Paris. 174 pp.
- Ursin, E. 1954. Efficiency of marine bottom samplers of the van Veen and Petersen types. *Meddelelser fra Danmarks Fiskeri- og Havundersøgelser. Ny Serie I (7):* 1-7.
- Valiela, I. 1984. *Marine ecological processes.* Springer-Verlag, New York. 546 pp.
- Vartdal, R. 1976. *Pontophilus norvegicus* (M. SARS) i Korsfjorden. Reproduksjon, vekst og energiinnhold. *Hovedfagsoppgave. Universitetet i Bergen. Biologisk Stasjon, Espeland.* 86 pp.
- Vathanaprida, C. 1966. On the life history of *Pandalina profunda* Holthuis in the Oslofjord. MS. 21 pp. + 12 fig. + 11 tab. Universitetet i Oslo. Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi.
- Vernberg, F. J. 1972. Dissolved gases - Animals. Pp. 1491-1526 in: *Marine Ecology. A comprehensive, integrated treatise on life in oceans and coastal waters.* Ed. by O. Kinne. Vol. I. *Environmental factors. Part 3.* Wiley-Interscience, London.
- Versvik, J. 1963. Undersøkelser over *Lycenchelys sarsi* i Oslofjorden, - og en sammenlikning med et materiale fra Barentshavet. *Hovedfagsoppgave. Universitetet i Oslo, Institutt for marinbiologi og limnologi, avd. marin zoologi og marin kjemi.* 68 pp.
- Wada, E. & A. Hattori 1991. *Nitrogen in the sea: Forms, abundances, and processes.* CRC Press, Boca Raton (Florida). 208 pp-
- Waterman, T. H. 1974. Underwater light and the orientation of animals. Pp. 415-443 in: N. G. Jerlov & E. Steemann Nielsen (eds) *Optical aspects of oceanography.* Academic Press, London. 494 pp.
- Weigelt, M. & H. Rumohr 1986. Effects of wide-range oxygen depletion on benthic fauna and demersal fish in Kiel Bay 1981-1983. *Meeresforschung* 31 (2): 124-136.
- Wiborg, K. F. 1940. The production of zooplankton in the Oslofjord in 1933-1934, with special reference to the copepods. *Hvalrådets Skrifter* 21: 1-85 + Tab. I-II.

- Wiborg, K. F. 1951. The whirling vessel. An apparatus for the fractioning of plankton samples. *Fiskeridirektoratets Skrifter. Serie Havundersøkelser* 9 (13): 1-16.
- Wiborg, K. F. 1954. Investigations on zooplankton in coastal and offshore waters of western and north-western Norway. With special reference to the copepods. *Fiskeridirektoratets Skrifter. Serie Havundersøkelser. Vol. XI. No. 1*: 1-246.
- Wigley, R. L. 1967. Comparative efficiencies of van Veen and Smith-McIntyre grab samplers as revealed by motion pictures. *Ecology* 48 (1): 168-169.
- Wilson, C. B. 1932. The Copepoda of the Woods Hole region, Massachusetts. *Bulletin of the United States Natural History Museum* 158: 1-635 + Pl. 1-41.
- With, C. 1915. Copepoda I. Calanoida Amphascandria. *The Danish Ingolf-Expedition III* (4): 1-260 + Pl. I-VIII.
- Wollebæk, A. 1903. Ræker og rækefisket. *Aarsberetning vedkommende Norges Fiskerier for 1903. 2^{det} Hefte*: 165-229.
- Wollebæk, A. 1906. Nogle statistiske Oplysninger om Fisket efter Dybvandsræker (*Pandalus borealis*) i Kristianiafjorden indenfor Drøbak. *Norsk Fiskeritidende* 1906. Nr. 11: 471-480.
- Wyrтки, K. 1962. The oxygen minima in relation to ocean circulation. *Deep Sea Research*, 9: 11-23.
- Zimmer, C. 1933. Mysidacea. *Die Tierwelt der Nord- und Ostsee Xg3*: 29-69. Leipzig.
- ZoBell, C. E. 1940. The effect of oxygen tension on the rate of oxidation of organic matter in sea water by bacteria. *Journal of Marine Research*, vol. 3, no 3: 211-223.

GRUPPER	ARTER/TAXA	G53	G59	G60	G61	G73	G81	G82	G84	G85	G86	G87	G88	G89	G90
	<i>Cheirocratus</i> sp.	0	0	0	0	0	0	0	6	0	0	0	0	0	0
	Gammaridae	0	0	0	0	0	0	0	6	3	0	0	0	0	0
	Photidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Podoceropsis sophiae</i>	0	0	0	0	0	5	0	0	0	0	0	0	0	0
	<i>Corophium bonelli</i>	2	1	1	0	0	5	5	11	0	0	3	2	12	0
	<i>Arrhis phyllonyx</i>	79	69	21	39	0	3	0	0	0	0	0	0	0	0
	<i>Monoculodes packardi</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Monoculodes tenuirostris</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Halicreion longicaudatus</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Halirages fulvocinctus</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Euphausiacea	<i>Meganyctiphanes norvegica</i>	0	1	0	5	0	1	0	11	0	0	0	1	0	0
	<i>Thysanoessa raschi</i>	0	0	0	2	0	261	0	6	0	0	0	0	0	7
	<i>Thysanoessa inermis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Decapoda	<i>Crangon allmani</i>	1	16	8	4	4	2	0	3	14	14	23	4	10	15
	<i>Pontophilus norvegicus</i>	13	16	51		4	3	0	6	2	2	5	1	1	9
	<i>Spirontocaris lilljeborgi</i>	0	5	1	1	0	2	0	0	4	2	1	1	3	9
	<i>Lobbeus polaris</i>	0	0	0	0	0	1	0	1	1	0	0	0	1	0
	<i>Pandalus borealis</i>	1	1	0	4	0	0	0	0	1	2	0	0	0	0
	<i>Pandalus propinquus</i>	0	1	0	0	0	0	0	0	12	0	0	0	0	0
	<i>Pandalus borealis</i> juv	1	1	0	0	0	0	0	206	188	19	0	42	38	197
	<i>Pandalina profunda</i>	3	1	0	30	4	27	1	86	40	224	108	7	125	547
	<i>Pasiphaea tarda</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Pandalus montagui</i>	0	0	0	1	0	0	0	0	0	0	0	0	0	0
Mollusca	Nudibranchiata	1	0	0	0	70	0	0	0	0	0	0	0	0	0
	<i>Delectopecten vitreus</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Pseudamussium septemradiatum</i>	2	0	1	1	0	0	0	0	0	0	0	0	0	0
	<i>Abra nitida</i>	0	0	0	0	0	0	0	0	560	0	0	2	0	0
	<i>Thyasira</i> sp	0	0	0	0	0	0	161	34	80	0	0	12	3	0
	<i>Bivalvia unident</i>	6	3	8	0	0	108	73	80	18	42	246	0	9	0
Echinodermata	Asteroida (larvae)	81	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ophiuroidea	4	3	6	0	19	0	0	40	45	425	246	5	34	0
	Echinoidea	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Holothuroidea	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tunicata	Ascidia	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pisces	<i>Myxine glutinosa</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Lycenchelys sarsi</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	<i>Rhinonemus cimrius</i>	0	0	0	0	0	0	0	0	1	0	0	0	0	0

GRUPPER	ARTER/TAXA	SD81	SD82	SD83	SD84	SD85	SD86	SD87	SD88	SD89
	<i>Cheirocratus</i> sp.	0	0	0	0	0	0	0	0	0
	Gammaridae	0	0	0	0	4	0	0	0	25
	Photidae	3	0	0	0	0	2	0	11	6
	<i>Podoceroopsis sophiae</i>	0	0	0	0	0	0	0	0	0
	<i>Corophium bonelli</i>	0	0	0	0	0	0	2	4	0
	<i>Arrhis phyllonyx</i>	0	0	0	0	0	0	0	0	0
	<i>Monoculodes packardi</i>	0	0	0	0	0	0	0	0	0
	<i>Monoculodes tenuirostratis</i>	0	0	0	0	0	0	0	0	0
	<i>Halicreion longicaudatus</i>	0	0	0	0	0	0	0	0	0
	<i>Halirages fulvocinctus</i>	0	0	0	0	0	0	0	0	0
Euphausiacea	<i>Meganyctiphanes norvegica</i>	5	1	6	1	0	6	8	7	84
	<i>Thysanoessa raschi</i>	13	0	1	3	0	1	1	0	45
	<i>Thysanoessa inermis</i>	0	0	0	0	1	1	0	0	0
Decapoda	<i>Crangon allmani</i>	3	0	0	0	0	2	2	1	19
	<i>Pontophilus norvegicus</i>	0	0	0	0	0	1	1	1	1
	<i>Spirontocaris lilljeborgi</i>	0	0	1	0	0	2	2	0	10
	<i>Lebbeus polaris</i>	0	0	0	0	0	0	0	0	2
	<i>Pandalus borealis</i>	0	0	3	0	0	1	1	1	0
	<i>Pandalus propinquus</i>	0	0	0	0	1	1	0	0	1
	<i>Pandalus borealis juv</i>	0	1	0	2	7	1	0	48	6
	<i>Pandalina profunda</i>	5	2	0	0	1	20	53	4	8
	<i>Pasiphaea tarda</i>	0	0	0	0	0	0	0	0	0
	<i>Pandalus montagui</i>	0	0	0	0	0	0	0	0	0
Mollusca	Nudibranchiata	0	0	0	0	0	0	0	0	0
	<i>Delectopecten vitreus</i>	0	0	0	0	4	0	0	1	11
	<i>Pseudamussium septemradiatum</i>	3	0	1	0	0	2	4	3	0
	<i>Abra nitida</i>	0	0	0	0	8	0	0	69	28
	<i>Thyasira</i> sp.	21	99	0	38	93	0	0	28	87
	<i>Bivalvia unident</i>	77	440	13	0	102	8	0	14	0
Echinodermata	Asteroida (larvae)	0	0	0	0	0	0	0	0	0
	Ophiuroidea	3	0	6	44	30	108	34	124	160
	Echinoidea	0	0	0	0	0	0	0	0	0
	Holothuroidea	0	0	0	0	0	0	0	0	0
Tunicata	Ascidia	0	0	0	4	17	2	0	1	1
Pisces	<i>Myxine glutinosa</i>	0	0	0	0	0	0	0	0	0
	<i>Lycenchelys sarsi</i>	0	0	0	0	0	0	0	0	0
	<i>Rhinonemus cimbricus</i>	0	0	0	0	0	1	1	1	0

GRUPPER	ARTER/TAXA	SS85	SS86	SS87	SS88	SS89	SS90
Foraminifera	Uidentifisert	36	71	0	9	13	30
Coelenterata	Tesseractaria muscosa	0	0	0	0	0	0
	Aglantha digitale	0	0	0	0	0	0
Chaetognatha	Sagitta elegans	4	6	0	6	4	0
	Eukrohnia hamata	0	3	0	0	0	0
Polychaeta	Syllis sp.	0	0	1	0	0	0
	Nereimyra punctata	0	42	0	0	0	0
	Ophiodromus flexuosus	0	0	0	0	0	0
	Antinoella sarsi	0	0	1	0	0	0
	Harmothoe spp.	0	0	0	0	0	0
	Leanira tetragona	0	0	0	0	0	0
	Pholoe minuta	0	3	0	0	0	0
	Phyllocoridae	0	0	0	0	0	0
	Nephtys spp.	0	0	0	4	0	0
	Lumbrineris spp.	44	0	0	0	0	0
	Sphaerodorum gracilis	0	0	0	0	0	0
	Spionidae	0	13	1	0	0	0
	Chaetozone setosa	0	0	0	0	0	0
	Flabelligeridae	0	0	0	0	0	0
	Scalibregmidae	0	0	0	0	0	0
	Ophelidae	0	0	0	0	0	0
	Capitella capitata	347	1017	1	0	0	0
	Ampharete baltica	0	0	0	0	0	0
	Ampharete falcata	0	0	0	0	0	0
	Amphiteis gunneri	0	0	0	0	0	0
	Ampharete sp.	0	0	0	0	0	0
	Melinna cristata	0	0	0	0	0	0
	Eupolymnia spp.	0	0	0	0	0	0
	Pista cristata	0	0	0	0	0	0
	Pectinaria spp.	0	0	0	0	0	0
	Thelepus spp.	0	0	0	0	0	0
	Trichobranchus spp.	0	0	0	0	0	0
	Terebellides stroemi	0	0	0	0	0	0
	Terebellidae sp.	0	0	0	0	0	0
	Sabellidae spp.	0	0	0	0	0	0
	Uidentifisert	0	10	0	0	4	0
	Hesionidae	0	0	0	0	0	0
	Aphroditidae unidentified	0	0	0	0	0	0
	Glyceridae	0	0	0	0	0	0
	Heteromastus sp.	0	0	0	0	0	0
Ostracoda	Philomedes brenda	0	0	0	0	0	0
	Philomedes liljeborgi	0	0	0	0	0	0
	Conchoecia elegans	0	0	0	0	0	0
	Conchoecia borealis	0	0	0	0	0	0
	Conchoecia obtusata	0	0	0	0	0	0
	Cytherella abyssorum	0	0	0	0	0	0
	Macrocypris minna	0	0	0	0	0	0
	Macrocypris angusta	4	0	0	0	0	0
	Elofonella concinna	0	0	0	0	0	0
	Eucytheridea punctillata	0	0	0	0	0	0
	Echinocythereis echinata	0	0	0	0	0	0
	Cythereis dunelmensis	0	0	0	0	0	0
	Hemicythere latimarginata	0	0	0	0	0	0
	Cythereis spp.	0	0	0	0	0	0
	Cytheropteron alatum	0	0	0	0	0	0
	Cythereis jonesi	0	0	0	0	0	0
	Cytherella abyssorum	0	0	0	0	0	0
Copepoda	Calanus finmarchicus	4	13	17	0	122	0
	Calanus helgolandicus	28	16	0	6	161	0
	Calanus hyperboreus	0	0	0	0	0	0
	Calanus sp. juv.	0	32	0	3	256	1015
	Pseudocalanus elongatus	0	3	0	0	4	0
	Bradydium bradyi	0	0	0	0	0	0
	Chiridius armatus	0	0	0	0	0	0
	Euchaeta norvegica	4	6	0	0	0	0
	Xanthocalanus fallax	0	0	0	0	0	0
	Xanthocalanus propinquus	0	0	0	0	0	0
	Xanthocalanus sp.	0	0	0	0	0	0
	Scolecithricella minor	0	6	0	0	0	0
	Diaxis hibernica	0	0	0	0	0	0
	Parastephus pallidus	0	0	0	0	0	0
	Tharybis macrophthalma	0	0	0	0	0	0
	Centropages hamatus	4	0	1	15	0	0
	Centropages typicus	0	0	0	15	83	61
	Temora longicornis	0	0	0	6	178	15
	Metridia longa	16	116	1	9	0	0
	Metridia lucens	0	0	0	0	0	0

GRUPPER	ARTER/TAXA	SS85	SS86	SS87	SS88	SS89	SS90
	<i>Pleuromamma robusta</i>	0	0	0	0	0	0
	<i>Heterorhabdus norvegicus</i>	0	0	0	0	0	0
	Unidentified copepodites	0	0	0	0	0	0
	<i>Harpacticoida</i>	0	0	0	0	0	0
	<i>Scottula inaequicornis</i>	0	0	0	0	0	0
	<i>Anomoclausia indrehusae</i>	0	0	0	0	0	0
Mysidacea	<i>Boreomysis arctica</i>	0	0	0	0	0	0
	<i>Erythrops serrata</i>	0	0	0	0	0	0
	<i>Amblyops abbreviata</i>	0	0	0	0	0	0
	<i>Mysideis insignis</i>	0	0	0	0	0	0
	<i>Mysidopsis didelphys</i>	0	0	0	0	0	0
Cumacea	<i>Leucon acutirostris</i>	0	0	0	0	0	0
	<i>Leucon nasicus</i>	0	0	0	0	0	0
	<i>Leucon</i> sp.	0	0	0	0	0	0
	<i>Eudorella emarginata</i>	0	0	0	0	0	0
	<i>Campylaspis rubicunda</i>	0	0	0	0	0	0
	<i>Campylaspis costata</i>	0	0	0	0	0	0
	<i>Campylaspis glabra</i>	0	0	0	0	0	0
	<i>Campylaspis</i> sp.	0	0	0	0	0	0
	<i>Diastylis lucifera</i>	0	0	0	0	0	0
	<i>Diastylis tumida</i>	0	0	0	0	0	0
	<i>Diastylis cornuta</i>	0	0	0	0	0	0
	<i>Diastylis echinata</i>	0	0	0	0	0	0
	<i>Diastylodes serrata</i>	0	0	0	0	0	0
	<i>Diastylodes biplicata</i>	0	0	0	0	0	0
	<i>Leptostylis longimana</i>	0	0	0	0	0	0
	<i>Eudorella truncatula</i>	0	0	0	0	0	0
Tanaidacea	Tanaidacea	0	0	0	0	0	0
Isopoda	<i>Ianira maculosa</i>	0	0	0	0	0	0
	<i>Pleurogonium inerme</i>	0	0	0	0	0	0
	<i>Macrostylis spinifera</i>	0	0	0	0	0	0
	<i>Ischnosoma bispinosum</i>	0	0	0	0	0	0
	<i>Desmosoma lineare</i>	0	0	0	0	0	0
	<i>Desmosoma armatum</i>	0	0	0	0	0	0
	<i>Desmosoma laterale</i>	0	0	0	0	0	0
	<i>Desmosoma</i> sp.	0	0	0	0	0	0
	<i>Munna</i> sp.	0	0	0	0	0	0
	<i>Echinopleura aculeata</i>	0	0	0	0	0	0
	<i>Munnopsis typica</i>	1	0	0	0	0	0
	<i>Eurycope cornuta</i>	0	0	0	0	0	0
	<i>Eurycope phalangium</i>	0	0	0	0	0	0
	<i>Eurycope</i> sp.	0	0	0	0	0	0
	<i>Ilyarachna longicornis</i>	0	0	0	0	0	0
Amphipoda	<i>Themisto abyssorum</i>	0	0	0	0	0	0
	Lysianassidae	0	0	0	0	0	0
	<i>Tryphosa h�rtingi</i>	0	0	0	0	0	0
	<i>Tryphosites longipes</i>	0	0	0	0	0	0
	<i>Harpinia crenulata</i>	0	0	0	0	0	0
	<i>Harpinia pectinata</i>	0	0	0	0	0	0
	Phoxocephalidae unident.	0	0	0	0	0	0
	<i>Ampelisca</i> spp.	0	0	0	0	0	0
	<i>Stegocephalus inflatus</i>	0	0	0	0	0	0
	<i>Stegocephalus</i> sp.	0	0	0	0	0	0
	<i>Andaniexis abyss</i>	0	0	0	0	0	0
	<i>Andaniopsis nordlandica</i>	0	0	0	0	0	0
	Amphilochidae	0	0	0	0	0	0
	<i>Astyra abyss</i>	0	0	0	0	0	0
	Stenothoidae	0	0	0	0	0	0
	Oediceridae	0	0	0	0	0	0
	<i>Perioculodes longimanus</i>	0	0	0	0	0	0
	<i>Synchelidium haplocheles</i>	0	0	0	0	0	0
	<i>Synchelidium tenuimanum</i>	0	0	0	0	0	0
	<i>Synchelidium intermedium</i>	0	0	0	0	0	0
	<i>Synchelidium</i> sp.	0	0	0	0	0	0
	<i>Westwoodilla caecula</i>	0	0	0	0	0	0
	<i>Bathymedon longimanus</i>	0	0	0	0	0	0
	<i>Bathymedon saussurei</i>	0	0	0	0	0	0
	<i>Bruzelia</i> spp.	0	0	0	0	0	0
	<i>Halice abyss</i>	0	0	0	0	0	0
	<i>Rhachotropis macropus</i>	0	0	0	0	0	0
	<i>Rhachotropis leucophthalma</i>	0	0	0	0	0	0
	<i>Apherusa bispinosa</i>	0	0	0	0	0	0
	<i>Apherusa</i> sp.	0	0	0	0	0	0
	<i>Melphidippa borealis</i>	0	0	0	0	0	0
	<i>Eriopisa elongata</i>	0	0	0	0	0	0
	<i>Cheirocratus intermedium</i>	0	0	0	0	0	0
	<i>Cheirocratus sundewalli</i>	0	0	0	0	0	0

GRUPPER	ARTER/TAXA	SS85	SS86	SS87	SS88	SS89	SS90
	<i>Cheirocratus</i> sp.	0	0	0	0	0	0
	Gammaridae	0	0	0	0	0	0
	Photidae	0	0	0	0	0	0
	<i>Podoceroopsis sophiae</i>	0	0	0	0	0	0
	<i>Corophium bonelli</i>	0	0	0	3	0	0
	<i>Arrhis phyllonyx</i>	0	0	0	0	0	0
	<i>Monoculodes packardi</i>	0	0	0	0	0	0
	<i>Monoculodes tenuirostris</i>	0	0	0	0	0	0
	<i>Halicreion longicaudatus</i>	0	0	0	0	0	0
	<i>Halirages fulvocinctus</i>	0	0	0	0	0	0
Euphausiacea	<i>Meganyctiphanes norvegica</i>	0	0	2	0	4	0
	<i>Thysanoessa raschi</i>	0	0	0	0	0	0
	<i>Thysanoessa inermis</i>	0	0	0	0	0	0
Decapoda	<i>Crangon allmani</i>	0	0	0	0	0	0
	<i>Pontophilus norvegicus</i>	0	0	0	0	0	0
	<i>Spirontocaris lilljeborgi</i>	0	0	0	0	0	0
	<i>Lebbeus polaris</i>	0	0	0	0	0	0
	<i>Pandalus borealis</i>	0	0	0	0	0	0
	<i>Pandalus propinquus</i>	0	0	0	0	0	0
	<i>Pandalus borealis</i> juv	0	0	0	0	0	0
	<i>Pandalina profunda</i>	0	0	0	0	0	0
	<i>Pasiphaea tarda</i>	0	0	0	0	0	0
	<i>Pandalus montagui</i>	0	0	0	0	0	0
Mollusca	<i>Nudibranchiata</i>	0	0	0	0	0	0
	<i>Delectopecten vitreus</i>	0	0	0	0	0	0
	<i>Pseudamussium septemradiatum</i>	0	0	0	0	0	0
	<i>Abra nitida</i>	0	0	0	0	0	0
	<i>Thyasira</i> sp.	0	0	0	0	0	0
	<i>Bivalvia unident</i>	0	0	2	0	0	0
Echinodermata	Asteroida (larvae)	0	0	0	0	0	0
	Ophiuroidea	0	0	0	0	0	0
	Echinoidea	0	0	0	0	0	0
	Holothuroidea	0	0	0	0	0	0
Tunicata	Ascidia	0	0	0	0	0	0
Pisces	<i>Myxine glutinosa</i>	0	0	0	0	0	0
	<i>Lycenchelys sarsi</i>	0	0	0	0	0	0
	<i>Rhinonemus cimbricus</i>	0	0	0	0	0	0

GRUPPE	KOMPONENT	AUTOR	ANMERKNING
Coelenterata	<i>Tesserogastria musculosa</i>	BEYER	
	<i>Aglantha digitale</i>	(O. F. MÜLLER)	
Chaetognatha	<i>Sagitta elegans</i>	VERRILL	
	<i>Eukrohnia hamata</i>	(MÖBIUS)	
	<i>Spadella cephaloptera</i>	(BUSH)	
Polychaeta	Aphroditidae	MALMGREN	
	<i>Harmothoë</i>	KINBERG	
	<i>Antinoëlla sarsi</i>	(KINBERG)	
	<i>Pholoë minuta</i>	(FABRICIUS)	
	<i>Leanira tetragona</i>	ÖRSTED	
	Phyllodocidae	WILLIAMS	
	Hesionidae	MALMGREN	
	<i>Nereimyra punctata</i>	(O. F. MÜLLER)	
	<i>Ophiodromus flexuosus</i>	(DELLE CHIAJE)	
	Syllidae	GRUBE	
	<i>Nephtys</i>	CUVIER	
	<i>Sphaerodorum gracilis</i>	(RATHKE)	Inkl. <i>S. flavum</i> ÖRSTED, kfr. FAUCHALD (1974).
	Glyceridae	GRUBE	
	<i>Lumbrineris</i>	BLAINVILLE	
	Spionidae	GRUBE	
	<i>Polydora ciliata</i>	(JOHNSTON)	
	<i>Spiophanes kröyeri</i>	GRUBE	
	<i>Chaetozone setosa</i>	MALMGREN	
	Flabelligeridae	SAINT-JOSEPH	
	<i>Scalibregma inflatum</i>	RATHKE	
	<i>Polyphysia crassa</i>	ÖRSTED	
	Opheliidae	MALMGREN	
	<i>Capitella capitata</i>	(FABRICIUS)	
	<i>Heteromastus</i>	MCINTOSH	
	<i>Pectinaria</i>	SAVIGNY	
	<i>Melinna cristata</i>	(M. SARS)	
	<i>Ampharete falcata</i>	ELIASON	
	<i>Ampharete baltica</i>	ELIASON	
	<i>Amphicteis gunneri</i>	(M. SARS)	
	Terebellidae	MALMGREN	
	<i>Pista cristata</i>	(O. F. MÜLLER)	
	<i>Eupolymnia</i>	VERRILL	
<i>Thelepus</i>	LEUCKART		
<i>Trichobranchus</i>	MALMGREN		
<i>Terebellides stroemi</i>	M. SARS		
Sabellidae	MALMGREN		
Ostracoda	<i>Philomedes brenda</i>	(BAIRD)	Syn.: <i>P. globosus</i> (Lilljeborg).
	<i>Philomedes lilljeborgi</i>	(G. O. SARS)	
	<i>Conchoecia elegans</i>	G. O. SARS	
	<i>Conchoecia borealis</i>	G. O. SARS	
	<i>Conchoecia obtusata</i>	G. O. SARS	
	<i>Cytherella abyssorum</i>	G. O. SARS	
	<i>Macrocypriis minna</i>	(BAIRD)	
	<i>Macrocypria angusta</i>	(G. O. SARS)	
<i>Eucytheridea punctillata</i>	(BRADY)		

GRUPPE	KOMPONENT	AUTOR	ANMERKNING
	<i>Hemicythere latimarginata</i>	(SPEYER)	
	<i>Echinocythereis echinata</i>	(G. O. SARS)	
	<i>Cythereis dunelmensis</i>	NORMAN	
	<i>Cythereis jonesi</i>	(BAIRD)	
	<i>Elofsonella concinna</i>	(JONES)	
	<i>Cytheropteron alatum</i>	G. O. SARS	
Copepoda	<i>Calanus finmarchicus</i>	(GUNNERUS)	
	<i>Calanus helgolandicus</i>	(CLAUS)	
	<i>Calanus hyperboreus</i>	KRÖYER	
	<i>Pseudocalanus elongatus</i>	BOECK	
	<i>Chiridius armatus</i>	(BOECK)	
	<i>Bradyidius bradyi</i>	(G. O. SARS)	
	<i>Euchaeta norvegica</i>	BOECK	
	<i>Xanthocalanus fallax</i>	G. O. SARS	
	<i>Xanthocalanus propinquus</i>	G. O. SARS	
	<i>Scolecithricella minor</i>	(BRADY)	
	<i>Diaixis hibernica</i>	(A. SCOTT)	
	<i>Parastephos pallidus</i>	G. O. SARS	
	<i>Tharybis macrophthalma</i>	G. O. SARS	
	<i>Centropages typicus</i>	KRÖYER	
	<i>Centropages hamatus</i>	(LILLJEBORG)	
	<i>Temora longicornis</i>	(MÜLLER)	
	<i>Metridia longa</i>	(LUBBOCK)	
	<i>Metridia lucens</i>	BOECK	
	<i>Pleuromamma robusta</i>	(DAHL)	
	<i>Heterorhabdus norvegicus</i>	(BOECK)	
	<i>Scottula inaequicornis</i>	G. O. SARS	
	<i>Anomoclausia indrehusae</i>	GOTTO	
Mysidacea	<i>Boreomysis arctica</i>	(KRÖYER)	
	<i>Erythrops serrata</i>	G. O. SARS	
	<i>Amblyops abbreviata</i>	(M. SARS)	
	<i>Mysideis insignis</i>	G. O. SARS	
	<i>Mysidopsis didelphys</i>	(NORMAN)	
Cumacea	<i>Leucon nasicus</i>	(KRÖYER)	
	<i>Leucon acutirostris</i>	G. O. SARS	
	<i>Eudorella emarginata</i>	(KRÖYER)	
	<i>Eudorella truncatula</i>	BATE	
	<i>Diastylis cornuta</i>	(BOECK)	
	<i>Diastylis lucifera</i>	(KRÖYER)	
	<i>Diastylis tumida</i>	(LILLJEBORG)	
	<i>Diastylis echinata</i>	BATE	
	<i>Diastylodes serrata</i>	(G. O. SARS)	
	<i>Diastylodes biplicata</i>	(G. O. SARS)	
	<i>Leptostylis longimana</i>	(G. O. SARS)	
	<i>Campylaspis rubicunda</i>	(LILLJEBORG)	
	<i>Campylaspis glabra</i>	G. O. SARS	
	<i>Campylaspis costata</i>	G. O. SARS	
Isopoda	<i>Ianira maculosa</i>	LEACH	
	<i>Munna</i>	BOECK	
	<i>Pleurogonium inerme</i>	G. O. SARS	

GRUPPE	KOMPONENT	AUTOR	ANMEKNING
	<i>Macrostylis spinifera</i>	G. O. SARS	
	<i>Ischnosoma bispinosum</i>	G. O. SARS	
	<i>Desmosoma lineare</i>	G. O. SARS	
	<i>Desmosoma laterale</i>	(G. O. SARS)	
	<i>Desmosoma armatum</i>	G. O. SARS	
	<i>Echinopleura aculeata</i>	(G. O. SARS)	
	<i>Munnopsis typica</i>	M. SARS	
	<i>Ilyarachna longicornis</i>	(G. O. SARS)	
	<i>Eurycope cornuta</i>	G. O. SARS	
	<i>Eurycope phalangium</i>	G. O. SARS	
Amphipoda	<i>Themisto abyssorum</i>	(BOECK)	
	Lysianassidae	DANA	
	<i>Tryphosa hörringi</i>	BOECK	
	<i>Tryphosites longipes</i>	(BATE)	
	Phoxocephalidae	G. O. SARS	
	<i>Harpinia pectinata</i>	G. O. SARS	
	<i>Harpinia crenulata</i>	BOECK	
	<i>Ampelisca</i>	KRÖYER	
	<i>Stegocephalus inflatus</i>	KRÖYER	
	<i>Andaniexis abyssi</i>	(BOECK)	
	<i>Andaniopsis nordlandica</i>	(BOECK)	
	Amphilochidae	G. O. SARS	
	<i>Astyra abyssi</i>	BOECK	
	Stenothoidae	STEBBING	
	Oediceridae	G. O. SARS	
	<i>Monoculodes packardii</i>	BOECK	
	<i>Monoculodes tenuirostratis</i>	BOECK	
	<i>Perioculodes longimanus</i>	BATE & WESTWOOD	
	<i>Synchelidium haplocheles</i>	(GRUBE)	= <i>S. brevicarpum</i> i SARS (1895), kfr. STEPHENSEN (1928).
	<i>Synchelidium tenuimanum</i>	NORMAN	= <i>S. haplocheles</i> i SARS (1895), kfr. STEPHENSEN (1928).
	<i>Synchelidium intermedium</i>	G. O. SARS	
	<i>Halicreion longicaudatus</i>	BOECK	
	<i>Westwoodilla caecula</i>	(BATE)	
	<i>Bathymedon longimanus</i>	(BOECK)	
	<i>Bathymedon saussurei</i>	(BOECK)	
	<i>Arrhis phyllonyx</i>	(M. SARS)	
	<i>Bruzelia</i>	BOECK	
	<i>Halice abyssi</i>	BOECK	
	<i>Rhachotropis macropus</i>	G. O. SARS	
	<i>Rhachotropis leucophthalma</i>	G. O. SARS	
	<i>Halirages fulvocinctus</i>	(M. SARS)	
	<i>Apherusa bispinosa</i>	(BATE)	
	Gammaridae	LEACH	
	<i>Melphidippa borealis</i>	BOECK	
	<i>Eriopisa elongata</i>	(BRUZELIUS)	
	<i>Cheirocratus sundewalli</i>	(RATHKE)	
	<i>Cheirocratus intermedius</i>	G. O. SARS	
	Photidae	BOECK	
	<i>Podoceropsis sophiae</i>	BOECK	
	<i>Corophium bonelli</i>	G. O. SARS	(MILNE-EDWARDS ?), kfr. STEPHENSEN (1928).

GRUPPE	KOMPONENT	AUTOR	ANMERKNING
Euphausiacea	<i>Meganyctiphanes norvegica</i>	(M. SARS)	
	<i>Thysanoessa raschi</i>	(M. SARS)	
	<i>Thysanoessa inermis</i>	(KRÖYER)	
Decapoda	<i>Pasiphaea tarda</i>	KRÖYER	
	<i>Spirontocaris lilljeborgi</i>	(DANIELSSEN)	
	<i>Lebbeus polaris</i>	(SABINE)	
	<i>Pandalus borealis</i>	KRÖYER	
	<i>Pandalus propinquus</i>	G. O. SARS	
	<i>Pandalus montagui</i>	LEACH	
	<i>Pandalina profunda</i>	HOLTHUIS	
	<i>Crangon allmani</i>	KINAHAN	
	<i>Pontophilus norvegicus</i>	(M. SARS)	
	<i>Macropipus holsatus</i>	(FABRICIUS)	
Nudibranchia	<i>Coryphella verrucosa</i>	(M. SARS)	
Bivalvia	<i>Pseudamussium septemradiatum</i>	(MÜLLER)	
	<i>Delectopecten vitreus var. abyssorum</i>	(M. SARS)	
	<i>Thyasira</i>	(LEACH) LAMARCK	
	<i>Abra nitida</i>	(MÜLLER)	
Ophiuroidea	<i>Ophiura sarsi</i>	LÜTKEN	
Pisces	<i>Myxine glutinosa</i>	L.	
	<i>Rhinonemus cimbricus</i>	(L.)	

Tekniske data for de aktuelle prøver. Lokalitet: Elle (Im). Dyp: Ca. 195 m.
Data for 1974 og 1975 er basert på FRITZVOLD (1981).

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 23	1953	17/6	11.45	608	0,90	
B 31	"	18/6	13.15	462	0,83	
B 36	"	19/6	11.10	260	0,51	
B 37	"	"	12.30	350	0,67	
Σ B 23-37				1680		
B 62	1959	4/6	12.45	1096	1,32	
B 69	"	5/6	12.10	901	1,27	
Σ B 62-69				1997		
B 80	1960	25/4	12.45	873	0,88	
B 112	1961	13/2	12.00	516	0,76	
B167	1962	10/11	11.25	2177	0,78	
B 178	1963	30/1	10.10	1591	0,86	
B 434	1971	4/11	12.00	1532	0,47	
B 471	1973	5/12	12.40	1063	1,15	
DB 3 - 10 8 prøver	1974	16/12	00.25- 21.25	8434	0,8-1,4 Gj.sn.= 1,12 (S.D.=0,21)	
DB 19 - 27 9 prøver	1975	16-17/6	00.20- 21.40	9700	0,7-1,4 Gj.sn.= 1,17 (S.D.=0,21)	
B 501	1981	10/8	13.00	693	0,30	Overfl.strøm snudde.
B 531	1982	1/7	15.15	1110	1,20	
B 548	1983	19/4	16.15	779	0,84	
B 552	"	26/10	12.10	761	0,85	
B 568	1984	8/8	12.00	1019	1,10	
B 602	1985	10/7	16.45	735	0,80	
B 638	1986	26/6	20.50	893	0,95	
B 655	1987	21/8	16.00	931	1,00	
B 667	1988	25/8	15.10	908	0,89	
B 671	1989	21/8	15.35	1205	1,30	
B 680	1990	16/8	12.00	712	0,74	
B 688	1992	1/9	10.30	1034	1,05	
B 700	1993	25/8	11.05	561	0,60	

Tekniske data for de aktuelle prøver. Lokalitet: Gråøyrenna (Gk). Dyp: Ca. 110 m.
Data for 1971 og 1972 er fra GJERMUNDSEN (1974). Data for mai 1973 er basert på HESTHAGEN & GJERMUNDSEN (1978) og data for desember 1974 og juni 1975 er basert på FRITZVOLD (1981).

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 24	1953	17/6	14.30	469	0,76	
B 25	"	"	15.35	275	0,42	Dyp 108 - 90 m
B 26	"	"	22.00	478	0,74	
B 28	"	18/6	09.15	866	0,98	
B 29	"	"	10.20	366	0,55	
B 30	"	"	11.15	542	0,84	
Σ B 24-30				2796		
B 44	1959	2/6	11.30	617	0,67	
B 45	"	"	13.15	839	1,29	
B 53	"	3/6	12.50	509	0,87	
B 54	"	"	14.00	572	0,93	
Σ B 44-54				2537		

Tekniske data for Gråøyrenna, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 76	1960	22/4	16.30	814	1,30	Nesten intet mudder.
B 107	"	12/12	14.25	715	1,10	
B 117	1961	14/2	13.40	698	1,08	Nesten intet mudder.
B 122	1962	8/1	12.15	610	0,99	
B 129	"	13/2	14.35	1826	1,00	
B 160	"	4/10	17.15	2297	0,82	
B 170	"	17/12	13.40	1926	0,69	
B 177	1963	29/1	12.50	1423	0,77	
B 190	"	4/5	10.55	1444	0,78	
B 417	1970	27/11	15.30	798	0,86	
B 435	1971	5/11	11.40	741	0,60	
B 455	1972	6/9	20.50	716	0,77	
B 456	"	"	21.55	766	0,83	
B 457 - 70 14 prøver	1973	21-22/5	12.40- 22.20	10261	0,4-1,4. Gj.sn.= 0,88 (S.D.=0,26)	
B 472	1973	5/12	14.55	778	0,87	
DB 11 - 18 8 prøver	1974	19/12	00.25- 21.20	7210	0,9-1,3. Gj.sn. = 1,11 (S.D.=0,15)	
DB 28 - 36 9 prøver	1975	19-20/6	00,10- 22.05	9302	0,7-1,7. Gj.sn.= 1,12 (S.D.=0,30)	
B 502	1981	10/8	14.55	1900	1,00	
B 530	1982	1/7	13.45	1100	1,19	Noe av prøven tapt.
B 547	1983	19/4	13.20	1000	1,08	Fanget også på opptur.
B 556	1984	10/1	15.10	960	1,04	
B 569	"	8/8	14.30	893	0,96	
B 603	1985	10/7	18.50	1019	1,10	
B 632	1986	25/6	12.30	673	0,73	
B 651	1987	19/8	18.20	787	0,85	
B 659	1988	23/8	11.10	1100	1,19	For lite sediment
B 672	1989	22/8	14.45	834	0,90	
B 683	1990	17/8	13.05	700	0,76	
B 689	1992	1/9	12.55	555	0,55	
B 699	1993	25/8	10.25	870	0,94	

Tekniske data for de aktuelle prøver. Lokalitet: Langårrenna (Fk). Dyp: 115 - 120 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 546	1983	19/4	12.15	1111	1,20	
B 557	1984	11/1	10.55	791	0,85	
B 571	"	9/8	09.20	736	0,78	Mest på ujevn bunn.

Tekniske data for de aktuelle prøver. Lokalitet: Sprodypet (Fl). Dyp: Ca. 160 - ca. 170 m.
Data for 1971 er fra GJERMUNDSSEN (1974).

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 159	1962	4/10	15.10	661	0,93	Sleden satte seg fast.
B 172	"	18/12	15.10	487	0,44	

Tekniske data for Sprodypet, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 179	1963	30/1	13.00	1147	0,86	
B 186	1963	20/3	11.50	1077	0,78	
B 193	"	6/5	16.55	979	1,02	
B 436	1971	5/11	13.30	705	0,62	
B 473	1973	6/12	14.15	1049	1,17	
B 506	1981	11/8	12.35	1759	0,95	
B 534	1982	2/7	15.20	561	0,91	160 - 125 m dyp.
B 545	1983	19/4	11.10	1204	1,30	
B 558	1984	11/1	13.15	655	0,71	
B 570	"	8/8	16.55	448	0,41	Til dels for nær land.
B 599	1985	9/7	18.50	982	1,10	158 - 149 m dyp.
B 631	1986	25/6	10.50	875	0,94	
B 650	1987	19/8	16.40	523	0,55	144 - 164 m dyp.
B 668	1988	26/8	09.15	1018	1,10	
B 698	1993	24/8	14.00	787	0,85	

Tekniske data for de aktuelle prøver. Lokalitet: Ytre Støadyp (El₂). Dyp: Ca. 145 m

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 544	1983	13/4	15.00	701	0,76	
B 576	1984	10/8	10.15	442	0,48	145 - 107 m. Masse grus og mudder i prøven.

Tekniske data for de aktuelle prøver. Lokalitet: Blåmagen (El₁). Dyp: Ca. 120 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 543	1983	13/4	13.55	656	0,71	
B 577	1984	10/8	11.30	559	0,50	

Tekniske data for de aktuelle prøver. Lokalitet: Svartedypet (Ek). Dyp: Ca. 100 m

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 130	1962	13/2	17.30	1816	0,98	
B 158	"	4/10	12.45	1540	0,91	Lite mudder i prøven.
B 176	1963	29/1	10.00	1098	0,85	
B 432	1971	3/11	15.05	1302	1,28	Lite mudder i prøven.
B 507	1981	11/8	15.05	1990	1,10	90 - 80 m dyp.
B 524	1982	30/6	11.00	926	1,00	
B 541	1983	13/4	11.05	813	0,88	Lite mudder i prøven.
B 553	"	27/10	11.00	796	0,86	Store gravende polychaeter.
B 572	1984	9/8	11.15	810	0,87	
B 596	1985	9/7	14.00	600	0,65	
B 628	1986	24/6	13.05	1205	1,30	
B 641	1987	17/8	14.05	1068	1,15	

Tekniske data for Svartedypet, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 658	1988	22/8	15.20	719	0,78	
B 673	1989	22/8	16.45	906	0,98	
B 690	1992	1/9	14.35	828	0,87	
B 697	1993	24/8	12.35	406	0,44	

Tekniske data for de aktuelle prøver. Lokalitet: Vesthullet (Ej). Dyp: Ca. 105 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 510	1981	12/8	10.40	1183	0,70	Ekstra meget mudder i prøven.
B 525	1982	30/6	12.20	415	0,45	
B 542	1983	13/4	12.10	443	0,48	
B 554	"	27/10	12.25	592	0,64	Grå vannhenter synlig til 14 m dyp. Store gravende polychaeter
B 561	1984	31/1	14.20	950	1,10	Mørkt mudder, løv og gress.
B 573	"	9/8	12.30	800	0,84	Kom inn mot bakken. Rykk i slepewiren. Mørkebrunt mudder.
B 595	1985	9/7	12.50	503	0,5	Stor mengde ascidier.
B 627	1986	24/6	11.50	856	0,84	
B 640	1987	17/8	12.35	687	0,74	
B 657	1988	22/8	12.20	442	0,48	Måtte snu p.g.a. oppgrunning. Ekstra meget mudder i prøv.
B 674	1989	22/8	18.50	979	1,00	Bunnen ikke så bløt som den har vært i senere år.
B 682	1990	17/8	11.35	893	0,96	
B 687	1992	31/8	15.50	898	0,99	Meget kraftig ekko (krill?) fra 70 m til bunn.
B 696	1993	24/8	11.30	400	0,43	Meget mudder, som er mørkere enn ved Steilene.

Tekniske data for de aktuelle prøver. Lokalitet: Steilene (Dk). Dyp: Ca. 100 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 13	1952	14/6	09.05	1074	0,87	
B 124	1962	8/1	19.20	608	0,82	Intet dagslys.
B 133	"	14/2	16.25	1559	1,01	
B 134	"	20/3	11.40	2099	1,06	
B 149	"	3/5	18.15	1498	0,81	
B 151	"	13/7	19.20	1596	0,86	Var litt innpå Morragrunnen, ca. 70 m dyp

Tekniske data for Steilene, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 152	"	21/8	11.20	1559	0,84	
B 157	1962	4/10	10.55	1305	0,70	Litt for langt øst med stein midt i draget.
B 164	"	9/11	11.55	2156	1,09	Lite mudder.
B 173	"	19/12	09.35	1509	0,81	
B 181	1963	31/1	11.50	1632	0,88	
B 184	"	19/3	11.25	1467	0,79	
B 194	"	7/5	09.55	1025	0,71	Satte seg fast i noe som ikke ga ekko.
B 199	"	18/6	10.35	1191	0,74	
B 207	"	10/7	14.10	1166	0,73	Lite mudder.
B 211	"	5/8	11.05	1553	0,84	Litt for langt over mot Morragrunnen.
B 224	"	3/9	16.35	2170	1,17	
B 234	"	27/11	11.40	1340	1,09	
B 243	1964	24/1	13.20	990	0,80	
B 249	"	12/3	11.25	1188	0,96	
B 257	"	27/4	18.25	849	0,92	
B 262	"	3/7	11.35	786	0,82	
B 270	"	1/9	14.55	874	0,94	Ikke helt lukket. Kom opp sidelengs.
B 278	"	12/11	10.55	812	0,88	
B 284	1965	2/2	10.10	587	0,59	Grunnere til slutt.
B 290	"	31/3	15.50	747	0,81	
B 302	"	25/6	16.00	1028	1,15	
B 306	"	27/8	15.25	705	0,76	
B 410	1970	28/4	12.00	930	0,35	Med kamera.
B 412	"	5/5	12.00	638	0,35	Med kamera.
B 421	"	4/12	14.35	759	0,61	
B 424	1971	19/8	15.00	1173	0,81	Med kamera.
B 431	"	2/11	12.10	759	0,63	Har dradd inn i bakken. 20 l mudder!
B 474	1973	6/12	16.40	846	0,90	
B 483	1976	7/1	15.10	825	0,91	
B 511	1981	12/8	13.03	2730	1,61	Kjørte seg til slutt fast i Morragrunnen.
B 523	1982	29/6	14.20	493	0,53	
B 540	1983	7/4	14.10	809	0,87	
B 555	"	27/10	13.55	760	0,82	
B 560	1984	31/1	12.10	771	0,83	Ingen levende dyr å se!
B 574	"	9/8	14.25	1005	1,09	Litt napping i wiren. Dyp 104 - 75 m!
B 594	1985	9/7	11.35	573	0,62	Var borti skråningen. Tallrike ascidier.
B 626	1986	24/6	10.30	847	0,90	
B 639	1987	17/8	11.00	884	0,95	
B 656	1988	22/8	09.55	624	0,67	
B 675	1989	23/8	09.25	741	0,80	
B 681	1990	17/8	10.10	971	0,90	
B 691	1992	4/9	15.50	487	0,48	Var for langt vest, på ujevn bunn for det meste.
B 695	1993	24/8	10.00	456	0,49	

Tekniske data for de aktuelle prøver. Lokalitet: Gåsøyrenna (Cl). Dyp: Ca. 70 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 156	1962	4/10	08.45	1281	0,69	Ujevn og hard bunn.
B 163	"	31/10	11.45	174		Bommen brakk. Ingen lukking.
B 174	"	19/12	13.10	1167	0,64	Tørnet i steinrøys.
B 185	1963	19/3	13.20	1212	0,65	
B 195	"	7/5	11.30	848	0,69	
B 198	"	17/6	17.40	2040	1,10	
B 206	"	10/7	12.30	1455	0,79	
B 216	"	6/8	13.50	1304	0,70	
B 223	"	3/9	15.00	1712	0,97	
B 233	"	27/11	10.35	807	0,87	
B 242	1964	24/1	12.15	475	1,03	Sleden kjørte seg fast.
B 248	"	12/3	10.00	740	0,60	Sleden gikk ujevnt.
B 256	"	27/4	17.15	714	0,72	Bare halvt lukket. Meget grus og stein.
B 263	"	3/7	12.45	567	0,61	
B 269	"	1/9	13.55	567	0,92	
B 277	"	12/11	09.50	857	0,93	
B 283	1965	1/2	15.15	552	0,60	
B 295	"	1/4	14.55	814	0,88	
B 303	"	25/6	18.00	645	0,99	
B 307	"	27/8	16.30	769	0,83	
B 430	1971	1/11	10.30	799	0,63	
B 475	1973	7/12	11.05	1028	1,11	

Tekniske data for de aktuelle prøver. Lokalitet: Lysakerfjorden (Bn). Dyp: Ca. 80 - ca. 70 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 162	1962	31/10	10.20	1229	0,90	
B 175	"	19/12	16.00	1859	1,00	
B 180	1963	31/1	09.30	1641	0,89	
B 183	"	18/3	14.55	1549	0,84	
B 189	"	3/5	15.00	Ca. 1370 m*	*	*) Vind på strømmåleren.
B 197	"	17/6	15.45	1772	1,25	
B 205	"	9/7	10.05	1661	0,90	Sleden sto fast noen sekunder midtveis.
B 219	"	7/8	13.55	1737	0,99	
B 222	"	3/9	13.10	1383	1,09	Grunnere ved lukking
B 232	"	27/11	09.10	1082	0,88	
B 241	1964	24/1	10.55	652	0,53	
B 247	"	10/3	16.05	988	0,80	
B 255	"	27/4	16.00	817	0,88	
B 264	"	3/7	14.05	547	0,59	
B 268	"	1/9	12.50	499	0,81	
B 274	"	10/11	14.50	923	1,00	
B 282	1965	1/2	14.00	491	0,53	
B 294	"	1/4	13.35	976	1,05	
B 299	"	24/6	14.35	886	0,96	
B 311	"	30/8	15.15	902	0,97	

Tekniske data for Lysakerfjorden, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 429	1971	29/10	15.05	1112	0,80	
B 476	1973	7/12	12.20	1032	1,11	Svært mørkt mudder. Død brisling, døde (opake) krill, ingen levende dyr.

Tekniske data for de aktuelle prøver. Lokalitet: Helviktangen (Cp). Dyp: Ca. 85 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 182	1963	18/3	12.55	1381	0,73	
B 240	1964	24/1	09,35	1186	0,96	For lite mudder.
B 246	"	10/3	14.50	1147	0,93	
B 254	"	27/4	14.55	877	0,95	
B 256	"	3/7	15.15	893	0,96	
B 267	"	1/9	11.15	551	0,89	Kom opp åpen.
B 273	"	10/11	13.45	816	0,88	
B 281	1965	1/2	12.50	928	0,94	For lite mudder.
B 293	"	1/4	12.30	1055	1,14	
B 298	"	24/6	13.30	1034	1,12	Fanget noe på nedtur.
B 310	"	30/8	16.15	666	0,72	
B 428	1971	29/10	12.35	710	0,77	Lite mudder.
B 477	1973	7/12	13.45	926	1,00	Svart mudder. Døde (opake) krill. Ingen dyr observert i beve- gelse. Rustne partier på sleden blitt svarte.

Tekniske data for de aktuelle prøver. Lokalitet: Kirkevikbanken (Dp). Dyp: Ca. 85 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 188	1963	3/5	12.25	1481	1,04	80 - 120m dyp. Noe tapt ved vasking.
B 196	"	17/6	13.45	1643	0,89	85 - 120 m dyp.
B 217	"	7/8	11.05	581	0,75	Fanget noe på nedtur.
B 221	"	3/9	11.45	364	0,69	Ujevn bunn. Bratt motbakke på slutten.
B 231	"	26/11	15.45	530	0,82	
B 237	1964	15/1	13.40	193	0,57	Sleden kjørte seg fast.
B 245	"	10/3	13.30	410	0,66	Bratt motbakke på slutten.
B 253	"	27/4	13.50	444	0,85	Motbakke på slutten.
B 261	"	3/7	09.30	282	0,76	
B 266	"	1/9	09.55	354	0,64	Svart mudder.
B 272	"	10/11	12.45	384	0,83	
B 280	1965	1/2	11.40	298	0,88	Ingen lukking. Fanget også på turen opp.
B 292	1965	1/4	11.25	535	0,94	For lite mudder.
B 297	"	24/6	12.25	440	0,71	Fanget noe på nedtur.

Tekniske data for Kirkevikbanken, fortsatt.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 309	1965	30/8	15.00	674	0,84	

B 188 og B 196 ble tatt sønnenfor, de øvrige nordenfor Husanberget.

Tekniske data for de aktuelle prøver. Lokalitet: Svartskog (Ep). Dyp: 150 - 155 m.

Prøve nr.	År	Dato	Tid (ca.)	Trekklengde m	Slepehastighet Knop	Anmerkninger
B 218	1963	7/8	12.10	497	0,80	
B 220	"	3/9	10.35	259	0,52	Lite mudder. Ikke råttent.
B 230	"	26/11	14.10	892	0,85	
B 236	1964	15/1	12.20	1188	0,96	Luktet H ₂ S. Meget løv
B 244	"	10/3	12.10	1089	0,88	
B 252	"	27/4	12.45	378	0,58	
B 271	"	10/11	11.40	848	0,92	
B 279	1965	1/2	10.35	851	0,89	
B 291	"	1/4	10.15	766	0,86	
B 296	"	24/6	11.15	871	0,94	Lite mudder.
B 308	"	30/8	13.40	703	0,76	Mudder svært slimet. Dyrene livløse.
B 427	1971	29/10	11.00	856	0,82	
B 478	1973	7/12	14.55	932	1,01	Wire svart.
B 514	1981	13/8	09.50	494	0,90	Svart mudder.
B 518	1982	28/6	12.15	442	0,50	Suppeaktig, svart, stinkende sediment.
B 535	1983	22/3	11.35	788	0,85	
B 589	1985	8/7	11.15	833	0,90	Sediment suppeaktig, svart og tynnt, men uten lukt av H ₂ S.
B 623	1986	23/6	12.20	790	0,80	
B 645	1987	18/8	13.25	815	0,88	Sediment suppeaktig, svart og tynnt, men uten lukt av H ₂ S. Løv. Levende bittesmå planktoncopepoder.
B 662	1988	24/8	10.35	876	0,89	Sediment suppeaktig, svart og tynnt. Intet synlig livstegn i planktonprøve fra 150 - 103 m dyp.
B 677	1989	23/8	13.30	605	0,67	Et snes levende <i>Oncaea</i> (eller <i>Oithona</i>) i planktonprøve fra 149 - 101 m dyp.
B 678	1990	15/8	11.40	336	0,7	Bunnsledenet fullt av svart suppe. Ikke lukket. Ytterst lite zooplankton i prøve fra 148 - 101 m dyp.

FIGURLISTE

- Fig. 1 Kart over Oslofjorden.
- Fig. 2 Kart over Indre Oslofjord med slededrag inntegnet samt dyp og oksygen-
innhold 1 m over bunnen i januar 1986.
- Fig. 3 Fotografi av bunnsleden med kortfattet beskrivelse.
- Fig. 4 Fotografi av fangst fra drag i dypet ved Torbjørnsskjær, del A.
- Fig. 5 " " " " " " " " " " " " B.
- Fig. 6 " " " " " " " " " Elle.
- Fig. 7 " " " " " " " " " Steilene.
- Fig. 8 " " " " " " " " " Lysakerfjorden.
- Fig. 9 " " " " " " " " " dypet ved Svartskog.
- Fig. 10 " " " " " " " " " ved Elle, Gråøy, Spro og Steilene, desember 1973.
- Fig. 11 " " " " " " " " " i Gåsøyrenna, Lysakerfjorden, Helvikdypet og ved
Svartskog, desember 1973.
- Fig. 12 " " " " " " " " " ved Steilene, juni 1965, oktober 1968, desember 1970
og desember 1973.
- Fig. 13 " " " " " " " " " april 1983, oktober 1983, januar 1984 og
august 1984, med angitt oksygeninnhold nær bunnen.
- Fig. 14 Antall taxa i utvalgte krepsdyrgrupper i prøver fra Elle (Im), Gråøyrenna (Gk),
Svartedypet (Ek), Vesthullet (Ej) og Steilene (Dk) i juni 1953 - april 1983.
Trekk lengder inntegnet.
- Fig. 15 Som Fig. 14 for august 1984 - august 1990.
- Fig. 16 Antall individer/100 m³ av de vanligste rekene ved Steilene ved forskjellige
anledninger fra juni 1952 til august 1993.
- Fig. 17 Samme som Fig. 16 ved Elle (Im), i Gråøyrenna (Gk), ved Spro (Fl), i Svarte-
dypet (Ek), Vesthullet (Ej) og ved Steilene (Dk) fra oktober-november 1962 til
januar 1984.
- Fig. 18A Som Fig. 17 fra august 1984 til august 1993.
- Fig. 18B Som Fig. 18A, men i forminskert målestokk for å få alt med.
- Fig. 19 Antall individer/100 m³ av *Tesserogastria musculosa*, *Bradyidius bradyi*,
Cumacea og Amphipoda samt slepehastighet og muddermengde i prøver fra
Steilene i juni 1952 og januar 1962 - august 1965. NB!: Logaritmisk skala, som
ikke har 0. Ingen funnet er tegnet som 0,60.
- Fig. 20 Individer/100 m³ av de samme dyr som i Fig. 19 ved forskjellige anledninger
fra november 1971 til august 1992.
- Fig. 21 Antall individer/100 m³ av de samme dyr som i Fig. 20 + Ostracoda ved
Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn), ved Helviktangen (Cp)
og ved Svartskog (Ep) i august 1963, juli 1964 og august 1965.
- Fig. 22 Antall individer/100 m³ av *Bradyidius bradyi*, *Tesserogastria musculosa*,
Philomedes brenda (en ostracode), Cumacea og Amphipoda ved Elle (Im),
i Gråøyrenna (Gk), ved Spro (Fl), i Svartedypet (Ek), Vesthullet (Ej), ved
Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn), ved Helviktangen (Cp)
og ved Svartskog (Ep) i oktober-november 1971. For *Philomedes* er intet tegnet
fra Elle da de 35 bunnlevende ostracodene som der ble funnet, ikke ble
identifisert til art. På de øvrige lokalitetene ble ingen ostracoder funnet.
- Fig. 23 Som Fig. 22, for desember 1973.
- Fig. 24 " " " " august 1981.
- Fig. 25 " " " " juni 1982.
- Fig. 26 " " " " april 1983.
- Fig. 27 " " " " august 1984.

- Fig. 28 Som Fig. 22 for juli 1985.
 Fig. 29 " " " " juni 1986.
 Fig. 30 " " " " august 1987.
 Fig. 31 " " " " " 1988.
 Fig. 32 " " " " " 1989.
 Fig. 33 " " " " " 1990.
 Fig. 34 Antall individer/100 m³ av polychaetene *Pholoë minuta*, *Antinoëlla sarsi*, *Nereimyra punctata*, *Polydora ciliata*, *Capitella capitata* og *Gyptis rosea* samt Nudibranchiata i oktober-november 1971 på lokalitetene som navngitt for Fig. 22.
 Fig. 35 Antall individer/100 m³ av polychaetene *Pholoë minuta*, *Ampharete* sp., *Antinoëlla sarsi*, *Chaetozone* sp., Scalibregmidae, Spionidae, *Nereimyra punctata* og *Capitella capitata* på utvalgte lokaliteter fra Elle til Svartskog i desember 1973.
 Fig. 36 Som Fig. 35 i august 1981.
 Fig. 37 " " " i juni 1982.
 Fig. 38 " " " i april 1983.
 Fig. 39 " " " i august 1984.
 Fig. 40 " " " i juli 1985.
 Fig. 41 " " " i juni 1986.
 Fig. 42 " " " i august 1987.
 Fig. 43 " " " i " 1988.
 Fig. 44 " " " i " 1989.
 Fig. 45 " " " i " 1990.
 Fig. 46A Antall taxa samt trekk lengder for prøver fra Elle og Gråøyrenna 1953 - 1990.
 Fig. 46B Antall individer/100 m³ totalt og derav foraminiferer i de samme prøver.
 Fig. 47A Som Fig. 46A for Svartedypet, Vesthullet og Steilene 1973 - 1990.
 Fig. 47B " " " 46B " " " " " " " " " " "
 Fig. 48 Diversitet ved Elle i august 1987, 1988, 1989 og 1990.
 Fig. 49 " i Gråøyrenna i juni 1953, juni 1959, april 1960 og februar 1961.
 Fig. 50 " i " i desember 1973, august 1981, juli 1985 og juni 1986.
 Fig. 51 " i " i august 1987, 1988, 1989 og 1990.
 Fig. 52 " i Svartedypet i august 1981, juni 1982, april 1983 og august 1984.
 Fig. 53 " i " i juli 1985, juni 1986, august 1987, 1988 og 1989.
 Fig. 54 " i Vesthullet i august 1981, juni 1982, april 1983 og august 1984.
 Fig. 55 " i " i juli 1985, juni 1986 og august 1987.
 Fig. 56 " i " i august 1988, 1989 og 1990.
 Fig. 57 " ved Steilene i august 1981, juni 1982 og april 1983.
 Fig. 58 " " " i " 1984, juli 1985, juni 1986 og august 1987.
 Fig. 59 " " " i " 1988, 1989 og 1990.
 Fig. 60 " " Svartskog i juli 1985, juni 1986 og august 1987.
 Fig. 61 Antall individer/100 m³ av *Macrocypriis minna*, *Macrocypria angusta*, *Rhachotropis macropus*, *Ilyarachna longicornis*, *Andaniopsis nordlandica* og *Arrhis phyllonyx* ved Elle 1953 - 1990.
 Fig. 62 Antall individer/100 m³ av *Xanthocalanus fallax* og *Xanthocalanus propinquus* ved Elle og i Gråøyrenna 1953 - 1990.
 Fig. 63 Antall individer/100 m³ av *Eudorella emarginata* og *Erythrope serrata* ved Elle og i Gråøyrenna 1953 - 1990.
 Fig. 64 Antall individer/100 m³ av *Bathymedon longimanus*, Ophiuroidea, *Diastylis lucifera*, *Diastylodes serrata*, *Arrhis phyllonyx* og *Metridia longa* i Gråøyrenna 1953 - 1990.

- Fig. 96 Vindobservasjoner fra Blindern oktober - mai 1960-69.
Fig. 97 " " " " " 1970-79.
Fig. 98 " " " " " 1980-89.
Fig. 99 " " " " " 1990-93.
Fig. 100 Individer/100 m³ av *Tesserogastria musculosa* ved Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn) og ved Helviktangen (Cp) 1962-65.
Fig. 101 Foto av bunnen i ca. 80 m dyp ved Steilene. April 1970.
Fig. 102 " " " i 73 m dyp i Lysakerfjorden. Oktober 1969.
Fig. 103 Bunnsleden med improvisert fotoarrangement på grunt vann. (Florida.)
Fig. 104 Gjenoppbygging av bunnfaunaen ved Steilene i 1962.
Fig. 105 Fotografi av fangst fra drag i Gråsøyrenna, november 1970.

- Fig. 65 Oksygeninnhold i dypvannet ved Spro (Fl), Steilene (Dk), i Lysakerfjorden (Bn), ved Helvik (Cp) og i Bunnefjord-dypet (Dp) 1962-65.
- Fig. 66A Temperaturen i 80 m dyp ved Steilene (Dk) 1973-93.
- Fig. 66B Oksygenkonsentrasjonen i 80 m dyp ved Steilene (Dk) 1973-83.
- Fig. 66C Oksygenkonsentrasjonen i 80 m dyp ved Steilene (Dk) 1973-93.
- Fig. 66D Oksygenekvivalenter i 145 - 150 m dyp ved Svartskog (Ep) 1973-93.
- Fig. 67 Oksygenkonsentrasjonen i 80 m dyp ved Steilene (Dk) 1966-71.
- Fig. 68 Oksygenkonsentrasjonen 1 m over bunnen ved Elle (Im), i Gråøyrenna (Gk), ved Spro (Fl), i Svartedypet (Ek), Vesthullet (Ej), ved Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn), ved Helviktangen (Cp) og Svartskog (Ep) i oktober-november 1971, mars-april 1983, oktober 1983, januar-februar 1984 og august 1984.
- Fig. 69 Oksygenkonsentrasjonen 1 m over bunnen i serier fra Elle til Svartskog i januar 1985, juli 1985, januar 1986, juni 1986 og august 1987.
- Fig. 70 Oksygenkonsentrasjonen 1 m over bunnen i serier fra Elle til Svartskog i august 1988, 1989, 1990, august-september 1992 og august 1993.
- Fig. 71 Opptellingstest. Middelet av 2/10 i prøvene fra Im, Gk, Ek, Ej, Dk og Ep i juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet. Polychaeta.
- Fig. 72 Som Fig. 71. Ostracoda & Chaetognatha.
- Fig. 73 " " " Copepoda & Ophiuroidea.
- Fig. 74 " " " Cumacea & Mollusca.
- Fig. 75 " " " Isopoda & Mysidacea.
- Fig. 76 " " " Amphipoda & Decapoda.
- Fig. 77 - 87: Antall individer/100 m³ av de vanligste faunakomponenter.
- Fig. 77 Ved Elle 1973 - 1990.
- Fig. 78 " " " - "
- Fig. 79 " " " - "
- Fig. 80 I Gråøyrenna 1973 - 1990.
- Fig. 81 I " " - "
- Fig. 82 I Svartedypet 1981 - 1989.
- Fig. 83 I " " - "
- Fig. 84 I Vesthullet 1981 - 1990.
- Fig. 85 I " " - "
- Fig. 86 Ved Steilene 1973 - 1990.
- Fig. 87 " " " - "
- Fig. 88 Slepehastighet, muddermengde i prøven samt ikke-planktoniske taxa.
- Fig. 89 Antall individer/100 m³ av *Diaixis hibernica* ved Steilene, i Gåsøyrenna og i Lysakerfjorden i 1962 - 1965.
- Fig. 90 Antall individer/100 m³ av Nudibranchiata ved Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn), ved Helviktangen (Cp), på Kirkevikbanken (Dp) og ved Svartskog (Ep) i 1962-63.
- Fig. 91 Som Fig. 90 for 1964-65, november 1971 og august 1993.
- Fig. 92 Antall individer/100 m³ av Nudibranchiata ved Elle (Im), i Gråøyrenna (Gk), ved Spro (Fl), i Svartedypet (Ek), i Vesthullet (Ej) og ved Steilene (Dk) 1962-63, 1971, 1973 og 1981-93.
- Fig. 93 Vertikalfordeling av oksygen, nitrat og ammonium i Bunnefjorden og ved Spro i april 1984.
- Fig. 94 Vertikalfordeling av temperatur, saltholdighet og tetthet - som σ_t - ved samme anledninger som i Fig. 93.
- Fig. 95 Ekkogrammer fra Bunnefjorden og Spro-dypet i april 1984 samt fra Spro-dypet og ved Steilene i august 1987.




Fig. 1. Kart over Oslofjorden med dybdekoter. Skrått merke på 59°00'N viser slededrag ved Torbjørnsskjær.




Fig. 2. Forenklet topografisk kart med slededrag og beliggenheten av VEAS kloakkrensning (Ej). Til høyre for lokalitetsnavnene er angitt vannets oksygeninnhold (ml/l) 7.-10 januar 1986 1 m over bunnen i dyp (m) som angitt nedenfor.

bo gl ar se di wi ha




Fig. 3. Beyers bunnslæde. Fig. 3. Beyer's epibenthic closing net.

bo = bolt som holder spannet på plass. *gl* = glider som, når den dras fremover, løfter armen *ar*, som da frigjør spannet, og videre strammer senene *se*, som snurper hovens innvendige kalv sammen. *di* = distansemåler. *wi* = wire som drar glideren fremover når ringen *ri* i hanefoten *ha* er frigjort ved hjelp av slippelodd fra fartøyet eller sprenget løs fra hanefoten hvis sleden har kjørt seg fast.

bo = pin holding the bucket in place. *gl* = sliding mechanism, which, when it is pulled forward causes the lever (*ar*) to pull out the pin and thus release the bucket. At the same time the diaphragm peripherally attached between the cylindrical and conical parts of the net is throttled by the nylon guts (*se*) which are attached to the sliding mechanism. *di* = flow meter. *wi* = wire which pulls the sliding mechanism forward when the ring (*ri*) attached with spun-yarn to the bridle (*ha*) is released by a messenger or torn away from the bridle if the gear has become stuck. The sliding mechanism becomes locked in its foremost position and the wire (*wi*) then released from it. The pull in the towing warp is then transferred to the back end of the frame, whereby the gear is turned. The propeller of the flow meter cannot rotate backward.



Fig. 4. Fotografi av fangst fra drag i dypet ved Torbjørnsskjær (Åo), august 1987; del A. Den øvrige del av samme prøve (del B) er vist i Fig. 5. Dyp = 465 m. Slept distanse = 1240 m. Fotokarets bredde = 15 cm.



Fig. 5. Fotografi av fangst fra drag i dypet ved Torbjørnsskjær (Åo), august 1987; del B. Den øvrige del av samme prøve (del A) er vist i Fig. 4. Dyp = 465 m. Slept distanse = 1240 m. Fotokarets bredde = 15 cm.



Fig. 6. Fotografi av fangst fra drag i dypet ved Elle (Im), august 1988.
Dyp = 200 m. Slept distanse = 908 m. Fotokarets bredde = 15 cm.
Oksygeninnhold 1 m over bunnen: 5,06 ml/l.



Fig. 7. Fotografi av fangst fra drag i dypet ved Steilene (Dk), august 1988.
Dyp = 100 m. Slept distanse = 624 m. Fotokarets bredde = 15 cm.
Oksygeninnhold 1 m over bunnen: 3,10 ml/l.



Fig. 8. Fotografi av fangst fra drag i Lysakerfjorden (Bn), august 1988.
Dyp = ca. 80 m. Slept distanse = 1200 m. Fotokarets bredde = 15 cm.
Oksygeninnhold 1 m over bunnen: 1,52 ml/l.




Fig. 9. Fotografi av fangst fra dypet ved Svartskog (Ep), august 1988.
Dyp = 155 m. Slept distanse = 376 m. Fotokarets bredde = 15 cm.
Oksygeninnhold 1 m over bunnen: 0,00 ml/l.



Fig. 10. Fotografi av fangster fra drag ved Elle (Im), Gråøy (Gk), Spro (Fl) og Steilene (Dk), desember 1973. Glassenes diameter = 11 cm og 10,5 cm. Kfr. Fig. 66B.



Fig. 11. Fotografi av fangster fra Gåsøyrenna (Cl), Lysakerfjorden (Bn), dypet ved Helviktangen (Cp) og ved Svartskog (Ep), desember 1973. Glassenes diameter = 10,5 cm.




Fig. 12. Fotografi av fangster i dypet ved Steilene (Dk), juni 1965, oktober 1968, desember 1970 og desember 1973. Glassenes diameter = 11 cm, 9,5 cm, 12 cm og 10,5 cm.



Fig. 13. Fotografi av fangster fra dypet ved Steilene (dk), april 1983, oktober 1983, januar 1984 og august 1984, med angitt oksygeninnhold i vannet nær bunnen. (Oksygenverdien for oktober 1983 = 0,32 ml/l.) Glassenes diameter = 10,5 cm.




Fig. 14. Antall taxa i krepsdyrgrupper fra Elle (1m) til Steilene (Dk). Copepoda: Bare de bunnlevende Calanoida. * = Ikke analysert. ** = Lite sediment. *** = Ingen prøve. **** = Ufullstendig prøve.




Fig. 15. Antall taxa i krepsdyrgrupper fra Elle (Im) til Steilene (Dk). Copepoda: Bare de bunnlevende Calanoidea. * = Ikke analysert. ** = Lite sediment. *** = Ingen prøve.




Fig. 16. Antall individer/100 m³ av de vanligste rekena ved Steilene (Dk). *** = Ingen prøve. 1966-69 = Minimumsverdier da slepedistansen ikke er godt nok kjent.





Fig. 17. Ind./100 m³ av de vanligste rekena fra Elle (1m) til Steilene (Dk). Verdier utenfor Dk i 1962 og alle i 1971 fra GJERMUNDSEN (1974).
 *** = Ingen prøve. ** = Lite sediment. **** = Ufullstendig prøve.





Fig. 18A. Antall individer/100 m³ av de vanligste rekeartene i observasjonsserier fra Elle (1m) ved Drøbak til Steilene (Dk). ** = Lite sediment. *** = Ingen prøve.





Fig. 18B. Samme som Fig. 18A, men i mindre målestokk for å få alt med.




Fig. 19. Individuer/100 m³ ved Steilene (Dk). 0,60 = Ingen prøve. *** = Ingen funnet. ** = Ingen funnet. * = Ingen funnet. Skala for slepehastighet og muddermengde til høyre. Data fra Beyer og Versvik (1968).




Fig. 20. Antall individer/100 m³ ved Steilene, Dk. 0,60 = Ingen funnet. * = Ikke analysert. *** = Ingen prøve. ** = Lite sediment.




Fig. 21. Antall individer/100 m³ i serier fra Steilene (Dk) til Svartskog (Ep) i Bunnefjorden. 0,60 = Ingen funnet. *** = Ingen funnet. * = Ikke analysert.




Fig. 22. Ind./100 m³ Okt.-Nov. 1971 fra Elle (Im) til Svartskog (Ep). Data fra GJERMUNDSEN (1974).




Fig. 23. Ind./100 m³ Desember 1973 fra Elle (Im) ved Drøbak til Svartskog (Ep) i Bunnefjorden.

Fig. 22 og 23: 0,60 = Ingen funnet. * = Ikke analysert. ** = Lite sediment. *** = Ingen prøve.





Fig. 24. Ind./100 m³ August 1981 fra Elle (Im) til Svartskog. 0,60 = Ingen funnet * = Ikke analysert.





Fig. 25. Ind./100 m³ Juni 1982. 0,60 = Ingen funnet. * = Ikke analysert. **** = Ufullstendig prøve.




Fig. 26. Ind./100 m³ April 1983 fra Elle (Im) til Svartskog (Ep). ** = Lite sediment.




Fig. 27. Ind./100 m³ August 1984. 1 ind. tegnet som 1,5 for å vises. *** = Ingen prøve.

Fig. 26 og Fig. 27: 0,60 = Ingen funnet. * = Ikke analysert.




Fig. 28. Ind./100 m³ Juli 1985 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.




Fig. 29. Ind./100 m³ Juni 1986 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.




Fig. 30. Ind./100 m³ Aug. 1987 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.




Fig. 31. Ind./100 m³ August 1988. 0,60 = Ingen funnet. * = Ikke analysert. ** = Lite sediment.




Fig. 32. Ind./100 m³ Aug. 1989 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert




Fig. 33. Ind./100 m³ Aug. 1990 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert

Fig. 32 og Fig. 33: *** = Ingen prøve.




Fig. 34. Ind./100 m³ Okt.-Nov. 1971 fra Elle (Im) til Svartskog (Ep). Data fra GJERMUNDSEN (1974).




Fig. 35. Ind./100 m³ Desember 1973 fra Elle (Im) ved Drøbak til Svartskog (Ep) i Bunnefjorden.

Fig. 34 og Fig. 35: 0,60 = Ingen funnet. * = Ikke analysert. ** = Lite sediment. *** = Ingen prøve.





Fig. 36. Ind./100 m³ Aug. 1981 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert





Fig. 37. Ind./100 m³ Juni 1982. 0,60 = Ingen funnet. * = Ikke analysert. **** = Ufullstendig prøve.





Fig. 38. Ind./100 m³ April 1983 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. ** = Lite sediment





Fig. 39. Ind./100 m³ Aug. 1984 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. *** = Ingen prøve.

Fig. 38 og Fig. 39: * = Ikke analysert.





Fig. 40. Ind./100 m³ Juli 1985 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.





Fig. 41. Ind./100 m³ Juni 1986 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.





Fig. 42. Ind./100 m3 Aug. 1987. 0,60 = Ingen funnet. Ca. 1 ind./100 m3 tegnet som 1,5 for å vises.





Fig. 43. Ind./100 m3 Aug. 1988 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. ** = Lite sediment.

Fig. 42 og Fig. 43: * = Ikke analysert.





Fig. 44. Ind./100 m3 Aug. 1989 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.





Fig. 45. Ind./100 m3 Aug. 1990 fra Elle (Im) til Svartskog (Ep). 0,60 = Ingen funnet. * = Ikke analysert.

Fig. 44 og Fig. 45: *** = Ingen prøve.




Fig. 46A. Søyler: Antall taxa. Kurve: Trekkklengde.




Fig. 46B. Totalt antall individer/100 m3 og derav Foraminifera.

Fig. 46A og Fig. 46B: * = Ikke analysert. ** = Lite sediment. **** = Ufullstendig prøve.




Fig. 47A. Søyler: Antall taxa. Kurve: Trekklengde.




Fig. 47B. Totalt antall individer/100 m3 og derav Foraminifera.

Fig. 47A og Fig. 47B: ** = Lite sediment. *** = Ingen prøve.




Fig. 48. Diversitet ved Elle i august 1987, august 1988, august 1989 og august 1990.




Fig. 49. Diversitet i Gråøyrenna i juni 1953, juni 1959, april 1960 og februar 1961.




Fig. 50. Diversitet i Gråøyrenna i desember 1973, august 1981, juli 1985 og juni 1986.




Fig. 51. Diversitet i Gråøyrenna i august 1987, august 1988, august 1989 og august 1990.




Fig. 52. Diversitet i Svartedypet i august 1981, juni 1982, april 1983 og august 1984.




Fig. 53. Diversitet i Svartedypet i juli 1985, juni 1986, august 1987, august 1988 og august 1989.




Fig. 54. Diversitet i Vesthullet i august 1981, juni 1982, april 1983 og august 1984.




Fig. 55. Diversitet i Vesthullet i juli 1985, juni 1986 og august 1987.




Fig. 56. Diversitet i Vesthullet i august 1988, august 1989 og august 1990.




Fig. 57. Diversitet ved Steilene i august 1981, juni 1982 og april 1983.




Fig. 58. Diversitet ved Steilene i august 1984, juli 1985, juni 1986 og august 1987.




Fig. 59. Diversitet ved Steilene i august 1988, august 1989 og august 1990.




Fig. 60. Diversitet ved Svartskog i juli 1985, juni 1986 og august 1987.





Fig. 61. Individuer/100 m3 ved Elle (1m). * = Ikke analysert. Nr. 1 og 2 = Ostracoder. Nr. 3, 5 og 6 = Amphipoder. Nr. 4 = Isopode.






Fig. 62. Antall ind./100 m³ av to bentiske copepoder ved Elle (1m) og i Gråøyrenna (Gk). * = Ikke analysert. ** = Mindre god bunnkontakt. **** = Ufullstendig prøve.

 Eudorella emarginata
  Erythrops serrata




Fig. 63. Ind./100 m3. * = Ikke analysert. ** = Mindre god bunnskontakt. **** = Ufullstendig prøve. 1974 og -75 fra FRITZVOLD (198); gjennomsnitt for 8 (1974) og 9 (1975) prøver fra hver lokalitet.





Fig. 64. Individuer/100 m³ i Gråøyrenna (Gk). * = Ikke analysert. ** = Dårlig bunnkontakt. **** = Ufullstendig prøve. Nr. 1 og 5 = Amphipoder. Nr. 3 og 4 = Cumacéer. Nr. 6 = Copepode.




Fig. 65. Oksygeninnhold i vannet nær bunnen ved Spro (Fl), Steilene (Dk), i Lysakerfjorden (Bn), ved Helviktangen og i Bunneford-dypet (Dp) 1962-65. (Fra Beyer & Versvik 1968.)




Fig. 66A. Temperaturen i 80 m dyp ved Steilene (DK) 1973 - 1993. Data fra NIVA.




Fig. 66C. Oxygenkonsentrasjonen i 80 m dyp ved Steilene (DK) 1973 - 1993. Data fra NIVA.




Fig. 66D. Oxygenekvivalenter i dypet ved Svartskog (Ep) 1973 - 1993. Data fra NIVA. Oxygenekvivalent -1 = 2 ml H₂S.




Fig. 66B. Oxygenkonsentrasjonen i 80 m dyp ved Steilene (DK) 1973-83. (Fra Magnusson, Kirkerud, Källqvist & Pedersen 1984.)

Oksygen




Fig. 67. Oksygen i m/l i 80 m dyp ved Steilene, Dk, fra NIVA's arkiv. April 1968: Lysakerfjorden (Bn, 80 m) da observasjon fra Dk mangler.

Oksygen ml/l




Fig. 68. Oksygen i ml/l 1 m over bunnen i serier fra Elle (Im) ved Drøbak til Svartskog (Ep) i Bunnefjorden. H2S = Hydrogensulfid.

Oksygen, ml/l




Fig. 69. Oksygen i ml/l 1 m over bunnen i serier fra Elle (Im) ved Drøbak til Svartskog (Ep) i Bunnefjorden.




Fig. 70. Oksygen i ml/l 1 m over bunnen i serier fra Elle (Im) ved Drøbak til Svartskog (Ep) i Bunnefjorden. H₂S = Hydrogensulfid.

× Polychaeta




Fig. 71. Middelen av 2/10 i prøvene fra Im, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.





Fig. 72. Middelet av 2/10 i prøvene fra lm, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.




Fig. 73. Middelen av 2/10 i prøvene fra lm, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.




Fig. 74. Middel av 2/10 i prøvene fra Im, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.




Fig. 75. Middelen av 2/10 i prøvene fra lm, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.




Fig. 76. Middelet av 2/10 i prøvene fra Im, Gk, Ek, Ej, Dk og Ep, Juli 1985 samt avvik fra dette middelet i 1/10 som % av middelet.





Fig. 77. Individuer/100 m³ ved Elle (1m). 0,60 = Ingen funnet. * = Bare delvis analysert.





Fig. 78. Individuer/100 m³ ved Elle (1m). 0,60 = Ingen funnet. * = Ikke analysert.





Fig. 79. Individuer/100 m³ ved Elle (1m). 0,60 = Ingen funnet. * = Ikke analysert.





Fig. 80. Individuer/100 m³ i Gråøyrenna (Gk). 0,60 = Ingen funnet. * = Bare delvis analysert. ** = Lite sediment. **** = Ufullstendig prøve.





Fig. 81. Individuer/100 m³ i Gråøyrenna (Gk). 0,60 = Ingen funnet. * = Ikke analysert. ** = Lite sediment. **** = Ufullstendig prøve.





Fig. 82. Individuer/100 m³ i Svartedypet (Ek). 0,60 = Ingen funnet. *** = Ingen prøve. ** = Lite sediment.





Fig. 83. Individuer/100 m³ i Svartedypet (Ek). 0,60 = Ingen funnet. *** = Ingen prøve.





Fig. 84. Individuer/100 m³ i Vesthullet (Ej). 0,60 = Ingen funnet. *** = Ingen prøve.





Fig. 85. Individuer/100 m³ i Vesthullet (Ej). 0,60 = Ingen funnet. *** = Ingen prøve.





Fig. 86. Individuer/100 m³ ved Steilene (Dk). 0,60 = Ingen funnet.





Fig. 87. Individuer/100 m³ ved Steielene (Dk). 0,60 = Ingen funnet.




Fig. 88. Slepefastighet, estimert muddermengde samt ikke-planktoniske taxa. * = Ikke analysert. *** = Ufullstendig prøve. **** = Ingen prøve.




Fig. 89. Ind./100 m³ av *Diatixis hibernica*. 0,60 = Ingen funnet. ** = Tallet for Lysakerfjorden noe for lite p. g. a. vind i måleren. Data fra Beyer og Versvik (1968).





Fig. 90. Nudibranchiata/100 m³ i serier fra Steilene (Dk) i Vestfjorden til Svartskog (Ep) i Bunnefjorden. 0,60 = Ingen funnet. Ingen søyle = Ingen prøve.







Fig. 91. Nudibranchiata/100 m³ i serier fra Steilene (Dk) i Vestfjorden til Svartskog (Ep) i Bunnefjorden. 0,60 = Ingen funnet. Ingen søyle = Ingen prøve. 1971 - data fra GJERMUNDSEN (1974).




Feb. Okt. Des. Jan. Mai Nov. Des. Aug. Juni Apr. Aug. Juli Juni Aug. Aug. Aug. Aug.
1962 **1963** **1971** **1973** **1981** **1982** **1983** **1984** **1985** **1986** **1987** **1988** **1989** **1990** **1991** **1992** **1993**
 Fig. 92. Nudibranchiata/100 m³ i serier fra Elle (Im) til Steilene (Dk). 0,60 = Ingen funnet. Ingen søyle = Ingen prøve, eller hvis *, ikke analysert. Data for 1962 og 1963 utenfor Steilene samt 1971-data fra GJERMUNDSEN (1974).



Bunnefjord 26.04.84

Spro 27.04.84

Fig. 93. Vertikalfordeling av oksygen (O₂) samt av nitrat (NO₃⁻) og ammonium (NH₄⁺) angitt ved den mengde nitrogen (N) som de representerer.



Bunnefjord 26.04.1984

Spro 27.04.1984

Fig. 94. Vertikalfordeling av temperatur (T), saltholdighet (S) og tetthet (σ_t).




Fig. 95. Ekkogrammer. A og B er fra Bunnefjorden. A: 26.4.1984, kl. 02¹⁵; B: 27.4.1984, ca. kl. 08⁴⁵. C og D er fra Spro. C: 27.4.1984, ca. kl. 16⁴⁰; D: 19.8.1987, ca. kl. 15⁴⁰. E er fra Steilene 19.8.1987, ca. kl. 12⁰⁰. Alle klokkeslett er norsk sommertid. 120 kHz svingerfrekvens ble benyttet ved alle registreringene unntatt høyre halvdel av B, der 38 kHz ble benyttet.





Fig. 96. Vind målt kl.00, 06, 12, 18 GMT på Blindern 1960-69. Produkt av frekvens og styrke i Beaufort. Data fra Meteorologisk Institutt.





Fig. 97. Vind målt kl. 00, 06, 12, 18 GMT på Blindern 1970-79. Produkt av frekvensprosent og styrke i Beaufort. Data fra Meteorologisk Institutt.





Fig. 98. Vind målt kl. 00, 06, 12, 18 GMT på Blindern 1980-89. Produkt av frekvensprosent og styrke i Beaufort. Data fra Meteorologisk Institutt.




Fig. 99. Vind målt kl. 00, 06, 12, 18 GMT på Blindern 1990-93. Produkt av frekvensprosent og styrke i Beaufort. Data fra Meteorologisk Institutt.




Fig. 100. Individuer/100 m³ ved Steilene (Dk), i Gåsøyrenna (Cl), Lysakerfjorden (Bn) og ved Helviktangen (Cp). Fra Beyer & Versvik 1968.



Fig. 101. Foto tatt med kamera montert på sleden i ca. 80 m dyp nær Storegrunnene ved Steilene (Dk) i april 1970.




Fig. 102. Foto tatt med kamera på sleden i 73 m dyp i Lysakerfjorden (Bn) i oktober 1969.




Fig. 103. Bunnsleden med improvisert fotoarrangement på grunt vann (Florida 1961).



Fig. 104. Totale prøver. Samtlige eksemplarer av *Pontophilus norvegicus* fra hver prøve i eget glass på toppen. De store glassenes diameter = 9,5 cm. Kfr. Fig. 16 og Fig. 65.



Fig. 105. Bunnsledefangst fra Gråøyrenna 27/11 - 1970. Dyp: 114 m. Slept distanse: 798 m. Glassets diameter = 12 cm. Merk forskjellen fra faunaen ved Steilene til samme tid, Fig. 12.

ENGLISH SUMMARY

INTRODUCTION. The subject of the present report is variation with time and locality in the fauna in the immediate vicinity of, on, and in the upper layer of the soft bottom of Oslofjorden, and the relations of these variations to pollution and deepwater exchanges. Since pollution effects on the fjord fauna are assumed to be of interest to many local inhabitants without expert knowledge of marine ecology, the Norwegian text has been written as a rather broad introduction to the subject.

TOPOGRAPHY, HYDROGRAPHY AND BIOCHEMICAL PROCESSES

A most important feature of the Oslofjord is its undulating bottom (Figs 1 & 2). Horizontal water movements are to a varying degree hampered by bottom ridges. The most significant of these ridges is the bar just inside Drøbak nearly blocking the western half of the sound and with a sill depth of 19.5 m in the eastern half.

Vertical stirring of the water masses is seriously hampered by a salinity-related density gradient (Fig. 94), which is significant even when the surface water temperature is at a minimum and its salinity has its maximum in late winter.

As only comparatively small amounts of freshwater are emptied into the fjord parts inside Drøbak estuarine circulation is only weakly developed there. The normal tidal range in the area is only 24 cm.

It follows that the considerable amounts of plant nutrients discharged to the fjord are not effectively diluted and cause intensive blooms of planktonic algae, which are the main cause of a pronounced reduction of light penetration. Because of this reduction, photosynthesis, and thus oxygen production, can hardly take place below the density discontinuity layer.

Decomposition of organic matter settling in the calm basins of the fjord inside Drøbak implies a consumption of oxygen which in the deeps nearly always exceeds the limited supply of oxygen.

Surface water is driven out of the fjord by northerly winds, which can lead to a marked lowering of the water level. As a consequence of the constriction of the fjord near Drøbak a pronounced upwelling of heavier water is then taking place just outside the Drøbak bar. This upwelling water will partly join the inflowing compensation current that will sooner or later be caused by pressure forces. Northerly winds are generally prevailing from October-November to April-May (Figs 96, 97, 98; exceptionally mild winters in the early -nineties, Fig. 99). During this season, inflowing water may, after turbulent mixing over the Drøbak bar still be heavy enough to sink to the bottom of the fjord inside.

During stagnation the density of the deepwater trapped in the inner fjord is gradually reduced by influence from above. This favours deepwater exchange. A preceding stay of the deepwater is, however, not an indispensable condition for its replacement, which is most distinctively revealed by a hump in the curve presenting variation with time of its oxygen content (Fig. 66B).

Exchange of the Bunnefjord deepwater is complicated both because of the long distance and many ridges separating it from the outer fjord, and because the surface water is trapped in the Bunnefjord during northerly winds. Several years in succession may pass without deepwater exchange there (Fig. 66D).

The resulting oxygen deficit makes the environment unsuitable for animal life, and organic matter accumulates on the bottom (Fig. 9). To a smaller and varying extent the same phenomena also occur in other localities of the inner fjord.

Although Gråøyrenna (Gk) is the outmost of our inner fjord localities new deepwater arrives late there and in comparatively small proportion due to the surrounding topography (Fig. 2; Fig. 68, Jan.-Feb. 1984). The oxygen loss is, however,

significantly smaller in Gråøyrenna than further in (Fig. 68, Aug. 84) because Gråøyrenna is farther away from big outlets of domestic sewage.

When and where conditions are acceptable for a versatile fauna, organic matter is both metabolized and cut into smaller parts by animals. This makes it increasingly accessible to bacterial decomposition, which in the presence of oxygen is more efficient than under anoxic conditions. It follows that a development towards a toxic and azoic bottom as well as towards a bottom with a rich and productive fauna is **self-propagating**. This can explain the enormous difference we find in the abundance and variety of living animals, respectively dead organic material, in different localities (Figs 4 - 9; Figs 10 - 11; Figs 12 & 105) as well as the dramatic changes with time observed in one and the same locality (Fig. 12; Fig. 16; Fig. 104).

CHARACTER OF THE BOTTOM. Although the bottom of the Inner Oslofjord basins is classified as a silt-clay sediment, it has a top layer which is extremely loose, has a very high water content and is featured by the presence of organic detritus and other organic matter. In the nineteesixties the thickness of this loose layer was often found to be 0.5 cm - 1 cm where the bottom was passably healthy, but in some places only 1 - 2 mm. With oxygen insufficiency the thickness of the loose layer can increase to 1 dm and more.

Partly in and mostly just below the loose layer we find - if there is a fauna or was one not too long ago - faecal pellets which in some places are so abundant as to completely dominate the structure of the sediment in a layer 1 - 3 inches thick. Most of the pellets have a length of about $\frac{1}{2}$ mm. Below, this layer gradually merges into the far more fine-grained, homogenous and compact clayey substance.

Our field experience indicated that the bottom of all the inner fjord had become much softer in the early nineteeeighties than it used to be.

METHODS. The framework and the plate of the gear used for collecting animals (Fig. 3) should preferably be made of acid resisting steel covered with a dark paint. All the mechanisms are made by Bergen Nautik AS, 50 Isdalstø, Norway.

The cylindrical part of the net is partly covered by a fine-meshed bib, which prevents small animals that have been skimmed off the bottom from falling through the coarse net. We have been using 0.5 mm mesh material in bib, diaphragm, conical part of net and bucket end cover.

The excentric position of the releasing mechanism makes the gear self-righting, and the water directly in front of the net is not disturbed by its presence. Fig. 101 and many other photographs show that the advancing sledge is not preceded by a wave disturbing the loose sediment, a small amount of which is scraped off and slides up along the sloping front of the gear. Occasionally crossing mud waves on the bottom (e. g. formed by otter boards, Fig. 102), the gear cannot work that smoothly.

Suitable length of the towing warp will depend on its weight. If the warp is too short or the towing velocity is too great the front of the gear may be lifted off the bottom and animals closely connected with the latter be missed. The amount of mud in the sample is estimated on board according to a coarse scale. Although most of them are small enough to pass through the meshes of our net the number of Foraminifera caught is used as a rough measure of the gear-bottom contact, and the relation between catches of Foraminifera and other fauna components indicates the connection between the latter and the substrate (Figs 77 - 87).

In order to facilitate comparison of the samples we relate the numbers of specimens found in each sample to the amount of water that would have passed through the net during the respective tow if the flow was completely unobstructed, and we use

100 m³ as a standard. Due to both flow reduction and animal avoidance of the gear all our estimates are smaller than the actual numbers in the field.

Our gear catches a much smaller fraction of the animals which are deeply rooted or hidden in the sediment than of those living detached in and on its uppermost layer and just above it. The reason why we have preferred to use the epibenthic net is that sampling of hyperbenthos too small for commercial trawls long has been a badly neglected enterprise. This is particularly unfortunate because the animals that our gear samples more efficiently than grab and dredge are closely related to fisheries, both directly (shrimp trawling) and as fish food, nutritious as they are, not being protected by thick calcium carbonate shells and not being hidden in the sediment.

Extensive experience gained both by others and by ourselves leave no doubt that variations and differences in our bottom net catches reflect corresponding variations and differences in the fauna. It is of particular value to the present studies that an epibenthic net frequently was used in Oslofjorden by G. O. SARS in last century - when pollution was only trifling.

Prior to microscopic examination a sample must be freed from mud. This is done by having the preserved sample in a jar (like jar No 2 in Fig. 12) the glass lid of which is replaced by a piece of 0.5 mm bolting cloth, and laying this jar on a pair of slightly sloping and slowly rotating rollers. A not too strong jet of tap water is directed towards the cloth.

After the bigger ones had been picked out, small animals were counted in 1/10 or 2/10 (occasionally 3/10) of the samples, which were fractioned by means of the apparatus described by WIBORG (1951). The usefulness of this procedure was tested and found satisfactory, other sources of inaccuracy taken into consideration. Only in two of the 258 cases constituting the basis of Figs 71 - 76 a component was found to be missing from one tenth of a sample when more than 5 specimens (7 and 17) were found in the other tenth.

No time was spent on species identification of Foraminifera and Copepoda Harpacticoida because of the great and varying proportion of these small animals that might have got lost through the meshes.

MATERIAL. Sampling in 1981 through 1993 was done with the intention of studying possible effects on the fauna of the sewage discharged from a new sewage purification plant (VEAS, Fig. 2), which after experimental operation in 1982 was in full work from the summer of 1983. The new sewerage system included transfer of part of the sewage that previously ended in the Bunnefjord and Oslo Harbour regions.

Positions of the sampling localities treated in the present report are shown in Fig. 1 and Fig. 2. All these localities have been, and many still are, used for shrimp (*Pandalus borealis*) trawling. Although the line between chart co-ordinates j and k is crossed by both Svartedypet and Vesthullet we invariably have used Ek to indicate the former and Ej to indicate the latter in order to avoid confusion.

Since we knew from earlier experience that very considerable changes in the local fauna from one year to the other could be caused by merely meteorological differences we have also made use of data previously obtained by means of the same gear from the same area both by others and by ourselves.

Time being a limiting factor the present report is based on analyses of a selected part of the material. In the store there are many irreplaceable samples that can be used to throw light upon special questions and expand our survey regionally. The contents of samples that so far only have been used for studying some selected components are not included in the appendix tables (Del II).

Zooplankton samples have also been taken from many of the localities by means of vertical hauls with a Nansen closing net. (Extensive use of the Nansen net

was made in the fjord sixty years ago, and later it has been used for the sake of comparison.) Zooplankton observations will be the subject of a future report.

At the localities inside Drøbak the oxygen content of the water was observed 1 m and 6 m above the bottom.

THE FAUNA VARIATIONS WITH TIME AND LOCATION that we have found are so great and consistent that inaccuracies and irregularities due to the imperfection of the methods cannot be kept responsible for the main features of these variations.

A sample obtained by means of the epibenthic net from a healthy soft bottom locality, in Norwegian coastal waters at depths at least from 60 to 700 m, is generally dominated by **crustaceans**, as illustrated by the photographs (Figs 4 & 5) of a sample from the mouth of Oslofjorden. This refers to biomass and number of specimens as well as to number of species and larger taxonomic groups. A sample obtained from a locality which is exposed to a heavy load of pollution is - if there are any animals at all - strongly dominated by **polychaetes** and may have a large number of specimens belonging to a small number of species (Fig. 8). With additional strain the numbers of both specimens and species are small, and in the deepest part of Bunnefjorden a complete absence of animals (Fig. 9) has been typical of recent years.

Large numbers, and even the same species, of **mussels** can be obtained from polluted as well as healthy areas (Figs 5, 6 & 7).

The 10 species and larger taxonomic units, excluding Foraminifera, Copepoda Harpacticoida and planktonic species, that were numerically most predominant at various localities are listed in the Tables 8 - 13 (Chapter 8.7). These tables show a gradual change from a dominance of Crustacea by Elle (Tab. 8) to a strong dominance of Polychaeta by Steilene (Tab. 12).

The ratio of these two groups was also subject to variations with time that are of great interest. In 1973 there was a pronounced increase in the number of polychaete taxa both by Elle and in Gråøyrenna (Tabs 8 & 9). The tendency of these polychaetes to reappear on the list was much greater in the Gråøyrenna case, which indicates that the Gråøyrenna bottom had been subject to a stronger influence.

In Svartedypet and Vesthullet (Tabs 10 & 11) the representation of polychaetes increased in 1984, whereas it was more than great already in 1981 by Steilene (Tab. 12). Coinciding with the increase in the numbers of Scalibregmidae and Spionidae in Vesthullet (Ej) and by Steilene (Dk) in 1984 there was an equally marked decrease in the numbers of the less pollution resistant polychaete *Pholoë minuta* (Fig. 39, cf Fig. 38).

A ranking was also undertaken with the planktonic components included, and their resulting representation appears from Tables 14 & 15. Planktonic components were particularly numerous by Elle and in Gråøyrenna in 1953, 1959 and 1960. They were found in much greater concentrations in the vicinity of the bottom than in the water column above, and they were found to be far more abundant along the bottom by Elle than in Gråøyrenna.

It should be noted that Tables 8 - 15 only indicate the relative importance of the various components in each case. Thus, due to the general paucity of the fauna by Steilene in 1981, *Euchaeta norvegica* obtained ranking number 7 (Table 14) with only 9 specimens per 100 m³ while *Tesserogastria musculosa* only obtained ranking number 10 with 265 specimens/100 m³ by Elle (Table 8).

Whereas the total number of taxa in the samples as a rule showed a pronounced correlation to the towed distance by Elle and in Gråøyrenna (Fig. 46A) this was not so at the localities further in (Fig. 47A), which is an indication of a more uniform fauna there. The particularly great variations found in the samples from Elle

(Figs 46A & 46B) are partly due to the absence of ridges that could prevent rapid and extensive supplies of water and animals from outside and probably also due to the fact that because of the wide space we have not towed our gear in exactly the same place each time. The bottom fauna may be influenced by a local variation in current pattern.

In agreement with the far more uniform nature of the pelagic realm than that of the benthic one the number of benthonic species within nearly all taxonomic groups is far greater than the number of planktonic species. As a consequence of this the species diversity curve for a sample obtained by means of an epibenthic net may be misleading if there is a great contribution of planktonic components. As an example can be mentioned the samples taken by Elle in June 1953, in which planktonic components made 76.3 % of the total number of specimens but only 17.2 % of the number of taxa, even though all the planktonic components with one exception (*Calanus finmarchicus* + *C. helgolandicus*) were counted as species whereas benthonic species in many cases were counted jointly. This was particularly the case with the foraminifera, which were all counted as one taxon. The diversity curve for Elle in 1953 (not shown) became lower along the path from 0 to 600 specimens than the SD81 curve in Fig. 52. The diversity curves presented in Fig. 48 are considered to be more suitable to characterize the fauna by Elle as compared to the fauna found at the localities inside the Drøbak bar, cf e. g. Fig. 51.

The scraping of sediment off the bottom can have two opposite effects on the diversity curve. A large number of Foraminifera, as for instance in the August 1990 sample from Elle (Fig. 46B), can - provided that the oxygen conditions in the sediment are satisfactory - be equivalent to a copious sample (Figs 77 & 78) with a high diversity (Fig. 48). A small amount of mud, as in the August 1989 sample from Elle, yielding a small number of Foraminifera (Fig. 46B), can have the same effect (Fig. 48) since the number of specimens can be subject to a stronger reduction than the number of taxa.

In Gråøyrenna highest diversity was found in 1953 (Fig. 49). The low profile of the 1960 curve is due to a strong predominance of *Calanus finmarchicus* and *C. helgolandicus* (counted together), which constituted 70.7 % of the total number of specimens. It was in 1973 that the lowest number of taxa was found by Elle as well as in Gråøyrenna (Fig. 46A), and this is responsible for the lacking rise in the right hand part of the diversity curve (Fig. 50).

The diversity curves from Svartedypet indicate a fauna depreciation **after 1981**. The low profile of the 1983 curve (Fig. 52) can be explained by insufficient gear-bottom contact. But the almost equally low profile of the 1984 curve is one of several symptoms of an impoverished fauna. Later improvements appear from Fig. 53.

The diversity curves from the localities Vesthullet and Steilene show great similarities for the same years, Fig. 54 being similar to Fig. 57, Fig. 55 to Fig. 58 and Fig. 56 to Fig. 59. In Vesthullet the diversity was lowest in 1984, when it was lower than it was by Steilene, and the same was also true of 1985 and 1986. Fig. 56 is indicative of a succeeding improvement of the fauna in Vesthullet. By Steilene a very low diversity was found in 1981. (An even lower diversity was found in the sample from 1973 in which *Calanus* sp. juv. made 78.7 % of the specimens.)

Turning to the fauna components, we see from Figs 14 & 15 an inwards reduction in the number of crustacean species which is quite consistent, with the exception of 1984, 1985 and 1986, when the fauna was no better in Vesthullet (Ej) than it was by Steilene (Dk). The observed species reduction was not due to (accidental) differences in the distance towed.

The difference between Elle and Gråøyrenna, i. e. outside and inside the Drøbak bar, is particularly conspicuous for the groups Isopoda + Tanaidacea and Ostracoda. Only in a few cases can such a difference be explained by the difference in

depth of the two localities. Many of the soft bottom crustacean species that we only have found outside the Drøbak bar, or not found at all, were found by G. O. SARS at 30 to 50 fathoms depth in the vicinity of Oslo (formerly named Christiania) a hundred years ago and more.

As far as the numbers of specimens of the various shrimp species are concerned, Fig. 16 and Figs 17 & 18 show sparse captures, or none at all, from the Vestfjord localities Svartedypet (Ek), Vesthullet (Ej) and Steilene (Dk) in 1971 and 1973 and also during the whole period 1981 - 1986/87. In Gråøyrenna (Gk) many shrimps were still found in 1971, but few in 1973 and in the period 1981 through January 1984. Later there was a remarkable increase in the shrimp fauna, originating in Gråøyrenna. This increase referred in particular to the small *Pandalina profunda* and the 0-group of *Pandalus borealis* (Fig. 16; Figs 18A & 18B), which are not so closely connected to the sediment as the crangonid shrimps. In 1992 and 1993, however, a noteworthy increase of the *Crangon allmanni* population occurred in Gråøyrenna. Further in this species did not seem to find the substrate satisfactory.

Figs 22 - 33 show decreasing numbers inwards of some other crustaceans as well as the epibenthic medusa *Tesserogastria musculosa*. A comparison with Fig. 19, Fig. 21 and Fig. 100 reveals that a dramatic reduction had taken place since the nineteen-sixties from Steilene and inwards, in spite of the fact that the deepwater exchange was even better in the nineteeneighties (Fig. 65; Figs 66B, -C, -D; Figs 68, 69 and 70).

From the Tables 4 and 5 (Ch. 8.4.5) it appears that the mysid *Erythroops serrata*, as well as some amphipods and cumaceans had disappeared more or less completely from the Vestfjord localities Svartedypet, Vesthullet and Steilene by August 1984, and that it took them several years to re-establish a population there, if at all.

Fig. 34 shows that many polychaetes, as opposed to the crustaceans, were found in increasing numbers inwards with maxima somewhere between Steilene (Dk) and Svartskog (Ep). *Pholoë minuta* had, however, a clearly different pattern of distribution. It evidently has a smaller tolerance of pollution than the others.

In connection with the repeated deepwater renewals in Bunnefjorden (Fig. 66D) some polychaetes had settled by Svartskog (Ep) in 1985 and 1986 (Figs 40 & 41). But for the rest there was not much to be found there (Figs 21 - 39; Figs 42 - 45). Samples and underwater video observations from the Helviktangen deep (Cp) give evidence of very miserable conditions even there. The transfer of sewage to VEAS had evidently not brought a sufficient relief to Bunnefjorden.

INDICATOR SPECIES. The most famous of all marine pollution indicators, the polychaete *Capitella capitata*, was the only benthonic animal that was found in considerable number in our innermost locality Svartskog (Ep) in Bunnefjorden. In Vesthullet (Ej) it occurred in considerable numbers in 1983 and 1984 (Figs 38 & 39), and in Svartedypet (Ek) it was only found in 1984.

Scalibregma inflatum (Scalibregmidae) also appeared to be a useful pollution indicator showing a distinct maximum of abundance in Svartedypet, Vesthullet and by Steilene in 1984 (Figs 38, 39 & 40, cf VEAS sewage discharge started in 1982-83).

The fact that nudibranchs, probably practically all of them belonging to the species *Coryphella verrucosa*, were found in greatest numbers (Figs 90 & 91) in Gåsøyrenna (Cl) and Lysakerfjorden (Bn), i. e. rather near Oslo, deluded BEYER (1968) into considering them, together with the polychaete families Spionidae and Hesionidae, characteristic of a heavily polluted bottom. We now think that the occurrence and absence of this nudibranch on the soft bottom is mainly determined by depth and distance from the slope, irrespective of pollution.

The Trachymedusa *Tesserogastria musculosa* seems to be indifferent to light. It normally stands or walks very slowly on the bottom and clings to the substrate if scared (HESTHAGEN 1971). It has never been found in plankton hauls, not even in vertical hauls starting with the weight on a bottom rich in *Tesserogastria*. In accordance with its light weight and its occurrence on the very top of the loose sediment layer *Tesserogastria* is easily caught in our epibenthic net. Being formerly found regularly in large numbers by Steilene (Fig. 100) the absence of *Tesserogastria* from this locality and its variable occurrence even further out in recent years is considered a reliable indication that conditions are not as they ought to be.

The calanoid copepod *Bradyidius bradyi* is also closely connected with the soft bottom. The very considerable and systematic variations of its occurrence and abundance (Table 3) show that this is another excellent indicator of the acceptability of the environment. *Bradyidius* appears to be somewhat more sensitive to pollution than *Tesserogastria*. Its scanty occurrence by Elle is due to its preference for moderate depths.

Since *Erythroops serrata* has been found at much greater depth elsewhere, depth is supposedly not responsible for the general absence of this mysid from Elle (Fig. 63). The great variation in abundance of *Erythroops* in Gråøyrenna (Gk) in 1971 - 1975 (Table 4) agrees with the simultaneous variation in the number of *Bradyidius* (Table 3). Later *Erythroops* has been by far the most successful of the two. A hypothetical explanation of this is that *Erythroops* with its extraordinarily long legs is better qualified to cope with the increasing softness of the substrate, which is also supposed to be the cause of the marked increase in abundance of *Erythroops* in Gråøyrenna and perhaps also of its appearance by Elle in the most recent years (Fig. 63).

Like *Erythroops* the cumacean *Diastylodes serrata* was few in number in Gråøyrenna in the ninetensixties and occurred in very much greater abundance thereafter (Fig. 64). In the nineteeneighties the location of its maximum abundance was even shifted from Gråøyrenna to Elle (Table 5).

A corresponding outwards shifting of maximum abundance is shown by the calanoid copepods *Xanthocalanus fallax* and *X. propinquus* (Fig. 62).

In 1953 and 1959 the amphipod *Arrhis phyllonyx* was found in greater abundance in Gråøyrenna (Fig. 64) than by Elle (Fig. 61). The same was also clearly the case in December 1974 and in June 1975 (FRITZVOLD 1981). In 1981 the ratio was opposite, and later this species has not been found in our samples from any of these two localities.

When the bottom by Steilene in 1962 became repopulated by *Pontophilus norvegicus* (Fig. 16) these were not a new generation that had been brought there as planktonic larvae by inflowing water, but they were one-year-old and older specimens (Fig. 104) which had migrated to the place on their own.

As can be seen from Fig. 17 and Table 2, *Pontophilus norvegicus* was far more abundant than *Crangon allmanni* in the inner fjord prior to 1970, whereas the ratio later has been the other way round. The reason(?) for this change should be sought through experiments. Both these species do well in aquaria.

The coldwater planktonic copepod *Calanus hyperboreus* has in our latitude a pronounced preference for depths greater than those found in the Inner Oslofjord, where it as a rule is only scantily represented. In the depths of the outer fjord it can be found in great abundance. In springtime young copepodites can be found in the upper 20 m outside Drøbak and become transported across the sill. In the inner fjord it then becomes a good indicator of the water transport, as in the year 1986 (Table 14, Ch. 8.7). Several months after the influx it was found in very much greater number in Gråøyrenna than in the other localities of the inner fjord, in spite of the more awkward

access to Gråøyrenna. We interpret this as a result of better possibility for survival in Gråøyrenna, and this is corroborated by its greater frequency of occurrence there.

CAUSES OF THE OBSERVED FAUNA VARIATIONS

The frequent observations in the nineteesixties show a clear correlation between the oxygen content of the water near the bottom (Fig. 65) and the fauna, with a reasonable response delay in the latter (Figs 16, 104, 19 & 100).

We do not have sufficient hydrographic or other data to explain the collapse in the fauna observed by Steilene in 1970 (Figs 16 & 12); but we do know that there was an extremely pronounced oxygen minimum in the density discontinuity layer which caused death of trapped fish. The importance of the distance from Oslo, and maybe also a protective effect of the ridge north of Gråøyrenna, is demonstrated by the rich fauna found in Gråøyrenna (Fig. 105) when it was depreciated by Steilene. It cannot be excluded that some of the animals found in Gråøyrenna were refugees from deserted areas further in. The fact that our observations from 1971 to 1993 show a significantly better fauna in Gråøyrenna than in Svartedypet (Figs 14, 15, 17, 18 & 22 - 33) although the oxygen conditions were hardly much better in the former locality (Figs 68 - 70) also indicates that Svartedypet is exposed to an unfavourable influence from the north which asserted itself also before the sewage was transferred to VEAS.

The reduced states of the fauna of Inner Oslofjord in October-November 1971 and December 1973 (Figs 14, 16, 10 & 11) are in agreement with the oxygen observations (Figs 68 & 66B). It was, however, surprising to find a distinct fauna depreciation in 1973 also by Elle, where a critically low oxygen content could not have occurred. A number of crustacean species had disappeared (Fig. 14) and the same was also true of the planktonic *Eukrohnia hamata*, which was found there on every other occasion. A hypothetic explanation is that sinking particles with a repulsive odour from the out-flowing polluted water had scared the animals away. The disappearance of the animals is thus not supposed to be due to lethality, but to a chemoreptory response (cf COOK & BOYD 1965).

In spite of the fact that, according to Fig. 66B, there was no year with particularly low oxygen content of the deepwater in the period 1976 - 1981 the fauna had still not recovered by 1981 (Fig. 16), and the big deepwater renewals that took place in 1982 and 1984 (Figs 66B & C, Fig. 68) did not bring fauna improvements like those we had in the nineteesixties (Figs 16, 100 & 104). Our explanation is that the increased discharge of sewage and repeated occurrence of oxygen shortage (1971, 1973, 1975) had accumulating effects in the sediment in the form of increased oxygen deficit and a looser consistency. There are reasons to believe that these changes were accompanied by repulsive odour. Moreover, the distances for the animals to migrate back had gradually increased, and the fauna reserves in adjacent areas had decreased.

The lacking recovery of the fauna in 1984 was certainly influenced by the poor oxygen conditions in the autumn of 1983 (Figs 66B & 68). Still there were changes in the fauna in Svartedypet and Vesthullet in 1982 - 1986 that are considered effects of the sewage discharge from VEAS. By means of underwater video an extraordinary abundance of particles were in October 1987 observed suspended in the water in the neighbourhood of the sewage outlet. That particle abundance had become a habitual phenomenon of the area was evident from the very dense population of the suspension feeding *Metridium senile* on the structures connected to the discharge pipe. *Metridium* was previously rare in that area. An abundance of *Ciona intestinalis* was also observed.

There are several indications that the improved sewage treatment and the consequent eutrophication reduction in recent years have introduced some improvements of the soft bottom fauna in Vestfjorden.

On the background of KONIECZNY's survey (1994) it can safely be concluded that the differences and changes in the fauna that we have observed were not caused by heavy metals, PCB, DDT or other organic toxicants in the sediment. Of our localities Elle, Gråøyrenna, Svartedypet, Vesthullet and Steilene it was the last one which had the poorest fauna and Elle by far the richest one. In Table 17 we have arranged an extract showing concentrations found in "our" localities of the toxicants considered to be of greatest concern. We see that the state of the sediment by Steilene was not inferior to that by Elle with respect to these toxicants. The reason is that by far the greater part of the toxicants are precipitated nearer the shore. Gråøyrenna is, however, considered to be in a danger zone because of its short distance from a basin where very large amounts of industrial wastes are discharged.

COMPARISONS WITH OTHER INVESTIGATIONS

Results obtained by HESTHAGEN (1971), GJERMUNDSSEN (1974), HESTHAGEN & GJERMUNDSSEN (1978) and FRITZVOLD (1981) using Beyer's epibenthic net in Oslofjorden fit nicely in between our own observations, and the present study has indeed derived great advantage from those reports.

In accordance with PETERSEN (1915) the fauna in our locality Torbjørnsskjær (Åo, 465 m depth) in the fjord mouth should undoubtedly represent an *Amphilepis* - *Pecten* community. According to THORSON (1968) this is a very poor community. This does not at all agree with our experience (cf Figs 4 & 5). Because of the pronounced increase of diluted pollution the fauna can be assumed to have become richer here than it was in 1914 (PETERSEN 1915, St. Kr. 350 - 410 m), like ROSENBERG & al. (1987) found to be the case in several other localities in the surroundings. The main reason for the discrepancy must, however, be the difference in sampling methods used, as anticipated by PETERSEN.

During their comparative studies of the catching ability of various sampling gear BRATTEGARD & HØISÆTER (1973) found that a large number of the epibenthonic medusa *Tesserogastria* was caught by Beyer's epibenthic net but none with a van Veen grab and some other gear. Nor has *Tesserogastria* been reported from the many samples taken in Oslofjorden by means of a Petersen grab, a Day grab or a dredge. These facts lend strong support to the view held by several authors that the uppermost layer of sediment as well as animals occurring in, on and next to it, are pushed away by the wave preceding a descending grab. The topmost loose sediment and small animals that are connected with it may also easily pass through the meshes of most dredges. These features and the far more limited digging behaviour of our gear are obvious reasons for the very considerable differences we see between our epibenthic net samples and samples obtained by means of grabs and dredges.

The shock wave preceding the Day grab used by ASCHAN & SKULLERUD (1990) seems the most likely reason why the tiny errant polychaete *Pholoë minuta* was extremely rare in their samples from Svartedypet, Vesthullet and Steilene in 1985 where and when we caught it in thousands. It may also be the reason why the very motile polychaete *Antinoëlla sarsi* is not even on the species lists of ASCHAN & SKULLERUD although we found it in all our samples from Vesthullet and nearly all our samples from Svartedypet and Steilene. The varying occurrence of the corpulent and sedentary Scalibregmidae in ASCHAN & SKULLERUD's (1990) material is, on the other hand, in very good agreement with our observations.

Using the same Petersen grab on as far as possible exactly the same spots as BROCH (1936) in 1933-34, STÅLESEN (1964) found that the fauna on the inside of the Drøbak bar in 1962 was clearly more marked by pollution than the fauna on the outside. STÅLESEN's data also indicate increased influence of pollution on both sides of the bar (cf BEYER 1967 a). Many of the sampling spots used by BROCH and

STÅLESEN are located in "our" localities Gråøyrenna and Elle, and STÅLESEN's observations are in very good agreement with our results.

The same can not be said about the results obtained by MIRZA & GRAY (1981), who used a dredge. On the basis of both log-normal distribution and factor analysis they concluded that the areas inside the Drøbak bar and south of the E/F boundary (in the present Fig. 2) as well as the area outside the Drøbak bar were unpolluted. One explanation of this serious discrepancy is probably that BROCH (1936), STÅLESEN (1962) and we have compared the bottoms of the various basins, whereas MIRZA & GRAY (1981) used random sampling. Unfortunately all their sampling stations on both sides of Håøy seem to have become located way up on the slope or on a shoal.

Moreover, the northern part of Vestfjorden was characterized by MIRZA & GRAY (1981) as slightly polluted ("Zone 2") whereas epibenthic samples for a long time have demonstrated a pronounced fauna reduction in this area.

Animals which in accordance with their mode of living and/or morphology cannot or do not leave the place when conditions are deteriorating have also developed a physiological capacity to tolerate reduced qualities of the environment, as for instance low oxygen concentration. As an example be mentioned the thick-shelled mussel *Arctica islandica*, which is too heavy to be collected in our epibenthic net. During an oxygen crisis in Lübecker Bay this was the only surviving macrobenthos species (SCHULZ 1969). For the same species THEEDE (1973) found a LT_{50} of 1400 hours in water with oxygen concentration < 0.15 ml/l and a temperature of 10° C. ROSENBERG & al. (1987) found that *Arctica islandica* was the predominating component in their grab samples from the fjord mouth (77 m depth, 49.5 % of the biomass) as well as by Steilene (100 m depth, 65.5 % of the biomass). Thus recording the local distribution of this species does not seem suitable for distinguishing different states of pollution influence. The same also applies to the small mussel *Thyasira flexuosa*.

In general, animals that cannot avoid being captured by a grab are also less sensitive to pollution than the more motile epibenthic species and hyperbenthos. With some exceptions the former group is dominated by polychaetes and mussels - as far as macrobenthos is concerned - and the latter by crustaceans. Because of these relations epibenthic net samples are by far better suited to demonstrate early pollution effects on the fauna and effects of moderate pollution influence than grab samples.

In 1994 COSTELLO & READ compiled data according to which crustaceans are far more sensitive to sewage sludge than both fish, molluscs and polychaetes over short time periods. The 96 h LC_{50} averages ranged from 20 % wet volume of sludge in seawater for polychaetes to 0.0003 % for shrimp larvae.

RECOMMENDATIONS

Because of the extensive damage that has been done to the soft bottoms of the inner Oslofjord and the poor state that is still found in the innermost parts, and because of the many incalculable pollution sources and the great variability and unpredictability of the meteorological conditions, which can be crucial, it is recommended that the efforts to improve the sewage treatment is continued.

It is also recommended to keep an eye on the development through taking samples of the fauna. Epibenthic net and shrimp trawls are best suited to give information on fauna components that do not have great pollution tolerance.

The existing knowledge represents an excellent basis and give rise to experiments testing the preferences, tolerances and aversion of some indicator species for various qualities of water and sediment including odour and suspended particles. Experimental studies concerning relations between species, i. e. competition and predation are also of great interest.

LIST OF TABLES IN THE TEXT

- Table 1 Numbers of specimens/100 m³ of shrimp species and oxygen concentration in the water 1 m over the bottom. *** = No oxygen record available.
- Table 2 Numeric relations of *Crangon allmanni* to *Pontophilus norvegicus* per 100 m³. **** = Samples on 10 occasions never gave more than 3 specim./100 m³ of any of them.
- Table 3 Numbers of *Bradyidius bradyi*/100 m³. Observations from Steilene and inwards in the nineteesixties are from BEYER & VERSVIK (1968) and from Svartedypet and outwards from GJERMUNDSEN (1974). 1971 observations are from GJERMUNDSEN (1974). The May 1973 figure is from HESTHAGEN & GJERMUNDSEN (1978) and is the average of 14 observations (S. D. = 256). 1974 and 1975 observations are from FRITZVOLD (1981). 1974 figures are averages of 8 samples from each locality (S. D. = 0 and 91), and 1975 figures are averages of 9 samples from each locality (S. D. = 2.6 and 180). * = Not analyzed. **** = Incomplete sample due to technical fault.
- Table 4 Numbers of *Erythrops* and some amphipods per 100 m³ at the various localities. * = Not analyzed. ** = Little sediment. *** = No sample. **** = Incomplete sample due to technical fault. 1971 figures are from GJERMUNDSEN (1974). The May 1973 figure is the average of 14 samples according to observations by HESTHAGEN & GJERMUNDSEN (1978). 1974 and 1975 figures are averages of 8 and 9 samples respectively from each locality according to observations by FRITZVOLD (1981).
- Table 5 Numbers/100 m³ of the most common cumaceans. Same arrangement as in Tab. 4.
- Table 6 Occurrence of maximum and minimum catch by Steilene and the apparent positive and negative or absent (Ingen) correlation of these to the amount of mud in the sample. () = Small effect.
- Table 7 Comparison of bottom fauna by Steilene (Dk) and Svartskog (Ep) when conditions were most favourable in Bunnefjorden. Only some of the components found by Steilene are included. The 1963 and 1965 data taken from BEYER & VERSVIK (1968).
- Table 8 Ranking of the ten most numerous fauna components on each occasion by Elle, planktonic components not included. 1 = most numerous, 2 = next etc.
- Table 9 Ranking of the ten most numerous fauna components on each occasion in Gråøyrenna, planktonic components not included. 1 = most numerous, 2 = next etc.
- Table 10 Ranking of the ten most numerous fauna components on each occasion in Svartedypet, planktonic components not included. 1 = most numerous, 2 = next etc.
- Table 11 Ranking of the ten most numerous fauna components on each occasion in Vesthullet, planktonic components not included. 1 = most numerous, 2 = next etc.
- Table 12 Ranking of the ten most numerous fauna components on each occasion by Steilene, planktonic components not included. 1 = most numerous, 2 = next etc.
- Table 13 Summary survey of the ten most numerous fauna components in each locality, planktonic components not included.
- Table 14 Occurrence and ranking of planktonic animals among the ten most numerous fauna components. 1 = most numerous, 2 = next etc. *** = No sample. * = Not analysed. Numbers in () give distribution groups according to text.
- Table 15 Numbers of planktonic species among the ten most numerous components. *** = No samples. * = Not analysed.
- Table 16 Numbers of specimens/100 m³ of Ophiuroidea. * = Not analysed. ** = Little sediment. *** = No sample. **** = Incomplete sample.
- Table 17 Selected data from KONIECZNY (1994). TTS = Total dry matter. TOC = Total organic carbon. Gjennomsnitt = Average. Rangering = Ranking. Tilstandsklasse = condition class. I = Good. II = Less good. III = Rather poor. IV = Poor. V = Very poor. - = Not measured.
- Table 18 Limit values of condition classes.

LIST OF FIGURES

General: * = Not analyzed. ** = Particularly small quantity of mud in the sample. *** = No sample. **** = Incomplete sample due to imperfect closing. In figures with logarithmic scale 0,60 means 0. (Im), (Dk) etc. = Chart coordinates according to Fig. 1 & Fig. 2.

- Fig. 1 Simplified topographic chart of Oslofjorden.
- Fig. 2 " " " " the inner part of Oslofjorden. Heavy lines indicate towing locations. Whole figures give depths in m, and figures with decimals give oxygen concentration in ml/l 1m above the bottom on January 7-9, 1986. VEAS = sewage treatment plant.
- Fig. 3 Beyer's epibenthic closing net. Photograph and short description.
- Fig. 4 Part of sample taken in August 1987 at 470 m depth W of Torbjørnsskjær (Åo). Towed distance = 1243 m. Width of glass vessel = 15 cm.
- Fig. 5 The remaining part of the same sample.
- Fig. 6 Sample taken in August 1988 by Elle (Im) at 200 m depth. Towed distance = 908 m. Oxygen concentration 1 m above the bottom = 5.06 ml/l.
- Fig. 7 Sample taken in August 1988 by Steilene (Dk) at 100 m depth. Towed distance = 624 m. Oxygen concentration 1 m above the bottom = 3.10 ml/l.
- Fig. 8 Sample taken in August 1988 in Lysakerfjorden (Bn) at about 80 m depth. Towed distance = 1200 m. Oxygen concentration 1 m above the bottom = 1.52 ml/l.
- Fig. 9 Sample taken in August 1988 by Svartskog (Ep) at 155 m depth. Towed distance = 376 m. Oxygen concentration 1 m above the bottom = 0.00 ml/l.
- Fig. 10 Samples taken in December 1973 by Elle, Gråøy (Gk), Spro (Fl) and Steilene. Bottle diameters = 11 cm and 10.5 cm.
- Fig. 11 Samples taken in December 1973 in Gåsøyrenna (Cl), Lysakerfjorden, by Helviktangen (Cp) and Svartskog. Bottle diameters = 10.5 cm.
- Fig. 12 Samples taken by Steilene in June 1965, October 1968, December 1970 and December 1973. Bottle diameters = 11 cm, 9.5 cm, 12 cm and 10.5 cm.
- Fig. 13 Samples taken by Steilene in April and October 1983 and January and August 1984. (Oxygen concentration in October 1983 = 0.32 ml/l.) Bottle diameters = 10.5 cm.
- Fig. 14 Numbers of taxa in selected groups of Crustacea from Elle (Im) to Steilene (Dk), 1953 - April 1983. Copepoda: Only benthic Calanoida. Trekklengde = Towed distance.
- Fig. 15 Same as Fig. 14, August 1984 - 1990.
- Fig. 16 Numbers of specimens/100 m³ of the most common shrimp species by Steilene 1952 - 1993. 1966-69: Minimum figures; towed distance not well enough known.
- Fig. 17 Same as Fig. 16 by Elle, in Gråøyrenna, by Spro, in Svartedypet, Vesthullet and by Steilene 1962 - January 1984.
- Fig. 18A Same as Fig. 17 August 1984 - 1993.
- Fig. 18B " " " " on a reduced scale.
- Fig. 19 Numbers of specimens/100 m³ by Steilene of *Tesserogastria musculosa*, *Bradyidius bradyi*, Cumacea and Amphipoda, scale on the left. Estimated quantities of mud and average towing velocities in knots, scale on the right. 1952 and 1962 - 1965.
- Fig. 20 Numbers/100 m³ by Steilene of same animals as in Fig. 19, 1968 - 1992.
- Fig. 21 Numbers of specimens/100 m³ of *Bradyidius bradyi*, *Tesserogastria musculosa*, Ostracoda and Amphipoda by Steilene (Dk), in Gåsøyrenna (Cl), Lysakerfjorden (Bn), by Helviktangen (Cp) and Svartskog (Ep) in August 1963, July 1964 and August 1965.

- Fig. 22 Numbers of specimens/100 m³ of *Bradyidius bradyi*, *Tesserogastria musculosa*, the ostracod *Philomedes brenda*, Cumacea and Amphipoda by Elle (Im), in Gråøyrenna (Gk), by Spro (Fl), in Svartedypet (Ek), Vesthullet (Ej) and the same localities as in Fig. 21 in October-November 1971. Benthic ostracods found by Elle were not identified to species (no column). In the other samples no ostracods were found.
- Fig. 23 Same as in Fig. 22 in December 1973.
- Fig. 24 " " " " " " August 1981.
- Fig. 25 " " " " " " June 1982.
- Fig. 26 " " " " " " April 1983.
- Fig. 27 " " " " " " August 1984.
- Fig. 28 " " " " " " July 1985.
- Fig. 29 " " " " " " June 1986.
- Fig. 30 " " " " " " August 1987.
- Fig. 31 " " " " " " 1988.
- Fig. 32 " " " " " " 1989.
- Fig. 33 " " " " " " 1990.
- Fig. 34 Numbers of specimens/100 m³ of Nudibranchiata and various polychaetes in October-November 1971 from the same localities as in Fig. 22.
- Fig. 35 Numbers of specimens/100 m³ of various polychaetes in December 1973.
- Fig. 36 Same as Fig. 35 in August 1981.
- Fig. 37 " " " " " " June 1982.
- Fig. 38 " " " " " " April 1983.
- Fig. 39 " " " " " " August 1984.
- Fig. 40 " " " " " " July 1985.
- Fig. 41 " " " " " " June 1986.
- Fig. 42 " " " " " " August 1987.
- Fig. 43 " " " " " " 1988.
- Fig. 44 " " " " " " 1989.
- Fig. 45 " " " " " " 1990.
- Fig. 46A Numbers of taxa (columns) and towed distances (lines) by Elle and in Gråøyrenna 1953 - 1990.
- Fig. 46B Total numbers of specimens and Foraminifera per 100 m³ by Elle and in Gråøyrenna 1953 - 1990.
- Fig. 47A Same as Fig. 46A for Svartedypet, Vesthullet and Steilene 1973 - 1990.
- Fig. 47B Same as Fig. 46B " " " " " " "
- Fig. 48 Diversity by Elle in August 1987, 1988, 1989 and 1990.
- Fig. 49 " in Gråøyrenna in June 1953, June 1959, April 1960 and February 1961.
- Fig. 50 " " " " December 1973, August 1981, July 1985 and June 1986.
- Fig. 51 " " " " August 1987, 1988, 1989 and 1990.
- Fig. 52 " " Svartedypet " " 1981, June 1982, April 1983 and August 1984.
- Fig. 53 " " " " July 1985, June 1986, August 1987, 1988 and 1989.
- Fig. 54 " " Vesthullet in August 1981, June 1982, April 1983 and August 1984.
- Fig. 55 " " " " July 1985, June 1986 and August 1987.
- Fig. 56 " " " " August 1988, 1989 and 1990.
- Fig. 57 " by Steilene in August 1981, June 1982 and April 1983.
- Fig. 58 " " " " " " 1984, July 1985, June 1986 and August 1987.
- Fig. 59 " " " " " " 1988, 1989 and 1990.
- Fig. 60 " " Svartskog in July 1985, June 1986 and August 1987.
- Fig. 61 Numbers of specimens/100 m³ of *Macrocyparis minna*, *Macrocypria angusta*, *Rhacotropis macropus*, *Ilyarachna longicornis*, *Andaniopsis nordlandica* and *Arrhis phyllonyx* by Elle 1953 1990.

- Fig. 62 Numbers of specimens/100 m³ of *Xanthocalanus fallax* and *X. propinquus* by Elle and in Gråøyrenna 1953 - 1990.
- Fig. 63 Numbers of specimens/100 m³ of *Eudorella emarginata* and *Erythrops serrata* by Elle and in Gråøyrenna 1953 - 1990.
- Fig. 64 Numbers of specimens/100 m³ of *Bathymedon longimanus*, Ophiuroidea, *Diastylis lucifera*, *Diastylodes serrata*, *Arrhis phyllonyx* and *Metridia longa* in Gråøyrenna 1953 - 1990.
- Fig. 65 Oxygen concentrations in the deeps by Spro (Fl), Steilene (Dk), in Lysakerfjorden (Bn), by Helviktangen (Cp) and in Bunnefjorden (Dp) 1962-65.
- Fig. 66A Temperature at 80 m depth by Steilene 1973-93.
- Fig. 66B Oxygen concentration at 80 m depth by Steilene 1973-83.
- Fig. 66C " " " " " " " " 1973-93.
- Fig. 66D " equivalents at 125 m depth by Svartskog (Ep) 1973-90.
- Fig. 67 Oxygen concentration at 80 m depth by Steilene 1966-71. April 1968: from 80 m depth in Lysakerfjorden (Bn) instead, since no observation from the Steilene station was available.
- Fig. 68 Oxygen concentration 1 m above the bottom in series from Elle (Im) to Svartskog (Ep) in October-November 1971, March-April 1983, October 1983, January-February 1984 and August 1984.
- Fig. 69 Same as Fig. 68 in January and July 1985, January and June 1986 and August 1987.
- Fig. 70 " " " " " August 1988, 1989, 1990, August-September 1992 and " 1993.
- Fig. 71 Counting test of samples from the localities Im, Gk, Ek, Ej, Dk and Ep in July 1985. Abscissa: Mean of 2/10. Ordinate: Deviation in 1/10 as % of the mean of 2/10. Polychaeta.
- Fig. 72 Same as Fig. 71. Ostracoda & Chaetognatha.
- Fig. 73 " " " " Copepoda & Ophiuroidea.
- Fig. 74 " " " " Cumacea & Mollusca.
- Fig. 75 " " " " Isopoda & Mysidacea.
- Fig. 76 " " " " Amphipoda & Decapoda.
- Fig. 77 - 87: Numbers of specimens/100 m³ of the commonest fauna components.
- Fig. 77 Elle 1973 - 1990.
- Fig. 78 " " - "
- Fig. 79 " " - "
- Fig. 80 Gråøyrenna 1973 - 1990.
- Fig. 81 " " " "
- Fig. 82 Svartedypet 1981 - 1989.
- Fig. 83 " " " "
- Fig. 84 Vesthullet 1981 - 1990.
- Fig. 85 " " " "
- Fig. 86 Steilene 1973 - 1990.
- Fig. 87 " " " "
- Fig. 88 Towing velocity in knots and estimated quantity of mud, left hand scale. Non-planktonic taxa, right hand scale.
- Fig. 89 Numbers of specimens/100 m³ of *Diaixis hibernica* by Steilene (Dk), in Gåsøyrenna (Cl) and Lysakerfjorden (Bn).
- Fig. 90 Numbers of specimens/100 m³ of Nudibranchiata by Steilene (Dk), in Gåsøyrenna (Cl), Lysakerfjorden (Bn), by Helviktangen (Cp), at Kirkevikbanken (Dp) and by Svartskog (Ep) in 1962-63. No column = No sample.
- Fig. 91 Same as in Fig. 90 for 1964-65, November 1971 and August 1993.

- Fig. 92 Numbers of specimens/100 m³ of Nudibranchiata by Elle (Im), in Gråøyrenna (Gk), by Spro (Fl), in Svartedypet (Ek), Vesthullet (Ej) and by Steilene (Dk) in 1962-63, 1971, 1973 and 1981-93. No column = No sample, or, with *, not analyzed.
- Fig. 93 Vertical distribution of oxygen, nitrate and ammonia in Bunnefjorden and by Spro in April 1984.
- Fig. 94 Vertical distribution of temperature, salinity and density - as σ_t - on the same occasions as in Fig. 93.
- Fig. 95 Echograms from Bunnefjorden and the Spro Deep in April 1984 and from the Spro Deep and by Steilene in August 1987.
- Fig. 96-99 Wind recorded at Blindern (Oslo) at 00, 06, 12 and 18 h. G. M. T. plotted as products of frequency and force in Beaufort.
- Fig. 96 October - May and the whole year, 1960-69.
- Fig. 97 As in Fig. 96. 1970-79.
- Fig. 98 " " " " 1980-89.
- Fig. 99 " " " " 1990-93.
- Fig. 100 Numbers of specimens/100 m³ of *Tesserogastria musculosa* by Steilene (Dk), in Gåsøyrenna (Cl), Lysakerfjorden (Bn) and by Helviktangen (Cp) 1962-65.
- Fig. 101 Photograph of the bottom at about 80 m depth by Steilene. April 1970.
- Fig. 102 " " " " " 73 m depth in Lysakerfjorden. October 1969.
- Fig. 103 The epibenthic closing net in shallow water with improvised arrangement for photography. (Florida.)
- Fig. 104 Fauna restoration by Steilene in 1962.
- Fig. 105 Sample taken in Gråøyrenna in November 1970. Depth = 114 m. Towed distance = 798 m. Bottle diameter = 12 cm. Note the difference from the sample taken by Steilene at about the same time and shown in Fig. 12.

NIVA



Norsk institutt for vannforskning

Postboks 173 Kjelsås, 0411 Oslo

Telefon: 22 18 51 00 Fax: 22 18 52 00

ISBN 82-577-2857-8