

Norwegian Pollution
Control Authority

Contractor

SFT Norway

Performing institute

NIVA

Report 623/95

Paris

Convention

Annual report on direct and riverine
inputs to Norwegian coastal waters during
the year 1994

A Principles, results and discussions

B Data report



The National Environmental
Monitoring Programme

Paris Convention

Annual report on direct and riverine inputs to
Norwegian coastal waters during the year 1994

- A Principles, results and discussion
- B Data report

Oslo, October 1995
Project manager: Gjertrud Holtan
Co-workers: Dag Berge
Hans Holtan
Terje Hopen

NIVA - REPORT

Norwegian Institute for Water Research



NIVA

Report No.:	Sub-No.:
O-90001	
Serial No.:	Limited distrib.:
3361	

Main Office	Regional Office, Sørlandet	Regional Office, Østlandet	Regional Office, Vestlandet	Akvaplan-NIVA A/S
P.O. Box 173, Kjelsås N-0411 Oslo Norway Phone (47) 22 18 51 00 Telefax (47) 22 18 52 00	Televeien 1 N-4890 Grimstad Norway Phone (47) 37 04 30 33 Telefax (47) 37 04 45 13	Rute 866 N-2312 Ottestad Norway Phone (47) 62 57 64 00 Telefax (47) 62 57 66 53	Thormøhlensgt 55 N-5008 Bergen Norway Phone (47) 55 32 56 40 Telefax (47) 55 32 88 33	Søndre Tollbugate 3 N-9000 Tromsø Norway Phone (47) 77 68 52 80 Telefax (47) 77 68 05 09

Report Title: Paris Convention Annual report on direct and riverine inputs to Norwegian coastal waters during the year 1994. A. Principles, results and discussion. B. Data report.	Date: 25.10.1995 Printed: NIVA 1995
Author(s): Gjertrud Holtan Dag Berge Hans Holtan Terje Hopen	Topic group: Geographical area: Norway
	Pages: 136 Edition: 175

Client(s): Norwegian Pollution Control Authority (SFT)	Client ref.:
---	--------------

Abstract:
 Riverine inputs of nutrients, selected heavy metals and organic micropollutants to Norwegian coastal waters from 10 main and 145 tributary rivers have been monitored during 1994. The inputs from rivers not monitored as well as direct discharges to marine waters along the coast from Sweden to Russia have been estimated. According to the results, total annual nutrient load to coastal waters from landbased sources, is approximately 4100 tonnes of phosphorus and 98100 tonnes of nitrogen. About 30 per cent of the phosphorus and 60 per cent of the nitrogen are inputs from the monitored rivers and tributaries. Most inputs of heavy metals are low, especially the riverine inputs of Cd, Pb and Hg. A few values of Cd and Pb are below the detection limits of the specific analysis, and about 50 % of the Hg-analyses except for the "Skagerrak-rivers", where more than 60 % of the Hg-values were above the detection limit. Most values of the different congeners of PCB were below the detection limit. The pesticide lindane is detected in most analyses in small amounts. Total load of this compound is estimated to about 89 kg. The largest yields from heavy metals comprise copper and zinc, with input estimates of 237 and 905 tonnes, respectively. Retention in the fjords is not included in the above mentioned values, which in several cases would reduce the actual load to open marine waters considerably.

4 keywords, Norwegian

1. Forurensningstilførsler
2. Norske kystområder
3. Elvetilførsler
4. Direkte tilførsler

4 keywords, English

1. Pollution inputs
2. Norwegian coastal waters
3. Riverine inputs
4. Direct discharges

Project manager

Gjertrud Holtan

Gjertrud Holtan

For the Administration

Dag Berge

Dag Berge

PREFACE

The report presents the data from the 1994 monitoring of waterborne pollutants, both riverine and direct discharges, to the Norwegian coastal waters. The study is part of a joint monitoring programme under the "Paris Convention for the prevention of Marine Pollution from Landbased Sources". The Norwegian contribution is administered by the Norwegian Pollution Control Authority (SFT) which has contracted the Norwegian Institute for Water Research (NIVA) to perform the actual investigations.

The 1994-investigation lasted from January throughout December. This report is the Norwegian part of the 1994 study, divided into two parts:

- A: Principles - Results and Discussion
- B: Data Report.

The Programme Committee has consisted of Dag Rosland (SFT), Dag Berge and Hans Holtan (NIVA). The practical investigation is coordinated, and performed by Gjertrud Holtan (NIVA). The calculations of all data has been performed by Terje Hopen (NIVA). The names of all participants are given in paragraph 5.

We would like to express our gratitude to all participants of the investigation, especially to the local fieldworkers for the collection and transport of the samples. The contact persons at the County Environmental Agencies and at the Municipalities of Oslo and Bærum are acknowledged for continuous support and goodwill. The contact persons at the Norwegian Water Resources and Energy Administration (NVE) and The Norwegian Meteorological Institute (DNMI), Per Lofberg and Stein Kristiansen, are acknowledged for their kind cooperation.

Report A:**TABLE OF CONTENTS**

PREFACE	2
TABLE OF CONTENTS	3
SUMMARY AND CONCLUSIONS	4
1. INTRODUCTION.....	6
2. RIVERSYSTEMS INCLUDED IN THE STUDY.....	7
2.1 General aspects	7
2.2 Riversystems monitored.....	8
2.3 Other riversystems included (tributaries).....	10
3. METHODOLOGY.....	11
3.1 Methodology for assessment of riverine inputs	11
3.2 Parameters monitored and analytical methods	12
3.2.1 Chemical parameters - detection limits and analytical methods	12
3.2.2 Method used to estimate flow rate	13
3.2.3 Calculation of annual load	19
3.3 Methodology for assessment of direct discharges to marine waters	20
3.3.1 Waste water treatment plants/sewage effluents.....	20
3.3.2 Industrial effluents	21
3.3.3 Other inputs.....	21
4. RESULTS AND DISCUSSION.....	22
4.1 Pollutants.....	22
4.2 1993-results and discussion.....	22
4.3 Mean annual runoff (1931-60, 1961-90) and "mean load"	27
4.4 Nutrient retention in fjords.....	27
4.5 Long term concentration data	28
5. REFERENCES.....	30
5.1 Project Personnel.....	30
5.2 Literature.....	31

Report B:

36

SUMMARY AND CONCLUSIONS

In 1988, the Paris Commission decided to launch a comprehensive annual monitoring programme covering inputs of selected pollutants to Convention Waters. The programme was to commence in 1990, and continue the following years (PARCOM, 10/3/2-E).

The purpose is to provide the Commission, in accordance with Article 17(B) of the Paris Convention, with an assessment of the waterborne inputs to Convention waters. Another objective of the programme is to control the fulfillment of The Ministerial Declaration of the North Sea.

In 1987 the ministers of the environment from 9 North Sea states agreed to "take effective national steps in order to reduce nutrient inputs into areas where these inputs are likely, directly or indirectly, to cause pollution, and to achieve a substantial reduction (of the order of 50%) in inputs of phosphorous and nitrogen to these areas between 1985 and 1995".

This Declaration applies for the coastal zone from the Norwegian-Swedish border to Lindesnes (the southern most point of Norway). Further discharges of selected micropollutants to the whole North Sea area are to be reduced by 50-70% depending on the micropollutant in question.

In this report the results (1994) are given for riverine inputs of 10 main rivers and 145 tributaries. Thus the active monitoring programme covers drainage from 75 per cent of the main land areas. For discharges entering directly into marine recipients, i.e. sewage and industrial effluents, estimates are based on numbers from effluent control programmes. Area runoff of Total phosphorus, Total nitrogen, phosphates, nitrates and ammonia from these coastal zones are estimated by use of area specific runoff coefficients.

Greatest emphasis with regard to accuracy has been given to the input estimate of the Skagerrak region, as this is considered the most susceptible part of the North Sea. The Skagerrak reception of Norway's total loads are 22 per cent of the phosphorus and 41 per cent of the nitrogen yield. In this region where 90 per cent of the area is river-monitored, more than 60 per cent of the P- and 75 per cent of the N-loads, are found in the riverine inputs.

According to the results of the 1994 investigation total annual nutrient loads to coastal waters from landbased sources in Norway are approximately 4100 tonnes of phosphorus and 98100 tonnes of nitrogen. Respectively 30 and 60 per cent of the grand total inputs of phosphorus and nitrogen are monitored in the main and tributary rivers. Riverine inputs of metals and micropollutants are low. A few of the concentrations found for Cd and Pb are lower than the detection limit requested from PARCOM. Therefore, two quantities have been estimated: one assuming that the true concentration is zero and the other assuming that the true concentration is the limit of detection. This provides maximum and minimum concentrations between which lies the true estimate. When evaluating inputs these data provide a basis for upper and lower estimates.

Inputs of cadmium are thus measured/calculated to be between 9.0 and 9.3 tonnes, mercury between 355 and 443 kg. The "below detection limit problem" also applies for the inputs of PCBs which are measured to be between 37.1 and 72.8 kg. The pesticide lindane was found in all analyses, but in small amounts. Assumably, lindane contamination in Norwegian rivers is mostly due to long range air pollution. Total load is estimated to about 89 kg. The largest share of heavy metals comprise copper and zinc, with input estimates of 237 and 905 tonnes, of which 79 and 85 % respectively, is river-monitored.

Retention of nutrients and micropollutants in the many threshold fjords of Norway is not included in the above given input figures. Estimates of retention of these substances would presumably reduce the actual input to open marine waters.

For most Norwegian rivers the input to the sea, show large annual variations due to differences in water discharge. In order to use the data as a control of the fulfillment of the Ministerial Declaration of the North Sea, the chemical data from 1994 are in addition "normalized", i.e. 1994 chemical concentrations in river water have been multiplied with normal annual runoff (LTA) in the period 1961-90 (main rivers) and 1931-60 (tributary rivers).

1. INTRODUCTION

At the eighth meeting of the Paris Commission (Madrid, June 1986) it was decided to carry out a pilot project to test methods for estimating transport of pollutants from rivers to marine areas. The Norwegian part of the pilot study comprised the two rivers Glomma and Skienselva (Fig. 1). The project was carried out from August 1986 till August 1987, and reported in October 1987 (Lingsten, 1987).

At the Tenth Meeting of the Paris Commission (Lisbon, June 1988) the principles for the comprehensive study on riverine inputs were accepted. It was then decided to commence the study with measurements carried out in 1990, and continue the work in the following years (PARCOM, 10/3/2-E).

The purpose is to provide the Commission, in accordance with Article 17(B) of the Paris Convention, with an assessment of the waterborne inputs to Convention waters. Besides riverine inputs, the information sought also relates to direct discharges.

The objectives of this study are the following:

- To give a quantitative assessment, as accurately as possible, of all riverborne and direct inputs of selected pollutants to Convention waters on an annual basis;
- to report these data annually to the Paris Commission and review them periodically with regard to determining trends;
- for each country, to aim at monitoring on a regular basis 90% of the inputs of each selected pollutant;
- to control that the objectives of the Ministerial Declaration* for reducing the loads of heavy metals, organic micropollutants and nutrients to the North Sea in an order of 50 per cent, between 1985 and 1995, are fulfilled.

The study is to be completed for each calendar year and submitted to PARCOM by the autumn following the year to which the data relate.

*

In 1987 the ministers of the environment from 9 North Sea states agreed to "take effective national steps in order to reduce nutrient inputs into areas where these inputs are likely, directly or indirectly, to cause pollution, and to achieve a substantial reduction (of the order of 50%) in inputs of phosphorous and nitrogen to these areas between 1985 and 1995".

This Declaration applies for the coastal zone from the Norwegian-Swedish border to Lindesnes (the southern most point of Norway). Further discharges of selected micropollutants to the whole North Sea area are to be reduced by 50-70% depending on the micropollutant in question.

2. RIVERSYSTEMS INCLUDED IN THE STUDY

2.1 General aspects

The length of the Norwegian mainland coast line including fjords and bays is 21347 km, and the length of the islands' coast line is 35662 km. (Table 1). Because of the length of the coast line, the great numbers of rivers, and retention processes in the fjords, in- and out-washing areas, monitoring of riverborne pollutants in Norway faces quite a few problems with respect to assessing their impact on coastal waters. Further, to measure 90 % of the load from the Norwegian rivers, a great number of rivers would have to be included, which would be extremely expensive. It was therefore decided that 8 of the major load bearing rivers should be monitored in accordance with the objectives of the comprehensive study. Further it was decided that 2 "unpolluted" rivers should be monitored at a reduced, but appropriate frequency. In these 10 rivers a number of investigations have been carried out during many years, and they have all been included in the National Monitoring Programme of Watercourses (SFT, 1980 - 1994).

These investigations have mainly concentrated on nutrients. Hence data on the load of the nutrients are satisfactory, while the data on heavy metals and organic micropollutants are rather insufficient. In addition to the ten rivers it was decided to estimate the load of 145 other rivers (tributaries) based on other 1994-monitoring programmes, and existing knowledge of the river systems concerned, supplemented with random samples taken in 1994.

The total drainage area of these monitored rivers is 229152 km², while the total area of mainland Norway is 323878 km² (Table 1). Totally 306747 km² of the drainage area is included in the investigation, of which 75 per cent is river monitored (Tables 4 and 5). It was of special importance to estimate the major loads to Skagerrak. In this region the monitored rivers and tributaries cover 94 per cent of the total area, whereof the main rivers alone 80 per cent.

The coastline is divided into subareas/-regions, comprising the drainage basins of the ten main rivers with tributaries, as shown in Fig. 1. Each subarea relates to one of the four discharge-/sub-regions (Fig. 1 and I.I-I.IV (Appendix I, Report B)) which are divided as follows (see also Table 4):

Barents Sea : From the Russian border (about 70°30'N, 70°30'E), to about 70°30'N, 21°E

Norwegian Sea: Southwards of 70°30'N, to about 62° N

North Sea : Southwards of 62° N, to the Swedish border

- Skagerrak : From Lindesnes (the southernmost point of Norway), about 57°44'N to Sweden about 58°58'N, 11°E.

Some key information about Norway and the adjacent oceans is given in Table 1.

Table 1. Norway: Population, areas and length of coastline.

<u>Population</u>	4.2 million
<u>Area:</u>	
- Mainland Norway	323878 km ²
- The whole country incl. Svalbard and Jan Mayen	386958 km ²
<u>Coastline:</u>	
- Length of the continental coastline	21347 km
- Not including fjords and bays	2650 km
- Length of the islands' coastline	35662 km

2.2 Riversystems monitored

The rivers chosen for the comprehensive study are the same as in 1990-1993 and presented in Table 2 and Fig. 1. The rivers marked 1 to 5, represent the major load bearing rivers in Norway. As mentioned in chapter 2.1, it was of special importance to estimate the major loads to Skagerrak (Fig. I.I, Appendix I, Report B) which is an important part of the North Sea, and considered to be most susceptible to pollutions. The five rivers (No 1 to 5) drain into the Skagerrak area. River Suldalslågen (No 7) and river Alta (No 10) are "unpolluted" riversystems where actual measurements have been carried out at a reduced frequency.

Orreelva (No 6) is draining the most intensive agriculture area in Norway especially concerning domestic animals (milk and meat production). Discharges from manure stores and silos together with area runoff from heavily manured fields are causing great problems (eutrophication, including toxic algal blooms) both in Orreelva and in the other water bodies in this area.

Orkla (No 8) is also draining agriculture area, but farming in this part of the country is rather extensive compared to the Orre area. More important in this area is the abandoned mines even if situated in the upper part of the watercourse. Several other rivers in this area also receive waste -water from abandoned mines (heavy metals).

Vefsna (No 9) is a watercourse where the runoff in periods is carrying quite a lot of suspended solids and as such is comparable with other rivers in this part of the country.

Fig. 1.

Norway. Main rivers and tributaries with drainage basins, the whole country divided into four discharge regions.

Source: Norwegian water resources and energy administration.
Design: NIVA.




Table 2. The main rivers with catchment areas and long term average flow (LTA) 1961-90.

No	River	Catchment area, km ²	LTA 1000 m ³ /day
		(at outlet)	(at outlet)
1	Glomma	41.918	61350
2	Drammenselva	17.034	28850
3	Numedalslågen	5.577	10200
4	Skienselva	10.772	23535
5	Otra	3.738	12870
6	Orreelva	105	335
8	Orkla	3.053	5710
9	Vefsna	4.122	15655
7	Suldalslågen	1.457	7420
10	Alta	7.373	7495
Total		95.149	

The ten water courses are all representing typical river systems in different parts of the country. As such they are very useful when estimating loads of comparable rivers, i.e corrections and adjustments in the estimates of load of tributaries, which are based on fewer data than the main rivers.

All these watercourses except Orreelva are regulated for hydroelectric power production.

2.3 Other riversystems included (tributaries)

In addition to the ten main rivers, it was determined to assess inputs from the same 145 river systems as in 1990 - 1993 (Fig. 1) using "best estimates" of concentrations and flows. In total all Norwegian rivers with catchment areas larger than 500 km², and several of the minor rivers (streams) also are included in the 1994 study. Some information about these rivers are shown in Tables 8.1-8.10 (Appendix VIII, Report B).

3. METHODOLOGY

3.1 Methodology for assessment of riverine inputs

In carrying out the Survey, the methodology described in the Commissions Document "Principles of the Comprehensive Study on Riverine Inputs" (1988 and 1993) and in the 1990-Report from Norway (Holtan et al., 1991*), was followed.

*In this document hereafter referred to as "The 1990-Report".

As for "Site selection" we refer to the abovementioned documents, but have chosen to repeat most of the text concerning "Sampling strategy and frequency", only with necessary adjustments.

Compared with the years 1990-1992 the programme was reduced in 1993, and also in 1994.

In all main rivers, except Suldalslågen and Alta it has been taken 12 samples at regular monthly intervals during the sampling period from January till December 1994 as prescribed in PARCOM 10/3/2/E.

For the "unpolluted" rivers Suldalslågen (No 7) and Alta (No 10), where, on the basis of existing knowledge, the concentration levels are very low, the requirement of 12 data sets per annum was found too stringent. These rivers were therefore sampled 4 times per annum. This sampling strategy should be sufficient enough to obtain a reliable estimate of the pollution load for these two rivers.

For the other rivers (tributaries), the concentrations are partly based on samples taken at the "standard" frequency (12, i.e. monthly, or more data sets per annum), which is the case in most rivers in the Skagerrak region. As for most rivers draining to the Norwegian Sea and the Barents Sea, the concentrations are based on measurements of samples taken at random (at least once) and compared with measurements from the last decade. With regard to the rest of the rivers, about 10 in the Orkla area, and nearly all draining to the Orre- and Suldalslågen areas, most data are from samples gathered in 1993.

For all main rivers except Suldalslågen and Alta the parameters lindane and PCBs have been monitored three times in 1994, in Suldalslågen and Alta, twice. For the other rivers these parameters and also Hg have been analyzed once in samples from the tributaries draining to the Skagerrak and up north from river Børselva in the Orkla area to river Grense Jacobselv in the Alta area (Appendix VIII, Report B). For the rest of the rivers the concentrations of lindane, PCBs and Hg are estimated on the basis of knowledge about the activity in the different drainage areas, the findings from the main rivers and samples/analyses from these areas in 1994 (paragraph 3.2.1).

The sample frequency for the main rivers is shown in Table 3.

Table 3. Sampling sites and frequency of the main rivers.

River/Location	J	F	M	A	M	J	J	A	S	O	N	D
Glomma at Sarpsfoss	x	x	x	x	x	x	x	x	x	x	x	x
Drammenselva upstr. outl.	x	x	x	x	x	x	x	x	x	x	x	x
Numedalslågen at Bommestad	x	x	x	x	x	x	x	x	x	x	x	x
Skienselva at Klosterf.	x	x	x	x	x	x	x	x	x	x	x	x
Otra upstream outlet	x	x	x	x	x	x	x	x	x	x	x	x
Orre upstream outlet	x	x	x	x	x	x	x	x	x	x	x	x
Orkla at Vormstad	x	x	x	x	x	x	x	x	x	x	x	x
Vefsna upstream Mosjøen	x	x	x	x	x	x	x	x	x	x	x	x
Suldalslågen upstream outlet		x				x		x		x		x
Alta upstream Alta			x			x		x		x		x

In 1994 the water samples were taken by local persons as in 1990 - 1993. The persons were carefully instructed in advance. The samples were sent to the laboratory at NIVA immediately after sampling, usually arriving a NIVA within 24 to 36 hours later.

3.2 Parameters monitored and analytical methods

3.2.1 Chemical parameters - detection limits and analytical methods

In 1994 the following parameters were monitored in accordance with the mandate: 4 nutrients (total phosphorus, orthophosphates, total nitrogen, ammonia and nitrates), 5 metals (copper, zinc, cadmium, lead and mercury), 1 pesticide (lindane) and a general parameter (suspended particulate matter, S.P.M.). PCBs were to be monitored on a voluntary basis for the 7 congeners (IUPAC numbers 28, 52, 101, 118, 138, 153 and 180).

As detailed information on methodology and obtainable limits of detection for all measured parameters were given in the 1990-Report, only new or improved methods will be described in this report.

As informed in the 1990-Report, the detection limits of the parameters Cd and Pb at the NIVA laboratory were above those requested from PARCOM. All Cd-, Pb- and also Cu- and Zn-samples from 1992 therefore have been analyzed at the Norwegian Institute for Air Research (NILU) where metal determinations are performed on an ICP-MS-instrument (NILU, 1990). On this instrument the recommended detection limits from Parcom (Cd: 10 ng/l, Pb: 0.1 µg/l) are obtainable.

According to PARCOM (10/3/2-E) it was necessary to choose an analytical method which would give at least 70 % of positive findings (i.e. above the detection limit).

As for Cd when determined on ICP-MS, more than 70 % of the findings in samples from the main rivers were positive (from 75 to 100%) and correspondingly in all main rivers for Pb (in 8 rivers 100%). As for the tributaries more than 70 % were above the detection limits both in the Cd- and Pb-samples.

From 1993 the limit of detection has been lowered from 2-1 ng/l (mercury) and from 0.05-0.03 ng/l (PCBs). This is a result of refinement and optimization of the methods. As for mercury more than 70% of the findings in the 1994-samples from three of the main rivers were above the detection limit, Glomma and Numedalslågen (83%) and Alta (75%). As for the findings from tributary rivers in the

Skagerrak area 17 of 25 river samples were higher than the detection limit (68%). As for the other main and tributary rivers, although better, we still had problems to obtain representative values for mercury, which during most of the investigation period were below the detection limit. This was also the case with PCBs. For these parameters most of the measured concentrations were extremely low, and certainly below "PARCOM-detection limits" (Appendix VII - VIII, Report B).

However, we assume that these difficulties do not affect the main results and conclusions of the 1994-study. In those cases where the results recorded were less than the limits of detection, two load quantities have been estimated, one assuming that the true concentration is zero and the other assuming that the true concentration is the limit of detection. This provides maximum and minimum concentrations within which the true estimate will fall. When used to evaluate inputs these data provide upper and lower boundaries for the estimate.

Occasionally on a voluntary basis, the metals arsenic, total chromium and nickel have been determined in some of the samples from the main and tributary rivers and are stipulated for others (Report B). These parameters also were determined on ICP-MS at NILU.

3.2.2 Method used to estimate flow rate

For the period 1931-60 the annual specific runoff from the total area of Norway is estimated at 42.9 l/s km². Expressed in volumetric units this amounts to 438 km³ water, which distributed over the whole country equals a mean runoff of 1350 mm. Mean annual runoff in Norway and from the sub-regions to the main surrounding seas for the period 1931-60 are shown in Table 4. For the main rivers mean annual runoff for the last LTA-period (1961-90) have been estimated. These values are preliminary and will be adjusted when edited and published by NVE, probably next year. For the main rivers mean annual runoff (1931-60 and 1961-90) together with annual runoff for the years 1985, 1990-1994 are shown in Fig. 2, mean annual and annual precipitation for the same stations and periods in Fig. 3. As for precipitation, normals for Norway based on the LTA-period 1961-90 were published in 1993 (DNMI, 1993).





Fig. 2 Main Rivers. Mean Annual Runoff (1931-60 and 1961-90) and Annual Runoff for the Years 1985, 1990 - 1994 (mill.cbm.).

Source: Norwegian Water Resources and Energy Administration



S = South E = East W = West N = North

Fig. 3 Main Rivers. Mean Annual Precipitation (1931-60 and 1961-90) at Stations near Outlet and Annual Precipitation in the Years 1985, 1990 - 1994 (mm/year).

Source: The Norwegian Meteorological Institute

Table 4. Mean annual runoff (1931-60) from the subregions to the main surrounding seas.
 (Fig. 1 and Appendix I, Report B).

Subregions	Main Seas	Drainage area, km ²	Runoff, mill. m ³
The Swedish border - Lindesnes	Skagerrak	98699	57934.47
Lindesnes - Stad	North Sea	138902	164875.88
Stad - the border of Finnmark	Norwegian Sea	94704	231928.67
Finnmark - the border of Sovjet	Barents Sea	73141	41462.90
Total		306747*	438267.45

* The difference between the total area and the area given in Table 1 is due to rivers which drain into the neighbouring countries (Sweden).

At a given location the runoff will change from one year to another and throughout the year. In natural river basins the seasonal variations will depend mostly on the distance from the coast, the altitude and the latitude, and on variations in precipitation. The mean discharge is determined both by the precipitation and the catchment area. Along the coast of Southern and Western Norway the summer low flows are usually dominant together with high runoff in autumn and winter. Thus although Western Norway has much more precipitation than Eastern Norway, its smaller catchment areas lead to much less absolute discharge in western rivers. In the central part of Southern Norway and in the Northern part of the country low water flows are typical both in summer and winter, whereas periods of higher runoff will appear during the snow melting period (spring and early summer). In late summer and in autumn the flow depends on the precipitation and may therefore vary considerably.

In all main rivers continuous observations of the rate of flow are collected. For most rivers these stations are located upstream the sampling stations (NVE, 1995). The additional water supplied is estimated using measured rainfall data from the local catchment areas (DNMI, 1995).

With regard to the river Orkla the runoff station was changed from Vormstad to Syrstad in 1993 as we were recommended by NVE to do so. According to NVE this station is more reliable than Vormstad.

For all main rivers, seasonal changes in runoff in the period 1961- 90, together with mean runoff in 1994, are shown in Fig. 4. In Fig. 5 monthly precipitation for the same period together with mean precipitation in 1994, are shown.

For the other rivers (tributaries) the runoff data partly are from continuous observations as the case is for most rivers in the Skagerrak area, and also for many of the rivers draining to the remaining North Sea. For the rest of the rivers mean runoff data (1931-60) and measured rainfall data (1994) are used for flow estimates.





Fig. 4 Seasonal Changes in Daily Runoff (m³/s).

Source: Norwegian Water Resources and Energy Administration



S = South E = East W = West N = North

Fig. 5 Monthly Changes in Precipitation (mm/month).

(m.o.h. = meters above sea level)

Source: The Norwegian Meteorological Institute

3.2.3 Calculation of annual load

The first of the 2 formulas given in the Paris Commission document and the 1990-Report (Holtan et al., 1991) was used for calculating loads for all main rivers and most of the larger rivers. The second formula was used where continuous records were not available.

For the other rivers, which have been monitored only once, the best available estimates of flow (catchment area multiplied by specific runoff adjusted for deviations from normal precipitation) and flow-weighted concentrations have been used to estimate contaminant loads.

Drainage basins to the different areas and regions (km^2 and per cent) is shown in Table 5.

Table 5. Drainage areas of monitored main and tributary rivers and Down Stream areas (km^2 and per cent monitored/estimated in each subarea and subregion).
(Fig. 1, Figs. I.I-I.V, Report B)

Sub-regions	Sub-areas	Drainage area of monitored rivers km^2		Down Stream areas km^2	Total km^2	Monitored %
		Main	Tributary			
Skagerrak	No 1: Glomma	41218	2389	2416	46023	94,8
	" 1: Inner Oslofjord		959	342	1301	73,7
	No 2: Drammenselva	17028	226	320	17614	98,2
	No 3: Numedalslågen	5513	1043	631	7187	91,2
	No 4: Skjenselva	10348	1200	1283	12831	90,0
	No 5: Otra	3730	9109	904	13743	93,4
Total		77837	14966	5896	98699	94,0
The remaining North Sea	No 6: Orre	105	7233	2513	9851	74,5
	No 7: Suldalslågen	1466	16205	12681	30352	58,2
Total		1571	23438	15194	40203	62,2
The Norwegian Sea	No 8: Orkla	2680	28118	17036	47834	64,4
	No 9: Vefsna	4113	23907	18850	46870	59,8
Total		6793	52025	35886	94704	62,1
The Barents Sea	No 10: Alta	7367	45155	20619	73141	71,8
Total		93568	135584	77595	306747	74,7

3.3 Methodology for assessment of direct discharges to marine waters

As the methodology for assessing direct discharges to marine waters is outlined in the 1990-Report (Holtan et al., 1991), and the same procedure is adopted for 1994, we refer to the above mentioned document for further information on this matter.

3.3.1 Waste water treatment plants/sewage effluents

The Central Bureau of Statistics (SSB) and SFT have jointly initiated annual registration of data from all waste water treatment plants in the country with a capacity of more than 50 population units (p.u.*). The data are updated each year by the County Environmental Agencies. The data program SSB-AVLØP has been installed at all county governors' environmental agencies, which are responsible for collecting data from the municipalities. The county environmental agencies then send the data to SSB on disc. Since 1994 onwards, the reporting system SSB-AVLØP has been extended also to include data on scattered settlements. Discharge figures from SSB-AVLØP are reported to NIVA. NIVA uses these figures in the model "TEOTIL" to calculate the total discharges of phosphorus and nitrogen to Norwegian coastal waters. The figures take into account retention in fjords and watercourses.

In Eastern and Southern Norway a large share of the municipal waste water is purified in "high grade" plants. These areas, as well as Sør-Trøndelag, are also the areas with the highest hydraulic capacity per inhabitant. For example, the plants serving Oslo/Akershus have a capacity of more than 1.5 p.u. per inhabitant. In Hordaland and northwards along the coast the greater part of the waste water is purified mechanically. Of a total of 1822 plants that were registered at the end of 1993, 16 had a hydraulic capacity of more than 50000 p.u. These plants account for almost half the total registered hydraulic capacity and load. Only 2 of these large plants are based on mechanical purification. Fjords are the recipients of the discharges from about 65 per cent of the total capacity of the plants (SSB, 1995).

Preferably, the annual loads from sewage effluents have been estimated as the product of annual flow and flow-weighted concentrations, which previously in particular has been the case for the sewage plants situated in the Skagerrak area (i.e. the area involved in the North Sea Agreement), but from 1994 has come into force for most plans.

For the rest of the municipal wastewater, the loads were estimated by multiplying the number of people with the coefficients listed.

For crude (untreated) sewage discharges, PARCOM (10/3/2-E, 1988), has recommended the following derived per capita loads to be used for nutrients:

	PARCOM:			NORWAY:		
BOD	0.063	kg	O/person/day	0.046	kg	O/person/day
COD				0.094	kg	O/person/day
TOC				0.023	kg	/person/day
SPM	0.063	kg	/person/day	0.042	kg	/person/day
Total N	0.009	kg	N/person/day	0.012	kg	N/person/day
Total P	0.0027	kg	P/person/day	0.0017	kg	P/person/day

* p.u. (population units) is the number of permanent residents plus the number of population equivalents (next page) in an area.

The Norwegian coefficients are based on recent studies of Norwegian sewerage districts. These data are also used to calculate pollutional loads from the different treatment plants, reduced by the removal efficiency of the treatment plants. Municipal sewage also includes a portion of industrial effluents. The fraction of the total person equivalents (p.e.) is proportioned between sewage and industrial wastewater according to the number of persons and the size of industrial effluents connected to each treatment plant.

For metals in sewage discharges the calculated loads are based on measured concentrations and flows in larger treatment plants in the Oslo part of the Glomma area. Metal inputs from the rest of the country are estimated from local knowledge (*) as follows:

Substance	Cu	Zn	Cd	Pb	Cr-T	Ni	Hg
mg/person-equivalent/day (p.e.)	30	35	0.2	1.0	2.9	5.0	0.10

The coefficients used in 1994 are the same as those used for calculations in 1992 and 1993. The coefficients are based on the results of an investigation on this topic launched by SFT (1993), i.e. monitoring at different types of treatment plants especially in the Oslo part of the Glomma area. Measured/estimated loads from sewage are shown in Appendix II, Report B.

Sources: Knutzen and Øren (1983), Myhrstad (1985), OVA (1995), SFT (1993), VEAS (1995),

3.3.2 Industrial effluents

Sampling frequency for industrial wastewater varies from weekly mixed samples to samples taken at random, but at least twice a year. Measured and estimated loads from industrial activities in the different areas are shown in Appendix III, Report B. According to SFT about 90 per cent of the industrial discharges (i.e. of the substances in question) are included in the total, and probably more for Total-P and Total-N.

3.3.3 Other inputs

(nutrients in area runoff from "Down Stream areas" of main and tributary rivers and rivers not monitored)

The pollution load model calculates the load from each pollution source by using area and activity specific load coefficients multiplied by areas (in square kilometres) of different categories and activity numbers, eg. population (Holtan and Åstebøl, 1990). The coefficients used are prepared according to precipitation, climate, vegetation and soil in the different areas.

To estimate load from agricultural land area runoff, coefficients in the range of 50-200 kg Total-P and 2000-6500 kg Total-N km²/year are used depending on point sources, location of the agricultural land in relation to major tributaries, and agricultural production intensity. Load from upland (remote unpolluted) areas were estimated by using export coefficients in the range of 4-6 kg Total-P and 200-600 kg Total-N km²/year. The highest values were used in areas most affected by long range pollution (acid rain) along the Southern and Western coast. The coefficients are based on mean annual runoff for the period 1931-60.

Total direct nutrient discharges (Down stream areas) are shown in Table I (Appendix I, Report B), in the different subareas (1-10) in Appendix VI, Report B. Direct discharge areas (km²) are shown in Table 5.

4. RESULTS AND DISCUSSION

4.1 Pollutants

Norwegian watercourses, coastal fjords and sea areas are recipients of various substances discharged from many different sources. The discharges may have widely different impacts of varying severity. SFT (Rensvik, 1990) considers that the most serious problems are connected to eutrophication (nutrient effluents and runoff), discharges of metals and organic micropollutants and acidification of water and soil.

In this investigation riverine and direct inputs of nutrients (P- and N-compounds), heavy metals, lindane and PCBs are measured or estimated. In addition to these contaminants the water is polluted by dissolved organic matter, especially from the pulp and paper industry and from municipal sewage, which also has been taken into account in this investigation.

SFT (Rensvik, 1990) has given first priority to eliminating the effluents of 13 of the substances classified as micropollutants, which are in use in Norway, as quickly as possible. Most of this pollution comes from industry, but other sources are municipal sewage, landfill leachate, and pesticide residues from agriculture.

Pollution by heavy metals is either due to discharges from industry, discharges from existing and abandoned mines, leaching from landfills polluted ground or atmospheric fallout. The municipal sewage is the source of several heavy metals. Long- range transboundary air pollution is another source of pollution both with respect to heavy metals and organic micropollutants.

Polychlorinated biphenyls (PCBs) are present in different industrial effluents and also released from discarded electrical equipment containing PCBs (e.g. transformer oil), when such equipment is unsatisfactorily stored or destroyed by incineration. There is no enterprise in Norway discharging PCBs regularly.

Major sources of phosphorus and nitrogen pollution are considered to be municipal sewage, agriculture and to a less degree, industry.

4.2 1994-results and discussion

The results given for riverine inputs (main rivers and tributaries) and discharges entering directly into marine recipients, are mainly based on monitoring data/direct measurements (paragraph 3.3).

Measured concentrations of the chemical parameters of the ten main rivers (1994), mean values, standard deviation and range are listed in Appendix VII, Report B. In the case of Cd, Pb, Hg and the different congeners of PCBs, where most of the rivers had concentrations below the respective detection limits, the concentrations are statistically treated as "limit-values". Total annual loads of the main rivers 1994 are shown in Appendix IV, Report B. Annual loads of nutrients and S.P.M. are also presented in Fig. 6. Total annual loads of the tributaries are shown in Appendix V, Report B. For the whole country, total annual loads (Direct discharges and Riverine inputs) are shown in Table I (Appendix I, Report B) and for the four subregions in Tables 1.1-1.4 (Appendix I, Report B), nutrients and S.P.M. also in Fig. 7.

As in 1990 -1993 the greatest emphasis with regard to accuracy has been given to the input estimate of the Skagerrak region, as this is considered the most susceptible part of the North Sea. The Skagerrak reception of total loads in 1994 were 22 per cent of the phosphorus and 41 per cent of the nitrogen yield.




Fig. 6 Main rivers. Nutrients and S.P.M. Total loads 1994 (A) and Total normalized loads (B) in the different rivers.

Whole columns = Total P / N / S.P.M.
Dark hatching = Phosphates / Nitrates




Fig. 7 Nutrients and S.P.M. Total and river discharges 1994 (A) and Total normalized loads (B) from mainland Norway to convention waters and the four subregions: I: Skagerrak, II: The remaining North Sea, III: The Norwegian Sea, IV: The Barents Sea.

Whole columns	= Grand total
Light hatching	= Direct discharges
Dark hatching	= Main and tributary rivers

In this region where 94 per cent of the area is river-monitored, more than 60 per cent of the P-load and 75 per cent of the N-load were found in the riverine inputs.

According to the results from the 1994 investigation, total annual nutrient load to coastal waters from landbased Norwegian sources, is approximately 4100 tonnes of phosphorus and 98100 tonnes of nitrogen (Fig. 7). About 30 per cent of the phosphorus and 60 per cent of the nitrogen yield were inputs from the monitored rivers and tributaries. The largest inputs of heavy metals were of copper and zinc, which in 1994 amounted to about 237 and 905 tonnes, of which 79 and 85 per cent respectively, were river monitored (Fig. 8).




Fig. 8 Cu and Zn. Total and river discharges 1994 (A) and Total normalized loads (B) from mainland Norway to convention waters and the four subregions:

I: Skagerrak, II: The remaining, North Sea, III: The Norwegian Sea, IV: The Barents Sea
 Whole columns = Grand total
 Light hatching = Direct discharges
 Dark hatching = Main and tributary rivers

Inputs of other metals and micropollutants were low. As the detection limits of the parameters Cd and Pb at the NIVA laboratory were above those requested from PARCOM, all analyses from 1992 and onwards of these substances are analyzed on an ICP-MS-instrument at NILU (paragraph 3.2). Still, a few of the concentrations found for these parameters also in 1994 were below the detection limits. Therefore two quantities have been estimated, one assuming that the true concentration was zero and the other assuming that the true concentration was the limit of detection. This provides maximum and minimum concentrations within which the true estimate will fall. When evaluating inputs these data provide upper and lower boundaries of the estimate.

Inputs of cadmium were measured/calculated to be between 9.0 and 9.3 tonnes, lead between 83.0 and 83.1 tonnes, mercury between 355 and 443 kg. The same "below detection limit problem" applies for the inputs of mercury, and also for PCBs which were measured to be between 37.1 and 72.8 kg. In Fig. 9 the lower and upper loads of these substances in the ten main rivers are presented. The pesticide lindane was found in most analyses, but in small amounts. The findings of this substance in Norwegian rivers are likely due to long range air pollution, as we often find lindane in runoff from areas where the compound has never been used (Olav Lodhe, State Plant Protection Agency, pers. comm.). Total load is estimated to about 89 kg.

In most areas except Skagerrak the riverine inputs, both of Total-P and S.P.M., were a little lower in 1994 than the year before, mainly due to precipitation/runoff-conditions (paragraph 4.3).




Fig. 9 Main rivers. Cd, Pb, Hg and PCBs. Total loads of the different rivers 1994 (lower and upper boundaries for the estimates).

Whole columns = upper boundary for the estimate
 Dark hatching = lower boundary for the estimate

4.3 Mean annual runoff (1931-60, 1961-90) and "mean load"

As mentioned (3.2.2) mean annual runoff for the last LTA-period (1961-90) has been estimated (fig. 2). For most main rivers the mean runoff 1961-90 is lower or in the same level as the 1931-60-period. However, these values are preliminary and may be adjusted when edited and published by NVE. As for precipitation, new normals were published in 1993.

Compared to Riverine Inputs to Marine Waters in 1990 - 1993, most calculated mean concentrations were in about the same level in 1994. Total flow for all "Skagerrak rivers" and also for Orreelva, and accordingly the calculated loads for most of the substances were higher in 1994 than the year before, except for the loads of total P and S.P.M. of Glomma and Orreelva. One of the reasons is probably fewer samples in 1994 (once monthly) than earlier in the flow period (every second week/weekly). As for the other main rivers, total flow was lower, with lower calculated loads for most substances as a result.

Annual variations in precipitation/runoff, erosion and seasonal activities of man in the drainage basins, strongly influence the mass transport in the watercourses. The transport values may vary considerably from one year to another. These variations are complicating the estimation of "normal transport values" i.e. mass transport in a "normal" year. Fig. 10 illustrates variations in annual runoff for the ten main rivers in 1985, 1990-1994, and annual variations in total discharge, of the nutrients total N and P for the same years.

In order to adjust the 1994 transport values to a "normal year", approximation have been made by multiplying weighted mean concentrations by mean runoff (LTA, 1961-90) - to normalize the concentrations is not possible. "Mean-values" (normals) for the ten main rivers and tributaries are given in Appendix X, Report B, where also total discharges to the sub regions are shown. In Fig. 6 - 7 the "normalized" nutrient transport values are compared with 1994-values.

During a normal winter the upland area of Norway together with the eastern part of the country show very little runoff. The frozen soil and snowcover protect vegetation and surface soil from erosion. But with the mild winters (1990-1993), soil especially in plowed field is exposed to more frequent and larger flood erosion also during the winter (eg. Glomma). As 1994 was a normal winter we suppose this is the main reason for lower concentrations/loads especially in Glomma, but also in Orreelva.

From experience we have learned that the product of weighted concentrations and mean runoff gives a relatively good estimate of the mass-transport in rivers, especially for the large rivers, and those not considerably exposed to erosion-matérial. In 1994 annual precipitation varied about normal or were a little higher in the Southern and South Western Norway, and a little lower in the rest of the country (Fig 3). On an annual basis also runoff varied about normal or were higher in most of the Southern and South Western Norway, but were somewhat lower in the rest of the country. The river Suldalslågen is recently regulated and has now considerably less annual water discharge than in the normal period (1931-60).

4.4 Nutrient retention in fjords

Both phosphorus and nitrogen retention in watercourses is taken into account in the calculation of the Norwegian contribution to marine pollution, but in a conservative way. However, no corrections are so far made for retention in fjords and other marine areas.

Considering the nutrient input to the open marine waters, one should also take into account retention in fjords, at least in well defined threshold fjords. As a result of high salinity compared to freshwater, marine waters have better conditions for sedimentation than lakes. For example clay settles very poorly

in lakes, but very efficient in fjords. In addition to temperature stratification, fjords also show salinity stratification, with light brackish water on top of heavier, saline waters. Thus the over all stratification in fjords is in most cases stronger than in lakes. This implies that particulate pollutants lost to deep waters by sedimentation have less chance to be brought back to the plankton producing layer than in lakes.

In addition, stronger stratification implies greater chance for oxygen depletion in deep waters, which in fact is seen in many sheltered Norwegian fjords. Theoretically this will improve the conditions for denitrification. However, this greater stability is often reduced by rougher physical conditions in fjords compared to lakes.

Retention in Norwegian fjords is very poorly studied by direct budget measurements. In the Drammensfjord Magnusson and Næss (1986) found that about 60% of the incoming phosphorus was retained in the fjord, while for nitrogen the retention was only about 15%. In the silled Inner Oslofjord preliminary calculations indicate that nitrogen retention is in the order of 30-50% (Baalsrud, K., and B. Bjerkeng, 1991).

Thus, nutrient retention in threshold fjords seems to be of the similar magnitude as we find in lakes, and it is likely to believe that retention can be estimated from the same type of models that applies for lakes. The general lack of calibration data on retention models in fjords implies that we find it to early, at this stage of knowledge, to include these correction in the Norwegian discharge budget. It should be kept in mind, however, that a significant part of the particulate pollutants, and pollutants with particle affinity, end up in fjord sediments and thus are prevented from reaching the coastal waters.




Fig. 10 Main rivers. Annual runoff and nutrient load in 1985, 1990-94.

5. REFERENCES

5.1 Project Personnel

At NIVA:

Dag Berge:	Member of the Programme Committee
Hans Holtan:	Member of the Programme Committee
Gjertrud Holtan :	Project manager

Contact with field workers/field work: G. Holtan, E. Bjerknes, B. Hals, H. Holtan, S. W. Johansen, E. Iversen.

Care of equipment, shipment of samples etc.: E. Bjerknes, B. Hals.

Analysis of PCBs and Lindane: E. Brevik, G. Sigernes, T. Tellefsen.

Analysis of Hg and Heavy Metals: H. Hovind, B. Lauritzen, M. Villø, R. Beba, L. Bryn.

Other chemical analyses: NIVAs Analytical Chemical Laboratory.

Data Collection/evaluation: G. Holtan, M. Grande, H. Holtan.

EDP/calculations/registrations: T. Hopen, L. Henriksen, G. Holtan, A. Lauritzen, T. Tjomsland.

Reporting work: G. Holtan, D. Berge, H. Holtan.

External analyses:

Heavy metals : O. Røyset, M. Vadset. NILU.

FIELD WORKERS:

(1) GLOMMA:	B. Moen. Fylkeslaboratoriet i Østfold. Moss.
(2) DRAMMENSELVA:	E. Iversen. NIVA. Oslo.
(3) NUMEDALSLÅGEN:	E. Løve. Næringsmiddeltilsynet. Larvik kommune.
(4) SKIENSVASSDRAGET:	A. Andersen. Vannlaboratoriet i Telemark. Skien
(5) OTRA:	M. Aadnevik. Ingeniørvesenet. Kr.-sand kommune.
(6) ORREELVA:	G. Undheim. Landbrukskontoret. Time kommune.
(7) SULDALSLÅGEN:	Ø. Vårvik. Suldal Elveeigarlag. Suldalsosen.
(8) ORKLA:	O. Lien. Rennebu.
(9) VEFSNA:	B. Hauan. Teknisk etat. Vefsn kommune. Mosjøen.
(10) ALTA:	P. Nilsen. Finnmarksforskning. Alta.

5.2 Literature


- Andersen, D.O., 1993: Mandalselva 1993. Rapport fra Agder distriktshøgskole, vannlaboratoriet. 24 s. + vedlegg.
- Baalsrud, K. og B. Bjerkeeng, 1991: Tiltaksanalyse for Indre Oslofjord. Brukerkrav. Siktedy og oksygen i dypvannet. En enkel lineær modell. NIVA-rapport O-90131 (l.nr. 2524). 79 s.
- Bjørklund, A., G.H. Johnsen, Å. Åtland og A. Kampestad, 1993: Overvåking av ferskvannsresipienter i Bergen kommune i 1992. Rådgivende Biologer A/S, Institutt for Miljøforskning. Rapport nr. 81. 168 s.
- Bloom, N.S. and E.A. Crecelius, 1983: Determination of Mercury in Sea-water at sub-nanogram per liter levels. Marine Chem. 14, pp. 49- 59.
- Brandrød, T.E. og M. Mjelde, 1992: Tilgroing med vannvegetasjon i terskelbasseng i Eksingedalselva, Hallingdalselva og Skjoma. Omfang, årsaker og tiltak. NIVA-rapport O-90136 (l.nr. 2826). 94 s.
- Brettum, P., 1990: Undersøkelser av Årdalsvatn i 1990. NIVA-rapport 0-90143 (l.nr. 2563). 32 s.
- Brun, P.F. og T. Haugen, 1990: Overvaking av fjordar og vassdrag i Møre og Romsdal 1986-88. Fylkesmannen i Møre og Romsdal. Rapport nr. 2/90. 101 s.
- Bærum kommune, 1995: 1994-Data fra Sandvikselva. Bærum (unpublished).
- DNMI, 1993: Nedbørnormaler 1961-90. DNMI. Oslo, 63 s.
- DNMI, 1995: Nedbørdata fra 1994. DNMI. Oslo (unpublished).
- Faafeng, B., 1995: Gjersjøens utvikling 1972-94 og resultater fra sesongen 1994. NIVA-rapport 0-70006 (in prep.).
- Faafeng, B., 1994: Årungen og Årungelva. NIVA-rapport (in prep.).
- Fagernes, K.E. og Å. Molversmyr, 1992: Vannkvalitet i Årosvassdraget 1984-1988. Rapport nr. 17. Fylkesmannen i Buskerud. Miljøvernnavdelingen. 25 s. + Vedlegg.
- Grande, M. og R. Romstad, 1994: Tiltaksorientert overvåking i Orkla, 1994. NIVA-rapport 0-800210 (l.nr. 3319) 53 s..
- Grande, M. og E.R. Iversen, 1992. Redusert vannføring i Årdalselva ved flytting av Tya kraftverk. Konsekvenser for forurensningssituasjonen. NIVA-rapport O-92183 (l.nr. 2815). 26 s.
- Haugen, T., 1994: Forurensningsundersøkelser i 12 vassdrag i Sør-Trøndelag - Mål for vannforekomstene og egnethet til bruk. Fylkesmannen i Sør-Trøndelag. Miljøvernnavdelingen, rapport nr. 2-94. 44 s.
- Hessen, D., E.-A. Lindstrøm og M. Mjelde, 1993: Storglomreguleringen. Undersøkelse av vannkjemi og vegetasjon. NIVA-rapport 0-901234. (l.nr. 2931). 77 s.
- Holtan, G., D. Berge, H. Holtan and T. Hopen, 1991: Paris Convention. Annual report on direct and riverine inputs to Norwegian coastal waters during the year 1990: A: Principles, results and discussions. SFT-report 452A/91. NIVA-report 0-90001/Serial No.: 2582. 43 pages.

- B: Data report. SFT-report 452B/91. NIVA-report 0-90001/Serial No.: 2577. 103 pages.
- Holtan, G., D. Berge, H. Holtan and T. Hopen, 1992: Paris Convention. Annual report on direct and riverine inputs to Norwegian coastal waters during the year 1991:
 A: Principles, results and discussions. SFT-report 488A/92. NIVA-report 0-90001/Serial No.: 2809. 40 pages.
 B: Data report. SFT-report 488B/92. NIVA-report 0-90001/Serial No.: 2777. 104 pages.
- Holtan, G., D. Berge, H. Holtan and T. Hopen, 1993: Annual report on direct and riverine inputs to Norwegian coastal waters during the year 1992. A. Principles, results and discussion. B. Data report. SFT-report 542/93. NIVA-report O-90001/Serial No.: 2964. 137 pages.
- Holtan, G., D. Berge, H. Holtan and T. Hopen, 1994: Annual report on direct and riverine inputs to Norwegian coastal waters during the year 1993. A. Principles, results and discussion. B. Data report. SFT-report 580/94. NIVA-report O-90001/Serial No.: 3162. 138 pages.
- Holtan, H. (red.), 1986: Norsk Vassdragsleksikon (utkast). Rapport fra Norsk Hydrologisk Komité, Oslo, 32 s. + vedlegg.
- Holtan, H. (red.), 1989: Vannkvalitetskriterier for ferskvann. Rapport TA-630 fra Statens forurensningstilsyn.
- Holtan, H., D. Berge and J. Molvær, 1990: Retention of nutrients in lakes and rivers with comments on retention in fjords. Paper prepared for PARCOM. NIVA-document 0-90145. 14 s. (unpublished).
- Holtan, H. og S.O. Åstebøl, 1990: Håndbok i innsamling av data om forurensningstilførsler til vassdrag og fjorder. Revidert utgave. NIVA-Jordforsk 0-89043/0-892301 (l.nr.2501). 53 s.
- Holtan, H., 1992. Overvåking av Aulielva 1991/1992. NIVA-rapport O-92111. (l.nr. 2796). 41 s.
- Holtan, H., 1994. Overvåking av Farris med tilløp 1993. NIVA-rapport O-91205. (l.nr. 3101). 20 s.
- Holtan, H., 1994: Konsentrasjon og transport av fosfor og nitrogen i Glomma 1978-1993. NIVA-rapport O-94031. (l.nr. 3000). 11 s.
- Hovind, H. og G. Holtan, 1994: Determination of unspecific organic substances in surface water - a comparison of different analytical parameters (in prep.).
- Ibrekk, H.O., H. Holtan, D. Berge, R. Gulbrandsen og K. Øren, 1991: Nordsjøplan. Vassdrag. NIVA-rapport 0-902302 (l.nr. 2628). 92 s.
- Indre Sogn Interkommunale Servicekontor, 1995: Vassdragsovervåking i Sogn og Fjordane 1994. ISIS-rapport/prosjekt nr. 474. 41 s + vedlegg.
- Innherred Kjøtt- og Næringsmiddelkontroll, 1994: Analysedata fra Snåsavassdraget 1993. Byveterinæren i Steinkjer. (Unpubl.).
- Iversen, E. R., 1993: Vannforurensninger fra nedlagte gruver. Del IV. NIVA-rapport 0-92152 (l.nr. 3045). 36 s.
- Johannessen, M. og K. Carm, 1992: Vannforurensning i Grenland. Endel resultater fra lokal miljøovervåking. Fylkesmannen i Telemark. Miljøvernavdelingen. Rapport nr. 11 - 1992. 21 s.

- Johnsen, G.H., 1992: Etterundersøkelse av Eidfjordvassdragene i Hordaland 1990-1991. Rådgivende Biologer A/S. Institutt for Miljøforskning. Rapport nr. 59. 59 s.
- Johnsen, G.H., G.B. Lehmann og A. Bjørklund, 1992: Tilstand og status for vann og vassdrag i Hordaland. Rådgivende Biologer A/S. Institutt for Miljøforskning. Rapport nr. 62. 75 s.
- Kaste, Ø., K.J. Aanes og E.-A. Lindstrøm, 1994: Otra 1993. Tiltaksorientert overvåking og konsekvensundersøkelse av industriutslipp. Overvåkingsrapport nr. 576/94. NIVA-rapport O-800208 (l.nr. 3109). 44 s.
- Kaste, Ø., K.J. Aanes og E.-A. Lindstrøm, 1995: Otra 1994. Tiltaksorientert overvåking og konsekvensundersøkelse av industriutslipp. Overvåkingsrapport nr. 606/95. NIVA-rapport O-800208 (l.nr. 3290). 42 s.
- Knutzen, J. og K. Øren, 1983: Miljøgifter i kommunalt avløpsvann - vurdering av effekter ved utslipp i sjøvann. Vann-3-83. pp. 292-305.
- Kroglund, F., P. Brettum og M. Lie, 1993: Kvinavassdraget. Resipientundersøkelse. NIVA-rapport O-91161 (l.nr. 2853). 46 s.
- Lindblad, V., 1994: Utslippskontroll av kloakkrenseanlegg i Østfold. Årsrapport for 1993. Rapport nr. 8/94. Fylkesmannen i Østfold. Miljøvernavdelingen. 102 s.
- Lingsten, L., 1987: Pilot Study on Riverine Inputs to Marine Waters. NIVA-report 0-86201. 37 s. (unpublished).
- Lindstrøm, E.A., 1994: Vurdering av vannkvalitet i Surna - basert på begroingsobservasjoner i 1993. NIVA-rapport O-93190. (l.nr. 2998). 17 s.
- Lindstrøm, E.A. og B. Relling, 1994: Overvåking av små og mellomstore landbruksforensede vassdrag i Møre og Romsdal. Undersøkelser i 1992 og 1993. NIVA-rapport O-94117. (in press).
- Løvstad, Ø., T. Hauger, P. Vallner og K. Bjørndalen, 1988: Vassdrag og kystområder. Overvåking 1987. Rapport nr. 6/88. Fylkesmannen i Østfold. Miljøvernavdelingen. 138 s. + vedlegg.
- Magnusson, J. og K. Næs, 1986: Basisundersøkelser i Drammensfjorden 1982-84: Delrapport 6: Hydrografi, vannkvalitet og vannutskiftning Overvåkingsrapport nr. 243/86. NIVA-rapport 8000315 (l.nr. 1892). 77 s.
- Moiseenko, T., M.Mjelde, T.E. Brandrud, P. Brettum, V. Dauvalter, L. Kagan, N. Kashulin, L. Kudriavtseva, A.Lukin, S. Sandimirov, T.S. Traaen, O. Vandysh and V. Yakovlev, 1994: Pasvik River Watercourse, Barents Region: Pollution Impacts and Ecological Responses. Investigations in 1993. INEP-NIVA-report O-93144. (l.nr. 3118). 87 s.
- Molværmyr, Å., S. Sanni og T. Tyvold, 1989: Basisundersøkelse av Figgjovassdraget 1984-1987. Rogalandsforskning. Rapport nr. RF-219/89. 79 s. + bilag.
- Molværmyr, Å. og S. Sanni, 1990: Tiltaksrettede undersøkelser i Ims-Lutsi vassdraget. Rogalandsforskning. Rapport nr. RF-171/90. 54 s. + vedlegg.
- Molværmyr, Å., 1992: Statusrapport for Orrevassdraget. Undersøkelser i 1990 og 1991. Rogalandsforskning. Rapport nr. RF-55/92. 61 s.

- Molversmyr, Å., 1993: Skas-Heigre kanalen. Undersøkelse av stofftransport. Årsrapport 1992. Rogalandsforskning. Rapport nr. RF-270/93. 18 s.
- Molvær, J. og A. Stigebrandt, 1991: Undersøkelse av eutrofiering i Grenlandsfjordene 1988-89. Delrapport 3. Vannutskiftning i fjordene. Overvåningsrapport nr. 450/91. NIVA-rapport 0-8000372, (l.nr. 2588). 45 s.
- Montaser, A. and D.W. Golightly, 1987: Inductively Coupled Plasmas in Analytical Chemistry. New York. VCH Publishers.
- Muladal, H. og T. Skotvold, 1993: Undersøkelse av forurensningstilstanden i Kåfjordvassdraget, Kåfjord kommune. Sluttrapport. Akvaplan-NIVA. Rapport nr. 325/3/93. 28 s.
- Myhrstad, A., 1985: Miljøgifter. Utslipp via kommunale anlegg. Rapport fra Elliot Strømme A.S. 13 s. + vedlegg.
- NILU, 1990: U-110. Forskrift for bestemmelse av elementer i vann med ICP-MS. 2 s.
- NVE, 1987: Avrenningskart over Norge. NVE. Oslo. 8 kartblad.
- NVE, 1990: Vassdragsregisterets kartbok. V 24. NVE. Oslo.
- NVE, 1995: Vannføringsdata fra 1994. NVE. Oslo (unpublished).
- Næringsmiddeltilsynet i Tønsberg, 1995: Aulielva 1994. Overvåningsrapport (in prep.).
- OSPAR, 1994. Oslo and Paris Conventions for the Prevention of Marine Pollution. Sixteenth Joint Meeting of the Oslo and Paris Commissions. Karlskrona 13-17 June 1994. OSPAR 16/4/8-E. 7 s.
- OVA, 1995: Bekkelaget renseanlegg. Årsrapport med nøkkeltall 1994. OVA-publikasjon. 12 s + vedlegg.
- Paris Commission, 1986: Eighth Meeting of the Paris Commission. Annex 4-6. Madrid 2-4 June 1986.
- Paris Commission, 1988: Tenth Meeting of the Paris Commission. PARCOM 10/3/2-E. Lisbon 15-17 June 1988.
- Paris Commission, 1993: Convention for the Prevention of Marine Pollution from Land-Based Sources. Fifth Meeting of the Ad hoc Working Group on Input Data. Parcom 5/6/1-E.
- Paulsen, L.I., 1994. Overvåking av vannkvaliteten i Årgårdsvassdraget 1993. Fylkesmannen i Nord-Trøndelag, Miljøvernavdelingen. Rapport 5-1994. 25 s.
- Rensvik, H. (red.), 1990: Vårt felles miljø. Oslo, SFT-rapport, 95 s.
- Schartau, A.K.L., 1995: Kjemisk overvåking av norske vassdrag - Elveserien 1994. NINA Oppdragsmelding (in press).
- Semb, R., 1992: Overvåking av vannkvaliteten i Drammenselva 1985-1991. Rapport nr. 22. Fylkesmannen i Buskerud. Miljøvernavdelingen. 74 s.
- Semb, R. 1993: Flerbruksplan for Mandalsvassdraget. Et notat om forurensningstilførsler og vannkvalitet. Fylkesmannen i Vest-Agder. Miljøvernavdelingen. 24 s.

- SFT, 1980-1994: The National Monitoring Programme of Water Courses. Monitoring Results. Annual Reports; TA-1980, 1981: TA-580, 1982: TA-588, 1983: TA-597, 1984: TA-607, 1985: TA-622, 1986: TA-628, 1987: TA-639, 1988: TA-629/1989, 1989: TA-711/1990.
- SFT, 1991: Næringsmiddelindustri. Stedfesting. Forurensningsproduksjon. Utslipp. Grøner, Rådgivende Ingeniører. Rapport nr. 28506. 11s. + vedlegg.
- SFT/NTNF, 1991: Karakterisering av nitrogenkomponenter i kommunalt avløpsvann. FAN - rapport R-4/91. 55 s. + vedlegg.
- SFT, 1993: Miljøgifter i kommunalt avløpsvann. 93:10. 52 s.
- SFT, 1994: Forurensning i Norge 1993. TA-1079/1994. 58 s.
- SFT, 1995: Overvåking av langtransportert forurenset luft og nedbør. Årsrapport 1994 (in press.).
- SSB, 1979, 1980, 1990-1995: Statistical Yearbooks. SSB. Oslo.
- SSB, 1994: Natural Resources and the Environment 1994, SSB, Oslo. 178 s.
- Stene-Johansen, S. og J.E. Samdal, 1994: Miljøundersøkelser i indre Oslofjord - Kartlegging av kilder. NIVA-rapport O-921312. (l.nr. 3291). 80 s.
- The County Environmental Agencies: Østfold, Oslo and Akershus, Buskerud, Vestfold, Telemark, Aust-Agder, Vest-Agder, Rogaland, Hordaland, Sogn og Fjordane, Møre og Romsdal, Sør-Trøndelag, Nord-Trøndelag, Nordland, Troms, Finnmark: Chemical data from monitored rivers in 1993 and/or monitored the last decade (mostly unpublished) and also information about treatment plants in the different counties.
- Tjomsland, T. og H.O. Ibrekk, 1992: TEOTIL. Modul for teoretisk beregning av fosfor- og nitrogentilførsler i Norge. NIVA-rapport 0-902301 (l.nr. 2786). 38 s.
- Traaen, T. og H. Huru, 1994: Vannkjemisk overvåking i Tanavassdraget 1988-1993. NIVA-rapport 0-88192 (l.nr. 3097). 37 s.
- Traaen, T., S. Rognerud og E. Steinnes, 1994: Forsuring og tungmetallforurensning i grenseområdene Norge/Russland. Årsrapport 1993. NIVA-rapport O-89187 (l.nr. 3074). 30 s.
- Traaen, T., 1995: Overvåking av Gaula. Vannkjemiske undersøkelser. Årsrapport 1994. SFT-rapport nr. 609/95, NIVA-rapport 0-90051 (l.nr. 3286). 48 s.
- Vallner, P., 1995: Overvåking av vassdrag og kystområder - Østfold. Fylkesmannen i Østfold. Miljøvernnavdelingen (in prep.).
- VEAS, 1995: Årsmelding 1994. Rapport fra Vestfjordens Avløpsselskap, 22 s.
- Welz, B., Melcher, M. Sinerius, H.W. og D. Maier, 1984: Picotrace Determination of Mercury using the Amalgamation Technique. At. Spectrosc. 1984, 5.
- Wivestad, T.M., 1994: Vassdragsundersøkelser i Lierelva 1992-1993. Rapport nr. 18. Fylkesmannen i Buskerud. Miljøvernnavdelingen. 25 s. + vedlegg.
- Wivestad, T.M., 1994: Vassdragsovervåking i Drammenselva 1992-1993. Fylkesmannen i Buskerud. Miljøvernnavdelingen, 59 s. + vedlegg.
- Wold, T., 1995: Vassdrag i Oslo. Status for elvene 1994. Rapport Miljøtilsyn, OVA. 124 s.

B Data report

CONTENTS	Page:
INSTRUCTIONS FROM PARCOM (Paragraph 1 - 3)	5
APPENDIX I : TOTAL DISCHARGES 1994 (Paragraph 4 - 6)	6
Table I Total discharges from mainland Norway to convention waters	7
Table 1.1 Total discharges to the Skagerrak region	8
Table 1.2 Total discharges to the remaining North Sea	10
Table 1.3 Total discharges to the Norwegian Sea region	12
Table 1.4 Total discharges to the Barents Sea region	15
APPENDIX II : SEWAGE EFFLUENTS FROM DOWN STREAM AREAS OF MAIN AND TRIBUTARY RIVERS AND RIVERS NOT MONITORED 1994 (Paragraph 7 - 8)	17
Table II Sewage effluents from mainland Norway to convention waters	18
Table 2.1 Sewage effluents to the Skagerrak region	19
Table 2.2 Sewage effluents to the remaining North Sea	20
Table 2.3 Sewage effluents to the Norwegian Sea region	21
Table 2.4 Sewage effluents to the Barents Sea region	22
APPENDIX III : INDUSTRIAL EFFLUENTS FROM DOWN STREAM AREAS OF MAIN AND TRIBUTARY RIVERS AND RIVERS NOT MONITORED 1994 (Paragraph 11 - 13)	23
Table III Ind. effluents from mainland Norway to convention waters	24
Table 3.1 Industrial effluents to the Skagerrak region	25
Table 3.2 Industrial effluents to the remaining North Sea	26
Table 3.3 Industrial effluents to the Norwegian Sea region	27
Table 3.4 Industrial effluents to the Barents Sea region	28
APPENDIX IV : MAIN RIVERINE INPUTS 1994 (Paragraph 14 - 16)	29
Table 4.1 Main riverine inputs. Glomma (1)	30
Table 4.2 Main riverine inputs. Drammenselva (2)	31
Table 4.3 Main riverine inputs. Numedalslågen (3)	32
Table 4.4 Main riverine inputs. Skienselva (4)	33
Table 4.5 Main riverine inputs. Otra (5)	34
Table 4.6 Main riverine inputs. Orreelva (6)	35
Table 4.7 Main riverine inputs. Suldalslågen (7)	36
Table 4.8 Main riverine inputs. Orkla (8)	37
Table 4.9 Main riverine inputs. Vefsna (9)	38
Table 4.10 Main riverine inputs. Alta (10)	39

APPENDIX V : INPUTS FROM TRIBUTARY RIVERS -94 (Paragraph 17-19)	40		
Table 5.1 Tributary rivers in the Sub-areas	(1-5). The Skagerrak reg. 41		
Table 5.2 Tributary rivers in the Sub-areas	(6-7). Remain. North Sea 42		
Table 5.3 Tributary rivers in the Sub-areas	(8-9). The Norwegian Sea 43		
Table 5.4 Tributary rivers in the Sub-area	(10). The Barents Sea 44		
APPENDIX VI : OTHER INPUTS 1994 (Paragraph 20)	45		
Table 6.1 Nutrients from "Down Stream Areas"	46		
APPENDIX VII : MAIN RIVERS 1 - 10. MEASURED CONCENTRATIONS 1994	47		
Table 7.1 Glomma	48		
Table 7.2 Drammenselva	49		
Table 7.3 Numedalslågen	50		
Table 7.4 Skienselva	51		
Table 7.5 Otra	52		
Table 7.6 Orreelva	53		
Table 7.7 Suldalslågen	54		
Table 7.8 Orkla	55		
Table 7.9 Vefsna	56		
Table 7.10 Alta	57		
APPENDIX VIII : TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994	58		
Table 8.1 Cond., Nutrients, Heavy metals, Suspended part.matter	59		
Table 8.2 Mercury, Lindane, PCBs			
APPENDIX IX : TRIBUTARY RIVERS. ANNUAL LOAD 1994	73		
Table 9.1 Cond., Nutrients, Heavy metals, Suspended part.matter	74		
Table 9.2 Mercury, Lindane, PCBs (limit-values = limit)	81		
(1) Glomma "tributaries"	:	Tista	- Hølenelva
(1) Inner Oslo-fjord	:	Årungelva	- Åroselva
(2) Drammenselva "tributary"	:	Lierelva	
(3) Numedalslågen "tributaries"	:	Sandeelva	- Farriselva
(4) Skienselva "tributary"	:	Tokkeelva	
(5) Otra "tributaries"	:	Gjerstade.	- Audna
(6) Orreelva "tributaries"	:	Lygna	- Ulla
(7) Suldalslågen "tributaries"	:	Saudaelva	- Hornindalselva
(8) Orkla "tributaries"	:	Ørstaelva	- Salsvatnelva
(9) Vefsna "tributaries"	:	Åbjøra	- Reisa
(10) Alta "tributaries"	:	Mattiselva	- Grense Jacobse.

APPENDIX X :	"MEAN" TOTAL DISCHARGES (Mean concentr. of main and trib.rivers multiplied with mean runoff 1961-90 (main rivers), 1931-60 (tributary rivers))	90
Table X	"Mean" total discharges from mainland Norway to convention waters	91
Table 10.1	"Mean" total discharges to the Skagerrak region	92
Table 10.2	"Mean" total discharges to the remaining North Sea	93
Table 10.3	"Mean" total discharges to the Norwegian Sea region	94
Table 10.4	"Mean" total discharges to the Barents Sea region	95
Table 10.5A	Main rivers. "Mean" load (Mean concentrations multiplied with mean runoff (1961-90)) (limit-values = limit)	96
Table 10.5B	Main rivers. "Mean" load (Mean concentrations multiplied with mean runoff (1961-90)) (limit-values = zero)	97
Table 10.6	The Skagerrak region. "Mean" inputs from tributary rivers in the Sub-areas (1-5)	98
Table 10.7	The remaining North Sea. "Mean" inputs from tributary rivers in the Sub-areas (6-7)	99
Table 10.8	The Norwegian Sea. "Mean" inputs from tributary rivers in the Sub-areas (8-9)	100
Table 10.9	The Barents Sea. "Mean" inputs from tributary rivers in the Sub-area (10)	101
FIGURES:		
Figure I.I	Main rivers and tributaries draining to The Skagerrak region of the North Sea	9
Figure I.II	Main rivers and tributaries draining to The remaining North Sea	11
Figure I.IIIA	Main rivers and tributaries draining to The Norwegian Sea (Southern part)	13
Figure I.IIIB	Main rivers and tributaries draining to The Norwegian Sea (Northern part)	14
Figure I.IV	Main rivers and tributaries draining to The Barents Sea	16

**CONVENTION FOR THE PREVENTION OF MARINE POLLUTION FROM LANDBASED
SOURCES
QUESTIONARY ACCORDING TO THE TENTH MEETING OF THE PARIS COMMISSION
LISBON 15 - 17 JUNE 1988.**

The purpose of this form is to provide the Commission, in accordance with Article 17(B) of the Paris convention, with an assessment of the waterborne inputs to Convention waters.

The form should be completed for each calendar year in retrospect and submitted to the Secretary by June following the year to which the data relate.

The information sought relates to inputs through direct discharges (questions 7 - 13) and riverine inputs (questions 14 - 19).

Some information on discharges other than those mentioned below are also attached (question 20).

Separate forms for the four single areas are filled in.

(1) **THE COUNTRY IS NORWAY**

(2) **LENGTH OF COASTLINE INCLUDING FJORDS AND BAYS IS 21347 KM**

(3) **NATURE OF THE RECEIVING WATER IS COASTAL**

APPENDIX I :	TOTAL DISCHARGES (Paragraph 4 - 6)	Page:
---------------------	---	--------------

Table I	Total discharges from mainland Norway to convention waters	7
Table 1.1	Total discharges to the Skagerrak region	8
Table 1.2	Total discharges to the remaining North Sea	10
Table 1.3	Total discharges to the Norwegian Sea	12
Table 1.4	Total discharges to the Barents Sea	15

Paragraph 4: Direct Discharges

Paragraph 5: Riverine Discharges

Paragraph 6: Grand Total

**Table I TOTAL DISCHARGES from MAINLAND NORWAY
to convention waters 1994 (Fig. I).**

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		1.2	4.0	*	3.7	*
Cadmium			4.3	**	3.8	**
Mercury		105	132	*	118	*
Mercury			199	**	140	**
Copper		49	95		92	
Zinc		140	379		386	
Lead		6.3	44.7	*	31.9	*
Lead			44.9	**	31.9	**
Arsenic		0.5	22.1	*	15.2	
Arsenic			22.1	**	15.2	
Cr-T		6.5	188.6	*	62.1	*
Cr-T			188.6	**	62.1	**
Ni		14.2	88.9	*	45.2	*
Ni			89.0	**	45.2	**
PCBs ***			0.54	*	36.52	*
PCBs			29.3	**	43.5	**
gamma-HCH			53		35	
NH4-N	1637	11263	1721		1143	15764
NO3-N	14020	194	17332		15659	47205
PO4-P	201	1163	201		180	1745
Total N	22364	17783	32584		25374	98106
Total P	766	2180	669		492	4107
S.P.M.		5675905	216064		248351	6140320
TOC		18053	172009		182927	372989
COD		215420				215420
AOX		210				210

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

Table 1.1 TOTAL DISCHARGES to The Skagerrak Region 1994 (Fig. I.I).

The Skagerrak Region with main rivers (1) Glomma, (2) Drammenselva, (3) Numedalslågen, (4) Skienselva, (5) Otra

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.09	0.9	*	3.3	*
Cadmium			0.9	**	3.4	**
Mercury		42.07	12	*	108	*
Mercury			18	**	126	**
Copper		20.65	9		57	
Zinc		19.07	103		306	
Lead		0.91	7.1	*	27.5	*
Lead			7.1	**	27.5	**
Arsenic		0.28	3.44	*	12.7	
Arsenic			3.44	**	12.7	
Cr-T		4.38	12.1	*	49.5	*
Cr-T			12.1	**	49.5	**
Ni		10.82	4.8	*	36.4	*
Ni			4.8	**	36.4	**
PCBs ***			0.13	*	35.73	*
PCBs			3.0	**	40.9	**
gamma-HCH			11.5		32	
NH4-N	158	4616	545		1003	
NO3-N	1560	150	4972		14653	
PO4-P	18	106	37		152	
Total N	2393	7497	7871		22861	
Total P	70	265	120		429	
S.P.M.		16466	31456		226229	
TOC		6838	50798		155354	
COD		117370				117370
AOX		210				210

Measurements below detection limits are treated in two ways :


*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

- 1 ØSTFOLD**
 1 Tista, Iddefj.
 2 Mosselva, Mossesundet Ø.
OSLO & AKERSHUS
 3 Hølenelva, Drøbaksundet Ø.
 4 Årungelva, I. Oslof.
 5 Gjersjøelva, I. Oslof.
 6 Ljanselva, I. Oslof.
 7 Loelva/Alna, I. Oslof.
 8 Akerselva, I. Oslof.
 9 Frognerelva, I. Oslof.
 10 Lysakerelva, I. Oslof.
 11 Sandvikselva, I. Oslof.
 12 Aroselva, I. Oslof.
- 2 BUSKERUD**
 13 Lierelva, Drammensfj. Ø.
- 3 VESTFOLD**
 14 Sandeelva, Sandebukta
 15 Aulielva, Tønsbergfj.
 16 Farriselva, Larvikfj.
- 4 TELEMARK**
 17 Tokkeelva, Kragerø
- 5 AUST-AGDER**
 18 Gjerstadelva, Søndeledf.
 19 Vegårdselva, Sandnesfj.
 20 Nidelva, Arendal
VEST-AGDER
 21 Tovdalselva, Topdalsfj.
 22 Søgneelva, Flekkerøy
 23 Mandalselva, Mannefj.
 24 Audna, Sniksfj.

Fig. I.1
 Main Rivers and Tributaries draining to
 The Skagerrak Region of The North Sea.



I North Sea/Skagerrak

**Table 1.2 TOTAL DISCHARGES to The Remaining North Sea
1994 (Fig. I.II).**

The North Sea Region with main rivers : (6) Orreelva, (7) Suldalslågen


Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.99	1.2	*	0.0	*
Cadmium			1.3	**	0.0	**
Mercury		42.39	32	*	0	*
Mercury			59	**	2	**
Copper		6.76	14		1	
Zinc		75.75	147		4	
Lead		4.59	9.6	*	0.4	*
Lead			9.7	**	0.4	**
Arsenic		0.00	5.4	*	0.1	
Arsenic			5.5	**	0.1	
Cr-T		1.34	39.3	*	2.7	*
Cr-T			39.3	**	2.7	**
Ni		2.19	6.3	*	2.0	*
Ni			6.3	**	2.0	**
PCBs ***			0.0	*	0.0	
PCBs			9.7	**	0.4	**
gamma-HCH			16.4		1	
NH4-N	579	3103	524		17	
NO3-N	5161	21	7674		377	
PO4-P	56	275	56		2	
Total N	8286	4742	11838		506	
Total P	198	537	188		7	
S.P.M.		2106041	39980		1971	
TOC		5287	69618		1399	
COD		38768				38768

Measurements below detection limits are treated in two ways:

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28, 52, 101, 118, 153, 138, 180



6 VEST-AGDER

- 25 Lygna, Lyngdalsfj.
- 26 Kvina, Fedafj.
- 27 Sira, Åna-Sira
- ROGALAND**
- 28 Sokndalselva, Sogndalsstr.
- 29 Hellelandselva, Egersund
- 30 Bjerkreimselva, Egersund
- 31 Håelva, Håtangen
- 32 Figgjo, Solavika
- 33 Ims-Lutsi, Høgsfj./Boknafj.
- 34 Oltedalselva, Høgsfj./Boknafj.
- 35 Dirdalselva, Høgsfj./Boknafj.
- 36 Frafjordelva, Frafj./Boknafj.
- 37 Espedalselva, Høgsfj./Boknafj.
- 38 Lyseelva, Lysefj./Boknafj.
- 39 Årdalselva, Årdalsfj./Boknafj.
- 40 Førreelva, Jøsenfj./Boknafj.
- 41 Ulla, Jøsenfj./Boknafj.
- 42 Saudaelva, Sandsfj./Boknafj.
- 43 Åbøelva, Saudafj./Boknafj.
- 44 Vikedalselva, Boknafj.

HORDALAND

- 45 Etneelva, Etnefj./Bømlafj.
- 46 Opo, Sørfj./Hardangerfj.
- 47 Tyssø, Sørfj./Hardangerfj.
- 48 Kinsø, Sørfj./Hardangerfj.
- 49 Veig, Eidf.v./Hardangerfj.
- 50 Bjoreia, Eidfj. v. Hardangerfj.
- 51 Sima, Eidfj./Hardangerfj.
- 52 Austdøla, Osafj./Eidfj.
- 53 Norddøla, Osafj./Eidfj.
- 54 Tysselva, Fusafj.
- 55 Oselva, Fusafj.
- 56 Bergsdalselva, Veafj./Herdlafl.
- 57 Vosso, Veafj./Sørfj.
- 58 Ekso, Osterfj.
- 59 Modalselva, Osterfj.

SOGN OG FJORDANE

- 60 Nærøyelva, Aurl.fj./Sognefj
- 61 Flåmselva, Aurl.fj./Sognefj.
- 62 Aurlandselva, Aurl.fj./Sognefj.
- 63 Erdalselva, Lærdalsfj./Sognefj.
- 64 Lærdalselva, Lærdalsfj./Sognefj.
- 65 Årdalselva, Årdalsfj./Sognefj.
- 66 Fortunelva, Lusterfj./Sognefj.
- 67 Mørkrisølva, Lusterfj./Sognefj.
- 68 Jostedøla, Lusterfj./Sognefj.
- 69 Årøyelva, Sognd.fj./Sognefj.
- 70 Sogndalselva, Sognd.fj./Sognefj.
- 71 Gaular, Dalsfj./Bufj.
- 72 Jølstra, Førdefj.
- 73 Nausta, Førdefj.
- 74 Oselva, Høydalsfj.
- 75 Hopselva, Hyefj./Nordfj.S.
- 76 Gjengedalselva, Hyefj./Nordfj.S.
- 77 Breimselva, Gløppenfj./Nordfj.S.
- 78 Oldenelva, Indre Nordfj.
- 79 Lænelva, Indre Nordfj.
- 80 Stryneelva, Indre Nordfj.
- 81 Hornindalselva, Nordfj.N.

Fig. I.II

Main Rivers and Tributaries draining to The Remaining North Sea.



Main Rivers



Tributaries



Areas not monitored

Table 1.3 TOTAL DISCHARGES to The Norwegian Sea 1994 (Fig. I.III).**The Norwegian Sea Region with main rivers (8) Orkla, (9) Vefsna**

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.14	1.8 *	0.3 *	2.2	tonnes
Cadmium			1.9 **	0.3 **	2.3	tonnes
Mercury	19.00		63 *	7 *	88	kg
Mercury			92 **	8 **	120	kg
Copper	21.58		53	32	107	tonnes
Zinc	44.71		110	74	228	tonnes
Lead	0.81		22.6 *	3.8 *	27.2	tonnes
Lead			22.6 **	3.9 **	27.3	tonnes
Arsenic	0.27		10.1 *	1.8	12.2	tonnes
Arsenic			10.1 **	1.8	12.2	tonnes
Cr-T	0.70		96.7 *	7.1	104.5	tonnes
Cr-T			96.7 **	7.1	104.5	tonnes
Ni	1.15		15.3 *	5.1 *	21.6	tonnes
Ni			15.4 **	5.1 **	21.6	tonnes
PCBs ***			0.4 *	0.4 *	0.8	kg
PCBs			13.2 **	1.5 **	14.8	kg
gamma-HCH			21.9	1	23	kg
NH4-N	808	3284	397	110	4599	tonnes
NO3-N	6289	22	4483	534	11328	tonnes
PO4-P	110	755	83	11	959	tonnes
Total N	10004	5195	10304	1612	27115	tonnes
Total P	410	1331	272	29	2041	tonnes
S.P.M.	1421222		113662	11509	1546393	tonnes
TOC		5588	27243	18614	51444	tonnes
COD		57745			57745	tonnes
AOX		1			1	tonnes

Measurements below detection limits are treated in two ways :***) Detection limit = Zero******) Detection limit = Limit******* the following congeners: IUPAC Nos. 28,52,101,118,153,138,180**




Fig. I.III A
Main Rivers and Tributaries draining to
The Norwegian Sea (Southern Part).






Table 1.4 TOTAL DISCHARGES to The Barents Sea 1994 (Fig. I.IV).**The Barents Sea Region with main river (10) Alta**

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.00	0.2	*	0.06	*
Cadmium			0.2	**	0.07	**
Mercury		1.35	25	*	3.36	*
Mercury			30	**	3.80	**
Copper		0.37	19		2.82	
Zinc		0.43	20		1.49	
Lead		0.01	5.4	*	0.19	*
Lead			5.4	**	0.19	**
Arsenic		0.00	3.1	*	0.62	
Arsenic			3.1	**	0.62	
Cr-T		0.04	40.5	*	2.73	*
Cr-T			40.5	**	2.73	**
Ni		0.06	62.4	*	1.74	*
Ni			62.4	**	1.74	**
PCBs ***			0.0	*	0.36	*
PCBs			3.4	**	0.65	**
gamma-HCH			3.4		0.11	
NH4-N	91	260.07	255		13.91	
NO3-N	1010	1.73	202		95.15	
PO4-P	18	27.56	25		14.84	
Total N	1681	349.25	2571		395.39	
Total P	88	46.44	89		26.78	
S.P.M.		2132176	30967		8642	2171785
TOC		340.44	24351		7560	32251
COD		1537.28				1537
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes
						tonnes

Measurements below detection limits are treated in two ways :***) Detection limit = Zero******) Detection limit = Limit******* the following congeners: IUPAC Nos. 28,52,101,118,153,138,180**

IV Barents Sea




Fig. I.IV
Main Rivers and Tributaries draining to
The Barents Sea.

**10 FINNMARK**

- 130 Mattiselva, Kåfj./Altafj.
- 131 Tverrelva, Altafj.
- 132 Repparfjordelva, Repparfj.
- 133 Stabburselva, I. Porsangen V.
- 134 Lakselva, I. Porsangen S.
- 135 Børselva, I. Porsangen Ø.
- 136 Mattusjákkka, I. Laksefj. V.
- 137 Storelva, I. Laksefj. V.
- 138 Soussjákkka, I. Laksefj. V.
- 139 Adamselva, I. Laksefj. Ø.
- 140 Tanavassdr., Tanafj. S.
- 141 Vesterelva, Syltefj.
- 142 V. Jakobselv, Y. Varangerfj.
- 143 Neiden Munkfj./Varangerfj.
- 144 Passvikelva, Bøkfj./Varangerfj.
- 145 Grense Jakobselv, Varangerfj.

**APPENDIX II : SEWAGE EFFLUENTS FROM DOWNSTREAM AREAS OF
MAIN AND TRIBUTARY RIVERS AND RIVERS NOT
MONITORED 1994 (Paragraph 7 - 8)**

		Page:
Table II	Sewage effluents from down stream areas of mainland Norway to convention waters 1994	18
Table 2.1	Sewage effluents to the Skagerrak region	19
Table 2.2	Sewage effluents to the remaining North Sea	20
Table 2.3	Sewage effluents to the Norwegian Sea region	21
Table 2.4	Sewage effluents to the Barents Sea region	22

Paragraph 7: Sewage effluents .

Paragraph 8: Measurements of calculation used - including information on the concentration upon which the measurement is based:

Paragraph 3.3 (Report A, 1991, 1992 - 1995)

Municipal sewage includes a portion of industrial effluents

Table II Sewage Effluents from down stream areas of mainland Norway to convention waters (1994).

Total quantity of substance discharged per year:

Regions:	I The Skagerrak	II The North Sea	III The Norwegian Sea	IV The Barents Sea	Sum
Substance:	Region	Sea	Sea	Sea	
Cd	74	40	34	2	151 kg
Hg	38	22	19	1	81 kg
Cu	10.8	6.0	5.2	0.4	22.4 tonnes
Zn	16.2	7.0	6.0	0.4	29.7 tonnes
Pb	640	201	172	12	1026 kg
Cr-T	3.1	0.6	0.5	0.0	4.3 tonnes
Ni	6.9	1.0	0.9	0.1	8.8 tonnes
PCBs					kg
gamma-HCH					kg
NH4-N	4616	3103	3284	260	11263 tonnes
NO3-N	150	21	22	2	194 tonnes
PO4-P	106	275	755	28	1163 tonnes
Tot-N	5796	4137	4379	347	14659 tonnes
Tot-P	176	458	1258	46	1938 tonnes
S.P.M.	8317	8233	9908	597	27054 tonnes
TOC	6771	4937	5564	340	17613 tonnes
COD	28853	22171	26002	1537	78564 tonnes

Table 2.1 Sewage Effluents to The Skagerrak Region (1994).

The Skagerrak region with sub-areas: (1) Glomma, (2) Drammenselva,
 (3) Numedalslågen, (4) Skienselva, (5) Otra

Sub-areas :	Total quantity of substance discharged per year:					Precision of the estimate of the load	
	1	2	3	4	5		
Substance:							
Cd	42	3	14	3	11	kg	%
Hg	21	2	8	2	6	kg	%
Cu	6.12	0.47	2.09	0.45	1.69	tonnes	%
Zn	10.74	0.55	2.44	0.53	1.97	tonnes	%
Pb	484	16	70	15	56	kg	%
Cr-T	2.62	0.05	0.21	0.05	0.17	tonnes	%
Ni	6.13	0.08	0.35	0.08	0.28	tonnes	
PCBs						kg	%
gamma-HCH						kg	%
NH4-N	2443	178	472	1048	475	tonnes	
NO3-N	135	1	3	7	3	tonnes	%
PO4-P	31	10	30	8	26	tonnes	%
Tot-N	2899	238	629	1397	634	tonnes	%
Tot-P	52	17	50	13	43	tonnes	%
S.P.M.	3591	410	1999	361	1956	tonnes	%
TOC	3714	285	1329	248	1195	tonnes	%
COD	15761	1208	5635	1107	5142	tonnes	%

Table 2.2 Sewage Effluents to The Remaining North Sea Region (1994).

The remaining North Sea Region with sub-areas: (6) Orreelva, (7) Suldalslågen

Sub-areas :	Total quantity of substance discharged per year:		Precision of the estimate of the load
	6	7	
Substance:			
Cd	12	28	kg
Hg	7	15	kg
Cu	1.86	4.18	tonnes
Zn	2.17	4.88	tonnes
Pb	62	139	kg
Cr-T	0.19	0.42	tonnes
Ni	0.31	0.70	tonnes
PCBs			kg
gamma-HCH			kg
NH4-N	1015	2088	tonnes
NO3-N	7	14	tonnes
PO4-P	52	223	tonnes
Tot-N	1353	2784	tonnes
Tot-P	86	372	tonnes
S.P.M.	2825	5407	tonnes
TOC	1592	3345	tonnes
COD	6924	15248	tonnes

Table 2.3 Sewage Effluents to The Norwegian Sea Region (1994).

The Norwegian Sea Region with sub-areas: (8) Orkla, (9) Vefsna

Sub-areas :	Total quantity of substance discharged per year:		Precision of the estimate of the load
	8	9	
Substance:			
Cd	18	16	kg
Hg	10	9	kg
Cu	2.73	2.43	tonnes
Zn	3.18	2.84	tonnes
Pb	91	81	kg
Cr-T	0.27	0.24	tonnes
Ni	0.45	0.41	tonnes
PCBs			kg
gamma-HCH			kg
NH4-N	1636	1648	tonnes
NO3-N	11	11	tonnes
PO4-P	173	582	tonnes
Tot-N	2181	2198	tonnes
Tot-P	288	970	tonnes
S.P.M.	5138	4770	tonnes
TOC	2933	2631	tonnes
COD	13837	12164	tonnes

Table 2.4 Sewage Effluents to The Barents Sea Region (1994).

The Barents Sea Region with sub-area: (10) Alta

Total quantity of substance discharged per year:		Precision of the estimate of the load
Sub-area :	10	
Substance:		
Cd	2	kg
Hg	1	kg
Cu	0.37	tonnes
Zn	0.43	tonnes
Pb	12	kg
Cr-T	0.04	tonnes
Ni	0.06	tonnes
PCBs		kg
gamma-HCH		kg
NH4-N	260	tonnes
NO3-N	2	tonnes
PO4-P	28	tonnes
Tot-N	347	tonnes
Tot-P	46	tonnes
S.P.M.	597	tonnes
TOC	340	tonnes
COD	1537	tonnes

APPENDIX III : INDUSTRIAL EFFLUENTS FROM DOWN STREAM AREAS OF MAIN AND TRIBUTARY RIVERS AND RIVERS NOT MONITORED 1994 (Paragraph 11 - 13)

Table III	Industrial effluents from down stream areas of mainland Norway to convention waters 1994	24
Table 3.1	Industrial effluents to the Skagerrak region	25
Table 3.2	Industrial effluents to the remaining North Sea	26
Table 3.3	Industrial effluents to the Norwegian Sea region	27
Table 3.4	Industrial effluents to the Barents Sea region	28

Paragraph 11: Industrial effluents ./.

Paragraph 12: Measurements of calculation used - including information on the concentration upon which the measurement is based:

Paragraph 3.3 (Report A, 1991, 1992 - 1995)

Paragraph 13: Any other relevant information (e.g. proportion of substance discharged as insoluble material):

A portion of industrial effluents is included in municipal sewage

Table III Industrial Effluents from down stream areas of mainland Norway to convention waters (1994).

Total quantity of substance discharged per year:

Regions:	I	II	III	IV	Sum
	The Skagerrak	The North Sea	The Norwegian Sea	The Barents Sea	
Substance:					
Cd	18	953	101	0	1072 kg
Hg	4	20	0	0	24 kg
Cu	9.83	0.71	16.41	0	27 tonnes
Zn	2.85	68.70	38.69	0	110.24 tonnes
Pb	274	4389	641	0	5303 kg
Arsenic	276	0	270	0	546 kg
Cr-T	1.28	0.74	0.19	0	2.21 tonnes
Ni	3.91	1.18	0.29	0	5.38 tonnes
PCBs					kg
gamma-HCH					kg
NO3-N					tonnes
PO4-P					tonnes
Tot-N	1700	606	816	2	3124 tonnes
Tot-P	89	79	73	1	242 tonnes
S.P.M.	8149	2097808	1411314	2131579	5648851 tonnes
TOC	67	350	23	0	440 tonnes
COD	88516	16597	31743	0	136856 tonnes
AOX	210	0	1	0	210 tonnes

Table 3.1 Industrial Effluents to The Skagerrak Region (1994).

The Skagerrak Region with sub-areas: (1) Glomma, (2) Drammenselva,
 (3) Numedalslågen, (4) Skienselva, (5) Otra

Sub-area :	Total quantity of substance discharged per year:					Precision of the estimate of the load	
	1	2	3	4	5		
Substance:							
Cd	10.15	0.00	5.80	0.26	1.30	kg	_____ %
Hg	3.60	0.00	0.00	0.01	0.00	kg	_____ %
Cu	8347	0	72	50	1365	kg	_____ %
Zn	1153	4	123	1256	318	kg	_____ %
Pb	86.5	0.0	20.2	53.0	114.0	kg	_____ %
Arsenic	0.0	0.0	0.0	0.0	276.0	kg	_____ %
Cr-T	1222.6	1.6	11.9	44.9	1.5	kg	_____ %
Ni	21.8	2.7	1966.2	11.8	1910.0	kg	_____ %
PCBs						kg	_____ %
gamma-HCH						kg	_____ %
NO3-N						tonnes	_____ %
PO4-P						tonnes	_____ %
Tot-N	166.3	16.4	303.8	1205.8	7.9	tonnes	_____ %
Tot-P	33.0	0.2	44.8	9.4	1.9	tonnes	_____ %
S.P.M.	3276	173	2195	2087	418	tonnes	_____ %
TOC	28.4	0.0	26.1	12.5	0.0	tonnes	_____ %
COD	55714	217	22013	10573	0	tonnes	_____ %
AOX	209.4	0.0	0.0	0.2	0.0	tonnes	_____ %

**Table 3.2 Industrial Effluents to The Remaining North Sea Region
(1994).**

The remaining North Sea Region with sub-areas: (6) Orreelva, (7) Suldalslägen

Sub-areas :	Total quantity of substance discharged per year:		Precision of the estimate of the load
	6	7	
Substance:			
Cd	0.00	952.99	kg
Hg	0.00	20.23	kg
Cu	6	709	kg
Zn	94	68602	kg
Pb	0.0	4389	kg
Arsenic	0.0	0.0	kg
Cr-T	231.2	505.6	kg
Ni	11.7	1166.7	kg
PCBs			kg
gamma-HCH			kg
NO3-N			tonnes
PO4-P			tonnes
Tot-N	46.0	559.6	tonnes
Tot-P	2.9	76.2	tonnes
S.P.M.	1759680	338128	tonnes
TOC	15.9	333.9	tonnes
COD	837	15759	tonnes
AOX	0.0	0.0	tonnes

Table 3.3 Industrial Effluents to The Norwegian Sea Region (1994).

The Norwegian Sea Region with sub-areas: (8) Orkla, (9) Vefsna

Sub-areas :	Total quantity of substance discharged per year:		Precision of the estimate of the load
	8	9	
Substance:			
Cd	55.12	46.20	kg
Hg	0.07	0.00	kg
Cu	1252	15162	kg
Zn	5020	33666	kg
Pb	220.9	420.2	kg
Arsenic	0.0	270.0	kg
Cr-T	153.6	32.0	kg
Ni	104.2	190.0	kg
PCBs			kg
gamma-HCH			kg
NO3-N			tonnes
PO4-P			tonnes
Tot-N	267.2	549.0	tonnes
Tot-P	46.5	26.6	tonnes
S.P.M.	152089	1259225	tonnes
TOC	0.0	23.5	tonnes
COD	31743.2	0.0	tonnes
AOX	0.7	0.0	tonnes

Table 3.4 Industrial Effluents toThe Barents Sea Region (1994).**The Barents Sea Region with sub-area: (10) Alta**

Total quantity of substance discharged per year:	Precision of the estimate of the load
Sub-area : 10	
Substance:	
Cd 0.00	kg _____ %
Hg 0.00	kg _____ %
Cu 0	kg _____ %
Zn 0	kg _____ %
Pb 0.0	kg _____ %
Arsenic 0.0	kg _____ %
Cr-T 0.0	kg _____ %
Ni 0.0	kg _____ %
PCBs	kg _____ %
gamma-HCH	kg _____ %
NO3-N	tonnes _____ %
PO4-P	tonnes _____ %
Tot-N 2.5	tonnes _____ %
Tot-P 0.5	tonnes _____ %
S.P.M. 2131579	tonnes _____ %
TOC 0.0	tonnes _____ %
COD 0.0	tonnes _____ %
AOX 0.0	tonnes _____ %

APPENDIX IV : MAIN RIVERINE INPUTS 1994 (Paragraph 14 - 16)			Page:
Table 4.1	Main riverine inputs. Glomma	(1)	30
Table 4.2	Main riverine inputs. Drammenselva	(2)	31
Table 4.3	Main riverine inputs. Numedalslågen	(3)	32
Table 4.4	Main riverine inputs. Skienselva	(4)	33
Table 4.5	Main riverine inputs. Otra	(5)	34
Table 4.6	Main riverine inputs. Orreelva	(6)	35
Table 4.7	Main riverine inputs. Suldalslågen	(7)	36
Table 4.8	Main riverine inputs. Orkla	(8)	37
Table 4.9	Main riverine inputs. Vefsna	(9)	38
Table 4.10	Main riverine inputs. Alta	(10)	39

Paragraph 14: Main Rivers /.

Paragraph15: Measurements of calculation used - including information on the concentration upon which the measurement is based:

Appendix VII (1-10) and Paragraph 3.2 (Report A, 1991, 1992 - 1995)

Paragraph 16: Any other relevant information (e.g. proportion of substance discharged as insoluble material):

Table 4.1 MAIN RIVERINE INPUTS 1994 (1) Glomma

Total volume: 57419 1000 m³/day Long term average flow (LTA): 60324 1000 m³/day
 Minimum flow: 24365 1000 m³/day LTA period : 1961 to 1990
 Maximum flow: 163296 1000 m³/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.10	12	0.00	0.44 µg/l	2.45 tonnes	YES	%
Cadmium **	0.10	12	0.01	0.44 µg/l	2.46 tonnes	_____	%
Mercury *	3.17	12	0.00	11.50 ng/l	79.04 kg	YES	%
Mercury **	3.33	12	1.00	11.50 ng/l	82.50 kg	_____	%
Copper	1.64	12	1.13	2.30 µg/l	37.26 tonnes	YES	%
Zinc	8.49	12	2.30	24.10 µg/l	202.14 tonnes	YES	%
Lead	0.89	12	0.15	2.08 µg/l	19.35 tonnes	YES	%
Arsenic	0.40	1	0.40	0.40 µg/l	8.38 tonnes	YES	%
Total Cr-T	1.31	1	1.31	1.31 µg/l	27.45 tonnes	YES	%
Ni	0.92	1	0.92	0.92 µg/l	19.28 tonnes	YES	%
PCBs *		3		ng/l	17.94 kg	NO	%
PCBs **		3		ng/l	19.39 kg	_____	%
gamma-HCH (lindane)	0.58	3	0.51	0.70 ng/l	11.63 kg	YES	%
Ammonia (NH4-N)	31.75	12	11.00	60.00 µg/l	557 tonnes	YES	%
Nitrates (NO3-N)	380.00	12	225.00	635.00 µg/l	7301 tonnes	YES	%
Orthoph. (PO4-P)	5.50	12	2.00	23.00 µg/l	102 tonnes	YES	%
Total N	573.75	12	385.00	910.00 µg/l	11470 tonnes	YES	%
Total P	13.75	12	7.00	38.00 µg/l	279 tonnes	YES	%
Susp. Part. Matter	7.10	12	1.85	29.70 mg/l	148111 tonnes	YES	%
TOC	3.70	1	3.70	3.70 mg/l	77545 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.2 MAIN RIVERINE INPUTS 1994 (2) Drammenselva

Total volume: 27353 1000 m³/day Long term average flow (LTA): 26743 1000 m³/day
 Minimum flow: 10109 1000 m³/day LTA period : 1961 to 1990
 Maximum flow: 75773 1000 m³/day

	Number of meas.	Mean	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.02	12	0.00	0.07 µg/l	0.23 tonnes	YES	%
Cadmium **	0.03	12	0.01	0.07 µg/l	0.25 tonnes		%
Mercury *	0.54	12	0.00	1.50 ng/l	5.24 kg	NO	%
Mercury **	1.13	12	1.00	1.50 ng/l	11.34 kg		%
Copper	0.85	12	0.30	1.40 µg/l	8.67 tonnes	YES	%
Zinc	3.24	12	2.00	5.70 µg/l	33.89 tonnes	YES	%
Lead	0.29	12	0.02	1.61 µg/l	2.71 tonnes	YES	%
Arsenic	0.17	4	0.08	0.37 µg/l	1.70 tonnes	YES	%
Total Cr-T	0.88	4	0.45	1.20 µg/l	8.85 tonnes	YES	%
Ni	0.59	4	0.45	0.91 µg/l	5.98 tonnes	YES	%
PCBs *		3		ng/l	6.19 kg	NO	%
PCBs **		3		ng/l	7.51 kg		%
gamma-HCH (lindane)	0.68	3	0.57	0.81 ng/l	6.77 kg	YES	%
Ammonia (NH4-N)	16.25	12	8.00	27.00 µg/l	151.64 tonnes	YES	%
Nitrates (NO3-N)	331.25	12	160.00	870.00 µg/l	3620.05 tonnes	YES	%
Orthoph. (PO4-P)	1.71	12	0.50	5.00 µg/l	16.95 tonnes	YES	%
Total N	512.92	12	375.00	1080.00 µg/l	5414.38 tonnes	YES	%
Total P	5.58	12	3.00	12.00 µg/l	58.95 tonnes	YES	%
Susp. Part. Matter	2.58	12	0.82	5.44 mg/l	27339 tonnes	YES	%
TOC	3.10	10	2.40	4.50 mg/l	31568 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.3 MAIN RIVERINE INPUTS 1994 (3) Numedalslågen

Total volume: 10644 1000 m3/day Long term average flow (LTA): 10082 1000 m3/day
 Minimum flow: 3093 1000 m3/day LTA period : 1961 to 1990
 Maximum flow: 57629 1000 m3/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.04	12	0.00	0.13 µg/l	0.17 tonnes	YES	%
Cadmium **	0.05	12	0.01	0.13 µg/l	0.18 tonnes	_____	%
Mercury *	4.71	12	0.00	25.50 ng/l	17.60 kg	YES	%
Mercury **	4.88	12	1.00	25.50 ng/l	18.04 kg	_____	%
Copper	1.04	12	0.30	3.81 µg/l	3.90 tonnes	YES	%
Zinc	5.42	12	1.70	11.60 µg/l	22.83 tonnes	YES	%
Lead	0.52	12	0.11	2.13 µg/l	2.01 tonnes	YES	%
Arsenic	0.19	1	0.19	0.19 µg/l	0.74 tonnes	YES	%
Total Cr-T	0.85	1	0.85	0.85 µg/l	3.30 tonnes	YES	%
Ni	0.62	1	0.62	0.62 µg/l	2.41 tonnes	YES	%
PCBs *		3		ng/l	3.86 kg	NO	%
PCBs **		3		ng/l	4.33 kg	_____	%
gamma-HCH (lindane)	0.44	3	0.30	0.64 ng/l	1.88 kg	YES	%
Ammonia (NH4-N)	28.08	12	5.00	54.00 µg/l	90.70 tonnes	YES	%
Nitrates (NO3-N)	250.17	12	59.00	730.00 µg/l	1030.55 tonnes	YES	%
Orthoph. (PO4-P)	3.13	12	0.50	7.00 µg/l	13.14 tonnes	YES	%
Total N	430.83	12	230.00	790.00 µg/l	1718.74 tonnes	YES	%
Total P	7.67	12	4.00	14.00 µg/l	32.95 tonnes	YES	%
Susp. Part. Matter	4.07	12	1.79	10.80 mg/l	17937 tonnes	YES	%
TOC	3.40	1	3.40	3.40 mg/l	13209 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.4 MAIN RIVERINE INPUTS 1994 (4) Skienselva

Total volume: 22438 1000 m³/day Long term average flow (LTA): 22611 1000 m³/day
 Minimum flow: 4320 1000 m³/day LTA period : 1961 to 1990
 Maximum flow: 49853 1000 m³/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.04	12	0.00	0.09 µg/l	0.33 tonnes	YES	%
Cadmium **	0.04	12	0.01	0.09 µg/l	0.33 tonnes		%
Mercury *	0.50	12	0.00	1.50 ng/l	3.98 kg	NO	%
Mercury **	1.08	12	1.00	1.50 ng/l	8.86 kg		%
Copper	0.55	12	0.30	1.20 µg/l	4.87 tonnes	YES	%
Zinc	2.55	12	1.91	3.60 µg/l	20.44 tonnes	YES	%
Lead *	0.17	12	0.00	1.01 µg/l	1.19 tonnes	YES	%
Lead **	0.17	12	0.02	1.01 µg/l	1.23 tonnes	YES	%
Arsenic	0.12	1	0.12	0.12 µg/l	0.98 tonnes	YES	%
Total Cr-T	0.57	1	0.57	0.57 µg/l	4.67 tonnes	YES	%
Ni	0.36	1	0.36	0.36 µg/l	2.95 tonnes	YES	%
PCBs *		3		ng/l	6.98 kg	NO	%
PCBs **		3		ng/l	7.98 kg		%
gamma-HCH (lindane)	1.05	3	0.83	1.31 ng/l	8.91 kg	YES	%
Ammonia (NH4-N)	15.67	12	6.00	39.00 µg/l	118.47 tonnes	YES	%
Nitrates (NO ₃ -N)	226.67	12	180.00	285.00 µg/l	1895.19 tonnes	YES	%
Orthoph. (PO ₄ -P)	0.88	12	0.50	2.00 µg/l	7.14 tonnes	YES	%
Total N	335.83	12	240.00	430.00 µg/l	2763.10 tonnes	YES	%
Total P	3.17	12	2.00	7.00 µg/l	26.95 tonnes	YES	%
Susp. Part. Matter	1.08	12	0.12	3.09 mg/l	9927 tonnes	YES	%
TOC	2.20	1	2.20	2.20 mg/l	18018 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.5 MAIN RIVERINE INPUTS 1994 (5) Otra

Total volume: 13734 1000 m3/day Long term average flow (LTA): 12841 1000 m3/day
 Minimum flow: 4398 1000 m3/day LTA period : 1961 to 1990
 Maximum flow: 39139 1000 m3/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.04	12	0.00	0.11 µg/l	0.17 tonnes	YES	%
Cadmium **	0.04	12	0.01	0.11 µg/l	0.18 tonnes	_____	%
Mercury *	0.42	12	0.00	1.00 ng/l	1.65 kg	NO	%
Mercury **	1.00	12	1.00	1.00 ng/l	5.01 kg	_____	%
Copper	0.50	12	0.10	0.90 µg/l	2.48 tonnes	YES	%
Zinc	5.04	12	3.61	7.50 µg/l	27.15 tonnes	YES	%
Lead	0.42	12	0.20	0.72 µg/l	2.23 tonnes	YES	%
Arsenic	0.17	4	0.09	0.27 µg/l	0.86 tonnes	YES	%
Total Cr-T	1.20	4	0.59	1.87 µg/l	5.26 tonnes	YES	%
Ni	0.95	4	0.51	1.48 µg/l	5.78 tonnes	YES	%
PCBs *		3		ng/l	0.76 kg	NO	%
PCBs **		3		ng/l	1.71 kg	_____	%
gamma-HCH (lindane)	0.83	3	0.55	1.13 ng/l	3.20 kg	YES	%
Ammonia (NH4-N)	13.33	12	5.00	30.00 µg/l	85.53 tonnes	YES	%
Nitrates (NO3-N)	143.58	12	87.00	225.00 µg/l	806.70 tonnes	YES	%
Orthoph. (PO4-P)	1.27	12	0.50	6.00 µg/l	12.83 tonnes	YES	%
Total N	282.92	12	225.00	360.00 µg/l	1495.06 tonnes	YES	%
Total P	5.25	12	3.00	9.00 µg/l	31.46 tonnes	YES	%
Susp. Part. Matter	3.76	12	0.67	21.10 mg/l	22915 tonnes	YES	%
TOC	3.26	12	2.30	4.70 mg/l	15014 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.6 MAIN RIVERINE INPUTS 1994 (6) Orreelva

Total volume: 282 1000 m3/day Long term average flow (LTA): 333 1000 m3/day
 Minimum flow: 29 1000 m3/day LTA period : 1961 to 1990
 Maximum flow: 782 1000 m3/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.06	12	0.00	0.32 µg/l	0.01 tonnes	YES	%
Cadmium **	0.06	12	0.01	0.32 µg/l	0.01 tonnes		%
Mercury *	0.50	12	0.00	5.00 ng/l	0.04 kg	NO	%
Mercury **	1.33	12	1.00	5.00 ng/l	0.12 kg		%
Copper	1.37	12	0.60	2.90 µg/l	0.14 tonnes	YES	%
Zinc	2.55	12	0.50	10.20 µg/l	0.36 tonnes	YES	%
Lead	0.35	12	0.10	0.95 µg/l	0.03 tonnes	YES	%
Arsenic	0.27	4	0.10	0.40 µg/l	0.03 tonnes	YES	%
Total Cr-T	2.58	4	1.13	4.64 µg/l	0.27 tonnes	YES	%
Ni	2.95	4	1.60	6.68 µg/l	0.31 tonnes	YES	%
PCBs *		3		ng/l	0.00 kg	NO	%
PCBs **		3		ng/l	0.02 kg		%
gamma-HCH (lindane)	1.29	3	0.46	2.94 ng/l	0.08 kg	YES	%
Ammonia (NH4-N)	45.08	12	7.00	120.00 µg/l	5.11 tonnes	YES	%
Nitrates (NO3-N)	757.92	12	1.00	1825.00 µg/l	102.15 tonnes	YES	%
Orthoph. (PO4-P)	9.25	12	2.00	32.00 µg/l	1.13 tonnes	YES	%
Total N	1352.1	12	610.00	2425.00 µg/l	163.58 tonnes	YES	%
Total P	38.92	12	25.00	60.00 µg/l	4.36 tonnes	YES	%
Susp. Part. Matter	4.91	12	2.84	7.79 mg/l	548 tonnes	YES	%
TOC	5.50	1	5.50	5.50 mg/l	567 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.7 MAIN RIVERINE INPUTS 1994 (7) Suldalslågen

Total volume: 4562 1000 m³/day Long term average flow (LTA): 7422 1000 m³/day
 Minimum flow: 1123 1000 m³/day LTA period : 1961 to 1990
 Maximum flow: 21168 1000 m³/day

	Number of meas.	Mean	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium	0.02	4	0.01	0.03 µg/l	0.04 tonnes	YES	%
Mercury *	0.50	4	0.00	2.00 ng/l	0.14 kg	NO	%
Mercury **	1.25	4	1.00	2.00 ng/l	1.74 kg		%
Copper	0.34	4	0.11	0.60 µg/l	0.64 tonnes	YES	%
Zinc	2.00	4	1.59	2.50 µg/l	3.56 tonnes	YES	%
Lead	0.14	4	0.05	0.27 µg/l	0.35 tonnes	YES	%
Arsenic	0.09	4	0.03	0.13 µg/l	0.12 tonnes	YES	%
Total Cr-T	1.36	4	0.54	2.26 µg/l	2.39 tonnes	YES	%
Ni	0.65	4	0.19	1.36 µg/l	1.67 tonnes	YES	%
PCBs *		2		ng/l	0.00 kg	NO	%
PCBs **		2		ng/l	0.35 kg		%
gamma-HCH (lindane)	0.83	2	0.81	0.84 ng/l	1.39 kg	YES	%
Ammonia (NH4-N)	7.25	4	5.00	13.00 µg/l	11.57 tonnes	YES	%
Nitrates (NO ₃ -N)	181.25	4	160.00	235.00 µg/l	275 tonnes	YES	%
Orthoph. (PO ₄ -P)	0.50	4	0.50	0.50 µg/l	0.83 tonnes	YES	%
Total N	227.50	4	190.00	275.00 µg/l	342 tonnes	YES	%
Total P	1.75	4	1.00	3.00 µg/l	2.57 tonnes	YES	%
Susp. Part. Matter	0.68	4	0.22	0.95 mg/l	1423 tonnes	YES	%
TOC	0.50	1	0.50	0.50 mg/l	833 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.8 MAIN RIVERINE INPUTS 1994 (8) Orkla

Total volume:	3652	1000 m3/day	Long term average flow (LTA):	5374	1000 m3/day		
Minimum flow:	1581	1000 m3/day	LTA period :	1961	to 1990		
Maximum flow:	17885	1000 m3/day					
	Number of meas.	Mean	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium	0.08	12	0.02	0.14 µg/l	0.10 tonnes	YES	%
Mercury *	0.46	12	0.00	1.50 ng/l	0.70 kg	NO	%
Mercury **	1.04	12	1.00	1.50 ng/l	1.38 kg		%
Copper	7.84	12	4.01	12.40 µg/l	10.83 tonnes	YES	%
Zinc	26.03	12	15.20	40.90 µg/l	36.44 tonnes	YES	%
Lead *	0.17	12	0.00	1.18 µg/l	0.26 tonnes	YES	%
Lead **	0.17	12	0.02	1.18 µg/l	0.27 tonnes		%
Arsenic	0.56	1	0.56	0.56 µg/l	0.75 tonnes	YES	%
Total Cr-T	1.15	1	1.15	1.15 µg/l	1.53 tonnes	YES	%
Ni	0.68	1	0.68	0.68 µg/l	0.91 tonnes	YES	%
PCBs *		3		ng/l	0.00 kg	NO	%
PCBs **		3		ng/l	0.28 kg		%
gamma-HCH (lindane)	0.19	3	0.15	0.23 ng/l	0.23 kg	YES	%
Ammonia (NH4-N)	9.67	12	5.00	19.00 µg/l	12.57 tonnes	YES	%
Nitrates (NO3-N)	215.50	12	100.00	395.00 µg/l	259 tonnes	YES	%
Orthoph. (PO4-P)	1.17	12	0.50	3.00 µg/l	1.46 tonnes	YES	%
Total N	331.75	12	211.00	435.00 µg/l	427 tonnes	YES	%
Total P	4.67	12	1.00	9.00 µg/l	6.37 tonnes	YES	%
Susp. Part. Matter	1.68	12	0.38	5.66 mg/l	2543 tonnes	YES	%
TOC	3.73	12	2.00	6.50 mg/l	4973 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.9 MAIN RIVERINE INPUTS 1994 (9) Vefsna

Total volume: 10992 1000 m³/day Long term average flow (LTA): 15620 1000 m³/day
 Minimum flow: 2843 1000 m³/day LTA period : 1961 to 1990
 Maximum flow: 79661 1000 m³/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.05	12	0.00	0.18 µg/l	0.15 tonnes	YES	%
Cadmium **	0.05	12	0.01	0.18 µg/l	0.16 tonnes		%
Mercury *	2.79	12	0.00	10.50 ng/l	5.89 kg	NO	%
Mercury **	3.13	12	1.00	10.50 ng/l	6.99 kg		%
Copper	5.03	12	1.20	10.35 µg/l	20.74 tonnes	YES	%
Zinc	9.54	12	1.20	19.51 µg/l	37.53 tonnes	YES	%
Lead	1.49	12	0.05	7.32 µg/l	3.59 tonnes	YES	%
Arsenic	0.27	4	0.11	0.36 µg/l	1.07 tonnes	YES	%
Total Cr-T	1.40	4	0.54	2.32 µg/l	5.60 tonnes	YES	%
Ni	1.00	4	0.37	1.97 µg/l	4.21 tonnes	YES	%
PCBs *		3		ng/l	0.43 kg	NO	%
PCBs **		3		ng/l	1.24 kg		%
gamma-HCH (lindane)	0.23	3	0.15	0.38 ng/l	1.16 kg	YES	%
Ammonia (NH4-N)	27.25	12	5.00	75.00 µg/l	97.26 tonnes	YES	%
Nitrates (NO ₃ -N)	108.33	12	39.00	325.00 µg/l	275.0 tonnes	YES	%
Orthoph. (PO ₄ -P)	2.67	12	0.50	5.00 µg/l	9.58 tonnes	YES	%
Total N	388.75	12	160.00	800.00 µg/l	1184 tonnes	YES	%
Total P	5.17	12	2.00	10.00 µg/l	22.48 tonnes	YES	%
Susp. Part. Matter	1.57	12	0.51	4.32 mg/l	8965 tonnes	YES	%
TOC	3.40	1	3.40	3.40 mg/l	13641 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

Table 4.10 MAIN RIVERINE INPUTS 1993 (10) Altaelva

Total volume: 5918 1000 m3/day Long term average flow (LTA): 7487 1000 m3/day
 Minimum flow: 2272 1000 m3/day LTA period : 1961 to 1990
 Maximum flow: 46829 1000 m3/day

	Mean	Number of meas.	Minimum concentr. during the year	Maximum concentr. during the year	Total quantity of substance discharged each year	Were 70 % of measurements above limit of detection ?	Precision of the estimate of the load
Cadmium *	0.03	4	0.00	0.06 µg/l	0.06 tonnes	YES	%
Cadmium **	0.04	4	0.01	0.06 µg/l	0.07 tonnes		%
Mercury *	1.38	4	0.00	2.50 ng/l	3.36 kg	NO	%
Mercury **	1.63	4	1.00	2.50 ng/l	3.80 kg		%
Copper	1.10	4	0.30	2.50 µg/l	2.82 tonnes	YES	%
Zinc	0.74	4	0.30	1.24 µg/l	1.49 tonnes	YES	%
Lead	0.09	4	0.05	0.19 µg/l	0.19 tonnes	YES	%
Arsenic	0.32	4	0.08	0.63 µg/l	0.62 tonnes	YES	%
Total Cr-T	1.20	4	0.15	2.12 µg/l	2.73 tonnes	YES	%
Ni	0.75	4	0.51	1.09 µg/l	1.74 tonnes	YES	%
PCBs *		2		ng/l	0.36 kg	NO	%
PCBs **		2		ng/l	0.65 kg		%
gamma-HCH (lindane)	0.05	2	0.05	0.05 ng/l	0.11 kg	YES	%
Ammonia (NH4-N)	6.00	4	5.00	9.00 µg/l	13.91 tonnes	YES	%
Nitrates (NO3-N)	48.50	4	23.00	84.00 µg/l	95.15 tonnes	YES	%
Orthoph. (PO4-P)	7.63	4	0.50	19.00 µg/l	14.84 tonnes	YES	%
Total N	178.75	4	155.00	210.00 µg/l	395.39 tonnes	YES	%
Total P	12.75	4	4.00	24.00 µg/l	26.78 tonnes	YES	%
Susp. Part. Matter	3.14	4	0.84	9.10 mg/l	8642 tonnes	YES	%
TOC	3.50	1	3.50	3.50 mg/l	7560 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

APPENDIX V : INPUTS FROM TRIBUTARY RIVERS 1994 (Paragraph 17 - 19) Page:

Table 5.1	Tributary rivers in the sub-areas	(1-5). The Skagerrak area	41
Table 5.2	Tributary rivers in the sub-areas	(6-7). Remain. North Sea	42
Table 5.3	Tributary rivers in the sub-areas	(8-9). The Norwegian Sea	43
Table 5.4	Tributary rivers in the sub-area	(10). The Barents Sea	44

Paragraph 17: Tributary rivers /.

Paragraph18: Measurements of calculation used - including information on the concentration upon which the measurement is based:

Appendix VIII and IX (1-10) and Paragraph 3.2 (Report A, 1991, 1992 - 1995)

Paragraph 19: Any other relevant information (e.g. proportion of substance discharged as insoluble material):

**Table 5.1 The Skagerrak Region. Inputs from tributary rivers 1994
in The Subareas (1-5).**

The Skagerrak Region with sub-areas: (1A) Glomma, (1B) Inner Oslofj., (2) Drammenselva,
(3) Numedalslågen, (4) Skienselva, (5) Otra

Sub-areas :	1A	1B	2	3	4	5	Were 70 % of measurements above the detection limit ?	Precision of the estimate of the load
Substance:								
Cd *	0.12	0.03	0.01	0.02	0.05	0.62 tonnes	YES	%
Cd **	0.12	0.03	0.01	0.02	0.05	0.62 tonnes		%
Hg *	0.63	0.74	0.24	0.10	1.08	9.51 kg	YES	%
Hg **	1.40	0.95	0.24	0.60	1.08	13.74 kg		%
Cu	1.0	0.6	0.2	0.8	0.6	6.0 tonnes	YES	%
Zn	8.6	2.3	2.0	8.8	8.2	73.1 tonnes	YES	%
Pb *	0.35	0.14	0.07	0.28	0.40	5.87 tonnes	YES	%
Pb **	0.35	0.14	0.07	0.28	0.40	5.87 tonnes		%
Arsenic *	0.35	0.13	0.05	0.16	0.34	2.42 tonnes	YES	%
Arsenic **	0.35	0.13	0.05	0.16	0.34	2.42 tonnes		%
Cr-T *	2.10	0.51	0.29	0.65	1.21	7.36 tonnes	YES	%
Cr-T **	2.10	0.54	0.29	0.65	1.21	7.36 tonnes		%
Ni *	1.11	0.54	0.28	0.63	0.31	1.95 tonnes	YES	%
Ni **	1.11	0.54	0.28	0.63	0.31	1.95 tonnes		%
PCBs *	0.00	0.13	0.00	0.00	0.00	0.00 kg	NO	%
PCBs **	0.24	0.20	0.05	0.12	0.23	2.16 kg		%
gamma-HCH	0.79	0.26	0.06	0.21	1.08	9.08 kg	YES	%
NH4-N	26	36	10	24	42	407 tonnes	YES	%
NO3-N	1001	447	210	599	244	2472 tonnes	YES	%
PO4-P	6	6	9	10	1	6 tonnes	YES	%
Total N	1205	603	265	903	433	4462 tonnes	YES	%
Total P	21	18	14	16	4	46 tonnes	YES	%
S.P.M.	3465	4598	3431	4808	1126	14029 tonnes	YES	%
TOC	8093	2586	821	2758	3680	32860 tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

**Table 5.2 The remaining North Sea Region. Inputs from tributary rivers 1994
in The Subareas (6-7).**

The remaining North Sea Region with sub-areas: (6) Orreelva, (7) Suldalslågen

Total quantity of substance discharged per year:			Were 70 % of measurements above the detection limit ?	Precision of the estimate of the load
	Sub-areas :	6 7		
Substance:				
Cd *	0.39	0.83	tonnes YES	%
Cd **	0.40	0.89	tonnes	%
Hg *	5.40	26.56	kg NO	%
Hg **	15.05	43.71	kg	%
Cu	4.1	9.9	tonnes YES	%
Zn	65.5	81.0	tonnes YES	%
Pb *	6.52	3.12	tonnes YES	%
Pb **	6.52	3.19	tonnes	%
Arsenic *	3.30	2.15	tonnes YES	%
Arsenic **	3.30	2.16	tonnes	%
Cr-T *	15.86	23.45	tonnes YES	%
Cr-T **	15.86	23.45	tonnes	%
Ni *	4.13	2.17	tonnes YES	%
Ni **	4.13	2.21	tonnes	%
PCBs *	0.00	0.00	kg NO	%
PCBs **	2.87	6.81	kg	%
gamma-HCH	8.00	8.40	kg YES	%
NH4-N	296	227	tonnes YES	%
NO3-N	3475	4199	tonnes YES	%
PO4-P	22	33	tonnes YES	%
Total N	5052	6786	tonnes YES	%
Total P	75	113	tonnes YES	%
S.P.M.	15627	24353	tonnes YES	%
TOC	38149	31468	tonnes YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

**Table 5.3 The Norwegian Sea Region. Inputs from tributary rivers 1994
in The Subareas (8-9).**

The Norwegian Sea Region with sub-areas: (8) Orkla, (9) Vefsna

Sub-areas :	8	9	Were 70 % of measurements above the detection limit ?		Precision of the estimate of the load
			tonnes	YES	
Total quantity of substance discharged per year:					
Cd *	1.45	0.35	tonnes	YES	%
Cd **	1.50	0.43	tonnes		%
Hg *	54.64	7.95	kg	NO	%
Hg **	64.26	27.89	kg		%
Cu	26.2	27.3	tonnes	YES	%
Zn	59.4	50.2	tonnes	YES	%
Pb *	8.84	13.75	tonnes	YES	%
Pb **	8.88	13.75	tonnes		%
Arsenic *	4.07		tonnes	YES	%
Arsenic **	4.08		tonnes		%
Cr-T *	42.58	54.08	tonnes	YES	%
Cr-T **	42.58	54.08	tonnes		%
Ni *	10.65	4.68	tonnes	YES	%
Ni **	10.67	4.68	tonnes		%
PCBs *	0.41	0.00	kg	NO	%
PCBs **	7.51	5.74	kg		%
gamma-HCH	13.64	8.25	kg	YES	%
NH4-N	256	140	tonnes	YES	%
NO3-N	3388	1096	tonnes	YES	%
PO4-P	55	28	tonnes	YES	%
Total N	7339	2964	tonnes	YES	%
Total P	181	91	tonnes	YES	%
S.P.M.	55226	58436	tonnes	YES	%
TOC	22867	4376	tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

**Table 5.4 The Barents Region. Inputs from tributary rivers 1994
in The Subarea (10).**

The Barents Sea Region with sub-area: (10) Alta

		Were 70 % of measurements above the detection limit ?		Precision of the estimate of the load
Total quantity of substance discharged per year:				
Sub-area :	10	tonnes	YES	%
Substance:		tonnes		%
Cd *	0.16	kg	NO	%
Cd **	0.23	kg		%
Hg *	25.47	tonnes	YES	%
Hg **	29.89	tonnes	YES	%
Cu	18.8	tonnes	YES	%
Zn	20.1	tonnes	YES	%
Pb *	5.41	tonnes	YES	%
Pb **	5.41	tonnes	YES	%
Arsenic *	3.07	tonnes	YES	%
Arsenic **	3.07	tonnes	YES	%
Cr-T *	40.53	tonnes	YES	%
Cr-T **	40.53	tonnes	YES	%
Ni *	62.44	tonnes	YES	%
Ni **	62.44	tonnes	YES	%
PCBs *	0.00	kg	NO	%
PCBs **	3.38	kg		%
gamma-HCH	3.41	kg	YES	%
NH4-N	255	tonnes	YES	%
NO3-N	202	tonnes	YES	%
PO4-P	25	tonnes	YES	%
Total N	2571	tonnes	YES	%
Total P	89	tonnes	YES	%
S.P.M.	30967	tonnes	YES	%
TOC	24351	tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

APPENDIX VI : OTHER INPUTS 1994 (Paragraph 20)	Page:
---	--------------

Table 6.1	Nutrients from "Down Stream Areas" of main and tributary rivers and rivers not monitored	46
-----------	--	----

Paragraph 20: Any available information on discharge through urban run-off - storm water overflow - polder effluents etc.:

"Background" is runoff from forested areas and highlands, including effect of acid precipitation

Agriculture runoff:

- "Area" is runoff from arable land
- "Point"-sources are drainage from silos, manures etc.

Paragraph 3.3 (Report A, 1991, 1992 - 1995)

**Table 6.1 Nutrients from "Down Stream Areas" of main and tributary rivers
and rivers not monitored 1994.**

Direct runoff of P and N :

Sub-areas:		Back-ground tonnes	Agriculture		
			Area tonnes	Point tonnes	Sum tonnes
1 Glomma	P	15.4	10.7	0.4	26.5
	N	266.4	518.9	6.4	791.6
	PO4-P	3.1	3.2	0.3	6.6
	NO3-N	159.8	363.2	0.8	523.8
	NH4-N	13.3	36.3	5.1	54.7
1 Inner Oslofjord	P	2.3	2.1	0.3	4.6
	N	27.2	90.0	2.7	119.8
	PO4-P	0.5	0.6	0.2	1.3
	NO3-N	16.3	63.0	0.3	79.6
	NH4-N	1.4	6.3	2.1	9.8
2 Drammenselva	P	1.4	2.1	0.0	3.5
	N	47.3	69.8	0.7	117.8
	PO4-P	0.3	0.6	0.0	0.9
	NO3-N	28.4	48.9	0.1	77.3
	NH4-N	2.4	4.9	0.5	7.8
3 Numedalslågen	P	4.9	10.5	0.2	15.5
	N	128.8	444.6	1.8	575.2
	PO4-P	1.0	3.1	0.1	4.2
	NO3-N	77.3	311.2	0.2	388.7
	NH4-N	6.4	31.1	1.4	39.0
4 Skienselva	P	6.8	2.0	0.1	8.9
	N	271.1	91.6	1.5	364.1
	PO4-P	1.4	0.6	0.1	2.0
	NO3-N	162.6	64.1	0.2	226.9
	NH4-N	13.6	6.4	1.2	21.1
5 Otra	P	7.0	3.6	0.3	10.9
	N	316.1	104.7	3.4	424.2
	PO4-P	1.4	1.1	0.2	2.7
	NO3-N	189.7	73.3	0.4	263.4
	NH4-N	15.8	7.3	2.7	25.9
6 Orreelva	P	23.0	43.6	4.2	70.9
	N	1238.6	1312.5	51.5	2602.6
	PO4-P	4.6	13.1	2.9	20.6
	NO3-N	743.2	918.8	6.2	1668.1
	NH4-N	61.9	91.9	41.2	195.0
7 Suldalslågen	P	58.0	62.2	7.2	127.5
	N	4284.0	1300.8	98.6	5683.4
	PO4-P	11.6	18.7	5.0	35.3
	NO3-N	2570.4	910.6	11.8	3492.8
	NH4-N	214.2	91.1	78.9	384.1
8 Orkla	P	131.1	141.2	14.1	286.5
	N	3501.0	3456.7	191.9	7149.6
	PO4-P	26.2	42.4	9.8	78.3
	NO3-N	2100.6	2419.7	23.0	4543.3
	NH4-N	175.0	242.0	153.6	570.6
9 Vefsna	P	83.6	32.5	7.2	123.3
	N	1920.5	829.7	104.6	2854.7
	PO4-P	16.7	9.8	5.0	31.5
	NO3-N	1152.3	580.8	12.5	1745.6
	NH4-N	96.0	58.1	83.7	237.8
10 Altaelva	P	86.1	1.6	0.6	88.2
	N	1617.9	54.9	7.9	1680.7
	PO4-P	17.2	0.5	0.4	18.1
	NO3-N	970.8	38.4	0.9	1010.1
	NH4-N	80.9	3.8	6.3	91.0

SUM	P	766	tonnes
SUM	N	22364	tonnes
SUM	PO4-P	201	tonnes
SUM	NO3-N	14020	tonnes
SUM	NH4-N	1637	tonnes

**APPENDIX VII : MAIN RIVERS 1 - 10. MEASURED CONCENTRATIONS
1994****Page:**

Table 7.1	Glomma	48
Table 7.2	Drammenselva	49
Table 7.3	Numedalslågen	50
Table 7.4	Skienselva	51
Table 7.5	Otra	52
Table 7.6	Orreelva	53
Table 7.7	Suldalslågen	54
Table 7.8	Orkla	55
Table 7.9	Vefsna	56
Table 7.10	Alta	57

Table 7.1 Measured concentrations - 1994

Watercourse :	Giomma												Min : 282 cbm/s													
	Annual flow				20958 mill. cbm				Max: 1890 cbm/s				282 cbm/s				Cr-T									
	Drainage area.				41218 sq.km				IUPAC NOS																	
Date	Q m3/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	Gam/HCH ng/l	HCH ng/l	PCB 28 ng/l	PCB 52 ng/l	101 ng/l	118 ng/l	138 ng/l	153 ng/l	180 ng/l	As µg/l	Cr-T µg/l	Ni µg/l
940121	451.9	4.96	7	3.0	555	40	1.13	6.37	0.10	0.54	2.57	<1.0	2.0	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.92		
940215	384.3	4.86	15	3	540	48	1.32	4.69	0.07	0.42	1.85	2.0	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.92			
940314	286.9	5.21	26	15.0	700	460	60	1.49	5.93	0.08	0.47	12.4	8.0	2.0	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.92		
940412	664.1	5.31	38	23	910	635	52	2.07	4.30	0.05	0.58	29.7	3.0	11.5	0.53	<0.03	0.17	0.33	0.3	0.3	0.22	0.07	0.07	0.07		
940510	1405	3.48	17	5.0	475	230	14	2.10	7.80	0.02	0.59	8.08	3.0	0.53	<0.03	0.17	0.33	0.3	0.3	0.22	0.07	0.07	0.07	0.07		
940610	1230	3.82	9	2.0	435	225	24	2.30	24.1	0.44	2.05	3.43	3.0	1.5	0.53	<0.03	0.17	0.33	0.3	0.3	0.22	0.07	0.07	0.07		
940714	950.0	4.31	7	2.0	395	235	11	1.60	2.30	0.03	0.15	3.89	<1.0	1.0	0.53	<0.03	0.17	0.33	0.3	0.3	0.22	0.07	0.07	0.07		
940818	717.4	22.1	8	2.0	385	235	16	1.60	17.9	0.17	0.81	4.55	2.0	1.5	0.51	<0.03	0.04	0.06	0.06	0.06	0.07	0.05	<0.03	0.40		
940915	871.1	4.47	13	2.0	740	530	20	2.00	7.30	0.08	0.62	7.63	3.5	2.0	0.51	<0.03	0.04	0.06	0.06	0.06	0.07	0.05	<0.03	0.40		
941012	569.8	4.32	8	3.0	480	320	27	1.20	4.50	0.02	0.08	2.81	2.5	1.5	0.51	<0.03	0.04	0.06	0.06	0.06	0.07	0.05	<0.03	0.40		
941110	461.2	4.68	8	3.0	565	425	26	1.20	5.90	<0.01	2.05	2.59	1.5	1.0	0.33	0.33	0.75	3.7	1.0	0.40	1.31	0.92	0.92			
941205	492.9	5.28	9	3	705	540	43	1.70	10.8	0.13	0.33	5.75	3.7	1.0	0.33	0.33	0.75	3.7	1.0	0.40	1.31	0.92	0.92			
Min:	286.9	3.48	7	2	385	225	11	1.13	2.3	0.01	0.15	1.85	3.7	1	0.51	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.92		
Max:	1405	22.1	38	23	910	635	60	2.3	24.1	0.44	2.08	29.7	3.7	11.5	0.70	0.03	0.17	0.33	0.3	0.3	0.22	0.07	0.4	0.92		
Aver.:	707.1	6.07	13.75	5.50	574	380	31.75	1.64	8.49	0.10	0.89	7.10	3.70	3.33	0.58	0.03	0.08	0.14	0.13	0.13	0.10	0.04	0.40	0.92		
St.dev.:	346.3	5.08	9.47	6.59	160	140	16.28	0.40	6.35	0.12	0.72	7.74	3.20	3.20	0.10	0.00	0.08	0.17	0.15	0.15	0.10	0.02	0.40	0.92		
Numb.:	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	3	3	1		

Table 7.3 Measured concentrations - 1994

Date	Q m3/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	HCH ng/l	Gamn ng/l	IUPAC NOS			
																	Min : 180 µg/l	Max : 180 µg/l	As µg/l	Cr-T µg/l
940117	97.8	3.35	6	4.0	400	235	43	0.47	3.47	0.04	0.19	3.08	1.5	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	
940217	80.4	2.65	4	3.0	335	140	30	0.39	3.02	0.11	0.15	1.79	7.5	7.5	7.5					
940316	69.2	4.12	8	5.0	590	435	50	3.81	11.6	0.06	2.13	4.28	10.80	1.5	0.20	10.80	1.5			
940417	154.7	4.21	14	7.0	790	730	26	1.13	7.36	0.13	0.20	0.20	0.20	0.20	0.20	0.20	0.20			
940509	254.4	2.25	8	3.0	340	160	9	0.70	6.80	0.02	0.39	4.84	10.0	2.0	0.64	<0.03	0.22	0.49	0.45	0.31
940607	132.1	1.89	9	3.0	230	109	5	1.40	9.40	0.02	1.77	3.10	2.0	2.0	0.64	0.22	0.49	0.43	0.45	0.31
940718	73.3	2.51	4	0.5	230	59	9	0.80	1.70	<0.01	0.15	2.71	<1.0	<1.0	<1.0	<1.0	<1.0			
940818	85.7	2.77	9	2.0	355	99	21	1.00	3.90	0.01	0.29	3.28	1.5	1.5	1.5	1.5	1.5			
940915	169.9	3.31	13	4.0	600	340	13	1.40	7.10	0.04	0.52	6.45	3.5	3.5	3.5	3.5	3.5			
941017	54.9	3.04	5	2.0	405	185	54	0.80	4.60	<0.01	0.23	4.12	25.5	25.5	0.38	<0.03	<0.03	<0.03	<0.03	
941108	103.7	3.14	6	2.0	460	245	34	0.30	2.90	<0.01	0.11	2.24	1.5	1.5	1.5	1.5	1.5			
941207	80.4	3.30	6	2.0	435	265	43	0.30	3.20	0.08	0.15	2.19	3.4	3.4	3.4	3.4	3.4			
Min:	54.93	1.89	4	0.5	230	59	5	0.3	1.7	0.01	0.11	1.79	3.4	1	0.30	0.03	0.03	0.03	0.19	0.85
Max:	254.4	4.21	14	7	790	730	54	3.81	11.6	0.13	0.21	25.5	10.8	10.8	0.64	0.03	0.22	0.49	0.45	0.31
Aver.:	113.0	3.05	7.67	3.13	431	250	28.08	1.04	5.42	0.05	0.52	4.07	3.40	4.07	0.44	0.03	0.09	0.18	0.16	0.12
S.dev.:	56.71	0.69	3.23	1.71	162	185	17.00	0.95	3.02	0.04	0.68	2.49	7.09	7.09	0.18	0.00	0.11	0.27	0.23	0.16
Numb.:	12	12	12	12	12	12	12	12	12	12	12	12	1	12	3	3	3	3	3	1

Table 7.4 Measured concentrations - 1994

Date	Q m3/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	Gamm ng/l	HCH ng/l	PCB ng/l	IUPAC NOS			
																		Min : 50 cbm/s			
Drainage area.	Watercourse :	Skienselva	Annual flow	8190 mill. cbm	10348 sq.km																
940120	231.0	2.25	2	1.0	335	240	11	0.32	2.27	0.08	0.06	0.36	1.5	<1.0	1.01	<0.03	<0.03	<0.03	<0.03	<0.03	
940216	200.0	2.13	2	1.0	240	230	9	0.30	1.91	0.04	0.05	0.32	<0.02	0.12	<1.0	1.64	<1.0	<0.03	<0.03	<0.03	
940316	190.0	2.28	2	1.0	330	240	9	0.69	2.10	0.05	<0.02	0.04	1.64	<1.0	0.02	0.82	<1.0	<0.03	<0.03	<0.03	
940412	332.0	2.33	3	2.0	380	285	9	0.31	2.10	0.02	0.04	1.64	<1.0	0.03	<0.02	0.82	<1.0	<0.03	<0.03	<0.03	
940519	473.0	2.30	3	0.5	350	255	11	1.10	2.10	0.03	<0.02	0.02	1.64	<1.0	0.03	0.82	<1.0	<0.03	<0.03	<0.03	
940613	262.0	2.13	4	1.0	335	225	6	1.20	2.70	0.02	0.11	0.88	<1.0	1.31	0.06	0.26	0.42	0.43	0.47	0.29	
940713	99.0	2.17	2	0.5	320	195	16	0.50	2.90	0.04	0.15	1.76	<1.0	0.02	0.83	<1.0	1.0	<0.03	<0.03	<0.03	
940818	137.0	2.08	2	0.5	300	180	16	0.50	2.20	0.02	0.12	0.83	<1.0	0.02	0.93	1.0	<0.03	<0.03	<0.03	<0.03	
940914	472.0	2.04	3	0.5	310	200	13	0.50	2.60	0.04	0.09	3.09	<1.0	0.04	0.93	1.0	<0.03	<0.03	<0.03	<0.03	
941014	165.0	2.07	4	1.0	430	215	39	0.40	3.60	0.08	0.24	1.69	<1.0	0.83	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	
941109	272.0	2.03	7	0.5	350	215	19	0.30	2.90	<0.01	0.11	0.66	1.5	<1.0	0.77	2.2	<1.0	<0.03	<0.03	<0.03	
941208	235.0	2.31	4	1.0	350	240	30	0.51	3.21	0.09	1.01	0.77	<1.0	0.02	0.83	1.0	<0.03	<0.03	<0.03	<0.03	
Min:	99	2.03	2	0.5	240	180	6	0.3	1.91	0.01	0.02	0.12	2.2	1	0.83	0.03	0.03	0.03	0.03	0.03	
Max:	473	2.33	7	2	430	285	39	1.2	3.6	0.09	1.01	3.09	2.2	1.5	1.31	0.06	0.26	0.42	0.43	0.47	
Aver.:	255.7	2.18	3.17	0.88	336	227	15.67	0.55	2.55	0.04	0.17	1.08	2.20	1.08	1.05	0.04	0.11	0.16	0.18	0.12	
St.dev.:	118.9	0.11	1.47	0.43	46	28	9.72	0.30	0.52	0.03	0.27	0.83	1.2	1.2	0.19	0.24	0.02	0.13	0.23	0.25	
Numb.:	12	12	12	12	12	12	12	12	12	12	12	12	12	12	1	12	3	3	3	1	

Table 7.5 Measured concentrations - 1994

Watercourse :	Otra										50.9 cbm/s									
	Annual flow					5013 mill. cbm					Min : 50.9 cbm/s									
	Drainage area.					3730 sq.km					Max: 453 cbm/s									
Date	Q m3/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	Gamm ng/l	PCB (The following Congeners)	IUPAC NOS	As µg/l	Cr-T µg/l	Ni µg/l
940119	168.8	2.49	3	1.0	265	149	18	0.42	4.55	0.06	0.28	0.67	2.4	<1.0			0.27	1.33	0.52	
940215	136.6	2.37	3	1.0	225	138	12	0.47	3.61	<0.01	0.20	1.41	2.3	<1.0			0.09	1.87	0.51	
940316	997.5	2.79	9	6.0	320	180	21	0.47	6.11	0.02	0.41	5.53	2.5	<1.0	0.55	<0.03	<0.03	<0.03	1.27	
940413	239.5	3.08	6	1.0	360	225	30	0.52	5.24	<0.01	0.49	1.79	3.5	<1.0			0.17	1.00	1.48	
940510	422.0	2.31	4	0.5	285	155	17	0.50	4.70	0.03	0.51	2.07	2.6	1.0			0.16	0.59	1.48	
940613	137.4	2.45	4	0.5	250	129	5	0.30	3.70	0.03	0.22	1.71	2.7	<1.0	1.13	<0.03	0.16	0.21	0.08	
940713	84.4	2.48	5	0.5	265	103	5	0.60	4.10	0.05	0.37	2.78	3.8	1.0						
940817	52.9	2.66	7	2.0	265	87	5	0.70	5.30	0.04	0.43	2.33	3.9	<1.0						
940914	216.2	2.34	6	0.5	320	124	8	0.90	7.50	0.08	0.72	2.86	4.7	1.0						
941012	109.5	2.19	4	0.5	255	113	13	0.10	4.20	0.02	0.29	1.21	3.0	1.0						
941109	126.1	2.32	5	0.5	315	160	6	0.60	6.10	0.03	0.51	1.64	3.9	1.0						
941207	219.4	2.25	7	1.2	270	160	20	0.40	5.40	0.11	0.64	21.1	3.8	<1.0						
Min:	52.88	2.19	3	0.5	225	87	5	0.1	3.61	0.01	0.2	0.67	2.3	1	0.55	0.03	0.03	0.03	0.51	
Max:	997.5	3.08	9	6	360	225	30	0.9	7.5	0.11	0.72	21.1	4.7	1	1.13	0.03	0.29	0.21	1.48	
Aver.:	242.5	2.48	5.25	1.27	283	144	13.33	0.50	5.04	0.04	0.42	3.76	3.26	1.00	0.83	0.03	0.07	0.12	0.95	
St.dev.:	256.4	0.25	1.82	1.56	38	37	8.03	0.20	1.14	0.03	0.16	5.60	0.77	0.00	0.29	0.00	0.08	0.15	0.50	
Numb.:	12	12	12	12	12	12	12	12	12	12	12	12	12	12	3	3	3	3	4	

Table 7.6 Measured concentrations - 1994
Watercourse: Orrelevå

Table 7.7 Measured concentrations - 1994
Watercourse : Suldalslägen

Annual flow		1665 mill. cbm		Min : 13 cbm/s		Max: 245 cbm/s		1457 sq.km		1665 mill. cbm		Min : 13 cbm/s		Max: 245 cbm/s		Drainage area.					
Date	Q m3/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	Hg ng/l	Gamm HCH ng/l	PCB (The following Congeners) IUPAC NOS ng/l	As ng/l	Cr-T µg/l	Ni µg/l	
940211	14.19	2.27	1	0.5	275	235	5	0.24	1.71	0.02	0.05	0.22	2.0	0.81	<0.03	<0.03	<0.03	0.13	1.21	0.19	
940629	196.1	1.85	1	0.5	190	160	6	0.40	2.20	0.03	0.27	0.95	<1.0	<1.0	<0.03	<0.03	<0.03	0.06	1.43	1.36	
940816	67.75	1.67	2	0.5	220	160	5	0.60	2.50	0.01	0.18	0.81	<1.0	<1.0	<0.03	<0.03	<0.03	0.12	2.26	0.62	
941010	57.23	5.65	3	0.5	225	170	13	0.11	1.59	0.01	0.07	0.74	0.5	<1.0	0.84	<0.03	<0.03	<0.03	0.03	0.54	0.43
Min:	14.19	1.67	1	0.5	190	160	5	0.11	1.59	0.01	0.05	0.22	0.5	1	0.81	0.03	0.03	0.03	0.03	0.54	0.19
Max:	196.1	5.65	3	0.5	275	235	13	0.6	2.5	0.03	0.27	0.95	0.5	2	0.84	0.03	0.03	0.03	0.03	0.13	1.36
Aver.:	83.81	2.86	1.75	0.50	228	181	7.25	0.34	2.00	0.02	0.14	0.68	0.50	1.25	0.83	0.03	0.03	0.03	0.03	0.09	0.65
Std.dev.:	78.34	1.88	0.96	0.00	35	36	3.86	0.21	0.43	0.01	0.10	0.32	0.50	0.50	1	4	2	2	0.05	0.71	0.50
Numb.:	4	4	4	4	4	4	4	4	4	4	4	4	4	4	1	4	2	2	2	4	4

Table 7.8 Measured concentrations - 1994

Date	Q m ³ /s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	HCH ng/l	Gamn ng/l	PCB (The following Congeners) ng/l	IUPAC NOS ng/l	As µg/l	Cr-T µg/l	Ni µg/l																				
Watercourse : Orkla						Annual flow						2872 sq.km						1333 mill. cbm																							
Drainage area						Min : 18.3 cbm/s						Max: 207 cbm/s						207 cbm/s																							
940124	33.69	6.10	3.0	3.0	300	170	12	4.46	20.9	0.04	0.40	1.07	2.0	<1.0																											
940223	20.28	7.50	2.0	1.0	365	265	13	4.01	15.2	0.04	0.04	0.38	2.4	<1.0																											
940319	20.28	7.40	4.3	1.7	285	260	9	4.45	18.2	0.14	<0.02	0.75	2.1	<1.0	0.23	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03																			
940412	21.69	9.30	9.0	2.0	435	193	16	12.4	35.3	0.07	0.02	2.31	5.3	<1.0																											
940516	50.42	6.30	7.0	1.0	300	150	6	12.0	39.9	0.12	1.18	5.66	4.7	1.0																											
940613	63.01	5.10	6.3	1.2	211	100	5	7.30	26.3	0.08	<0.02	1.45	3.9	<1.0	0.18	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03																			
940716	53.72	7.10	6.0	1.0	320	180	5	8.90	16.6	0.05	0.08	2.99	2.9	1.0																											
940825	29.40	5.20	1.0	1.0	300	115	5	7.20	18.0	0.02	0.09	1.31	6.5	1.5																											
940913	23.88	7.10	6.4	0.5	315	348	5	5.90	21.7	0.09	0.16	1.42	4.7	1.0																											
941018	64.23	6.70	3.0	0.5	355	200	19	8.80	40.9	0.09	0.02	1.39	3.7	1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03																			
941116	23.13	9.10	4.0	0.5	430	395	13	11.2	36.0	0.06	0.02	0.69	3.5	<1.0																											
941212	44.13	7.10	4.0	0.6	365	210	8	7.40	23.3	0.11	<0.02	0.69	3.1	<1.0																											
Min:	20.28	5.1	1	0.5	211	100	5	4.01	15.2	0.02	0.38	2	1	0.15	0.03	0.03	0.03	0.03	0.03	0.56	1.15	0.68																			
Max:	64.23	9.3	9	3	435	395	19	12.4	40.9	0.14	1.18	5.66	6.5	1.5	0.23	0.03	0.03	0.03	0.03	0.56	1.15	0.68																			
Aver.:	37.32	7.00	4.67	1.17	332	216	9.67	7.84	26.03	0.08	0.17	1.68	3.73	1.04	0.19	0.03	0.03	0.03	0.03	0.56	1.15	0.68																			
St.dev.:	16.93	1.29	2.31	0.75	63	88	4.85	2.92	9.46	0.04	0.34	1.45	1.37	0.14	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00																			
Numb.:	12	12	12	12	12	12	12	12	12	12	12	12	12	12	3	3	3	3	3	1	1	1																			

Table 7.9 Measured concentrations - 1994

Table 7.10 Measured concentrations - 1994

Annual flow	2160 mill. cbm										Min : 26.3 cbm/s									
	Drainage area.										Max: 542 cbm/s									
Date	Q m³/s	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	TOC mg/l	Hg ng/l	Gamm HCH ng/l	PCB (The following Congeners) ng/l	IUPAC NOS ng/l	As µg/l	Cr-T µg/l	Ni µg/l
940320	31.18	17.40	11	7.0	190	84	5	0.78	1.24	0.06	0.05	1.31	1.5	0.05	<0.03	0.06	0.11	0.07	0.05	0.46
940606	54.20	5.53	12	4.0	210	23	9	2.50	0.50	0.01	0.07	9.10	2.5	0.05	<0.03	0.06	0.11	0.07	0.12	2.12
940817	34.59	6.11	4	0.5	155	26	5	0.80	0.30	<0.01	0.06	0.84	1.5	<0.03	<0.03	<0.03	0.08	0.08	0.86	1.09
941028	30.66	21.40	24	19.0	160	61	5	0.30	0.90	0.06	0.19	1.29	3.5	<1.0	0.05	<0.03	<0.03	<0.03	0.15	0.53
Min:	30.66	5.53	4	0.5	155	23	5	0.3	0.3	0.01	0.05	0.84	3.5	1	0.05	0.03	0.03	0.03	0.08	0.51
Max:	54.2	21.4	24	19	210	84	9	2.5	1.24	0.06	0.19	9.1	3.5	2.5	0.05	0.03	0.06	0.11	0.63	2.12
Aver.:	37.7	12.61	12.75	7.63	179	49	6.00	1.10	0.74	0.04	0.09	3.14	3.50	1.63	0.05	0.05	0.05	0.07	0.05	1.09
St.dev.:	11.2	8.01	8.30	8.04	26	29	2.00	0.96	0.42	0.03	0.07	3.98	4	1	0.63	0.03	0.04	0.04	0.04	0.75
Numb.:	4	4	4	4	4	4	4	4	4	4	4	4	4	4	2	2	2	2	2	0.28

APPENDIX VIII : TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994			Page:
Table 8.1	Cond., Nutrients, Heavy metals, Suspended part.matter		59-65
Table 8.2	Mercury, Lindane, PCBs		67-73

(1) Glomma	"tributaries"	:	Tista	- Hølenelva
(1) Inner Oslo-fjord		:	Årungelva	- Åroselva
(2) Drammenselva	"tributary"	:	Lierelva	
(3) Numedalslågen	"tributaries"	:	Sandeelva	- Farriselva
(4) Skienselva	"tributary"	:	Tokkeelva	
(5) Otra	"tributaries"	:	Gjerstade.	- Audna
(6) Orreelva	"tributaries"	:	Lygna	- Ulla
(7) Suldalslågen	"tributaries"	:	Saudaelva	- Hornindalselva
(8) Orkla	"tributaries"	:	Ørstaelva	- Salsvatnelva
(9) Vefsna	"tributaries"	:	Åbjøra	- Reisa
(10) Alta	"tributaries"	:	Mattiselva	- Grense Jacobse.

Table 8.1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

Tributary Rivers. Mean Concentrations 1994.

County	Watercourse	Parameters (mean values)																
		Runoff data				Discharge				Sampling station								
		Drainage area	Outlet	Samp.	Disch. station	Normal	1994	Cond	Tot-P	PO4-P	Tot-N	NO3-N	Cu	Zn	cd	Pb	S.P.M.	
Vest-Agder (5.)	Tovdalselva, Topdalsfl. Søgneelva, Flekkerøy Mandalselva, Mannefj. Audna, Sniksfj. Lygna, Lyngdalsfl. Kulna, Fedafj. Sira, Ana-Sira	1856	1854	1794	33.9	31.3	33.9	2.38	4.0	0.5	467	178	44	0.4	7.5	0.07	1.19	
		204	192	38.0	41.8	38.0	8.03	6.0	2.0	1025	800	56	0.7	10.5	0.11	0.41	1.56	
		1809	1800	1740	46.0	50.0	47.6	50.0	2.41	5.0	0.5	394	214	39	0.4	5.3	0.03	1.03
		450	400	59	45.0	51.8	50.4	5.27	4.0	0.5	615	430	41	0.4	7.7	0.06	0.48	
		664	660	266	48.0	61.6	57.9	71.5	3.24	4.0	0.5	453	244	30	0.2	10.5	0.06	1.19
		1445	1140	1140	57.6	18.1	57.6	18.1	3.95	5.0	1.0	350	170	24	0.2	6.6	0.01	1.71
		1916	1872	1872	59.4	57.6	59.4	57.6	2.49	3.0	0.5	265	155	30	0.1	4.4	0.01	0.38
		293	107	51.1	56.7	51.1	5.37	5.0	2.0	370	275	30	0.1	8.3	0.01	0.25	0.80	
		241	240	194	57.5	60.4	71.1	71.6	4.61	7.0	2.0	450	330	12	0.1	6.6	0.01	0.73
		705	704	633	77.7	69.3	86.4	78.0	3.70	5.0	0.5	462	372	12	0.1	4.8	0.01	0.45
Rogaland (6.)	Sokndalselva, Sogndalsstr. Hellelandselva, Egersund Bjerkelandselva, Egersund Hæiva, Håtangen Fliggo, Solavika Ims-Lutsi, Høgsfj.Boknafj. Oltedalse., Høgsfj.Boknafj. Dirdalse., Høgsfj.Boknafj. Frafjorde., Frafj. Boknafj. Espedalsel., Høgsfj.Boknafj. Lysee., Lysefj.Boknafj. Ardalse., Ardalafj.Boknafj. Førree., Jøsenfj.Boknafj. Ulla., Jøsenfj.Boknafj. Sauda., Saudafj.Boknafj. Abøelva., Saudafj.Boknafj. Vikedalse., Boknafj.	127	127	127	34.9	40.9	34.9	40.9	6.63	8.0	0.5	716	545	18	0.5	2.7	0.02	0.14
		101	129	70.0	76.3	70.0	3.62	15.0	2.0	396	290	29	0.3	4.3	0.03	0.21	0.70	
		158	95	83.0	89.6	83.0	2.74	3.0	0.5	322	232	11	0.2	2.6	0.02	0.40	0.26	
		178	124	94.4	101.0	94.4	2.49	4.0	0.5	261	176	13	0.2	2.7	0.03	0.42	0.42	
		138	124	90.0	94.5	90.0	2.31	3.0	0.5	255	195	9	0.3	2.8	0.06	0.27	0.51	
		182	46	74.0	77.7	74.0	2.50	3.0	1.0	310	200	9	1.0	3.0	0.06	0.30	0.51	
		519	516	501	81.4	85.4	81.4	2.76	11.0	6.0	0.5	252	249	7	0.3	3.6	0.05	1.01
		163	163	85.8	90.1	85.8	2.30	3.0	0.5	320	245	13	0.4	2.7	0.08	0.30	0.43	
		393	385	83.4	87.6	83.4	2.32	3.0	0.5	220	155	7	0.3	2.7	0.08	0.30	0.43	
		353	353	85.0	93.5	85.0	3.49	3.0	0.5	880	750	3	0.7	21.0	0.11	0.35	0.33	
(7.)		82	82	85.0	93.5	85.0	1.58	3.0	0.5	240	165	8	0.1	1.3	0.01	0.11	0.33	
		118	117	80.0	103.1	80.0	2.75	3.0	0.5	283	166	5	0.4	3.3	0.04	0.19	1.17	

Table 8.1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data												Parameters (mean values)													
		Drainage area				Discharge				gauging station				Cond mS/m			Tot-P µg/l		Tot-N µg/l		NH4-N µg/l		Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l
		Outlet sq.km	Sampi. Disch. station sq.km	Disch. gaug. station sq.km	Sampling station I/s sq.km	Normal 1994 I/s sq.km	Normal 1994 Normal I/s sq.km	Normal 1994 Cond mS/m	Normal 1994	Tot-P µg/l	Tot-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l										
Hordaland (7.)	Eneelva, Etnefj. Børmlaflj. Opo, Sørfj. Hardangerfj. Tysso, Sørfj. Hardangerfj. Kinsø, Sørfj. Hardangerfj. Veig, Eldfjv. Hardangerfj. Bjorela, " Hardangerfj. Sima, Eldfj. Hardangerfj. Ausdøla, Osafj. Eldfj. Norddøla, Osafj. Eldfj. Tysseelva, Fusafj. Oselva, Fusafj. Bergsdalselva, Veafj. Herdlaflj. Vosso, Veafj. Sørfj. Eitso, Osterfj. Modalselva, Osterfj.	252 482 388 281 496 592 145 131 40 240 109 50 198 1492 414 385	250 464 385 281 496 592 128 130 39 240 108 91.7 198 1102 400 384	127 79.3 79.3 91.2 46.0 26.0 69.2 74.6 82.0 85.0 91.7 92.6 80.0 58.2 86.2 95.5	46.8 93.1 93.1 79.3 59.3 28.6 76.8 82.0 82.0 93.5 91.7 86.4 76.5 96.5 105.0	58.4 115.0 96.0 79.3 46.0 26.0 69.2 74.6 74.6 85.0 92.6 80.0 58.2 86.2 95.5	3.10 1.54 1.54 1.78 41.8 26.0 2.21 1.28 1.28 2.02 3.04 0.91 76.5 86.2 95.5	4.5 2.5 2.0 2.0 3.5 2.04 3.5 2.04 2.02 3.04 10.0 0.5 1.38 2.0 1.82	0.5 0.5 0.5 0.5 0.5 0.5 0.5 2.0 3.0 0.5 10.0 0.5 0.5 0.5 2.0 0.5	402 200 96 145 190 180 95 203 209 203 300 137 128 164 225	330 101 63 46 46 190 95 148 182 116 300 85 77 112 166	14 108 63 46 46 190 95 148 182 116 300 85 77 112 166	0.3 6 0.1 5 5 0.1 0.1 5 5 7.0 0.5 0.5 0.5 0.5 0.5	2.7 3.1 2.7 1.4 0.5 0.1 0.5 0.2 0.1 10.0 0.1 0.2 0.5 1.0 5.0	0.04 0.04 <0.02 0.01 0.01 <0.01 0.01 0.01 <0.01 0.01 0.01 0.1 0.2 <0.01 0.01 0.01	0.09 0.20 0.11 0.05 0.05 0.07 0.07 0.04 0.02 0.30 0.12 0.56 0.70 0.50											

Table 8.1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data						Parameters (mean values)											
		Drainage area		Sampling station		Discharge gauging station		Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l	
		Outlet sq.km	Sampi. sq.km	Disch. station sq.km	Normal l/s sq.km	1994 l/s sq.km	Normal 1994												
Sogn og Fjordane (7.)	Nærøye,, Aurl. f., Sognefj. Flåmse,, Aurl. f., Sognef. Aurlandv. Aurl. f., Sognef. Erdalse,, Lærd. f., Sognefj. Lærdalsv. Lærd. f., Sognefj. Ardalsv., Ardalsf. Sognefj. Fortunv., Lustefj. Sognef. Mørkrivs., Lustefj. Sognef. Jostedøla, " Sognef. Areøe,, Sogn.d. f., Sognef. Sogndalsæ, " Sognef. Gaular, Dalsfj. Bufj. Jølstra, Førdefj. Nausta, Førdefj. Oselva, Høydalsfj. Høse,, Hyef. Nordfj. S Glengedalsæ, " Nordfj. S Breimsv., Gloppefj. " Oldene,, Indre Nordfj. Løneniva, Indre Nordfj. Stryne, Indre Nordfj. Hornindalsæ, Nordfj. N	290	290	267	59.5	66.6	59.5	3.34	2.5	0.5	170	126	5	0.1	0.6	0.02	0.03		
		280	275	275	52.4	70.4	52.4	70.4	2.26	1.5	0.5	137	103	5	0.1	0.9	0.02	<0.02	
		800	799	762	48.6	53.9	48.6	53.9	1.44	2.0	0.5	122	86	5	0.1	1.1	0.02	0.10	
		138	138	30.0	33.3	30.0	33.3	1.43	2.0	0.5	122	85	5	0.1	0.9	<0.01	0.04	0.45	
		1184	1172	30.0	22.1	30.0	22.1	1.88	4.0	0.5	212	126	5	0.2	1.1	<0.02	0.02	1.18	
		989	989	44.9	48.1	44.9	48.1	1.21	4.2	1.7	174	85	8	0.9	0.8	<0.01	0.07	2.97	
		508	508	367	51.0	56.1	51.0	56.1	1.11	2.5	1.0	137	103	5	0.2	0.9	0.02	0.08	
		282	282	203	54.7	57.5	54.7	57.5	1.28	1.5	0.5	137	103	5	0.1	0.6	0.02	<0.02	
		865	864	57.3	68.0	76.2	68.0	1.46	3.5	2.0	160	122	5	0.3	1.7	0.02	0.17	1.14	
		449	446	384	77.2	86.5	77.2	86.5	1.58	2.5	0.5	120	59	6	0.2	1.5	0.02	0.03	0.46
		175	172	111	66.1	77.8	66.1	77.8	1.48	5.5	3.0	225	160	5	0.2	1.1	0.10	0.06	0.77
		627	625	505	79.3	93.8	79.3	93.8	1.54	4.0	1.0	213	118	5	0.4	1.4	<0.01	0.14	0.36
		714	709	384	74.3	82.5	74.3	82.5	1.93	4.3	2.0	203	111	17	0.1	2.0	0.01	0.04	0.54
		277	273	232	81.7	101.2	81.7	101.2	1.82	7.0	2.0	181	85	8	0.1	1.0	0.02	0.11	0.84
		287	285	225	78.7	86.6	78.7	86.6	2.36	4.5	0.5	369	79	11	0.1	2.1	0.09	0.13	0.31
		73	73	161	75.0	82.5	75.0	82.5	1.71	3.0	1.0	126	90	5	0.1	0.9	0.02	0.11	0.41
		170	168	161	75.0	82.5	75.0	82.5	1.56	4.0	1.0	123	64	11	0.3	2.2	0.08	0.19	0.11
		636	634	585	68.0	75.5	68.8	75.5	1.93	6.0	0.5	335	143	11	0.1	1.6	0.07	0.13	0.51
		226	225	214	70.1	78.5	70.1	78.5	2.03	6.3	0.5	265	136	6	0.1	0.5	0.01	0.03	0.59
		261	260	234	65.0	69.4	65.0	69.4	1.95	9.0	1.5	180	103	6	0.2	0.6	0.08	0.02	0.81
		532	530	493	60.2	66.2	60.2	66.2	1.80	9.3	4.0	195	75	5	0.1	0.5	<0.01	0.02	8.66
		428	424	378	58.1	63.9	58.1	63.9	1.98	3.5	2.0	170	75	11	0.3	1.4	0.01	0.17	1.55

Table 8-1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Parameters (mean values)																	
		Runoff data				Discharge				Sampling									
		Drainage area	Outlet	Sampi.	Disch.	Sampling station	gauging station	Normal	1994	Cond	Tot-P	PO4-P	Tot-N	NO3-N	Cu	Zn	Cd	Pb	S.P.M.
sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	Normal	1994	Normal	1994	mS/m	µg/l	µg/l	µg/l	µg/l	µg/l	µg/l	µg/l	mg/l	
Møre og Romsdal (8)	Ørstaæ., Ørstafj.	160	155	70.0	73.5	70.0	73.5	3.05	16.0	8.0	359	195	14	0.5	2.7	0.04	0.17	2.93	
	Valldøla, Nordafj. Storfj.	359	357	60.0	62.4	60.0	62.4	2.01	3.0	0.5	170	108	5	0.2	0.5	<0.01	0.05	0.77	
	Rauma, Romsdalsfj. Moldefj.	1202	1190	1142	32.8	34.4	32.8	1.96	2.5	0.5	104	59	5	0.3	0.9	0.15	0.16	0.37	
	Isa, Isfj. Moldefj.	175	175	89	57.0	58.0	57.0	2.56	3.0	0.5	160	90	11	0.2	0.7	0.05	0.51	0.51	
	Eira, Erest. Moldefj.	1119	1119	1085	34.8	36.2	34.8	2.61	2.5	0.5	175	112	5	0.4	3.5	0.03	0.13	0.26	
	Littlandsfj., Sunndalsfj.	359	330	41.0	43.0	41.0	43.0	1.08	2.0	0.5	51	15	5	0.3	0.6	0.01	0.32	0.29	
	Driva, Sunnd. fj. Tingvollsfl.	2487	2435	2435	27.9	28.7	27.9	3.82	3.0	0.5	265	185	11	0.5	0.7	0.03	0.06	0.65	
	Ulvåa, Alvundsfj.	199	199	207	57.0	58.7	60.7	1.95	3.5	1.0	137	76	5	0.3	0.9	0.08	0.10	0.09	
	Toåa, Todalsfj.	251	251	207	58.5	60.3	58.5	1.51	4.0	1.0	117	21	5	0.2	0.6	<0.01	0.04	0.39	
	Surna, Surnadalsfj.	1200	1200	1125	48.0	50.0	49.3	2.57	8.4	3.0	349	175	5	0.3	0.7	<0.02	0.57	0.57	
Sør-Trøndelag (8)	Bøvra, Hamnesfj. Halsafj.	243	243	196	55.0	57.2	55.0	2.78	3.0	0.5	250	140	8	0.6	1.7	<0.01	0.07	1.01	
	Børse, Gaulosen Tr.h.fj.	110	100	30.0	28.5	30.0	30.0	10.30	10.0	2.0	730	475	3	1.4	1.1	0.01	0.14	4.74	
	Vigta, Gaulosen Tr.h.fj.	150	150	30.0	28.5	30.0	30.0	12.00	23.0	12.0	415	210	11	2.2	3.8	<0.01	0.59	25.90	
	Gaulosen Tr.h.fj.	3659	3650	3062	26.4	20.6	26.4	20.6	6.14	5.0	1.0	235	102	3	1.6	2.6	0.01	0.14	1.57
	Nidelva, Trondheimsfj.	3110	3100	3049	35.5	27.8	35.5	27.8	4.03	7.0	0.5	210	59	3	1.0	2.7	0.01	0.29	0.54
	Homla, Stjørd. fJ. Tr.h.fj.	157	157	30.0	29.0	30.0	29.0	7.31	11.0	4.0	365	51	47	4.3	6.0	<0.01	0.71	0.98	
	Stjørdalsv., " Tr.h.fj.	2117	2117	1863	38.5	35.9	38.5	3.56	5.0	3.0	245	132	3	1.7	2.9	<0.01	0.13	3.82	
	Gråe, " Tr.h.fj.	93	93	25.0	25.0	28.6	44.5	35.8	6.02	4.0	3.0	865	660	6	1.6	1.3	0.01	0.33	2.13
	Verdalsvassdr., Tr.h.fj.	1472	1472	898	40.0	29.0	33.6	32.6	5.62	12.0	5.0	280	126	7	1.3	3.9	0.60	0.82	0.82
	Figgafj.Leksdalse, Tr.h.fj.	282	282	178	30.0	29.0	30.0	25.4	35.1	25.1	4.93	4.0	0.5	250	10	1.1	1.3	0.01	0.24
Nord-Trøndelag (8)	Snåsavassdr., Trondhfj.	2125	2125	35.1	25.4	35.1	35.1	25.1	4.93	4.0	0.5	310	175	7	0.7	1.0	0.01	0.14	6.27
	Argårdseiva, Namsfj.	543	510	238	41.0	50.9	48.3	10.80	37.0	21.0	540	147	84	1.4	2.7	0.01	0.36	6.92	
	Namsen, Namsfj. Ø	6277	5718	43.4	44.5	43.4	44.5	2.62	3.0	0.5	132	39	6	0.6	1.5	0.01	0.18	1.10	
	Salsvatneiva, Follafl.	432	422	59.7	59.4	59.7	59.4	5.06	1.0	0.5	125	64	3	1.2	1.4	0.01	0.18	0.44	

Table 8.1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data						Parameters (mean values)											
		Drainage area			Discharge			Sampling station	gauging station	Cond	Tot-P	PO4-P	Tot-N	NO3-N	NH4-N	Cu	Zn	Cd	Pb
		sq.km	sq.km	sq.km	Normal	1994	Normal												
Nordland (9.)	Abjøra, Blindalsfj. S	526	520	384	80.2	59.0	80.2	59.0	3.99	5.0	1.0	114	24	3	0.9	0.5	0.01	0.29	2.77
	Skjæra, Vefsenvf. S	104	104	98	41.3	36.3	41.3	36.3	7.41	9.0	6.0	225	59	40	0.5	1.0	0.01	0.14	0.88
	Fusta, Vefsenvf. N	544	543	520	63.4	57.0	63.4	57.0	3.81	3.0	0.5	120	19	10	0.4	1.4	<0.01	0.17	1.20
	Dreyla, Vefsenvf. N	177	176	98	65.0	58.5	65.0	58.5	3.96	5.0	2.0	111	34	3	0.5	1.6	0.01	0.23	3.07
	Ressåga, Sørflj.	20922	20877	1880	45.4	51.0	45.4	51.0	5.27	4.0	1.0	140	34	11	0.7	4.9	0.02	0.32	2.31
	Bjerka, Sørflj.	385	385	273	55.4	48.7	55.4	48.7	2.61	3.0	1.0	66	19	3	0.4	0.5	0.01	0.07	1.43
	Dalselva, Ranafj. N	211	211	129	39.5	35.5	39.5	35.5	2.53	1.0	3.0	87	19	3	0.6	1.2	0.02	0.23	1.95
	Ranavassdraget, Ranafj. N	3847	3846	1892	44.9	50.0	44.9	50.0	3.90	3.0	0.5	104	51	3	0.4	1.1	<0.01	0.98	1.62
	Fykanåga, Glomfjord	297	297	243	103.7	100.0	103.7	100.0	2.91	3.0	2.0	85	40	7	0.5	1.0	0.01	0.30	2.01
	Belare,, Belarflj. Nordflj.	1064	875	797	45.1	30.3	45.1	30.3	3.67	2.0	1.0	89	39	3	0.6	1.0	0.01	0.30	2.55
	Saltidalsvassdr., Saltidflj.S	1544	1543	1168	32.1	26.6	32.1	26.6	2.79	4.0	3.0	96	54	3	0.5	1.5	0.02	1.08	5.35
	Sultfjellevassdr., Saltidflj.S	1028	800	791	44.0	36.0	44.0	36.0	42.00	1.0	0.5	96	44	7	13.9	11.4	0.07	2.17	0.65
	Kobbe,, Leirflj. Sørflolda N	405	386	66.9	62.7	66.9	62.7	1.50	1.0	0.5	60	26	3	0.3	1.5	0.01	0.39	0.29	0.53
	Skjøna, Oftofl. S	845	840	797	36.3	29.0	36.3	29.0	1.81	2.0	0.5	68	10	3	0.5	1.4	0.03	0.21	0.53

Table 8.1 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

county	Watercourse	Runoff data										Parameters (mean values)										
		Drainage area					Discharge					Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l
		Outlet sq.km	Sampi. station	Disch. gaug. station	Sampling station	Normal I/s sq.km	1994 Normal I/s sq.km	Normal I/s sq.km	1994 Normal I/s sq.km	Cond mS/m	Tot-P µg/l	PO4-P µg/l	Tot-N µg/l	NO3-N µg/l	NH4-N µg/l	Cu µg/l	Zn µg/l	Cd µg/l	Pb µg/l	S.P.M. mg/l		
Troms (9.)	Spanselva, Astafj. Vågsfj. Salangse., Astafj. Vågsfj.	142	533	50.0	51.0	50.0	40.9	41.5	39.5	7.25	2.0	0.5	89	39	3	0.5	1.2	0.09	0.67	1.02		
	Rossfjorde., Maangen	539	539	40.9	51.0	40.9	39.5	28.7	28.1	39.5	7.27	3.0	0.5	89	44	3	0.3	0.5	<0.01	0.21	0.84	
	Målselv., Målselvfj. "	196	190	39.5	41.5	39.5	41.5	26.3	27.8	27.8	6.88	4.0	0.5	155	5	16	0.4	0.5	<0.01	0.12	0.68	
	Bardue., Målselvfa.	3299	3206	3118	28.7	28.1	28.7	26.3	27.7	28.0	6.13	4.0	1.0	119	39	3	0.5	0.7	0.01	0.19	2.53	
	Nordkjøsøya, Balsfj.	2906	2906	2049	27.8	27.8	27.8	27.7	27.7	28.0	6.13	4.0	1.0	119	39	3	0.5	0.7	0.01	0.19	2.53	
	Signaldalselva, Lyngen V	191	191	415	27.7	28.0	27.7	27.7	27.7	28.0	4.31	3.0	0.5	78	34	3	0.5	0.5	<0.01	0.03	0.90	
	Skibotnøya, Lyngen V	473	467	415	27.7	28.0	27.7	27.7	27.7	28.0	4.29	10.0	7.0	90	54	3	0.9	1.0	<0.01	0.25	13.40	
	Kåfjordelva, Lyngen Ø	770	770	724	18.0	16.8	18.0	18.0	18.0	18.0	2.99	2.0	0.5	96	44	7	0.6	0.5	0.01	0.03	0.95	
	Reisa, Reisatj.	358	358	348	20.0	23.0	20.0	23.0	20.0	23.0	4.34	1.0	0.5	147	105	3	1.1	0.6	0.01	0.02	0.28	
	2702	2702	16.0	13.6	16.0	13.6	16.0	13.6	16.0	13.6	3.76	2.0	0.5	114	49	10	1.4	1.9	0.03	0.18	1.33	
	Mattiselva, Kåfj. Altafj.	325	325	319	26.5	21.2	26.5	26.5	21.2	26.5	2.75	3.0	0.5	90	13	6	0.6	0.5	<0.01	0.16	0.87	
	Tverrelva, Altafj.	234	233	15.1	12.0	15.1	12.0	15.1	12.0	15.1	4.27	4.0	2.0	140	45	7	0.6	0.5	0.01	0.10	0.55	
	Reparfjordv., Repparfi.	1050	1089	25.0	22.5	25.0	22.5	25.0	22.5	25.0	4.50	2.0	0.5	119	24	8	0.4	0.5	<0.01	0.14	0.38	
Finnmark (10.)	Stabburse., I. Porsangen V	1108	1102	870	18.3	15.2	18.3	15.2	18.3	18.3	3.02	2.0	0.5	104	24	9	0.4	1.3	0.09	2.33	0.74	
	Lakse., Indre Porsangen S	1532	941	15.9	12.8	15.9	12.8	15.9	12.8	15.9	4.41	5.0	1.0	134	14	3	1.0	1.0	0.01	0.74	2.41	
	Børselva, Indre Porsangen Ø	883	883	863	29.8	29.0	29.8	29.0	29.8	29.8	3.74	1.0	0.5	60	14	10	0.2	1.2	0.01	0.16	0.62	
	Mattusjakkå, I. Laksefj. V	101	101	101	22.8	20.5	22.8	20.5	22.8	22.8	4.01	3.0	1.0	69	14	5	0.6	0.4	<0.01	0.16	0.95	
	Storelva, Indre Laksefj. V	690	690	760	21.9	19.0	19.9	19.9	19.9	19.9	2.10	1.0	0.5	75	3	0.5	3.9	0.04	0.03	0.22	0.10	
	Soussjakkå, I. Laksefj. V	92	92	102	25.3	22.8	22.8	22.8	22.8	22.8	4.68	1.0	0.5	75	34	3	0.3	1.6	0.02	0.13	0.29	
	Adamsselva, I. Laksefj. Ø	705	705	760	19.9	15.0	19.9	15.0	19.9	19.9	6.14	2.0	0.5	105	5	11	0.6	1.5	0.03	0.16	0.59	
	Tanavassdraget, Tanatj. S	16389	15713	14169	11.5	9.2	11.5	9.2	11.5	11.5	3.96	6.0	2.0	165	10	20	0.7	0.5	0.01	0.32	1.58	
	Vesterelva, Syltefj.	469	469	79	34.6	31.6	34.6	31.6	34.6	34.6	8.55	4.0	1.0	60	5	3	0.4	0.5	0.03	0.09	0.22	
	V. Jakobse., Y. Varangerfj.	627	627	239	18.1	14.5	18.1	14.5	18.1	18.1	5.72	4.0	2.0	135	5	14	0.4	0.7	0.01	0.10	0.19	
	Pasvilkie., Bektfj. Varangfj.	18494	18400	16175	9.3	10.2	9.3	10.2	9.3	10.2	4.37	8.0	2.0	165	5	21	2.1	1.9	<0.01	0.21	3.29	
	Neiden, Munkfj. Varangfj.	2960	2960	2911	9.8	8.8	9.8	8.8	9.8	9.8	2.95	3.0	1.0	185	34	3	0.7	1.0	<0.01	0.24	0.96	
	Grense Jakobse., Varangfj.	234	234	234	18.0	18.0	18.0	18.0	18.0	18.0	4.20	3.0	0.5	175	10	27	2.6	1.7	0.02	1.35	0.85	

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data						Parameters (mean values)												
		Discharge			Gauging station			IUPAC NOS												
		Drainage area	Sampi. Disch. station	Sampling station	Normal	1994	Hg	PCB (The following Congeners)	28	52	101	118	138	153	180	TOC	DOC	Cr-T	NI	As ug/l
sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	mg/l	mg/l	ug/l	ug/l	ug/l
Østfold (1.)	Tista, Iddefj. Mosseselva, Mossesundet Ø	1588 690	1582 689	14.4 14.5	15.5 15.4	14.4 14.5	<1.0 1.50	0.62 <0.03	<0.03 <0.03	<0.03 <0.03	<0.03 <0.03	<0.03 <0.03	<0.03 <0.03	<0.03 <0.03	6.70 7.20	1.69 1.93	0.79 1.09	0.27 0.34		
Oslo & Akershus (1.)	Helenelva, Drøbakseendet Ø Arungeliva, I. Oslofj. Glersjøelva, I. Oslofj. Ljanselva, I. Oslofj. Loelva/Alna, I. Oslofj. Akerselva, I. Oslofj. Frognerelva, I. Oslofj. Lysakerelva, I. Oslofj. Sandvikselva, I. Oslofj. Åroselva, I. Oslofj.	137 52 86 42 75 227 23 178 223 113	121 50 85 41 69 225 20 173 187 109	14.0 13.0 14.0 13.0 13.0 22.2 20 173 18.4 17.0	13.3 12.4 13.3 13.0 13.9 12.6 20.9 27.1 18.0 16.3	15.5 14.5 15.4 14.5 13.9 12.6 20.9 <1.0 18.4 17.0	2.50 0.58 0.60 0.52 0.42 0.84 0.62 0.55 1.50 0.21	<0.03 <0.03 <0.03 <0.03 0.28 0.03 <0.03 <0.03 0.03 <0.03	<0.03 <0.03 <0.03 <0.03 0.19 0.03 0.11 <0.03 <0.03 <0.03	<0.03 <0.03 <0.03 <0.03 0.27 0.04 0.11 <0.03 0.04 <0.03	<0.03 <0.03 <0.03 <0.03 0.19 0.04 0.10 <0.03 0.05 <0.03	<0.03 <0.03 <0.03 <0.03 0.29 0.04 0.11 <0.03 0.05 <0.03	<0.03 <0.03 <0.03 <0.03 0.22 0.05 0.11 <0.03 0.07 <0.03	9.90 2.80 4.90 6.20 4.80 3.80 4.10 5.20 5.20 5.20	2.81 2.26 2.01 1.45 <0.5 1.06 1.48 0.76 0.77 0.77	2.66 2.79 1.68 1.01 0.50 0.71 1.87 0.79 1.06 1.13	0.59 0.91 0.21 0.03 0.35 0.19 0.49 0.08 0.34 0.34			
Buskerud (2.)	Lierelva, Drammensfj. Ø	266	222	18.6	28.8	18.6	0.26	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.40	1.19	1.17	0.19		
Vestfold (3.)	Sandeelva, Sandebukta Aulielva, Tønsbergfj. Farriselva, Larvikfj.	193 363 491	190 362 491	17.0 14.9 21.6	16.2 14.6 21.4	17.0 14.9 21.6	1.00 <1.0 <1.0	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	3.40	1.12	0.29	0.31		
Telemark (4.)	Tokkeelva, Kragerø	1238	1200	26.7	28.6	26.7	1.00	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.60	1.26	0.83	0.44		
Aust- Agder (5.)	Gjerstadelva, Søndeledfj. Vegårdselva, Sandnesfj. Nidelva, Arendal	419 457 4025	414 429 4020	291 291 3956	27.0 29.3 31.3	27.0 30.8 29.8	1.50 1.00 <1.0	1.00 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	<0.03 <0.03 <0.03	4.60 4.83 2.85	0.51 0.95 0.51	1.14 0.49 0.11	0.31 0.19 0.18		

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data										Parameters (mean values)												
		Drainage area			Sampling station			Discharge gauging station				PCB (The following Congeners)				IUPAC NOS								
		sq.km	sq.km	sq.km	Normal	1994	Normal	I/s	sq.km	I/s	sq.km	1994	Hg	Gamma HCH	28 ng/l	52 ng/l	101 ng/l	118 ng/l	138 ng/l	153 ng/l	180 ng/l	TOC mg/l	DOC mg/l	Cr-T mg/l
Vest-Agder (5)	Tovdalselva, Topdalsfj. Segneelva, Flekkerøy	1856	1854	1794	33.9	31.3	33.9	<1.0	1.00	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.78	0.84	0.22	0.35	0.33
	Mandalselva, Mannefj. Audna, Sniksfj.	1809	1800	1740	46.0	50.0	47.6	50.0	1.50	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.80	1.76	0.85	0.19	0.19
	Lygna, Lyngdalsfj. Kynna, Fedafj.	450	400	59	45.0	50.4	51.8	1.00	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.87	0.66	0.07	1.41	0.13
	Sira, Ana-Sira	1445	1140	1140	57.6	18.1	57.6	<1.0	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.90	1.82	0.27	0.15	0.39
	Sokndalselva, Sogndalsstr. Hellelandselva, Egersund	294	293	107	51.1	56.7	51.1	<1.0	0.90	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.70	0.85	0.47	0.21	0.21
	Bjerkreimselva, Egersund	241	240	194	57.5	60.4	71.1	71.6	1.00	0.90	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	7.75	0.49	0.51	0.87	0.87
	Hælva, Håtangen	705	704	633	77.7	69.3	86.4	78.0	1.00	0.90	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	7.75	0.87	0.87	0.87	0.87
	Figgjo, Solavika	165	160	135	46.9	45.4	46.9	45.4	<1.0	0.81	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	5.10	0.87	0.87	0.15	0.15
	Ims-Lutsl. Høgsfj.Boknafj.	229	218	135	50.0	52.0	50.0	<1.0	0.80	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.90	1.75	0.41	0.25	0.24
	Oltedals-, Høgsfj.Boknafj.	127	127	127	34.9	40.9	34.9	40.9	<1.0	0.80	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.40	4.05	0.25	0.25	0.24
(7)	Dirdalse-, Høgsfj.Boknafj.	102	101	129	70.0	76.3	70.0	<1.0	0.80	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.40	2.56	0.14	0.16	0.16
	Frafjorde., Frafl. Boknafj.	158	158	95	83.0	89.6	83.0	<1.0	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.95	1.68	0.03	0.13	0.13
	Espedalsde., Høgsfj.Boknafj.	178	178	124	94.4	101.0	94.4	<1.0	0.60	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.30	1.47	0.03	0.11	0.11
	Lyssefj.Boknafj.	182	182	46	74.0	77.7	74.0	<1.0	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.10	1.86	0.02	0.09	0.09
	Lysee., Ardalsfj.Boknafj.	519	516	501	81.4	85.4	81.4	2.00	0.36	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.60	0.32	0.06	0.03	0.03
	Førre., Jøssenfj.Boknafj.	163	163	65.8	90.1	85.8	65.8	1.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.07	2.01	0.31	0.38	0.38
	Ulla., Jøsenfj.Boknafj.	393	393	83.4	87.6	83.4	<1.0	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.90	1.08	0.05	0.11	0.11
	Saudae., Saudafj.Boknafj.	353	353	82	82	85.0	93.5	85.0	1.00	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.90	0.32	0.06	0.03	0.03
	Aabelsø, Aabelsøfj.Boknafj.	118	117	80.0	103.1	60.0	103.1	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.40	1.05	0.12	0.07	0.07
	Vikedalselv.,																			0.90	0.95	0.05	0.05	0.05

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data										Parameters (mean values)										
		Drainage area			Sampling station			Discharge gauging station				IUPAC NOS										
		sq.km	sq.km	sq.km	Normal	1994	Normal	Hg	HCH	PCB (The following Congeners)	28	52	101	118	138	153	180	TOC mg/l	DOC mg/l	Cr-T ug/l	Ni ug/l	As ug/l
(7.)	Etnelva, Etnefj. Bømlafj. Opo, Sørfl. Hardangerfj.	252	250	127	48.8	58.4	96.0	115.0	<1.0	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.53	1.15	0.15	0.10	
	Tysso, Sørfl. Hardangerfj.	482	480	464	79.3	93.1	2.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.70	1.00	0.05	0.10	
	Kinsø, Sørfl. Hardangerfj.	388	385	407	79.3	91.2	79.3	1.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.30	0.85	0.05	0.03	
	Veig, Eldfjv. Hardangerfj.	281	281	232	46.0	59.3	46.0	59.3	<1.0	0.33	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.50	0.35	<0.01	0.10
	Bjorela, " Hardangerfj.	496	496	386	41.8	46.2	41.8	1.00	0.23	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.10	0.60	0.15	0.01
	Sima, Eldfj. Hardangerfj.	592	592	592	26.0	28.6	26.0	0.23	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.10	0.60	0.15	0.01
	Austdøla, Osafj. Eldfj.	145	145	128	69.2	76.8	69.2	<1.0	0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.30	0.90	0.02	0.03
	Norddøla, Osafj. Eldfj.	131	130	89	74.6	82.0	74.6	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.20	0.03	0.03	0.15
	Tysseelva, Fusafj.	40	39	89	74.6	82.0	74.6	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.20	0.65	0.80	0.60
	Oseleva, Fusafj.	240	240	50	91.7	92.6	91.7	92.6	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.79	1.40	0.05	0.10
	Bergsdalselva, Veafj. Herdlaflj.	109	108	198	80.0	86.4	80.0	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.42	1.40	0.05	0.10
	Vosso, Veafj. Sørfl.	1492	1465	1102	58.2	76.5	58.2	76.5	2.50	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.90	1.15	0.20	0.02
	Ekso, Osterfj.	414	400	342	86.2	96.5	86.2	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.90	0.85	<0.01	0.10
	Modalselva, Osterfj.	385	384	248	95.5	105.0	95.5	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.24	0.05	0.01	0.15

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Parameters (mean values)																									
		Runoff data				Discharge				IUPAC NOS								DOC	Cr-T	Ni	As						
		Outlet	Sampl.	Disch.	gaug.	station	Sampling	station	gauging	Normal	1994	Normal	1994	I/s sq.km	I/s sq.km	I/s sq.km	I/s sq.km	mg/l	ug/l	ug/l	ug/l						
Sogn og Fjordane (7.)		sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	Normal	1994	Normal	1994	Hg	Hg	HCH	28	52	101	118	138	153	180	DOC	Cr-T	Ni	As
	Nærøye,, Aurl. fl. Sognefj. Flåmse,, Aurl. fl. Sognefj. Aurlandv. Aurl. fl. Sognefj. Erdalset., Lærd. fl. Sognefj. Lærdalsv. Lærd. fl. Sognefj. Ardalsv., Ardalsfl. Sognefj. Fortunv., Lusterfl. Sognefj. Mørkrivs., Lusterfl. Sognefj. Jostedøla, " Sognefj. Arøye., Sognd. fl. Sognefj. Sogndalsæ., " Sognefj. Gaular, Dalsfl. Buifl. Jølstra, Førdefl. Nausta, Førdefl. Oselva, Høydalsfl. Hopse., Høyfl. Nordfl. S. Gjengedalsæ., " Nordfl. S. Breimse., Gløppenfl. " Oldens., Indre Nordfl. Loenelva, Indre Nordfl. Stryneee., Indre Nordfl. Hornindalsæ., Nordfl. N. 428	290	290	267	59.5	66.6	59.5	1.00	0.20	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.40	1.35	0.10	0.10	
		280	275	52.4	70.4	52.4	70.4	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.30	0.70	0.15	0.05	
		800	799	762	48.6	53.9	48.6	<1.0	0.12	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.40	0.50	0.03	<0.01	
		138	138	30.0	33.3	30.0	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.90	0.55	0.05	0.20	
		1184	1172	30.0	22.1	30.0	22.1	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	1.51	0.15	0.05	0.05	
		989	989	44.9	48.1	44.9	48.1	3.00	0.25	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.76	0.40	0.10	0.03	
		508	508	367	51.0	56.1	51.0	2.00	0.28	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.30	0.15	0.10	0.02	
		282	282	203	54.7	57.5	54.7	1.00	0.28	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.40	0.30	0.02	<0.01	
		865	864	573	68.0	76.2	68.0	<1.0	0.28	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.40	0.40	0.50	0.10	
		449	446	384	77.2	86.5	77.2	<1.0	0.28	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.60	0.65	0.03	0.03	
		175	172	111	66.1	77.8	66.1	77.8	<1.0	0.28	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	2.90	0.55	0.05	0.02
		627	625	505	79.3	93.8	79.3	93.8	1.00	0.36	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	1.57	0.40	0.10	0.10
		714	709	384	74.3	82.5	74.3	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	1.10	0.95	0.03	0.05	
		277	273	232	81.7	101.2	81.7	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	1.58	1.20	0.10	0.03	
		287	285	225	78.7	86.6	78.7	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.60	2.50	1.90	0.03	
		73	73	161	75.0	82.5	75.0	1.00	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.70	1.49	0.40	0.05	
		170	168	161	75.0	82.5	75.0	2.50	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.70	0.80	0.10	0.05	
		636	634	585	68.0	75.5	68.8	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.70	0.80	0.05	0.02	
		226	225	214	78.5	77.1	78.5	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.75	0.75	0.05	<0.01	
		261	260	234	65.0	69.4	65.0	69.4	1.00	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.80	0.80	0.50	0.05
		532	530	493	60.2	66.2	60.2	66.2	1.00	0.30	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	0.80	0.80	0.10	0.05
		428	424	378	63.9	58.1	63.9	58.1	1.00	0.40	<0.03	<0.03	<0.03	<0.03	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	1.20	1.30	0.10	0.15

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data												Parameters (mean values)																	
		Drainage area			Sampling station			Discharge gauging station			1994			Hg			PCB (The following Congeners)			IUPAC NOS			DOC			Cr-T			Ni		
		Outlet	Sampl.	Disch.	station	gaug.	station	Normal	1994	Normal	sq.km	sq.km	l/s sq.km	Normal	1994	Normal	ng/l	ng/l	ng/l	ng/l	ng/l	ng/l	mg/l	mg/l	mg/l	ug/l	ug/l	ug/l			
Nordland (9.)	Abjøra, Blindsightj. S	526	520	384	80.2	59.0	80.2	59.0	<1.0	0.30	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.52	0.31	0.14				
	Skjerva, Vefsentj. S	104	104	98	41.3	36.3	41.3	36.3	2.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.61	0.03	0.26				
	Fista, Vefsentj. N	544	543	520	63.4	57.0	63.4	57.0	1.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.85	0.21	0.25				
	Drevia, Versentj. N	177	176	98	65.0	58.5	65.0	58.5	<1.0	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.71	0.16	0.19				
	Røssåga, Sørkj.	2092	2087	1880	45.4	51.0	45.4	51.0	1.00	0.30	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.50	0.11	0.07				
	Bjerkå, Sørkj.	385	385	273	55.4	48.7	55.4	48.7	1.00	0.30	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.91	0.09	0.07				
	Dalselva, Ranafj. N	211	211	129	39.5	35.5	39.5	35.5	<1.0	0.25	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.39	0.24	0.08				
	Ranavassdraget, Ranafj. N	3847	3846	1892	44.9	50.0	44.9	50.0	<1.0	0.25	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.32	0.22	0.09				
	Fykanåga, Glomfjord	297	297	243	103.7	100.0	103.7	100.0	<1.0	0.25	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.32	0.22	0.22				
	Belare., Belafj. Nordfl.	1064	875	797	45.1	30.3	45.1	30.3	<1.0	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.60	1.99	0.17					
	Saltdalsvassdr., Salidflj. S	1544	1543	1168	32.1	26.6	32.1	26.6	<1.0	0.70	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.39	0.11	0.26				
	Sulttelimavassdr., Saltedflj. S	1028	800	791	44.0	36.0	44.0	36.0	1.00	0.74	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	19.50	0.22	1.74				
	Kobbe., Leirfi. Sørfolda N	405	405	386	66.9	62.7	66.9	62.7	<1.0	0.50	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.29	0.35	0.09				
	Skjoma, Ofotflj. S	845	840	797	36.3	29.0	36.3	29.0	1.00	0.40	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.78	0.37	0.11				

Table 8.2 TRIBUTARY RIVERS. MEAN CONCENTRATIONS 1994.

County	Watercourse	Runoff data						Parameters (mean values)										
		Discharge			gauging station			IUPAC NOS										
		Drainage area	Sampling station	Normal 1994 l/s sq.km	Hg ng/l	PCB 28 ng/l	The following Congeners 118 ng/l	DOC mg/l	Cr-T ug/l	Ni ug/l	As ug/l							
Troms (9)	Spanselva, Astafj. Vågsfj. Salangs., Astafj. Vågsfj. Rossfjorde., Målangen Måse., Måselvfl. ” Bardue., Måselvra Nordkioselva, Balafj. Signaldaiselva, Lyngen V Skitbonelva, Lyngen Kafjordelva, Lyngen Ø Reisa, Relesfj.	142	142	533	50.0	51.0	50.0	<1.0	0.35	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.11	0.21	0.04
		539	539	533	40.9	51.0	40.9	<1.0	0.35	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.14	0.11	1.75
		196	190	39.5	41.5	39.5	2.00	0.33	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.01	0.21	0.14
		3239	3200	3118	28.7	28.1	28.7	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.95	0.02	0.07
		2906	2906	2049	28.3	27.8	28.3	<1.0	0.20	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.95	0.02	0.07
		191	191	415	27.7	28.0	27.7	<1.0	0.10	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.98	0.14	0.12
		473	467	415	27.7	28.0	27.7	<1.0	0.10	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.18	0.47	0.09
		770	770	724	18.0	16.8	18.0	1.50	0.05	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	0.95	0.66	0.09
		358	348	348	20.0	23.0	20.0	23.0	1.00	0.05	<0.03	<0.03	<0.03	<0.03	<0.03	0.92	0.34	0.12
		2702	2702	2702	16.0	13.6	16.0	<1.0	0.05	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.16	0.15	0.13
Finmark (10)	Mattiselva, Kåfj. Altafj. Tverrelva, Altafj. Repparfjordv., Repparfj. Stabburse., I. Porsangen V Lakse., Indre Porsangen S Børselva, Indre Porsangen Ø Matusjäkka, I. Laksetj. V Storelva, Indre Laksetj. V Soussjäkka, I. Laksetj. V Adamselva, I. Laksetj. Ø Tanavassdraget, Tanafj. S Vesterelva, Sytten. V. Jakobse., Y.Varangerfj. Påssvike., Bøkfj.Varangfj. Neiden, Munkfj. Varangfj. Gjense Jakobse., Varangfj.	325	325	319	26.5	21.2	26.5	<1.0	0.10	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.15	1.37	0.12
		234	233	233	15.1	12.0	15.1	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.65	1.34	0.15
		1090	1089	1089	25.0	22.5	25.0	2.00	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.95	1.50	0.10
		1108	1102	870	18.3	15.2	18.3	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.40	2.44	1.46
		1532	941	941	15.9	12.8	15.9	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.51	0.88	0.06
		883	883	863	29.8	29.0	29.8	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.81	0.31	0.11
		101	101	101	22.8	20.5	22.8	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.27	0.52	0.11
		690	690	760	21.9	19.0	19.9	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	1.27	0.52	0.11
		92	92	102	25.3	22.8	22.8	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.11	1.18	0.11
		705	705	760	19.9	15.0	19.9	<1.0	0.15	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	4.45	1.13	0.20
		16389	15713	14169	11.5	9.2	11.5	2.50	0.11	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.36	0.82	0.15
		469	469	79	34.6	31.6	34.6	<1.0	0.03	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	4.95	1.01	0.42
		627	627	239	18.1	14.5	18.1	1.00	0.30	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.06	1.44	0.13
		18404	18400	18175	9.3	10.2	9.3	2.00	0.32	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.38	8.77	0.27
		2960	2960	2911	9.8	8.8	9.8	<1.0	0.35	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	2.04	1.32	0.10
		234	234	234	18.0	18.0	18.0	3.00	0.35	<0.03	<0.03	<0.03	<0.03	<0.03	<0.03	3.49	7.40	0.27

APPENDIX IX : TRIBUTARY RIVERS. ANNUAL LOAD 1994			Page:
Table 9.1	Cond., Nutrients, Heavy metals, Suspended part.matter		75-81
Table 9.2	Mercury, Lindane, PCBs *(Detection limit = limit)		83-89

(1) Glomma	"tributaries"	:	Tista	- Hølenelva
(1) Inner Oslo-fjord		:	Årungelva	- Åroselva
(2) Drammenselva	"tributary"	:	Lierelva	
(3) Numedalslågen	"tributaries"	:	Sandeelva	- Farriselva
(4) Skienselva	"tributary"	:	Tokkeelva	
(5) Otra	"tributaries"	:	Gjerstade.	- Audna
(6) Orreelva	"tributaries"	:	Lygna	- Ulla
(7) Suldalslågen	"tributaries"	:	Saudaelva	- Hornindalselva
(8) Orkla	"tributaries"	:	Ørstaelva	- Salsvatnelva
(9) Vefsna	"tributaries"	:	Åbjøra	- Reisa
(10) Alta	"tributaries"	:	Mattiselva	- Grense Jacobse.

* Measurements below detection limits are treated in two ways:
 "Detection limit = Zero", and "Detection limit = limit". This concerns the substances Cd, Pb, Hg and PCBs. In Tables 9.1-9.2 as well as in Tables 5.1-5.4 both "zero- and limit-values" are shown.

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Runoff data										Parameters (mean values)															
		Drainage area		Sampling station		Discharge gauging station		1994		Cond mS/m		Cu		Zn		C d		P b		S.P.M.		H g limit kg					
Outlet	Sampi. station	Disch. gaug. station	Normal	1994	Normal	1/s sq.km	1/s sq.km	1/s sq.km	1/s sq.km	Tot-P	P04-P	Tot-N	NO3-N	NH4-N	Tons	Tons	Tons	Tons	Tons	Tons	zero tons	limit tons	zero tons	limit tons	t.tons	zero kg	H g limit kg
Østfold (1.)	Tista, Iddefj. Mosselvå, Mossesundet	1588 690	1582 689	14.4 14.5	15.5 15.4	14.4 14.5	15.5 15.4	14.4 14.5	15.5 15.4	2.3 1.7	6.24 9.8	9.0 3.35	7.73 3.36	0.54 0.37	7.11 1.14	0.077 0.033	0.201 0.124	1.24 1.57	0.000 0.502	0.773 0.502	0.127 0.124	0.023 0.023	0.65 0.65	0.127 0.127	0.127 0.127		
Oslo & Akershus (1.)	Hølenelva, Drøbak sundet Ø Arungelva, I. Oslofj. Gjerseløva, I. Oslofj. Lianseleva, I. Oslofj. Loelva/Alna, I. Oslofj. Akerselva, I. Oslofj. Frognerelva, I. Oslofj. Lyseakerelva, I. Oslofj. Sandvikselva, I. Oslofj. Åreselva, I. Oslofj.	137 52 86 42 75 227 225 23 178 223 113	121 50 85 41 69 225 225 20 173 187 109	14.0 13.0 14.0 13.0 13.0 17.5 17.5 20 16.8 18.4 17.0	13.3 12.4 13.3 13.9 22.2 12.6 12.6 15.0 27.1 18.0 16.3	14.4 13.0 14.0 13.0 13.0 12.6 12.6 20.9 16.8 18.4 17.0	15.5 12.4 13.3 13.9 22.2 12.6 12.6 20.9 27.1 18.0 16.3	14.4 12.4 13.3 13.9 22.2 12.6 12.6 20.9 27.1 18.0 16.3	15.5 14.5 15.4 15.4 22.2 13.0 13.0 21.0 27.1 18.0 16.3	2.4 0.8 0.5 1.1 6.7 1.6 1.6 1.0 2.4 1.8 2.1	19.40 26.70 18.25 24.00 31.00 5.80 5.80 1.0 7.40 13.20 13.70	1.7 0.8 0.5 1.1 2.3 0.2 0.2 0.5 0.6 0.8 0.8	151 69 59 25 96 58 58 18 67 123 65	14.67 3.42 1.39 2.73 87 5.10 5.10 7 52 96	0.05 0.02 0.05 0.01 0.16 0.31 0.06 0.07 0.05 0.01 0.38	0.31 0.04 0.01 0.01 0.01 0.78 0.06 0.07 0.35 0.13 0.16	0.008 0.002 0.001 0.001 0.001 0.008 0.008 0.000 0.000 0.001 0.001	0.023 0.003 0.004 0.010 0.022 0.032 0.032 0.003 0.013 0.038 0.013	0.201 0.124 0.010 0.010 0.022 0.34 0.34 0.12 0.53 0.30	1.24 1.57 0.10 0.10 2.08 0.179 0.179 0.046 0.148 0.000	0.000 0.029 0.036 0.054 0.242 0.179 0.179 0.046 0.148 0.056						
Vestfold (3.)	Lierelva, Drammensfl. Ø Sandeelva, Sandebukta Aulleva, Tønsbergfl. Farriselva, Larvikfl.	309 193 363 491	266 190 362 491	18.6 17.0 14.9 21.6	222 16.2 14.6 21.4	28.8 17.0 14.9 21.6	18.6 17.0 14.9 21.6	14.4 12.30 17.22 4.01	9.2 0.8 8.1 2.4	13.50 125 542 236	265 102 358 139	10.39 5.44 13.33 4.97	0.19 0.12 0.18 0.50	2.01 0.014 0.87 1.66	0.075 0.072 0.045 0.003	3.43 0.49 0.45 0.166	0.242 0.097 4.15 0.166	0.242 0.097 0.167 0.331									
Telemark (4.)	Tokkeelva, Kragerø	1238	1200	26.7	28.6	26.7	28.6	4.3	0.5	433	244	42.21	0.65	8.23	0.054	0.400	1.13	1.082	1.082								
Aust-Agder (5.)	Gjerstadelva, Søndeledfl. Vegårdselva, Såndnesfl. Nidelva, Arendal	419 457 4025	414 429 4020	291 291 3956	27.0 29.3 29.8	27.8 30.8 31.3	27.0 30.8 31.3	2.2 2.5 15.9	3.00 3.02 2.17	176 171 1452	98 85 138.88	16.70 20.00 20.00	0.25 0.25 28.57	0.269 0.183 0.278	0.036 0.029 0.278	0.544 0.417 0.2024	0.544 0.417 0.2024										

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Runoff data						Parameters (mean values)												
		Drainage area			Discharge			Tot-P	PO4-P	Tot-N	NO3-N	NH4-N	Cu	Zn	Cd	Pb	S.P.M.	Hg		
		Outlet	Samp. station	Disch. gauging station	Sampling station	Normal 1994 l/s	Normal 1994 mS/m													
		sq.km	sq.km	sq.km	sq.km	l/s sq.km	l/s sq.km	tons	tons	tons	tons	tons	tons	tons	tons	tons	tons	t.tons	kg	
Vest-Agder (5.)	Tordalselva, Tordalsfj.	1856	1854	1794	33.9	31.3	2.38	7.3	0.9	855	326	80.52	0.73	13.73	0.128	1.226	2.18	3.660	3.660	
	Søgneelva, Flekkerøy	204	192	38.0	41.8	38.0	8.03	1.5	0.5	259	202	14.17	0.18	2.66	0.028	0.104	0.39	0.000	0.253	
	Mandalselva, Mannefj.	1809	1800	1740	46.0	50.0	47.6	50.0	2.41	14.2	1.4	1118	607	110.69	1.14	15.04	0.085	1.760	2.92	
	Audna, Sniksfj.	450	400	59	45.0	50.4	51.8	5.27	2.5	0.3	391	273	26.07	0.25	4.90	0.038	0.305	0.77	0.636	4.257
	Lygna, Lyngdalsfj.	664	660	266	48.0	61.6	57.9	71.5	3.24	5.1	0.6	581	313	38.46	0.26	13.46	0.077	0.808	1.53	
	Kvina, Fedafj.	1445	1140	1140	57.6	18.1	57.6	18.1	3.95	3.3	0.7	228	111	15.62	0.13	4.29	0.007	0.560	1.11	
	Sira, Ana-Sira	1916	1872	1872	59.4	57.6	59.4	57.6	2.49	10.2	1.7	901	527	102.01	0.34	14.96	0.034	1.666	1.29	
	Sokndalselva, Sogndalsstr.	294	293	107	51.1	56.7	51.1	53.7	2.6	1.0	194	144	15.72	0.05	4.35	0.005	0.131	0.42	0.000	
	Helelandselva, Egersund	241	240	194	57.5	60.4	71.1	71.6	4.61	3.2	0.9	206	151	5.49	0.05	3.02	0.005	0.174	0.33	
	Birkelandselva, Egersund	705	704	633	77.7	69.3	86.4	78.0	3.70	7.7	0.8	711	572	18.46	0.15	7.39	0.015	0.323	0.457	
Rogaland (6.)	Hælvela, Håtangen	165	160	135	46.9	45.4	46.9	45.4	11.30	5.5	2.5	337	244	14.43	0.02	0.89	0.002	0.064	0.38	
	Figgjo, Solavikta	229	218	135	50.0	52.0	50.0	10.40	6.4	3.2	390	277	23.59	0.25	2.11	0.004	0.004	0.232	0.232	
	Ims-Lutsi, Høgsfj.Boknafj.	127	127	127	34.9	40.9	34.9	40.9	6.63	1.3	0.1	117	89	2.95	0.08	0.44	0.003	0.003	0.023	
	Oltedalsse., Høgsfj.Boknafj.	102	101	129	76.3	70.0	3.62	3.6	0.5	96	70	7.05	1.05	0.07	0.007	0.051	0.174	0.15	0.164	
	Dirdalse., Høgsfj.Boknafj.	158	158	95	83.0	89.6	83.0	2.74	1.3	0.2	144	104	4.91	0.09	1.16	0.009	0.079	0.179	0.12	
	Frafjordse., Frafj. Boknafj.	178	178	124	94.4	101.0	94.4	2.49	2.3	0.3	148	100	7.37	0.11	1.53	0.017	0.238	0.24	0.000	
	Espedalsse., Høgsfj.Boknafj.	138	138	124	90.0	94.5	90.0	2.31	1.2	0.2	105	80	3.70	0.12	1.15	0.025	0.111	0.111	0.411	
	Lysee., Lysefj.Boknafj.	182	182	46	74.0	77.7	74.0	2.50	1.3	0.4	138	89	4.01	0.45	1.34	0.027	0.134	0.23	0.000	
	Ardalse., Ardalsfj.Boknafj.	519	516	501	81.4	85.4	81.4	2.76	15.3	8.3	293	222	15.29	1.39	5.00	0.069	1.404	1.404	0.446	
	Farese., Jøsenfj.Boknafj.	163	163	163	85.3	90.1	85.8	2.30	1.4	0.2	117	115	3.24	0.14	0.42	0.000	0.093	0.093	0.463	
(7.)	Ula., Jøsenfj.Boknafj.	393	393	385	83.4	87.6	83.4	2.32	3.3	0.5	347	266	14.11	0.43	2.93	0.087	0.326	0.47	0.000	
	Sædæ., Sædæfj.Boknafj.	353	353	82	93.5	85.0	85.0	3.49	3.1	0.5	916	827	3.12	0.73	21.86	0.104	0.114	0.36	1.041	
	Abælvæ., Abælvæfj.Boknafj.	118	117	80.0	103.1	80.0	103.1	2.75	1.9	1.1	108	63	4.18	0.15	0.31	0.002	0.027	0.027	0.000	
	Vikedalsse., Vikedalsfj.															0.015	0.072	0.45	0.380	

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Runoff data						Parameters (mean values)														
		Drainage area			Discharge			Tot-P	PO4-P	Tot-N	NO3-N	NH4-N	Cu	Zn	Cd	Pb	S.P.M.	Hg				
		Outlet	Sampi.	Disch.	Sampling station	gauging station	Normal															
		sq.km	sq.km	sq.km	Normal	1994	l/s sq.km	l/s sq.km	l/s sq.km	1994	Normal	l/s sq.km	Cond	tons	tons	tons	tons	tons	kg			
(7.)	Hordaland Etheelva, Etnefj. Bømlafj. Opo, Særfj. Hardangerfj. Tysso, Særfj. Hardangerfj. Kinsø, Særfj. Hardangerfj. Veig, Eidsfjv. Hardangerfj. Bjoreia, " , Hardangerfj. Sima, Eidsfj. Hardangerfj. Austdøla, Osafj. Eldfj. Norddøla, Osafj. Eldfj. Tysseliva, Fusafj. Ossefja, Fusafj. Bergsdalselva, Veafj. Herdalsfj. Vosso, Veafj. Særfj. Eltsø, Osterfj. Modalselva, Osterfj.	252	250	127	48.8	58.4	96.0	115.0	3.10	2.1	0.2	185	152	6.45	0.14	1.24	0.018	0.041	0.33	0.000	0.460	
		482	480	464	79.3	93.1	79.3	93.1	1.54	3.5	0.7	247	152	8.46	0.14	4.37	0.056	0.282	1.18	2.819	2.819	
		388	385	407	79.3	91.2	79.3	91.2	1.78	2.2	0.6	221	112	5.54	0.11	2.99	0.044	0.044	0.12	1.107	1.107	
		281	281	232	46.0	59.3	46.0	59.3	2.01	1.1	0.3	50	33	2.63	0.05	0.74	0.005	0.005	0.026	0.000	0.525	
		496	496	386	41.8	46.2	41.8	46.2	2.04	2.5	0.4	105	33	3.61	0.07	0.36	0.000	0.007	0.051	0.23	0.723	
		592	592	592	26.0	28.6	26.0	28.6	2.04	2.1	0.3	101	25	2.67	0.05	0.27	0.000	0.005	0.037	0.17	0.534	
		145	145	128	69.2	76.8	69.2	76.8	2.21	1.2	0.2	63	33	1.76	0.04	0.18	0.014	0.014	0.000	0.21	0.351	
		131	130	89	74.6	82.0	74.6	82.0	1.28	1.0	0.7	68	50	1.68	0.07	0.27	0.000	0.003	0.013	0.05	0.000	
		40	39	39	74.6	82.0	74.6	82.0	8.52	0.3	0.1	21	18	0.50	0.01	0.05	0.000	0.001	0.002	0.03	0.000	
		240	240	85.0	93.5	85.0	93.5	2.02	2.1	0.4	144	82	4.25	0.35	1.91	0.00	0.007	0.184	0.184	0.708	0.708	
		109	108	50	91.7	92.6	91.7	92.6	3.04	3.2	2.2	95	56	1.58	0.32	3.15	0.003	0.003	0.095	0.095	0.315	
		198	198	80.0	86.4	80.0	86.4	1.91	1.6	0.3	74	46	3.24	0.05	1.13	0.016	0.016	0.065	0.065	0.25	0.000	0.539
		1492	1465	1102	58.2	76.5	58.2	76.5	1.38	7.1	1.8	452	272	28.27	0.71	4.59	0.071	0.071	0.283	1.98	8.836	8.836
		414	400	342	86.2	96.5	86.2	96.5	1.92	4.9	2.4	224	136	6.09	1.22	0.012	0.012	0.122	0.122	0.000	0.217	
		385	384	248	95.5	105.0	95.5	105.0	1.82	5.1	0.6	286	211	6.36	1.27	6.36	0.013	0.013	0.127	0.64	0.000	1.272

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

Table 9.1 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Runoff data						Parameters (mean values)													
		Drainage area			Discharge gauging station			Tot-P	PO4-P	Tot-N	NO3-N	NH4-N	Cu	Zn	Cd	Pb	S.P.M.	Hg limit			
		Outlet	Samp. station	Disch. gaug. station	Sampling station	Normal	1994														
(9.)	Abøra, Bindalsfj. S	526	520	384	80.2	59.0	80.2	3.99	4.8	1.0	110	23	2.90	0.87	0.48	0.010	0.281	2.68	0.000	0.968	
	Skjerva, Vefsenvfj. S	104	104	98	41.3	36.3	41.3	7.41	1.1	0.7	27	7	4.76	0.06	0.12	0.011	0.017	0.10	0.238	0.298	
	Fusta, Vefsenvfj. N	544	543	520	63.4	57.0	63.4	3.81	2.9	0.5	117	19	9.76	0.39	1.37	0.000	0.010	0.166	1.17	0.976	
	Drevia, Versevff. N	177	176	98	65.0	58.5	65.0	3.96	1.6	0.6	36	11	0.97	0.16	0.52	0.003	0.075	1.00	0.000	0.325	
	Røsståga, Sørff.	2092	2087	1880	45.4	51.0	45.4	51.0	5.27	13.4	3.4	470	114	36.92	2.35	16.45	0.067	1.074	1.074	7.75	3.357
	Bjerkå, Sørff.	385	385	273	55.4	48.7	55.4	2.61	1.8	0.6	51	11	1.77	0.24	0.30	0.006	0.041	0.85	0.591	0.591	
	Dalselva, Ranafj. N	211	211	129	39.5	35.5	39.5	2.53	0.2	0.7	21	4	0.71	0.14	0.28	0.005	0.005	0.46	0.000	0.236	
	Ranavassdraget, Ranafj. N	3847	3846	1892	44.9	50.0	44.9	50.0	3.90	18.2	3.0	631	309	18.19	2.43	6.67	0.000	0.061	5.943	5.943	
	Fykanåga, Glomfjord	297	297	243	103.7	100.0	103.7	2.91	2.8	1.9	60	37	6.56	0.47	0.94	0.009	0.009	0.281	1.88	0.000	
	Belare, Belafj. Nordff.	1064	875	797	45.1	30.3	45.1	30.3	3.67	1.7	0.8	74	33	2.51	0.50	0.84	0.008	0.008	0.251	2.13	0.000
	Saldalsvassdr., Salidff. S	1544	1543	1168	32.1	26.6	32.1	2.79	5.2	3.9	124	70	3.88	0.65	1.94	0.026	0.026	1.398	1.398	6.92	
	Sultielmavassdr., Saltff. ffl	1028	800	791	44.0	36.0	44.0	42.00	0.9	0.5	87	40	6.36	12.62	10.35	0.064	0.064	1.971	0.59	0.908	
	Kobbe, Leirff. Sørifolda N	405	405	386	66.9	62.7	66.9	1.50	0.8	0.4	48	21	2.40	0.24	1.20	0.008	0.008	0.312	0.312	0.23	
	Skjoma, Ofotff. S	845	840	797	36.3	29.0	36.3	1.81	1.5	0.4	52	8	2.30	0.38	1.08	0.023	0.023	0.161	0.161	0.768	

Table 9.1 TRIBUTARY RIVERS: ANNUAL LOAD 1994.

County		Watercourse		Runoff data								Parameters (mean values)									
				Discharge				gauging station				Cond mS/m				Cu					
				Outlet station	Sampling station	Gauging station	Normal 1994 I/s sq.km	Tot-P tons	P O4-P tons	Tot-N tons	NO3-N tons	Cu tons	Zn tons	Cd zero tons	Pb limit tons	S.P.M. zero t/tons	Hg zero kg	Limit kg			
Troms (9.)	Spanselva, Astafj., Vågsfj., Salangsæ., Astafj., Vågsfj., Rossfjorde., Målangen Målse., Målselvflj., Bardlie., Målselvåa Nordkjøselva, Balsflj. Signaldalselva, Lyngen V Skibotnelva, Lyngen Kåfjordelva, Lyngen Ø Reisa, Reisafj.	142	142	533	50.0	51.0	50.0	7.25	0.5	0.1	20	9	0.69	0.11	0.27	0.021	0.153	0.23	0.000	0.228	
		539	539	533	40.9	51.0	40.9	7.27	2.6	0.4	77	38	2.60	0.26	0.43	0.000	0.182	0.73	0.000	0.867	
		196	190	39.5	41.5	39.5	8.88	1.0	0.1	39	1	3.98	0.10	0.12	0.000	0.002	0.030	0.030	0.17	0.497	
		3239	3200	3118	28.7	28.1	28.7	6.13	11.3	2.8	337	111	6.51	1.42	1.99	0.028	0.539	7.17	0.000	2.836	
		2906	2906	2049	28.3	27.8	28.3	6.13	10.2	2.5	303	99	7.64	1.27	1.78	0.025	0.484	6.45	0.000	2.548	
		191	191	415	27.7	28.0	27.7	4.31	0.5	0.1	13	6	0.51	0.08	0.08	0.000	0.002	0.005	0.015	0.000	0.169
		473	467	415	27.7	28.0	27.7	4.29	4.1	2.9	37	22	1.24	0.37	0.41	0.000	0.004	0.103	0.553	0.000	0.412
		770	770	724	18.0	16.8	18.0	2.99	0.8	0.2	39	18	2.86	0.24	0.20	0.004	0.012	0.012	0.39	0.612	0.612
		358	358	348	20.0	23.0	20.0	23.0	0.3	0.1	38	27	0.78	0.29	0.16	0.003	0.005	0.005	0.07	0.000	0.260
		2702	2702	16.0	13.6	16.0	3.76	2.3	0.6	132	57	11.59	1.62	2.20	0.035	0.035	0.209	0.209	1.54	0.000	1.159
		325	325	319	26.5	21.2	26.5	2.75	0.7	0.1	20	3	1.30	0.13	0.11	0.000	0.002	0.035	0.19	0.000	0.217
		234	234	233	15.1	12.0	15.1	4.27	0.4	0.2	12	4	0.62	0.05	0.04	0.001	0.001	0.009	0.005	0.000	0.088
		1090	1089	25.0	22.5	25.0	4.50	1.5	0.4	92	19	6.18	0.31	0.39	0.000	0.008	0.108	0.29	0.545	1.545	
(10.)	Mattuslåkka, I. Laksflj. V Storelva, Indre Lakesflj. V Soussjåkka, I. Laksflj. V Adamseva, I. Laksflj. Ø Tanavassdræget, Tanafj. S Vesterelva, Syteflj. V Jakobse, Y Varangerflj. Passyve, Bøkfj. Varangflj. Neldsen, Munkflj. Varangflj. Grense Jakobse, Varangflj.	1108	1102	870	18.3	15.2	18.3	3.02	1.1	0.3	55	13	4.75	0.21	0.69	0.048	0.048	1.231	0.39	0.000	0.528
		1533	1532	941	15.9	12.8	15.9	4.41	3.1	0.6	83	9	1.86	0.62	0.62	0.006	0.006	0.458	1.49	0.000	0.618
		883	883	663	29.8	29.0	29.8	3.74	0.8	0.4	48	11	8.08	0.16	0.97	0.008	0.008	0.129	0.50	0.000	0.808
		101	101	101	22.8	20.5	22.8	4.01	0.2	0.1	4	1	0.33	0.04	0.03	0.000	0.001	0.010	0.010	0.000	0.065
		690	690	760	21.9	19.0	19.9	2.10	0.4	0.2	37	31	1.24	0.21	1.61	0.017	0.017	0.215	0.04	0.000	0.413
		92	92	102	25.3	22.8	22.8	4.68	0.1	0.0	5	2	0.20	0.02	0.11	0.001	0.009	0.009	0.02	0.000	0.066
		705	705	760	19.9	15.0	19.9	6.14	0.7	0.2	35	2	3.67	0.20	0.50	0.010	0.010	0.053	0.20	0.000	0.333
		16389	15713	14169	11.5	9.2	11.5	3.96	27.4	9.1	83	46	91.18	3.19	2.28	0.046	0.046	1.459	7.20	11.397	11.397
		469	469	79	34.6	31.6	34.6	8.55	1.9	0.5	28	2	1.40	0.19	0.23	0.014	0.014	0.042	0.10	0.000	0.467
		627	627	239	18.1	14.5	18.1	5.72	1.1	0.6	39	1	4.01	0.11	0.20	0.003	0.003	0.029	0.05	0.029	0.287
		18404	18400	18175	9.3	10.2	9.3	4.37	47.3	11.8	1095	30	124.29	12.43	11.25	0.000	0.059	1.243	19.47	11.837	11.837
		2960	2960	2911	9.8	8.8	9.8	2.95	2.5	0.8	152	28	2.46	0.58	0.82	0.000	0.008	0.197	0.79	0.000	0.821
		234	234	234	18.0	18.0	18.0	4.20	0.4	0.1	23	1	3.59	0.35	0.35	0.003	0.003	0.23	0.179	0.11	0.398

Page:**Table 9.2**

83-89

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Parameters (mean values)															As limit tons	
		PCB (The following Congeners) IUPAC NOS																
		Gamma HCH kg	28 zero kg	52 limit kg	101 zero kg	118 limit kg	138 zero kg	153 limit kg	180 zero kg	Sum : PCB kg	TOC t tons	Cr-T zero tons	Ni limit zero tons	Cr-T zero tons	Ni limit zero tons			
Vest-Agder (5.)	Tovdalselva, Tøpdalsfj.	1.830	0.000	0.055	0.000	0.055	0.000	0.055	0.000	0.055	0.000	0.055	0.000	0.384	6.92	1.54	0.40	0.64
	Søgneelva, Flekkerøy	0.202	0.000	0.008	0.000	0.008	0.000	0.008	0.000	0.008	0.000	0.008	0.000	0.053	0.96	0.45	0.22	0.08
	Mandalselva, Mannenfj.	1.987	0.000	0.085	0.000	0.085	0.000	0.085	0.000	0.085	0.000	0.085	0.000	0.596	8.15	1.87	0.20	0.54
	Audna, Sniksfj.	0.318	0.000	0.019	0.000	0.019	0.000	0.019	0.000	0.019	0.000	0.019	0.000	0.134	1.84	0.90	0.08	0.25
	Lygna, Lyngdalsfj.	0.641	0.000	0.038	0.000	0.038	0.000	0.038	0.000	0.038	0.000	0.038	0.000	0.269	4.24	2.33	0.35	0.19
	Kvina, Fedafj.	0.325	0.000	0.020	0.000	0.020	0.000	0.020	0.000	0.020	0.000	0.020	0.000	0.137	2.54	0.93	0.03	0.21
	Sira, Ana-Sira	1.700	0.000	0.102	0.000	0.102	0.000	0.102	0.000	0.102	0.000	0.102	0.000	0.714	5.44	2.79	0.24	0.48
	Sokndalselva, Sogndalsst.	0.472	0.000	0.016	0.000	0.016	0.000	0.016	0.000	0.016	0.000	0.016	0.000	0.110	0.84	0.42	1.50	0.04
	Helelandselva, Egersund	0.411	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.096	0.78	0.39	0.21	0.10
	Birkelandselva, Egersund	1.385	0.000	0.046	0.000	0.046	0.000	0.046	0.000	0.046	0.000	0.046	0.000	0.323	11.92	0.75	0.78	1.34
Rogaland (6.)	Håelva, Håtangen	0.186	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.048	1.17	0.20	0.20	0.03
	Figgjo, Solavika	0.286	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.075	1.39	0.63	0.15	0.05
	Ims-Lutsi, Høgsfj.Boknafj.	0.131	0.000	0.005	0.000	0.005	0.000	0.005	0.000	0.005	0.000	0.005	0.000	0.034	0.56	0.66	0.04	0.04
	Oltedalse., Høgsfj.Boknafj.	0.194	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.051	0.34	0.62	0.03	0.04
	Dirdalse., Høgsfj.Boknafj.	0.313	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.094	0.42	0.75	0.01	0.06
	Frafjorde., Fraflj.Boknafj.	0.340	0.000	0.017	0.000	0.017	0.000	0.017	0.000	0.017	0.000	0.017	0.000	0.119	0.74	0.83	0.02	0.06
	Espedalsse., Høgsfj.Boknafj.	0.206	0.000	0.012	0.000	0.012	0.000	0.012	0.000	0.012	0.000	0.012	0.000	0.086	0.45	0.76	0.01	0.04
	Lysee., Lysefj.Boknafj.	0.178	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.013	0.000	0.094	0.71	0.14	0.03	0.01
	Ardalase., Ardalsfj.Boknafj.	0.500	0.000	0.042	0.000	0.042	0.000	0.042	0.000	0.042	0.000	0.042	0.000	0.292	1.49	2.79	0.43	0.53
	Fareæ., Jøsenfj.Boknafj.	0.185	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.014	0.000	0.097	0.88	0.50	0.02	0.05
(7.)	Ula., Jøsenfj.Boknafj.	0.543	0.000	0.033	0.000	0.033	0.000	0.033	0.000	0.033	0.000	0.033	0.000	0.228	4.23	0.35	0.07	0.03
	Saudaæ., Saudafj.Boknafj.	0.208	0.000	0.031	0.000	0.031	0.000	0.031	0.000	0.031	0.000	0.031	0.000	0.219	2.50	1.09	0.12	0.07
	Abøelva, Sauðafj.Boknafj.	0.048	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.051	0.23	0.01	0.01	0.01
	Vikedalsse., Boknafj.	0.076	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.011	0.000	0.080	0.37	0.82	0.13	0.08

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Parameters (mean values)																
		PCB (The following Congeners) IUPAC NOS																
Gamma	28	52	101	118	138	153	180	Sum : PCB	TOC	Cr-T	Cr-zero	Ni	Aszero	Aslimit	As tons			
HCH	zero	limit	zero	limit	zero	limit	zero	kg	kg	kg	kg	kg	kg	kg	kg	t.tons	t.tons	t.tons
kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg
Sogn og Fjordane (7.)	Nærøye., Aurl. fl. Sognefj. Flåmse., Aurl. fl. Sognefj. Aurlandy. Aurl. fl. Sognefj. Erdalse., Lærd. fl. Sognefj. Lærdalsv. Lærd. fl. Sognefj. Ardalsv., Ardalsfl. Sognefj. Fortunv., Lustefj. Sognefj. Mørkrivs., Lustefj. Sognefj. Jostedøla, " Sognefj. Arsv. Sogn. fl. Sognefj. Sogndalsse, " Sognefj. Gaujar, Dalsfl. Buff. Jøstra, Førdefl. Nausla, Førdefl. Oselva, Høydalsfl. Hopse., Hyet, Nordfl. S Glengedalse,, Nordfl. S Bremlse., Gløppenfl. " Oldene., Indre Nordfl. Lønnevat, Indre Nordfl. Strynese., Indre Nordfl. Hornindalsse., Nordfl. N	0.122 0.122 0.163 0.022 0.163 0.375 0.252 0.143 0.581 0.341 0.118 0.666 0.553 0.261 0.234 0.057 0.131 0.453 0.167 0.171 0.332 0.342	0.000 0.000	0.018 0.018 0.041 0.004 0.025 0.045 0.027 0.015 0.062 0.036 0.013 0.055 0.055 0.055 0.026 0.023 0.006 0.013 0.045 0.017 0.017 0.033 0.026	0.000 0.000	0.018 0.018 0.041 0.004 0.025 0.045 0.027 0.015 0.062 0.036 0.013 0.055 0.055 0.055 0.026 0.023 0.006 0.013 0.045 0.017 0.017 0.033 0.026	0.000 0.000	0.018 0.018 0.041 0.004 0.025 0.045 0.027 0.015 0.062 0.036 0.013 0.055 0.055 0.055 0.026 0.023 0.006 0.013 0.045 0.017 0.017 0.033 0.026	0.000 0.000	0.128 0.128 0.041 0.004 0.025 0.045 0.027 0.015 0.062 0.036 0.013 0.055 0.055 0.055 0.026 0.023 0.006 0.013 0.045 0.017 0.017 0.033 0.026	0.24 0.18 0.54 0.004 0.000 0.045 0.027 0.015 0.062 0.036 0.013 0.055 0.055 0.055 0.026 0.023 0.006 0.013 0.045 0.017 0.017 0.033 0.026	0.82 0.43 0.68 0.004 0.000 0.14 0.18 0.20 0.12 0.12 0.14 0.27 0.20 0.15 0.15 0.17 0.13 0.13 0.18 0.11 0.11 0.11 0.11	0.06 0.09 0.04 0.01 0.01 0.12 0.13 0.15 0.04 0.04 0.15 0.09 0.01 0.01 0.21 0.18 0.18 0.02 0.03 0.03 0.06 0.09	0.06 0.03 0.01 0.03 0.04 0.15 0.09 0.02 0.10 0.04 0.05 0.02 0.01 0.01 0.21 0.18 0.18 0.01 0.03 0.03 0.06 0.09				

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

County	Watercourse	Parameters (mean values)														Cr-T	TOC	N	As		
		PCB (The following Congeners) IUPAC NOS														zero	zero	zero	zero	zero	zero
		Gamma	28	52	101	118	138	153	180	Sum : PCB	Cr-T	N	As	zero							
		HCH	zero	limit	zero	limit	zero	limit	zero	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg
		kg	kg	kg	kg	kg	kg	kg	kg												
(9.)	Abjøra, Blindselvflj. S	0.290	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.203	2.44
	Skjerva, Vefsenvflj. S	0.048	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.004	0.000	0.025	0.31
	Fusta, Vefsenvflj. N	0.390	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.029	0.000	0.205	1.81
	Drevla, Vefsenvflj. N	0.130	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.010	0.000	0.068	0.56
	Røssåga, Sørflj.	1.007	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.101	0.000	0.705	5.03
	Blerka, Sørflj.	0.177	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.018	0.000	0.124	0.54
	Dalselva, Ranafj. N	0.059	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.007	0.000	0.050	0.33
	Ranavassdraget, Ranafj. N	1.516	0.000	0.182	0.000	0.182	0.000	0.182	0.000	0.182	0.000	0.182	0.000	0.182	0.000	0.182	0.000	0.182	0.000	1.274	8.00
	Fykanåga, Glomiford	0.234	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.028	0.000	0.197	1.24
	Belare., Belarfj. Nordflj.	0.418	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.025	0.000	0.176	2.17
	Saltdalsvassdr., Saltdflj. S	0.906	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.039	0.000	0.272	3.09
	Sulttelmavassdr., Saltdflj. S	0.672	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.027	0.000	0.191	17.71
	Kobbe., Leirfj. Sørfolda N	0.400	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.024	0.000	0.168	1.03
	Skjoma, Otoffl. S	0.307	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.023	0.000	0.161	0.60

Table 9.2 TRIBUTARY RIVERS. ANNUAL LOAD 1994.

APPENDIX X :	"MEAN" TOTAL DISCHARGES (Mean concentrations of main and tributary rivers multiplied with mean runoff 1961-90 (main rivers), 1931-60 (tributary rivers).	Page:
Table X	"Mean" total discharges from mainland Norway to convention waters	91
Table 10.1	"Mean" total discharges to the Skagerrak region	92
Table 10.2	"Mean" total discharges to the remaining North Sea	93
Table 10.3	"Mean" total discharges to the Norwegian Sea region	94
Table 10.4	"Mean" total discharges to the Barents Sea region	95
Table 10.5A	Main rivers. "Mean" load (Mean concentrations multiplied with mean runoff, 1961-90) *(Detection limit = limit)	96
Table 10.5B	Main rivers. "Mean" load (Mean concentrations multiplied with mean runoff, 1961-90) *(Detection limit = zero)	97
Table 10.6	The Skagerrak region. "Mean" inputs from tributary rivers in the Sub-areas (1-5)	98
Table 10.7	The remaining North Sea. "Mean" inputs from tributary rivers in the Sub-areas (6-7)	99
Table 10.8	The Norwegian Sea region. "Mean" inputs from tributary rivers in the Sub-areas (8-9)	100
Table 10.9	The Barents Sea region. "Mean" inputs from tributary rivers in the Sub-area (10)	101

- * Measurements below detection limits are treated in two ways:
 "Detection limit = Zero", and "Detection limit = limit". This concerns the substances Cd, Pb, Hg and PCBs. In Table 10.5A the "limit-values" are shown, in Table 10.5B the "zero-values" are presented.

**Table X "MEAN" TOTAL DISCHARGES from MAINLAND NORWAY
to convention waters (Mean runoff 1961 - 90, main rivers,
1931 - 60, tributary rivers (Fig. I)).**

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		1.2	4.3	*	3.7	*
Cadmium			4.6	**	3.7	**
Mercury	105	135	*	108	*	347
Mercury		203	**	135	**	442
Copper	49	101		102		252
Zinc	140	382		390		912
Lead	6.3	48.8	*	36.5	*	91.6
Lead		48.9	**	36.5	**	91.7
Arsenic	0.5	23.2		16.4	*	40.1
Arsenic		23.2		16.4	**	40.2
Cr-T	6.5	199.6	*	67.2	*	273.2
Cr-T		199.6	**	67.2	**	273.2
Ni	14.2	87.2	*	46.1	*	147.5
Ni		87.2	**	46.1	**	147.5
PCBs ***		0.5	*	32.1	*	32.6
PCBs		29.7	**	34.4	**	64.1
gamma-HCH		55		37		92
NH4-N	1637	11263	1730	1344	15974	tonnes
NO3-N	14020	194	17100	16416	47730	tonnes
PO4-P	201	1163	199	199	1763	tonnes
Total N	22364	17783	32791	26850	99788	tonnes
Total P	766	2180	672	510	4128	tonnes
S.P.M.		5675905	219552	240780	6136236	tonnes
TOC		18053	173	192090	210316	tonnes
COD		215420			215420	tonnes
AOX		210			210	tonnes

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

**Table 10.1 "MEAN" TOTAL DISCHARGES to The Skagerrak Region
(Mean runoff 1961 - 90, main rivers,
1931 - 60, tributary rivers (Fig. I.I)).**

The Skagerrak Region with main rivers (1) Glomma, (2) Drammenselva, (3) Numedalslågen,
(4) Skienselva, (5) Otra

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.09	0.8	*	3.1	*
Cadmium			0.8	**	3.1	**
Mercury	42.07	12	*	87	*	141
Mercury		17	**	109	**	168
Copper	20.65	9		54		83
Zinc	19.07	99		276		395
Lead	0.91	6.9	*	27.0	*	34.8
Lead		6.9	**	27.0	**	34.8
Arsenic	0.28	3.3	*	12.6	*	16.2
Arsenic		3.3	**	12.6	**	16.2
Cr-T	4.38	11.5	*	49.6	*	65.6
Cr-T		11.6	**	49.6	**	65.6
Ni	10.82	4.6	*	34.9	*	50.3
Ni		4.6	**	34.9	**	50.3
PCBs ***		0.1	*	26.4	*	26.5
PCBs		2.8	**	27.6	**	30.4
gamma-HCH		11		33		44
NH4-N	158	4616	523	1129		6426
NO3-N	1560	150	4690	14659		21059
PO4-P	18	106	32	158		314
Total N	2393	7497	7494	22709		40093
Total P	70	265	109	427		871
S.P.M.		16466	28662	217490		262618
TOC		6838	49	153780		160666
COD		117370				117370
AOX		210				210

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

**Table 10.2 "MEAN" TOTAL DISCHARGES to The Remaining North Sea
 (Mean runoff 1961 - 90, main rivers,
 1931 - 60, tributary rivers (Fig. I.III)).**

The North Sea Region with main rivers : (6) Orreelva, (7) Suldalslågen

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.99	1.1	*	0.1	*
Cadmium			1.2	**	0.1	**
Mercury		42.39	28	*	0	*
Mercury			54	**	3	**
Copper		6.76	13		1	
Zinc		75.75	144		6	
Lead		4.59	10.2	*	0.4	*
Lead			10.3	**	0.4	**
Arsenic		0.00	5.7	*	0.3	*
Arsenic			5.7	**	0.3	**
Cr-T		1.34	37.4	*	4.0	*
Cr-T			37.4	**	4.0	**
Ni		2.19	6.0	*	2.1	*
Ni			6.0	**	2.1	**
PCBs ***			0.0	*	0.0	*
PCBs			9.1	**	0.6	**
gamma-HCH			16		2	
NH4-N	579	3103	521		24	
NO3-N	5161	21	7386		583	
PO4-P	56	275	53		2	
Total N	8286	4742	11467		782	
Total P	198	537	181		9	
S.P.M.		2106041	39171		2440	
TOC		5287	72		2024	
COD		38768				38768

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

****) Detection limit = Limit**

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

**Table 10.3 "MEAN" TOTAL DISCHARGES to The Norwegian Sea
(Mean runoff 1961 - 90, main rivers,
1931 - 60, tributary rivers (Fig. I.III)).**

The Norwegian Sea Region with main rivers (8) Orkla, (9) Vefsna

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.14	2.1	*	0.4	*
Cadmium			2.3	**	0.4	**
Mercury	19.00		67	*	17	*
Mercury			99	**	19	**
Copper	21.58		60		44	
Zinc	44.71		119		105	
Lead	0.81		25.6	*	8.8	*
Lead			25.6	**	8.8	**
Arsenic	0.27		10.9	*	2.6	*
Arsenic			10.9	**	2.6	**
Cr-T	0.70		106.2	*	10.2	*
Cr-T			106.2	**	10.2	**
Ni	1.15		16.9	*	7.0	*
Ni			16.9	**	7.0	**
PCBs ***			0.4	*	5.2	*
PCBs			14.0	**	5.6	**
gamma-HCH			24		2	
NH4-N	808	3284	414		173	
NO3-N	6289	22	4795		1039	
PO4-P	110	755	87		18	
Total N	10004	5195	11046		2868	
Total P	410	1331	288		39	
S.P.M.	1421222		119930		12242	
TOC		5588	30		26691	
COD		57745				57745
AOX		1				1

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

**Table 10.4 "MEAN" TOTAL DISCHARGES to The Barents Sea
(Mean runoff 1961 - 90, main rivers,
1931 - 60, tributary rivers (Fig. I.IV)).**

The Barents Sea Region with main river (10) Alta

Substance:	Area runoff	Direct Discharges	Tributary Inputs	Main Riverine Inputs	Grand Total	
Cadmium		0.00	0.2	*	0.11	*
Cadmium			0.3	**	0.11	**
Mercury		1.35	28	*	4.11	*
Mercury			33	**	4.11	**
Copper		0.37	19		3.02	
Zinc		0.43	21		2.03	
Lead		0.01	6.1	*	0.25	*
Lead			6.1	**	0.25	**
Arsenic		0.00	3.2	*	0.88	*
Arsenic			3.2	**	0.88	**
Cr-T		0.04	44.4	*	3.29	*
Cr-T			44.4	**	3.29	**
Ni		0.06	59.8	*	2.06	*
Ni			59.8	**	2.06	**
PCBs ***			0.0	*	0.49	*
PCBs			3.7	**	0.66	**
gamma-HCH			4		0.14	
NH4-N	91	260	273		16	640
NO3-N	1010	2	229		134	1375
PO4-P	18	28	27		21	94
Total N	1681	349	2784		491	5305
Total P	88	46	95		35	264
S.P.M.		2132176	31789		8608	2172572
TOC		340	23		9595	9958
COD		1537				1537

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

*** the following congeners: IUPAC Nos. 28,52,101,118,153,138,180

Table 10.5 A Main rivers. "Mean" load (Mean concentrations multiplied with mean runoff 1961-90) (Limit-values=limits).

Watercourse	Runoff data				Discharge				Parameters (mean values)									
	Drainage area		Sampl. station	Disch. gaug. station	Sampling station		gauging station		Cond	Tot-P	PO4-P	Tot-N	NH4-N	Cu	Zn	Cd limit tons	Pb limit tons	S.P.M. t.tons
	Outlet	sq.km			Normal	1994	Normal	1994										
	sq.km	sq.km	sq.km	sq.km	I/s	sq.km	I/s	sq.km	ms/m	tons	tons	tons	tons	tons	tons	tons	tons	tons
Glomma, Hvaler-Singlefj.	41918	41218	40221	16.5	15.7	16.9	16.1	6.07	295.98	117.96	12311	8150	686.32	35.17	182.09	2.145	19.088	152.28
Drammensvassdr. Dr.fj. V	17034	17028	16020	17.1	17.5	18.2	18.6	3.58	51.42	15.61	4711	3039	146.92	7.81	29.75	0.275	2.663	23.69
Numedalslågen, Larvikfj.	5577	5513	5197	21.2	22.3	21.2	22.3	3.05	28.38	11.43	1589	921	103.20	3.83	19.98	0.184	1.917	15.00
Skimensvassdr, Grenlandsfj.	10772	10348	10348	25.3	25.1	25.3	25.1	2.18	26.42	7.27	2774	1874	132.10	4.54	21.05	0.330	1.404	8.92
Otra, Kr.Sandsfj.	3738	3730	3668	39.8	42.6	39.8	42.6	2.48	24.58	6.09	1325	674	60.86	2.34	23.60	0.187	1.966	17.60
Orreleva, Ørrestrand	105	105	54	36.7	31.1	40.7	34.6	20.48	4.73	1.12	164	92	5.47	0.17	0.31	0.007	0.043	0.60
Suldalsl.,Sandsfj.Boknafj.	1457	1457	1457	59	36	59	36	2.86	4.74	1.36	618	491	18.98	0.92	5.42	0.054	0.380	1.84
Orkla, Orkdalsfj.Tr.h.fj.	3053	2872	2247	21.7	14.7	21.7	14.7	7.00	9.18	2.30	653	425	19.65	15.41	51.16	0.157	0.334	3.30
Vefsna, Verfsenfj. S	4122	4113	3323	43.9	30.9	43.9	30.9	6.40	29.44	15.20	2215	615	153.74	28.64	54.32	0.285	8.484	8.94
Altælva, Altafj.	7373	7367	6257	11.8	9.3	11.8	9.3	12.61	34.95	20.92	491	134	16.45	3.02	2.03	0.110	0.247	8.61

Watercourse	Parameters (mean values)								PCB (The following Congeners) IUPAC NOS								SUM : kg			
	Hg		Gamma		28.		52		101		118		138.		153.		180		kg	
	limit	kg	Hg	HCH	limit	kg	limit	kg	limit	kg	limit	kg	limit	kg	limit	kg	limit	kg		
Glomma, Hvaler-Singlefj.	69.276	12.440	0.643	1.501	2.788	2.574	2.574	1.930	0.429	12.440	79.356	28.096	19.732	8.579	19.732	1.561	5.418			
Drammensvassdr. Dr.fj. V	9.183	6.244	0.275	0.551	1.010	0.918	0.918	0.735	0.184	4.591	28.466	8.081	5.418	1.561	5.418	0.700	2.285			
Numedalslågen, Larvikfj.	17.655	1.622	0.111	0.258	0.590	0.516	0.553	0.369	0.147	2.543	12.532	3.133	3.133	2.972	2.972	0.991	4.706			
Skimensvassdr, Grenlandsfj.	8.256	8.669	0.165	0.743	1.156	1.156	1.321	0.826	0.330	5.697	18.164	4.706	4.706	4.448	4.448	0.796	5.618			
Otra, Kr.Sandsfj.	4.682	3.886	0.140	0.234	0.515	0.468	0.468	0.328	0.140	2.294	15.262	0.668	0.668	0.358	0.358	0.033	0.358			
Orreleva, Ørrestrand	0.122	0.157	0.004	0.004	0.004	0.004	0.004	0.004	0.004	0.004	0.026	0.026	0.026	0.026	0.026	0.026	0.026			
Suldalsl.,Sandsfj.Boknafj.	2.711	2.250	0.081	0.081	0.081	0.081	0.081	0.081	0.081	0.081	0.569	1.355	3.687	3.687	1.762	1.762	0.244			
Orkla, Orkdalsfj.Tr.h.fj.	1.965	0.373	0.059	0.059	0.059	0.059	0.059	0.059	0.059	0.059	0.413	7.331	2.260	2.260	1.336	1.336	1.101			
Vefsna, Verfsenfj. S	16.855	1.310	1.765	1.310	0.740	0.342	0.285	0.285	0.456	0.182	19.360	7.972	5.694	5.694	1.537	1.537	0.877			
Altælva, Altafj.	4.112	0.137	0.082	0.082	0.082	0.164	0.110	0.082	0.055	0.082	0.658	9.595	3.290	3.290	2.056	2.056	0.877			

Table 10.5 B Main rivers. 'Mean' load (Mean concentrations multiplied with mean runoff 1961-90) (Limit-values=zero).

Watercourse	Runoff data										Parameters (mean values)											
	Drainage area		Sampling station		Discharge gauging station		Sampling station		Discharge		Tot-P		PO4-P		Tot-N		NO3-N		NH4-N			
	Outlet Samp!.	station	Disch. gaug. station	Normal	1994	Normal	1994	I/s sq.km	Cond	ms/m	tons	tons	tons	tons	tons	tons	tons	tons	Cu	Zn	Cd zero tons	Pb zero tons
sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km	sq.km
Gloomma, Hvaler-Singlefj.	41918	41218	40221	16.5	15.7	16.9	16.1	6.07	295.98	117.96	12311	8150	686.32	35.17	182.09	2.145	19.088	152.28				
Drammensvassdr, Dr.fj. V	17034	17028	16020	17.1	17.5	18.2	18.6	3.58	51.42	15.61	4711	3039	146.92	7.81	29.75	0.275	2.663	23.69				
Numedalslügen, Larvikfj.	5577	5513	5197	21.2	22.3	21.2	22.3	3.05	28.38	11.43	1589	921	103.20	3.83	19.98	0.184	1.917	15.00				
Skagensvassdr, Grenlandsfj.	10772	10348	10348	25.3	25.1	25.3	25.1	2.18	26.42	7.27	2774	1874	132.10	4.54	21.05	0.330	1.404	8.92				
Otra, Kr. Sandsfj.	3738	3730	3668	39.8	42.6	39.8	42.6	2.48	24.58	6.09	1325	674	60.86	2.34	23.60	0.187	1.966	17.60				
Oreelva, Øresondøen	105	105	54	36.7	31.1	40.7	34.6	20.48	4.73	1.12	164	92	5.47	0.17	0.31	0.007	0.043	0.60				
Suldalsfj., Sandsfj., Boknafj.	1457	1457	1457	59	59	59	59	3.6	2.86	4.74	1.36	618	491	18.98	0.92	5.42	0.054	0.380	1.84			
Orka, Orkafj., Tr.h.fj.	3053	2872	2247	21.7	14.7	21.7	14.7	7.00	9.18	2.30	653	425	19.65	15.41	51.16	0.157	0.334	3.30				
Vefna, Vefsnesfj. S	4122	4113	3323	43.9	30.9	43.9	30.9	6.40	29.44	15.20	2215	615	153.74	28.64	54.32	0.285	8.484	8.94				
Altælva, Altafj.	7373	7367	6257	11.8	9.3	11.8	9.3	12.61	34.95	20.92	491	134	16.45	3.02	2.03	0.110	0.247	8.61				

Parameters (mean values)

Watercourse	Parameters (mean values)																
	Hg			Gamma			PCB (The following Congeners)			IUPAC NOS			TOC	Cr-T	Ni	As	
	zero	HCH	kg	zero	zero	kg	28	52	101	118	138	153	180	SUM : zero	zero	zero	zero tons
Gjømma, Hvaler-Singlefj.	69.276	12.440	0.000	1.501	2.788	2.574	1.930	0.429	11.796	79.356	28.096	19.732	8.579				
Drammensvassdr., Dr.fj. V	0.000	6.244	0.000	0.551	1.010	0.918	0.918	0.735	0.184	4.316	28.466	8.081	5.418	1.561			
Numedalslågen, Larvikfj.	17.655	1.622	0.000	0.258	0.590	0.516	0.553	0.369	0.147	2.433	12.532	3.133	2.285	0.700			
Skiensvassdr., Grenlandsfj.	0.000	8.669	0.165	0.743	1.156	1.156	1.321	0.826	0.330	5.697	18.164	4.706	2.972	0.991			
Otra, Kr.Sandsfj.	0.000	3.886	0.000	0.234	0.515	0.468	0.468	0.328	0.140	2.154	15.262	5.618	4.448	0.796			
Orreleva, Orresanden	0.000	0.157	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.668	0.314	0.358	0.033			
Suldsfj., Sandsfj.Boknafj.	0.000	2.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	1.355	3.687	1.762	0.244			
Orkla, Orkdalsfj.Tr.h.fj.	0.000	0.373	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	7.331	2.260	1.336	1.101		
Vefsna, Vefsnfj. S	16.855	1.310	1.765	1.310	0.740	0.342	0.285	0.285	0.456	5.182	19.360	7.972	5.694	1.537			
Altaielva, Altaij.	4.112	0.137	0.000	0.082	0.164	0.110	0.082	0.055	0.055	0.000	0.493	9.595	3.290	2.056	0.877		

**Table 10.6 The Skagerrak Region. "Mean" inputs from tributary rivers in
The Sub-areas (1 - 5)
(Mean concentrations 1994 multiplied with mean runoff, 1931-60)**

The Skagerrak Region with sub-areas: (1A) Glomma, (1B) Inner Oslofj., (2) Drammenselva,
(3) Numedalslågen, (4) Skienselva, (5) Otra

Sub-areas :	Total quantity of substance discharged per year:						Were 70 % of measurements above the detection limit ?	Precision of the estimate of the load
	1A	1B	2	3	4	5		
Substance:								
Cd *	0.11	0.03	0.01	0.02	0.05	0.60	tonnes	YES _____ %
Cd **	0.11	0.03	0.01	0.02	0.05	0.60	tonnes	_____ %
Hg *	0.61	0.70	0.16	0.10	1.01	9.37	kg	YES _____ %
Hg **	1.32	0.85	0.16	0.61	1.01	13.38	kg	_____ %
Cu	0.9	0.5	0.1	0.8	0.6	5.7	tonnes	YES _____ %
Zn	8.0	2.4	1.3	9.1	7.7	70.6	tonnes	YES _____ %
Pb *	0.33	0.14	0.05	0.29	0.37	5.68	tonnes	YES _____ %
Pb **	0.33	0.14	0.05	0.29	0.37	5.68	tonnes	_____ %
Arsenic *	0.33	0.13	0.03	0.16	0.31	2.35	tonnes	YES _____ %
Arsenic **	0.33	0.13	0.03	0.16	0.31	2.35	tonnes	_____ %
Cr-T *	1.97	0.51	0.19	0.66	1.13	7.08	tonnes	YES _____ %
Cr-T **	1.97	0.53	0.19	0.66	1.13	7.08	tonnes	_____ %
Ni *	1.05	0.51	0.18	0.65	0.29	1.90	tonnes	YES _____ %
Ni **	1.05	0.51	0.18	0.65	0.29	1.90	tonnes	_____ %
PCBs *	0.00	0.11	0.00	0.00	0.00	0.00	kg	NO _____ %
PCBs **	0.23	0.16	0.03	0.13	0.21	2.08	kg	_____ %
gamma-HCl	0.74	0.25	0.04	0.22	1.01	8.80	kg	YES _____ %
NH4-N	25.77	33.77	6.71	24.33	39.41	392.7	tonnes	YES _____ %
NO3-N	953	408	136	612	227	2354	tonnes	YES _____ %
PO4-P	5.5	4.6	5.9	10.1	0.5	5.5	tonnes	YES _____ %
Total N	1145	558	171	922	404	4293	tonnes	YES _____ %
Total P	20	15	9	16	4	44	tonnes	YES _____ %
S.P.M.	3319	3660	2216	4919	1051	13497	tonnes	YES _____ %
TOC	8	2	1	3	3	32	tonnes	YES _____ %

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

**Table 10.7 The remaining North Sea. "Mean" inputs from tributary rivers
in The Subareas (6-7).**

(Mean concentrations 1994 multiplied with mean runoff, 1931-60)

The remaining North Sea Region with sub-areas: (6) Orreelva, (7) Suldalslågen

Sub-areas :	6	7	Were 70 % of measurements above the detection limit ?		Precision of the estimate of the load
			tonnes	YES	%
Substance:					
Cd *	0.38	0.73	tonnes	YES	_____ %
Cd **	0.38	0.79	tonnes	YES	_____ %
Hg *	5.39	22.63	kg	YES	_____ %
Hg **	16.04	38.13	kg	YES	_____ %
Cu	4.2	8.8	tonnes	YES	_____ %
Zn	71.8	72.3	tonnes	YES	_____ %
Pb *	7.49	2.74	tonnes	YES	_____ %
Pb **	7.49	2.80	tonnes	YES	_____ %
Arsenic *	3.85	1.86	tonnes	YES	_____ %
Arsenic **	3.85	1.88	tonnes	YES	_____ %
Cr-T *	17.00	20.39	tonnes	YES	_____ %
Cr-T **	17.00	20.39	tonnes	YES	_____ %
Ni *	4.03	1.93	tonnes	YES	_____ %
Ni **	4.03	1.96	tonnes	YES	_____ %
PCBs *	0.00	0.00	kg	NO	_____ %
PCBs **	3.09	6.02	kg	YES	_____ %
gamma-HCl	8.56	7.40	kg	YES	_____ %
NH4-N	321.42	199.96	tonnes	YES	_____ %
NO3-N	3639	3747	tonnes	YES	_____ %
PO4-P	23.0	29.7	tonnes	YES	_____ %
Total N	5409	6057	tonnes	YES	_____ %
Total P	80	101	tonnes	YES	_____ %
S.P.M.	17313	21858	tonnes	YES	_____ %
TOC	44	28	tonnes	YES	_____ %

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit

**Table 10.8 The Norwegian Sea. "Mean" inputs from tributary rivers
in The Subareas (8-9).
(Mean concentrations 1994 multiplied with mean runoff, 1931-60)**

The Norwegian Sea Region with sub-areas: (8) Orkla, (9) Vefsna

Total quantity of substance discharged per year:			Were 70 % of measurements above the detection limit ?	Precision of the estimate of the load
	Sub-areas :	8 9		
Cd *	1.77	0.38	tonnes YES _____	%
Cd **	1.82	0.47	tonnes _____	%
Hg *	58.95	8.22	kg YES _____	%
Hg **	68.94	29.58	kg _____	%
Cu	29.3	31.1	tonnes YES _____	%
Zn	65.8	52.9	tonnes YES _____	%
Pb *	10.70	14.86	tonnes YES _____	%
Pb **	10.74	14.86	tonnes YES _____	%
Arsenic *	4.59	6.34	tonnes YES _____	%
Arsenic **	4.61	6.34	tonnes YES _____	%
Cr-T *	45.46	60.77	tonnes YES _____	%
Cr-T **	45.46	60.77	tonnes YES _____	%
Ni *	11.80	5.08	tonnes YES _____	%
Ni **	11.82	5.08	tonnes YES _____	%
PCBs *	0.39	0.00	kg NO _____	%
PCBs **	7.94	6.09	kg YES _____	%
gamma-HCl	15.02	9.00	kg YES _____	%
NH4-N	267.36	146.19	tonnes YES _____	%
NO3-N	3641	1154	tonnes YES _____	%
PO4-P	56.8	30.3	tonnes YES _____	%
Total N	7931	3115	tonnes YES _____	%
Total P	193	95	tonnes YES _____	%
S.P.M.	57568	62361	tonnes YES _____	%
TOC	24	6	tonnes YES _____	%

Measurements below detection limits are treated in two ways :

***) Detection limit = Zero**

****) Detection limit = Limit**

**Table 10.9 The Barents Sea. "Mean" inputs from tributary rivers
in The Sub-area (10).**

(Mean concentrations 1994 multiplied with mean runoff, 1931-60)

The Barents Sea Region with sub-area: (10) Alta

		Were 70 % of measurements above the detection limit ?		Precision of the estimate of the load
Total quantity of substance discharged per year:				
Sub-area :	10	tonnes	YES	%
Substance:		tonnes	_____	%
Cd *	0.19	kg	YES	%
Cd **	0.26	kg	_____	%
Hg *	27.51	tonnes	YES	%
Hg **	32.62	tonnes	YES	%
Cu	19.0	tonnes	YES	%
Zn	20.6	tonnes	YES	%
Pb *	6.14	tonnes	YES	%
Pb **	6.14	tonnes	_____	%
Arsenic *	3.21	tonnes	YES	%
Arsenic **	3.21	tonnes	_____	%
Cr-T *	44.40	tonnes	YES	%
Cr-T **	44.40	tonnes	_____	%
Ni *	59.76	tonnes	YES	%
Ni **	59.76	tonnes	_____	%
PCBs *	0.00	kg	NO	%
PCBs **	3.69	kg	_____	%
gamma-HCl	3.51	kg	YES	%
NH4-N	272.65	tonnes	_____	%
NO3-N	229	tonnes	YES	%
PO4-P	27.3	tonnes	YES	%
Total N	2784	tonnes	YES	%
Total P	95	tonnes	YES	%
S.P.M.	31789	tonnes	YES	%
TOC	23	tonnes	YES	%

Measurements below detection limits are treated in two ways :

*) Detection limit = Zero

**) Detection limit = Limit