


REPORT SNO 3719-97

**P**lanning of an Integrated  
Acidification Study and  
Survey on Acid Rain  
Impacts in China

Final Report

***Acid  
Rain  
Research***

REPORT 48/97



**Main Office**

P.O. Box 173, Kjelsås  
N-0411 Oslo  
Norway  
Phone (47) 22 18 51 00  
Telefax (47) 22 18 52 00

**Regional Office, Sørlandet**

Televeien 1  
N-4890 Grimstad  
Norway  
Phone (47) 37 29 50 55  
Telefax (47) 37 04 45 13

**Regional Office, Østlandet**

Sandvikaveien 41  
N-2312 Ottestad  
Norway  
Phone (47) 62 57 64 00  
Telefax (47) 62 57 66 53

**Regional Office, Vestlandet**

Nordnesboder 5  
N-5005 Bergen  
Norway  
Phone (47) 55 30 22 50  
Telefax (47) 55 30 22 51

**Akvaplan-NIVA A/S**

Søndre Tollbugate 3  
N-9000 Tromsø  
Norway  
Phone (47) 77 68 52 80  
Telefax (47) 77 68 05 09


<b>Title</b> Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China. Final Report.	<b>Serial No.</b> 3719-97	<b>Date</b> 98.01.26
	<b>Report No.</b> <b>Sub-No.</b> 48/97        O-97061	<b>Pages</b> <b>Price</b> 77 pp + Appendix
<b>Author(s)</b> Espen Lydersen, Valter Angell (NUPI), Odd Eilertsen (NINA), Thorjörn Larssen (UiO), Jan Mulder (NISK), Ivar P. Muniz (NINA), Hans Martin Seip (UiO), Arne Semb (NILU), Rolf D. Vogt (UiO), Per Aagaard (UiO).	<b>Topic group</b> Acid Rain	<b>Distribution</b> 250
	<b>Geographical area</b> China	<b>Printed</b> NIVA

<b>Client(s)</b> World Bank and NORAD	<b>Client ref.</b>
--	--------------------


**Abstract**

The PIAC-project has been a multidisciplinary survey on acid rain in China. One goal has been to document effects of airborne acidifying compounds on vegetation, soil, soil- and surface- water and aquatic biota. Other goals have been to exchange knowledge between Chinese and Norwegian scientists, as well as visiting research sites in highly polluted areas in China and evaluating their need of support in a future collaborative monitoring and research program. Besides doing field measurements, we have collected samples at more than 20 sites in three areas in China, i.e. Chongqing, Guiyang and Guangzhou. Data from this survey may also be used in the RAINS-Asia model work, where more data are needed to validate and improve this model. Our survey documents negative effects of air pollution on all ecosystem levels investigated. The air concentration of sulfur in the urban and nearby areas is very high. The concentration of volatile organic compounds is generally high, which means that increased NO<sub>x</sub> emissions in coming years may create increasing O<sub>3</sub> problems. Reduced photosynthetic activity in some plants is documented, as well as soil and surface water acidification. Aquatic biota also reflects the acidification status of the surface waters investigated. However, the degree of damage in these regions is difficult to assess, since too few sites are incorporated in the survey. Surface water acidification is currently not a major environmental problem in China and will presumably not become one during the next decades. This report also contains a status report about acidification in China (in Appendix A) as well as a proposed framework for a monitoring program based on Norwegian experiences (in Appendix D).

<b>4 keywords, Norwegian</b> 1.    Luft forurensning 2.    Forsuring 3.    Økosystem 4.    Kina	<b>4 keywords, English</b> 1.    Air pollution 2.    Acidification 3.    Ecosystems 4.    China
---	---

  
 Espen Lydersen (NIVA)  
 Project co-ordinator

ISBN 82-577-3287-7

  
 Bjørn Olav Rosseland  
 Head of Research Department

Planning of an Integrated Acidification Study and  
Survey on Acid Rain Impacts in China

PIAC

## Preface

This report is prepared for the Norwegian Agency for Development Co-operation (NORAD) and Environmental and Natural Resource Division, Asia Technical Department, the World Bank (WB-ASTEN), as the main product from the joint WB/NORAD funded PIAC-project (*Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China*). According to the contract we have also made a report "Acid rain and its effects in China". The report is based on information available in printed form, basically from international and Chinese scientific journals about acidification status and impacts of acid rain in China. This report is enclosed as Appendix A. The main report contains results from our 1 month travel in China during April/May 1997 and presents data from visited sites as well as a general impression from meetings with Chinese scientific and management institutions. A proposed framework on monitoring and research in China is presented in Appendix D.

The Norwegian project participants are: Dr. Arne Semb, Norwegian Institute for Air Research; Professor Hans M. Seip, Associate professor Rolf D. Vogt and Research Fellow Thorjørn Larssen, Department of Chemistry, University of Oslo; Professor Per Aagaard, Department of Geology, University of Oslo; Dr. Ivar P. Muniz and Senior Research Fellow Odd Eilertsen, Norwegian Institute for Nature Research; Dr. Espen Lydersen, Norwegian Institute for Water Research; Dr. Jan Mulder, Norwegian Institute for Forest Research; Senior Research Fellow Valter Angell, Norwegian Institute for International Affairs. Dr. Espen Lydersen is the technical leader of both the WB and NORAD funded parts of the project, while Valter Angell is the administrative coordinator between WB-ASTEN and the scientific Norwegian group.

We gratefully acknowledge Dr. Tang Dagang, Director of Atmospheric Environment Institute at the China Research Academy of Environmental Science (CRAES), Beijing, who was our excellent guide during the trip in China. His excellent English, his administrative talent and humor were decisive for the success of the trip. Further, we would like to thank Counselor (Environmental Affairs) Gunnar Mathisen at the Royal Norwegian Embassy in Beijing for valuable support during the planning phase and the dinner sponsored by the Norwegian Embassy. We also want to thank Director Zhang Yutian, Director of International Co-operation Division at CRAES, who together with Dr. Tang was responsible for the implementation of our China visit and the Chinese-Norwegian seminar on acid rain held at the National Environmental Pollution Agency (NEPA) in Beijing.

Selection of actual sites in China was made in cooperation with Dr. Marcus Amman (IIASA), Prof. Leen Hordijk (Project manager RAINS-Asia), Dr. Jean-Paul Hettelingh (IIASA, RIVM) and local Chinese scientists. During our travel to these regions, we met a lot of friendly people and we were excellently treated. Therefore, we are most grateful to everyone, in particular Senior Engineer Zhao Dawei, Chongqing Institute of Environmental Science in Chongqing, Dr. Xiong Ji Ling, Director at the Guizhou Institute of Environmental Sciences in Guiyang and Professor Zhou Guoyi, South China Institute of Botany, Chinese Academy of Sciences, Guangzhou. We feel that our visit in China was very successful. We experienced the gradient from extremely beautiful nature to highly polluted sites. We have established numerous important contacts with open-minded Chinese scientists and people from Chinese environmental management institutions and politicians. This will obviously be of major importance for a successful implementation of a larger project on acidification in China.

Oslo, 1. September 1997

Espen Lydersen

---

---

# Contents

<b>Executive summary</b>	<b>6</b>
<b>1. Introduction</b>	<b>9</b>
<b>2. Material and methods</b>	<b>11</b>
2.1 Air measurements - analyses	11
2.2 Vegetation measurements - analyses	12
2.3 Soil and soil water measurements - analyses	12
2.4 Surface water measurements - analyses	13
2.5 Invertebrate measurements - analyses	13
<b>3. Data from the Chinese sites</b>	<b>15</b>
3.1 Chongqing area	16
3.1.1 Nanshan	16
3.1.2 Tie Shan Ping	19
3.1.3 Simian Shan	24
3.2 Guiyang area	26
3.2.1 Liu Chong Guang	27
3.2.2 Leishan	33
3.3 Guangzhou area	36
3.3.1 Dinghushan	37
3.3.2 Heshan	40
3.3.3 Guangzhou Botanical Garden	42
3.3.4 Baiyun Shan	43
3.3.5 Liu Xi River	45
3.4 Volatile organic compounds in air	47
3.5 Intercalibration of water analyses	49
3.6 Data Synthesis	51
3.6.1 Deposition and precipitation chemistry	51
3.6.2 Soil and soil water	52
3.6.3 Vegetation	56
3.6.4 Surface water	57
3.6.5 Aquatic invertebrates	59
<b>4. Other information related to acid rain in China</b>	<b>62</b>
4.1 The Chinese-Norwegian seminar	62
4.2 The RAINS-Asia model	62
4.3 Management and scientific relationships	63
4.4 Acid rain research activity in China	65
4.5 Conclusions from the Forth Expert Meeting on Acid Deposition in East Asia.	66
<b>5. Main conclusions and recommendations</b>	<b>67</b>
5.1 Main findings	67
5.2 A framework for further cooperation	70

---

<b>6. References</b>	<b>73</b>
<b>7. Appendix</b>	<b>77</b>
Appendix A	
Appendix B	
Appendix C	
Appendix D	

## Executive summary

### Introduction

Economic and social development in China is intimately linked to production of energy through combustion of fossil fuels, primarily by using high sulfur coal. The resulting emissions of sulfur dioxide cause serious air pollution in urban areas and relatively high concentrations of air pollutants and acid rain in many rural areas, affecting the environments. Future developments will involve further building of electric power plants and increased industrial activities. This development, together with a strong increase in the use of motor vehicles, may result in rapid increases in the emission of nitrogen oxides. This may increase the photochemical production of ozone on a regional scale, as has been observed both in Europe and in North America. As urban air pollution problems are being solved, emissions of fly ash and dust may also decrease, causing a decrease in the emissions of basic particles that has to some extent reduced the effects of acid deposition.

Three following issues are addressed in this report:

- How and to what extent are natural environments in China affected by acid rain ?
- To what extent may the RAINS Asia model be useful in China in simulating present and forecasting future effects of acid rain ?
- How may China benefit from foreign assistance in addressing its acid rain problems ?

Effects of acid deposition are clearly seen on biota, and in water and soil chemistry in China. Current emission intensities and deposition rates in some areas are much higher than in Europe and North America where most earlier studies of acid precipitation and the effects have been carried out. However, since the climate and the character of the natural environments in China are vastly different from conditions in the western countries affected by acid rain, the knowledge can not be applied directly. The survey has increased the scientific basis for translating the experience with acid rain in Europe and North America to China.

The RAINS-Asia model is a valuable tool, particularly for determining the relationship between economic activity, emissions and deposition rates; and for identifying potential problem areas. It should be realized, however, that the limited geographical resolution is a problem when working in areas where a large fraction of airborne deposits derives from sources within a few tens of kilometers, and with strong deposition gradients due to topographic conditions. The critical loads derived for the RAINS-Asia model on the basis of European experience are also very crude and their value as a predictive tool needs to be qualified. There is a strong need for further refinement and validation of the model. In order for the model to become an instrument in Chinese abatement policy, transfer of ownership to Chinese users appears to be necessary.

The understanding of the nature and the impacts of acid rain in China is growing. Abatement strategies have been introduced and important policy instruments are being developed. But, even though the Chinese authorities have most of the qualifications needed for implementing adequate control measures, there still appear to be gaps in the scientific knowledge about air pollution effects in China. Monitoring activities in China need to be improved, and monitoring of air pollution and acid deposition should be coordinated with effect studies and monitoring of vegetation, soil and runoff water. In order to provide a sound scientific basis for control measures and planning of future developments involving emissions of acidifying substances, the Chinese may find it beneficial to

exploit foreign experience and expertise, methodologies and "state of the art" equipment. This can be achieved through cooperation with bilateral and multilateral development agencies.

## Key findings

Findings from the PIAC-survey relate to assessments of ambient air quality, vegetation including forests, soils, water chemistry and aquatic biota:

- \* Precursors for the formation of O<sub>3</sub> and other photooxidants occur at levels which are high compared to European conditions. Their presence is caused both by regional emissions and by long range transport, and the concentration levels suggest that increased emissions of nitrogen oxides (NO<sub>x</sub>) may lead to seriously increased O<sub>3</sub>-concentrations, which may further damage vegetation and reduce yields in agriculture.
- \* Severe effects of acid deposition in China relate to soils. The investigated top and sub soils appear prone to acidification in the sense that acidity and aluminum concentrations in soil water increase. In areas with high deposition the conditions may cause damage to forests. Deeper soils and bedrock have in general higher neutralizing capacities.
- \* Reduced photosynthesis in plants and changes in the chemical composition in plant tissues in urban and suburban areas and necrosis of plant tissues may be attributable to high levels of sulfur dioxide (SO<sub>2</sub>), photochemical ozone (O<sub>3</sub>), lack of phosphorous or high aluminum content in soil water.
- \* Some water bodies in regions with high loading show typical signs of being acidified, i.e. high acidity or weakened natural buffering systems and elevated levels of strong acid anions and toxic aluminum compounds. Sensitive waters low in dissolved salts and hence poorly buffered, were found in some sparsely populated rural areas with low acid loading at present.
- \* Ecological effects of acidification have been observed on surface water organisms. Many of our observations resemble those of acidified systems in Norway and other countries in Europe and North America, i.e. that the diatom communities in streams are clearly associated with the level of water acidity and invertebrate abundance and biodiversity are lower in low pH streams.

The total information collected and evaluated by the PIAC team confirms that China has serious air pollution problems, not only in the urban areas and in the immediate surroundings of large conurbations, but also at considerable distances from emission sources. The problems are complex, involving damage to human health, materials, and terrestrial and aquatic ecosystems. In addition to the compounds usually involved in acid rain, photochemical ozone formation may also be a serious problem in the future. Scientific understanding of these problems requires close cooperation between different disciplines with quite different academic traditions. We have found the Chinese scientists knowledgeable and well informed, also with respect to acid rain issues. However, to make full use of previous studies in other countries, more bilateral or multilateral scientific cooperation is needed. In order to be able to run long-term monitoring studies with permanent measuring programs, it is essential to provide Chinese scientists with more resources, including equipment and training.



## The RAINS-Asia model

The RAINS-Asia model may become a valuable tool for understanding acid rain in Asia and for developing mitigation strategies in China. Scientists and administrators at many of the institutions and institutes visited had been introduced to the RAINS model. They expressed strong interest in using the model both for research and for policy-related activities.

The RAINS-Asia formulations are based on a similar model for Europe, (RAINS-Europe) which includes the Critical Loads Concept and different options and energy pathways. The PIAC team would like to stress that there are some obvious problems related to the applied spatial and temporal scales. The model forecasts become inaccurate on small spatial scales (m/km), because of the large heterogeneity in sensitivity. Likewise, the criteria for assessing soil sensitivity based on soil types and bedrock sensitivities to anthropogenic inputs have barely been studied for Chinese conditions. In this connection it is important, as our surveys have repeatedly reminded us, to consider the extensive and multiple use of forests and vegetation as part of local subsistence agriculture or forestry. Throughout the millennia, the ecosystems have been impoverished by gathering of edible plants, grazing of domestic animals, collection of litter and organic material as firewood or as soil fertilizing material.

The model does not, at this stage, pretend to cover the relationships between soils, freshwater chemistry and biotic responses, like the RAINS-Europe model. Therefore, further work is necessary in order to make the RAINS-Asia model a more effective tool in future planning.

Another set of issues is related to the question of transferring the "ownership" of the Rains model to the users in China. Several Chinese researchers expressed interest to have access to the modalities of the model in order to refine its operation and improve the quality of the data used. The authorities were particularly interested to incorporate their own emission data and emission forecasts.

## Concluding remarks

We recommend that Chinese authorities and scientific institutions embark upon a joint process of creating, and partly funding, a nationwide network of cooperating scientists and administrators with the overall objective to lay the foundations for sound federal and provincial policies regarding air pollution.

# We recommend that the World Bank assists in funding R&D projects needed in order to develop Critical Load criteria and response models for vegetation, soils, freshwater and aquatic and terrestrial biota.

# We suggest that bilateral cooperation is established in a 5 year Chinese-Norwegian monitoring and research project which includes transfer of expertise, training and technologies to Chinese counterparts in order to (a) develop a national network for integrated monitoring of air pollution and acid rain of high international standard and (b) promote basic and applied research into the cause and effects of such pollution on Chinese environments as well as forecasting future changes.

More details about the scientific conclusions and the proposed framework for a monitoring/research and training program on acid rain in China are presented in Chapter 5 (**Main conclusions and recommendations**) and Appendix D (**Proposed framework for future research, monitoring and training program**).

# 1. Introduction

The energy consumption in China has increased annually by 5.3% over the period 1980-1991 (Byrne et al., 1996) and coal accounts for more than 75% of the commercial energy production. The average sulfur content of the coal consumed is 1.2%, but in the Sichuan and Guizhou provinces the average S-content is 2.8% and 3.2%, respectively. The nitrogen emissions in China are dominated by NH<sub>3</sub> from fertilizer and domestic animal waste (Zhao and Wang, 1994; Galloway et al., 1996). Commercial fertilizers account for about 80% of China's total 20 million tons per year of nitrogen mobilization to the atmosphere.

Acid rain was recognized as a potential environmental problem in China in the late 1970s and early 1980s (Zhao and Sun, 1986; Zhao et al., 1988; Wang et al., 1997a; Wang et al., 1996). Although the SO<sub>2</sub> emissions have shown a lower growth rate during the last years due to cleaner technology used in power plants and boilers, the emissions are still increasing. Published reasonable scenarios have predicted that the Chinese SO<sub>2</sub> emissions will continue to increase for years. At present there are several acid rain projects under implementation in China. These projects are included in the National Ninth Five-Year Plan for Environmental Protection and the Long-Term Targets for the Year 2010. According to this plan the amount of sulfur dioxide emissions will be reduced from 22.3 million tons in 1995 to 22.0 million tons in the year 2000, a reduction of 1.5%. Even if China manages to reduce the SO<sub>2</sub>-emissions in the nearest future, the nitrogen (NH<sub>3</sub> and NO<sub>x</sub>) emissions are expected to increase significantly in the coming decades due to increased use of fertilizer and the fast increasing number of motor vehicles in the cities. This means that air pollution problems are expected to increase in China. Consequently, the country is likely to face increasing health problems and material damage (corrosion), reduced crop production, and negative impacts at several ecosystem levels such as vegetation (including forest), soil- and surface water biology. Preliminary calculations, e.g. by the RAINS-Asia model, suggest that the critical loads of forest ecosystems are seriously exceeded in many areas already (Hettelingh et al., 1995). However, there is an urgent need for improving the critical load estimates. More details about the acidification status and impacts of acidification in China are presented in Appendix A.

Most of the knowledge of effects of acid deposition has been supplied from studies in Europe and North America. Due to the severe impacts of acidification in these regions, scientists have shown increasing concerns about the situation in China and in nearby countries. Already in 1988, Professor Hans M. Seip at the University of Oslo established contacts with Chinese scientists, and Seip and co-workers have collaborated with several Chinese groups since then (Seip et al., 1995). In particular this has resulted in detailed studies in small catchments close to Guiyang and Chongqing. Acid rain research has a long tradition in Norway and is among the most successful examples of interdisciplinary research carried out in this country. A broad international contact net has also been established, and Norwegian scientists have been involved in many international research activities related to acidification and its effects on terrestrial and aquatic ecosystems. One of the main goals of this project was to transfer our experience and knowledge to Chinese colleagues. Concurrently, our cooperation will allow us to learn from Chinese experience and it will broaden our understanding of the effects of acidification because the ecosystems are quite different from those considered in the past. The differences between Chinese and European and North American ecosystems may reduce the applicability of the most used acidification models. At present there are no reliable estimates of critical loads for China. Assessment of critical loads for different ecosystems in China is necessary before effective emission reductions plans can be made.

Experience from acidification research in Europe and North America shows that critical loads of pollutants may vary depending on the target studied. An optimal strategy for air pollution control can

therefore only be obtained if environmental problems are treated in an integrated way. This implies that effects on ecosystems, human health, materials and climate must be considered at the same time (see e.g. Aunan et al., 1995).

We are convinced that problems related to emissions of acidifying compounds are severe in China and, if proper measures are not taken, very likely will get worse. Accordingly we conclude that a highly interdisciplinary research program is necessary as a basis for action. Consequently, an important aspect of this project is a tight link between different scientific disciplines. At present China primarily focuses on the health and corrosion problems caused by air pollutants. Our project will focus on terrestrial and aquatic ecosystems, because as China follows the same emission strategy as the Western world, negative impacts are to be expected.

In the pilot project, PIAC (*Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China*), proposed to NORAD and WB, we had six major objectives:

- 1) Establish contacts between scientific institutions and environmental authorities in China and Norway.
- 2) Obtain an overview of existing knowledge and activities within acidification research in China at both local and national level, and if necessary contribute with our knowledge and experience.
- 3) Evaluate needs for education, training, analytical tools, spare parts etc. for implementation of the action plan for monitoring.
- 4) Evaluation of the acidification problems in China.
- 5) Collect and analyze samples as an aid in getting quantitative information on acid rain and acidification in China. The data are also relevant for the RAINS-Asia modeling work.
- 6) Based on the pilot project, propose an action plan for monitoring and research in China in close cooperation with Chinese partners.

This action plan should be useful for our Chinese counterparts when they apply for economic support to NORAD/WB.

In order to obtain these goals we have had frequent contacts with Chinese colleagues since late 1996. Accordingly, we arranged a one day seminar in Beijing together, and thereafter implemented a one month travel in China (April/May 1997) including visits to potential sites for future monitoring. During this travel, a lot of measurements and samples were taken. In addition we received much information from Chinese scientists, environmental institutions and industrial companies. Most of this information is presented in this report.

## 2. Material and methods

Three regions in China have been visited: the Chongqing area, the Guiyang area and Guangzhou area. In Chongqing we visited three main sites: *Nanshan*, *Tie Shan Ping* and *Simian Shan*. In Guiyang we visited two major sites: *Liu Chong Guang* and *Leishan*; and in Guangzhou another four sites: *Dingushan*, *Heshan*, *Baiyun Shan* and *Liu Xi River* catchment. Where possible, measurements were conducted and samples were collected at the sites. Totally 27 different locations were visited. An overview of the main sites and the sampling/measuring program is given in Table 2.1. More details are presented in Appendix B, while all data from the survey are presented in Appendix C.

**Table 2.1.** Overview of sites where samples were collected during the tour. N: Needle samples; PEA: Plant Efficiency Activity measurements

		Soil No. of plots	Soil water	Surface water	Vegetation
Chongqing	Nanshan East	1	No	No	PEA
	Nanshan West	1	No	No	No
	Tie Shan Ping	7 <sup>1</sup>	Yes	Yes	N, PEA
	Simian Shan	1	No	Yes	PEA
Guiyang	Liu Chong Guang	7 + 13 <sup>2</sup>	Yes	Yes	N, PEA
	Leishan	3	No	Yes	PEA
Guangzhou	Dingushan	3	No	Yes	N, PEA
	Heshan	1	No	Yes	N, PEA
	Baiyun Shan	1	No	Yes	N, PEA
	Guangzhou Bot. G.	No	No	No	PEA
	Liu Xi River	1	No	Yes	N, PEA

1. Studied by Zhao Dawei, Chongqing Inst. of Environmental Science and Monitoring

2. The 7 plots are included in the study by Larssen et al. (1997). In addition soils from 12 locations along a transect and one additional plot were sampled.

### 2.1 Air measurements - analyses

Measurements of the airborne concentrations of sulfur dioxide, nitrogen oxides, ozone and particulate matter require heavy instrumentation and are not easy to perform on a field trip. Moreover, since concentrations vary from day to day, the results of such measurements are difficult to interpret in the absence of long data series and supplementary information, e.g. in the form of air mass trajectories and detailed emission inventories. Information on the concentration levels of sulfur dioxide in the cities and at some of the measurement sites was supplied by our Chinese colleagues. Precipitation chemistry data for these sites were also received. These data give information not only on the acidity of the precipitation, but also on the general air quality.

The only direct air quality measurements carried out by the group were determinations of the concentrations of light hydrocarbons and carbonyl compounds (aldehydes and ketones). The sampling procedure followed the EMEP Manual (Hanssen et al., 1996). Table C.16. gives the sampling sites, dates and time of sampling. The samples were analyzed by gas chromatography and high performance liquid chromatography at NILU. Some estimates of atmospheric residence times of different

compounds are presented in Table C.17 (Appendix). All samples were taken some distance away from major roads and motor vehicles. Contamination sources were generally absent. However, some of the sites were more close to urban centers as described under site description. The concentrations of these compounds are indicative of the influence of combustion sources on the air quality at the sampling sites, and also of the photochemical reactivity and the ozone formation potential. The data may be directly compared with corresponding data from Europe.

## 2.2 Vegetation measurements - analyses

Chlorophyll emits a red fluorescence when excited by visible light. When a leaf is illuminated at constant intensity it will fluoresce at a steady level. However, if a leaf is kept in darkness for several minutes and then is brightly illuminated, fluorescence rapidly rises to a peak level and then gradually decays to a low level ( $F_0$ ) level. The maximum achievable fluorescence level at a given light intensity is known as  $F_M$ . The difference between  $F_M$  and the low level signal ( $F_0$ ) is the variable component of fluorescence ( $F_V$ ). The ratio  $F_V/F_M$  has been shown to be proportional to the quantum yield of photochemistry and shows a high correlation with the quantum yield of net photosynthesis of intact leaves (Björkman & Demmig 1987).

Recently, chlorophyll fluorescence has been used to study cellular processes other than photosynthesis, in particular, the response of plants to environmental stress. A number of stress factors cause a progressive decline in  $F_V/F_M$ . Changes in chlorophyll fluorescence may well occur before any physical signs of deterioration are evident. Early indications of this type allow the collection of valuable data on the onset of stress conditions and tolerance threshold values. One of the objectives in the PIAC project was to make a screening for stress tolerance by chlorophyll fluorescence, by studying the  $F_V/F_M$  ratio, on different species at geographically separated localities in a gradient from central to south-east China.

In our study a portable Plant Efficiency Analyzer (PEA) was used for measuring chlorophyll fluorescence in the field. Early detection of reduced photosynthetic capacity may be achieved non-destructively by measurement of chlorophyll fluorescence using a PEA.

Tree damage parameters, including crown density, and crown color, have been estimated. Furthermore, samples of fresh needles from Masson pine and Chinese fir have been collected and analyzed for the content of plant nutrients, aluminum and several heavy metals. The chemical composition of the needles was compared to soil chemical parameters, which were determined at the same site. All chemical analyses of plant samples were conducted at the Norwegian Forest Research Institute (NISK), according to accredited methods.

The main purpose of these analysis was to evaluate the possible direct effects of air pollutants and indirect effects of soil acidification on trees and ground vegetation.

## 2.3 Soil and soil water measurements - analyses

Soil samples were analyzed at the Department of Chemistry, University of Oslo (cation exchange characteristics) and at the Department of Soil and Water Sciences, Agricultural University of Norway (C and N analysis). The soils' bulk density, porosity and moisture characteristics were determined at NISK and the soil mineralogy was determined at Department of Geology, University of Oslo. In some cases, published results from soil analyses at Chinese institutes are presented. Data on soil water

---

chemistry from Tie Shan Ping are provided by Dr. Zhao Dawei, Chongqing Institute of Environmental Science and Monitoring (CIESM). Soil water chemistry data in the Liu Chong Guang site are from Larssen et al. (1997).

The following soil analyses have been conducted: Total content of carbon and nitrogen, bulk density, porosity and water retention in soil, effective cation exchange capacity ( $CEC_E$ ), base cation saturation (BS), grain size and mineralogy. Fractionation of aluminum in soil water has been conducted, and soil water samples have been analyzed for all major anions and cations.

Parameters essential in evaluating effects of acid rain and for the critical load estimates connected to RAINS-Asia have been determined. Based on the soil chemical and physical analyses, soil and soil water sensitivity to acidification can be estimated. Mechanisms for interactions between soil and soil water can be discussed where soil water data, including Al-fractionation, exist. Furthermore, the content of major plant nutrients in the soil, important for forest health, can be assessed.

## **2.4 Surface water measurements - analyses**

Water pH, conductivity and temperature measurements were conducted directly in field. Water samples were collected for major chemical analyses at the Norwegian Institute for Water Research (NIVA) and for trace-elements and heavy metals at the Norwegian Institute for Air Research (NILU). Both NIVA and NILU are accredited laboratories.

On the basis of our analyses and existing Chinese data on surface water chemistry, it should be possible to document water acidification and to assess the resistance or buffering capacity of the different water bodies in light of likely increases in acidic inputs. Based on these analyses we are also able to give a first estimate of the water chemical conditions in relation to critical loads for some fish species according to our knowledge of critical loads for some European and North American fish species. The data from our China tour are not sufficient to assess the critical loads for the different water bodies in relation to the current sulfur and nitrogen deposition. This will be one goal in a future project on acid rain in China.

## **2.5 Invertebrate measurements - analyses**

The visited localities are all very small first order streams, some of which probably periodically dry out. Fish, crayfish, larger forms of snails and stream invertebrates were absent in all the actual streams. Macro-invertebrates in adjacent soils (earthworms, snails and woodlice) were also extremely scarce and difficult to collect in adequate sizes and numbers. Measurements of physiological (fish) and biochemical (soil and freshwater invertebrates) indicators of the level of stress in individual specimens, as proposed in the project description, have therefore not been performed. Consequently the sampling of stream invertebrates, microalgae and diatoms was merely qualitative. These groups of organisms are frequently used in Europe and North America to indicate the level of freshwater acidification through shifts in community composition.

The invertebrates were collected by using a fine meshed dipnet and/or by collecting specimens on stones from the streambed. When specimens were scarce or seemed absent from these samples, diatom samples (see below) were later examined for the presence of invertebrates. The field samples were preserved in ethylalcohol and later examined under the microscope. Only major taxonomic groups have been identified and registered. To determine which species are present in our samples,

there is a need for establishing contacts with Chinese taxonomic expertise on stream invertebrates. The data sets have been used to assess the more general relationship between stream fauna and environmental variables like water chemistry.

Diatoms, like invertebrates, occur in many habitats, and are also frequently used in the context of freshwater acidification, for monitoring and for comparative purposes. Their most widely sampled habitat in streams is the surface of upwards facing stones, submerged in the stream. Samples were taken by removing them from surfaces of several closely spaced stones by brushing into a tray, decanted into plastic vials and a pooled sample was preserved with Lugols Iodine. The material consists of *12 samples* of such epilithon material collected from various stream localities. The diatom samples have been processed using standard protocols and analyzed by Professor R. Battarbee at the Environmental Change Research Center (ECRC), UK. The samples were rapidly screened in the microscope and identification at this stage has been restricted in most cases to the generic level. Based on these results and known pH preferences, the diatom flora for each site have also been used to assess its relationship with ambient water quality.

### 3. Data from the Chinese sites

In China 3 different classes of air pollution standards are used. Class III is for industrial areas, Class II for urban residential areas, while Class I is for specially protected areas and nature. The annual average standard for SO<sub>2</sub> is 60 µg m<sup>-3</sup> for Class II. (WHO gives 40-60 µg m<sup>-3</sup>). The National standard for SO<sub>2</sub> for 24-hour averaging time is 150 µg m<sup>-3</sup> for Class II and 50 µg m<sup>-3</sup> for Class I.

The acidity (pH) of the precipitation is governed by a balance between acidic and basic components. Emissions of sulfur dioxide (SO<sub>2</sub>) and nitrogen oxides (NO<sub>x</sub>) are main contributors to the acids while basic dust (both man-made and natural) containing Ca and Mg tends to neutralize the precipitation (see e.g. Hedin et al., 1994). Emissions of ammonia (NH<sub>3</sub>) will neutralize the precipitation, but may cause acidification of soils and waters when ammonium is oxidized to nitrate.

Direct damage to forest vegetation is usually caused by short-term exposure to concentrations of several hundred micrograms of sulfur dioxide during a few hours, while frequent exposure to somewhat lower concentrations may result in general weakening of the plants and needle loss in the case of conifers. Conifers are generally susceptible to SO<sub>2</sub> and HF damage.

When discussing the sensitivity to acid deposition at the various sites, one must distinguish between further acidification of soil, soil water and streamwater. Defining soil acidification as a decrease in base saturation, the most sensitive soils are those with low cation exchange capacity and moderate BS. Soil water acidification is likely to occur where BS is low.

For water chemistry the acid neutralization capacity (ANC) is an expression of base cations (ΣCa, Mg, Na, K) in relation to strong acid anions (ΣSO<sub>4</sub>, Cl, NO<sub>3</sub>), i.e.  $ANC = \Sigma Ca, Mg, Na, K - \Sigma SO_4, Cl, NO_3$  (in µeq L<sup>-1</sup>). This variable gives a measure of the acidification status and the water's ability to resist acidification. In Norway and Sweden countermeasures like liming of acidic waters are never conducted in surface water with ANC > 50 µeq L<sup>-1</sup>. Lakes with ANC > 100-150 µeq L<sup>-1</sup> are assessed to be well buffered towards acidification. No fish damage is documented in Norway in lakes with annual average ANC > 20 µeq L<sup>-1</sup>. The fish incorporated in the Norwegian surveys (salmonids) are among the most sensitive species in the world to acid water.

Al<sub>a</sub> and Al<sub>i</sub> are two fractions of aluminum. Al<sub>a</sub> primarily represents the total concentrations of low molecular weight Al forms, while Al<sub>i</sub> is the inorganic fraction of Al<sub>a</sub> and represents the acute toxic forms of Al. The molar ratio Al<sub>i</sub>/BC<sup>2+</sup>, where BC<sup>2+</sup> = Ca<sup>2+</sup> + Mg<sup>2+</sup>, is an important variable, especially in soil water. It has been suggested that values > 1 may cause forest damage (Sverdrup and de Vries, 1994). The molar Ca/Al ratio in needles is often used as a diagnostic criteria to indicate the risk for aluminum damage in forest trees (e.g. Cronan and Grigal, 1995). A value < 12.5 is believed to indicate a 50 % risk for aluminum damage. However, it is not known if this threshold value also applies for the Chinese tree species investigated in our survey.

The evapotranspiration is high in the places we visited. This implies that ion concentrations increase relative to concentrations in precipitation.



### 3.1 Chongqing area

The Chongqing region is within the most air-polluted area in China. In the center of the city of Chongqing, average pH in rain is about 4.1, but pH increases to about 4.4 in the nearby rural areas. Predominant wind directions are north and north-east and rain pH varies according to emission sources and wind direction. The concentration of  $\text{SO}_2$  is 200 - 400  $\mu\text{g m}^{-3}$ . The Chongqing area has a subtropical climate. Average air temperature is 18°C, with an average humidity of 80%.

The soils in the Chongqing area are very variable. Three soil types predominate (Professor Ma, Vice-director of forestry Bureau of Chongqing Municipality and Professor at Southwest China Normal University, personal com.):

- 1) Non-acidic, well buffered soils developed from purple mudstone and shale (with 6-17%  $\text{CaCO}_3$  covering about 81% of the area. These soils are mostly used for agriculture and relatively little for forestry (some bamboo).
- 2) Acidic yellow soils derived from gray and yellow sandstone. These soils have little or no  $\text{CaCO}_3$  and cover about 17% of the area. Yellow soils have a pH between 4 and 5 and the base saturation is generally low. This means that these soils and/or the soil waters are sensitive to acidification. Normal pH in yellow soil water is about 4.3. About 80% of the local forests occur on yellow soil and these forests are dominated by Masson pine. Soil depth in Masson pine forest is 20-80 cm.
- 3) Soils derived from limestone containing more than 17%  $\text{CaCO}_3$  cover about 3.3% of the area,

Chongqing city covers an area of 23 000  $\text{km}^2$ , and the forest stand is about 12 mill  $\text{m}^3$ , covering 17% of the area. The annual growth is 0.8 mill  $\text{m}^3$  and 0.5 mill  $\text{m}^3$  is cut down annually for private use and because of diseases/insect attacks etc. Forest harvesting is strictly regulated by the authorities. The primary function of forests in Chongqing is to reduce soil erosion in the steep terrain. They are also used for recreational purposes. At the end of the 1950's, when coal combustion increased markedly, significant reductions in forest growth were reported in the area.

According to Professor Ma, the predominant forest vegetation is Masson pine ( $\approx 68\%$  of the forest), evergreens ( $\approx 13\%$ ), Chinese fir ( $\approx 10\%$ ) and Cypress ( $\approx 6\%$ ). Large parts of the forest were deforested in 1958 and 1962 during the "The Great Leap Forward", a common situation over large areas of China. The forests of today are mainly planted, and accordingly nearly 40 years old.

#### 3.1.1 Nanshan

##### Site description

Nanshan is a hilly area just outside the center of Chongqing city (29°32'N, 106°39'E). Annual precipitation is about 1100  $\text{mm yr}^{-1}$ , and almost 60% falls during April-September. Maximum intensity of rain is about 100  $\text{mm day}^{-1}$ . Annual evapotranspiration is about 900  $\text{mm}$  (Professor Ma, personal comm.), while annual mean temperature is about 17-18°C. We visited both the east and west side of the mountain.

In the 1980's, forest dieback occurred in a 700 ha area at Nanshan. Some of this has been reported at the Rio Conference. Also foliar damage, including yellow needle tips (likely chlorosis), has been observed. Possible reasons for the dieback have been discussed in several scientific papers, also internationally. There seems to be general agreement that insect pest (e.g. bark beetles) was the direct cause of tree death, but that the trees were already stressed from pollution. Professor Ma believes that

indirect effects through soils are more important than direct effects through SO<sub>2</sub>. He suggested toxic Al and nutrient problems (Mg shortage) as probable causes, while other scientists like Bian and Yu (1992) assumed the direct effects from SO<sub>2</sub> and hydrogenfluoride (HF) to be the most important reasons.

On the east side of the Nashan mountain, facing away from the city, we observed scattered signs of damage to the most dominating tree species (i.e. Masson pine) and some bryophytes species. The forest vegetation showed a considerable crown thinning because of such damage to Masson pine. The second most dominating tree species (i.e. Chinese fir) and a number of vascular plants showed better vitality. In areas with thin soils (20-50 cm) coniferous forests predominate while deciduous trees populate areas with thicker soils (> 50 cm). Soils are generally acidic yellow mountain soil with a thin O horizon on top of an approximately 3 cm deep A horizon. Below the A horizon is a B horizon with high clay content (Bt).

On the west side of the hill, facing Chongqing city, extensive dieback of Masson pine occurred in the mid-1980's. According to our guide, a number of other sensitive species had disappeared from this part of the mountain, and been replaced by more tolerant species. Currently, some deciduous trees and vascular plants seem to manage the levels of air pollution in the area, while others showed stress symptoms. On the west side of Nanshan the soils are considerably more sandy than on the east side, but also here they classify as yellow mountain soils.

### Air pollution

Precipitation chemistry (Table 3.1) and concentrations of some pollutants in air have been extensively studied in the Chongqing area, see Zhao et al. (1994) and Xue and Schnoor (1994). The sulfate concentrations in precipitation are very high, while the nitrate values are still fairly low. Of the cations, calcium and ammonium show the highest values. The former dominates in the urban areas, the latter in the rural areas. In addition to deposition by precipitation, dry deposition of both sulfur dioxide and particles should be considered. Annual SO<sub>2</sub> deposition in this area could easily amount to several grams per m<sup>2</sup> (i.e. > 10 g S m<sup>-2</sup>). The SO<sub>2</sub>-gas concentrations in Chongqing are about 30 times greater than those measured in the Bavarian and Black Forests of Germany (Blank, 1985; Blank et al., 1988), and 3-4 times greater than those reported from the most polluted regions in the prior Czechoslovakia (Moldan and Schnoor, 1992).

**Table 3.1.** *Precipitation chemistry (µeq L<sup>-1</sup>) in the Chongqing area. The three upper rows correspond to two urban (U) and one suburban (SU) station (for the years 1982-86), based on Xue and Schnoor (1994). The two lower rows are from one urban and one rural (R) station (for 1987-89), from Zhao et al. (1994).*

	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	NH <sub>4</sub> <sup>+</sup>	H <sup>+</sup>	pH	SO <sub>4</sub> <sup>2-</sup>	Cl <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	F <sup>-</sup>
U	187	26	14	17	126	72	4.14	387	26	27	18
U	170	23	15	8	152	51	4.29	328	25	30	22
SU	125	18	13	15	136	33	4.48	228	19	26	31
U	125	31	17	17	123	77	4.11	299	30	23	35
R	74	15	22	17	116	47	4.33	200	21	20	33

## Soil

At both sites the effective cation exchange capacity ( $CEC_E$ ) in the mineral soil is relatively high, presumably due to the high clay content. On the east side the soils are highly acidic with a base saturation in the A and B horizon below 10% (see table C.1). High clay content was especially noticeable at the east side site, showing a somewhat higher  $CEC_E$  than the west side. The exchange sites are dominated by aluminum so that the base saturation (BS) is low, rendering the soil water sensitive to acid deposition, despite the high  $CEC_E$ . C/N ratios decrease from 15.0 in the A horizon to 10.5 in the B (Table C.1). These C/N ratios are rather low, compared with values reported for forest soils in temperate and boreal regions. The low C/N ratios suggest that the availability of N is high.

On the west side of Nanshan we sampled the soil close to one of the remaining Masson pines. At this site the soils have a considerably higher base saturation in the topsoil (A-horizon) than in the deeper B-horizon, and the values are higher in both horizons than on the east side (Table C.1). The higher base saturation on the west side, facing the local pollution sources in Chongqing, may be due to the elevated input of alkaline dust. However, also the input of base cations from mineralizing woody debris after the dieback may have contributed. The C/N ratio (Table C.1) is extremely low in the A horizon (11.9) suggesting that the organic matter is well decomposed and that nitrogen is highly available. Leaching of nitrate from this site may be considerable, but unfortunately no data are available.

## Surface water

At the east side of Nanshan, there was one small artificial pool, but this was considered to be too much affected by human activity. No water body was present at the Nanshan west site. We also visited a second catchment on the hillside facing westwards towards Chongqing, but at a lower altitude than Nanshan west. In this catchment, there were two small streams originating as groundwater springs, one clearwater stream with pH 4.3 and one with pH 7. The latter probably contained high concentrations of heavy metals, presumably iron, since significant amounts of red precipitate (ochre) were found in the stream. We did not collect samples from this site for further analysis, because we did not find this catchment applicable as a future monitoring site. Thus, no water body was found suitable for monitoring.

Xue and Schnoor (1994) sampled two lakes in the area (Table 3.2). The concentrations of ions in Lake I are so extreme that we will not discuss these data further. Even in Lake II, the concentrations of ions are very high, and the acid neutralizing capacity (ANC) of the water is  $822 \mu\text{eq L}^{-1}$ .

The ANC of the lake is a result of neutralization and buffering of strong acids in the atmosphere (primarily  $\text{H}_2\text{SO}_4$ ) by dry-deposited compounds (e.g.  $\text{CaCO}_3$  dust) and weathering and cation exchange reactions in the watersheds. The calcium concentration in Lake II is close to saturation concentration for calcium in surface water in equilibrium with  $\text{CO}_2$  in air. Such high concentrations of calcium are normally found only in calcite areas, where acidification is unimportant. Some surface waters in Nanshan Mountain were also analyzed by Zhao Dawei and coworkers (Chongqing Inst. of Environmental Science and Monitoring). These data have not been published, but show the same picture, i.e. high concentrations of calcium and sulfate and high ANC and pH. This does not rule out the possible presence of other harmful compounds, in particular heavy metals, since the area is highly polluted by atmospheric inputs of  $\text{SO}_2/\text{H}_2\text{SO}_4$  and dust. Such compounds should be analyzed in the future, because they may have negative impacts on aquatic life. The concentration of dissolved organic carbon in the surface waters analyzed is high, and much higher than the other surface waters within our China survey.

**Table 3.2.** Water chemistry in two Nanshan Mountain lakes near Chongqing, sampled 4 times during 1987-88 (Xue and Schnoor, 1994). All concentrations in  $\mu\text{eq L}^{-1}$ , except for DOC (Dissolved organic carbon) which is in  $\text{mg C L}^{-1}$ .

Lake	pH	H <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	SO <sub>4</sub> <sup>2-</sup>	Cl <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	F <sup>-</sup>	ANC	DOC
I	8.46	0.004	3336	1010	855	115	1934	1241	17	16.3	2026	13.1
II	6.79	0.16	1450	364	205	118	979	216	86	0	822	9.3

### Vegetation

Nanshan Mountain is an area of dramatic forest decline, and the Masson pine forests at the top of this mountain are dying (Xue and Schnoor, 1994). Photosynthesis activity was only measured on the east side of the mountain. The few remaining specimens were so damaged that it was not possible to measure photosynthesis activity on the west side. For Masson pine the mean value of three measurements from six ramets was 0.75, the lowest value both within the Sichuan province and the other 12 studied areas. The standard deviation of these measurements was 0.043, the highest registered in our material (Table C.20). The  $F_V/F_M$  ratio measured on the tree showing the best condition was 0.80, while the lowest  $F_V/F_M$  ratio was 0.69. This may indicate that some of the trees have more sub-optimal growth conditions, due to more air pollution and/or less available soil nutrients, than others. For Chinese fir, the other dominant coniferous tree species, the mean value was 0.80 and the standard deviation very low (0.008), indicating that Chinese fir is a non-sensitive species compared to Masson pine. For a number of randomly selected vascular plants the measurements showed more or less the same picture as for Chinese fir.

Because Nanshan did not appear to be one of the potential monitoring sites, we have not carried out chemical analysis of needles.

### 3.1.2 Tie Shan Ping

#### Site description

The Tie Shan Ping area (29°38'N, 106°41'E) is approximately 25 km north-east from Chongqing center, at an elevation of about 450 meters. The catchment visited is located within a protected area. The forest area is partly protected, but heavily used for recreational purposes. It consists of 1200 ha forest where Masson pine is the predominant tree species. Large parts of the forest were logged in 1958 and 1962. Later forest has been replanted. The area receives about 1 million Yuan annually from tourists, while 0.1 million Yuan come from governmental funding. The administrative management of the forest is undertaken by Jiang Bei District, Agricultural Dept., Chongqing.

The main soil type is yellow mountain soil. The soils are reported to be homogenous and they are relatively rich in finer particles (i.e. < 63  $\mu\text{m}$ , see Table C.15). In general the soils are 50 - 80 cm deep with a thin A horizon (about 3 cm) on top of a B1 (3 - 13 cm depth) and a B2 horizon.

During previous co-operation between CIESM and the UiO, equipment for sampling precipitation, throughfall and soilwater has been installed at seven plots within the catchment. Throughfall and soil water chemistry data given in this report are unpublished results from these plots provided by Dr. Zhao Dawei at CIESM.

### **Air pollution**

The site appears to be downwind of Chongqing industrial areas, and may therefore experience quite high airborne concentrations of sulfur dioxide and other pollutants, particularly in the winter season. Inputs by dry deposition may be similar to or higher than the wet deposition.

pH in volume weighted precipitation is low with values during winter months dropping to 4.6 in open air and below 4 in throughfall (annual avg. 5.0 and 4.1 respectively). Due to high concentrations of calcium, the lime potential ( $L_p = \text{pH} - \frac{1}{2}\text{p}(\text{Ca}+\text{Mg})$ ) in precipitation was 3.0, dropping to 2.5 in the throughfall. Ammonium deposition is significant, i.e. 32  $\mu\text{M}$  in throughfall (0.29 g N  $\text{m}^{-2}\text{yr}^{-1}$ ). Fluoride concentrations in throughfall are also relatively high (up to 14.4  $\mu\text{M}$ ).


### **Soil and soil water**

The soils at Tie Shan Ping are rather dense with bulk densities increasing from 1.25  $\text{kg dm}^{-3}$  in the A horizon to 1.34  $\text{kg dm}^{-3}$  in the B2 horizon (Table C.4). Soil porosity is 50% and 47% in the A and B horizon, respectively. The generally high densities and low porosities are mainly due to a large fraction of fine secondary silt and clay material ( $< 63 \mu\text{m}$ ). The grain fraction less than 63  $\mu\text{m}$  decreases somewhat down into the profile. Increasing grain size with depth is common in soils developed through chemical weathering. Water retention characteristics are similar for the A and B horizon and typical for clay rich soils. The soils are mainly composed of quartz (74%) and clay minerals (14%). The remaining material being both plagioclase and K-feldspar (Table C.15). Moderate and similar amounts of plagioclase and K-feldspar probably reflect the generally low amounts of these minerals found in the indigenous sandstone bedrock. Water contents in the A horizon decrease from 36% at field capacity to 22% at wilting point (Table C.5). This suggests that the upper 60 cm of the soil, which is assumed to be the rooting zone, has a storage capacity of water available for plants of only 69 mm. This indicates that the site is rather drought sensitive.

The C/N ratio of soil organic matter decreases from 16.1 in the A to 8.1 in the B2 horizon (Table C.1). These values are rather low and suggest that that nitrogen availability for the vegetation is good and that nitrate is likely to leak from the soils causing high concentrations in soil water and surface water (see below). The total store of nitrogen in the soil is estimated at 5100  $\text{kg ha}^{-1}$  (from the soil surface to 28 cm depth; Table C.4).

Average soil water chemistry data from the A and B horizon in the seven plots are presented in Figure 3.1 and Table C.9. Charge balance calculations on the soil water data show that there is no clear bias, though the agreement is not good; average deviation was 7.5% in excess of anions, with a standard deviation of 25.0% (see figure 3.1). Results from an intercalibration between CIESM and University of Oslo indicate fairly good agreement, but there is a pertinent need for further intercalibration studies. See Chapter 3.5 for further discussion of the intercalibration data.

The major anion in soil water is as usual sulfate, though also nitrate concentration is high at this site. The high nitrate values suggest that ammonium from deposition may oxidize and leach out causing high nitrate and even significant ammonia concentrations in surface water (see below). As in deposition, fluoride concentrations are relatively high. Large spatial and temporal variations in pH and  $\text{Al}_i$  are observed. In plot 2, pH is generally low (median 4.2) and  $[\text{Al}_i]$  is high, especially in the B-horizon. Lowest acidity was found in plot 1, which is a groundwater spring, feeding water into a lake.



**Figure 3.1.** Soil water chemistry in the various plots at the Tie Shan Ping catchment. Left and right bars represent cations and anions respectively. Al eq denotes charge equivalence of aluminum species.

Where the  $Al_i$  concentrations are high, the  $Al_i$  over  $Ca+Mg$  molar ratio also becomes high, despite high  $BC^{2+}$  concentrations. Hydrated  $Al^{3+}$  was the dominating Al species in soil water accounting for more than 50% of the total, except in the groundwater spring (plot 1; figure 3.1). Soil samples were collected at plot 2, but the results are not yet available.

The  $Al^{3+}$  concentration appears not to follow a gibbsite mineral ( $Al(OH)_3$ ) solubility, instead we observe the typical increasing undersaturation with decreasing pH. Soil solutions were close to saturation with jurbanite ( $AlSO_4OH$ ). This has also been observed elsewhere where the sulfate concentrations are high (Vogt et al., 1994). The lime potential through the soil profile does not change significantly relative to the throughfall deposition, except for an increase in the "spring plot" (i.e. plot 1: 3.3 and 3.0 in A- and B-horizon respectively). The water at plot 1 has much lower anion concentrations relative to the other soil plots. This may be due to sulfate adsorption/reduction and nitrate uptake. Several observations show clearly that neutralization processes occur in deeper soil layers or in the sandstone bedrock, which usually is fairly permeable due to cracks and fissures. While pH in throughfall is about 3.0, the average pH in soil water in all plots except plot 1 (the ground water spring) is 4.0 and 4.2 in the A and B horizons, respectively. The increase is greater for plot 1, where pH in both soil horizons is close to that in surface water (i.e. about 5). Furthermore, the alkalinity of soil water is about -100 and -50  $\mu eq L^{-1}$  in the A- and B-horizons respectively for all plots except plot 1 where the values are +10 and -5  $\mu eq L^{-1}$ . Only a few samples from plot 6 and 7 have been analyzed.

### Surface water

The water bodies investigated within Tie Shan Ping area have low pH and ANC (average ANC values vary from about -20 to -90  $\mu eq L^{-1}$ ). The concentrations of  $Al_i$  are relatively high (Table 3.3). Sulfate is the dominating anion, but considerably lower than in soil water and also much lower than in surface water in the Nanshan Mountain area. However, the dissolution of base cations by cation exchange and weathering reactions in Tie Shan Ping (see above) is too low to prevent negative ANC-values in surface waters in the area. The concentration of total organic carbon in the waters was too low to affect the water pH. However, when we visited the catchment April 18 (1997), the streams had only minor discharge. We have no data for the water chemistry at high flow.

**Table 3.3.** Water chemistry of waterbodies in the Tie Shan Ping area. The rows marked I, II and III show average values for sites within the monitored catchment, sampled during March to December 1995 ( $n = 9$ ), while A and B are grab samples from two small streams within the same area taken April 18, 1997. TOC: Total organic carbon.

Site	pH	$Ca^{2+}$	$Mg^{2+}$	$Na^+$	$K^+$	_____			ANC	$Al_a$	$Al_i$	TOC
						$\mu eq L^{-1}$						
I	4.84	217	151	42.3	9.2	268	96.5	77.1	-22.1	230	200	
II	5.15	263	171	63.7	17.7	425	65.9	84.0	-59.1	330	240	
III	4.95	277	155	52.5	20.6	464	44.1	88.5	-92.0	460	440	
A	5.34	298	128	43.9	16.4	408	59.2	40.7	-21.0	127	121	0.54
B	4.89	341	136	54.8	23.8	475	79.0	57.5	-56.0	204	196	0.64

The concentrations of heavy and trace metals in the waterbodies are not extreme (Table C.7). One exception is manganese (Mn) at site B, which had the highest concentration ( $632 \mu\text{g Mn L}^{-1}$ ) of all water bodies investigated. Zhao Dawei and coworkers at CIESM have also documented high concentrations of manganese, i.e. from  $417 \mu\text{g Mn L}^{-1}$  at site I to  $538 \mu\text{g Mn L}^{-1}$  at site III. Manganese is not a very toxic metal, but the concentrations in the surface water in Tie Shan Ping are very high. Manganese interferes in the determination of aluminum. This has not been corrected for due to lack of Mn data for the other sites. At site B this has led to a 3.5% overestimate of Al. The concentration of iron is close to the median concentration of this element in our survey. At site B, we observed a redbrown precipitate on the stream bed, probably indicating high levels of iron and/or manganese.

### **Vegetation**

The trees at Tie Shan Ping appear less defoliated than at Nanshan, although Masson pine on exposed ridges showed occasionally low crown densities. This was reflected by the chlorophyll fluorescence, showing a mean  $F_V/F_M$  ratio of 0.82 (maximum value: 0.84, minimum value: 0.77). The standard deviation of these measurements, 0.025, was considerably lower than in Nanshan, indicating more stable and better growth conditions. Chinese fir, on the other hand, shows the opposite picture: The mean  $F_V/F_M$  ratio was 0.80, which was slightly lower than at Nanshan. It is difficult to explain this difference. But we cannot preclude differences in local climate and nutrient conditions.

Chemical analyses of needles (Table C.3) indicate that phosphorus nutrition is sub-optimal for both Masson pine and Chinese fir. Nitrogen contents in the needles indicate that this nutrient is in good supply reflecting the low C/N ratios in soils and high nitrate values in soil water. The content of Ca in needles is quite different for Masson pine and Chinese fir, being highest in the latter. By contrast the Al content is lowest in the fir needles. In Masson pine the Ca/Al ratio in the needles even approaches 12.5, which is believed to indicate a 50% risk for aluminum damage.

### **Invertebrates and diatoms**

Diatom samples were taken below the confluence of the two small streams (A and B, Table 3.3). This station had pH 4.95 and conductivity  $66.4 \mu\text{S cm}^{-1}$  at  $16.6^\circ\text{C}$ . This sample is of low floristic diversity, containing several small *Pinnularia spp.*, *Eunotia spp.* and *Brachysira serians*, indicating a strongly acidic environment.

We observed no stream invertebrates at this site through visual inspection of the substrate. But by later examination of the diatom samples we observed a few (15) very small Diptera (midge) larvae, one pupae and one little Trichoptera (caddisfly) larvae. This signifies a poor fauna with very low species diversity which probably is related to the ambient water quality. The hydrological effects of the upstream dams are not known, but even if the stream periodically carries very little water, this alone cannot explain the absence of major common invertebrate groups.



### 3.1.3 Simian Shan

#### Site description

The Simian Shan (or Four Side Mountain) is located about 110 km south of Chongqing (28°40'N, 106°22'E) and about 200 km from Guiyang. The sampling site was a few hundred meters behind the Forest Management Department, in the catchment of a small stream named Piao Chang Qoy, located about 1100 m a.s.l. Annual mean temperature in the area is 15-17°C. Air temperature is normally 3-5°C during winter, but temperatures  $\leq 0^\circ$  are not uncommon. Annual precipitation is about 1200 mm. Deciduous forest is the dominating forest type, but there are also areas with conifers.

The area under the Forest Management Department administration is 40 000 ha. The major soil type is yellow mountain soil derived from sandstone. Average soil pH representative for this area was reported to be 4.2 in the A-horizon, 4.6 in the B-horizon and 4.5 in the C-horizon. Average organic matter contents were 81%, 2.2% and 2.2% in the A, B, and C horizon, respectively. The information was presented by a representative from the Forest Management Department. At the site sampled by us, there was a few cm thick forest floor on top of a 4 cm deep A horizon. Below the A is a B1 (6 cm deep) and a B2 horizon. More information about the Simian Shan Site is given in Zhao et al. (1994).

#### Air pollution

According to Zhao et al. (1994), the volume weighted pH of rain in this area is about 4.4, the sulfate concentration in precipitation about 130  $\mu\text{eq L}^{-1}$ , and wet deposition about 2.3 g S  $\text{m}^{-2}$ . Some data (20 samples) presented to us by the representative of the Forest Management Dept., collected during a short period in the summer, showed high pH in precipitation, varying from pH 6.2 to 6.9. Sulfur dioxide concentrations were a few micrograms per  $\text{m}^3$  (cf. Lei et al., 1997).

#### Soil

The soil plot sampled at Simian Shan was dominated by Chinese fir and broadleaf trees. C/N ratios in soil organic matter are rather low, being 17.0 in the A horizon and 15.2 in the B2 (Table C.1). Nitrogen stores in the upper 18 cm of the soil (including A, B1 and B2 horizon) are estimated at 3800 kg  $\text{ha}^{-1}$  (Table C.4). The soils at Simian Shan have a relatively low bulk density increasing from 0.57 kg  $\text{dm}^{-3}$  in the A horizon to 0.92 kg  $\text{dm}^{-3}$  in the B. Soil porosity is 69% and 58% in the A and B horizon, respectively (see Table C.15). More than 80% of the soil material is quartz, which is the highest amount among our sampled sites. Furthermore, the soils do not contain the secondary mineral kaolinite. Instead the clay minerals are mainly residuals as illite and chlorite. This may indicate that the soils are relatively young. A lack of plagioclase mineral in these soils may be due to the sandstone bedrock being previously depleted of plagioclase. The mineralogy does not change much with depth.

The  $\text{CEC}_E$  is high in the topsoil horizons (Table C.1). This is probably caused by fairly high organic matter content probably due to rather slow organic breakdown in the relatively cold climate (see above). Also the B-horizon has a relatively high  $\text{CEC}_E$  considering that the soil material consisted mainly of quartz sand. The soils are among the most acidic ones sampled on our trip through China, with base saturation values decreasing slightly down into the profile from 8.5% in the A horizon to 6.4% in the B2. These low BS values suggest that soil water at this site may be easily acidified even though the  $\text{CEC}_E$  is high. Increased acid deposition is likely to result in elevated concentrations of toxic aluminum in soil water. Despite the acid soils the streamwater was not acid (see below). This is likely due to deep hydrological flowpaths, passing soils or bedrock richer in base cations. The high

alkalinity in the stream sample collected further downstream even indicates the presence of some carbonates in the bedrock.

Water contents in the A horizon decrease from 40% at field capacity to only 9% at wilting point. In the upper 60 cm this soil has a storage capacity of water available to plants of 165 mm, indicating that this site is far less drought sensitive than Tie Shan Ping.

### Surface water

Water samples were taken from the Piao Chang Qoy stream, between two small artificial dams stocked with carp (site A), and from another stream further down the road draining a small valley where rice and vegetables were grown (site B). The data from Zhao Dawei and coworkers at CIESM presented in Table 3.4 are from sites located at somewhat higher altitudes. Even though the Simian Shan area is located 110 km south of Chongqing, the concentration of sulfate in runoff is relatively high, probably mainly due to the substantial S-deposition (see above). Since the concentrations of base cations are even higher than of sulfate, the ANC of surface water is high, and  $Al_i$  negligible. The concentrations of trace- and heavy metals are generally low (Table C.7), but the concentrations of titanium ( $80 \mu\text{g Ti L}^{-1}$ ) and vanadium ( $0.80 \mu\text{g V L}^{-1}$ ) at site B are the highest, and the concentration of strontium ( $125 \mu\text{g Sr L}^{-1}$ ) the second highest recorded in this survey. Titanium is normally closely associated with iron (Fe), and even Fe is at relatively high levels. These elements, especially Fe and Ti, are presumably primarily associated with clay particles and organic compounds, and not dissolved in the water as true ions. The turbidity, and therefore concentration of particles in this stream is relatively high (Table C.6), explaining the high Fe and Ti.

**Table 3.4.** Average concentrations of some chemical compounds in surface water at the Simian Shan Mountain located from 1000-1350 m a.s.l. Data set I gives average values for three sites based on unpublished data from Zhao Dawei and coworkers at CIESM, while the data denoted A and B are from sites collected during our survey April 19-20, 1997.

Data set	pH	$Ca^{2+}$	$Mg^{2+}$	$Na^+$	$K^+$	$SO_4^{2-}$	$Cl^-$	$NO_3^-$	ANC	$Al_a$	$Al_i$	TOC
		$\mu\text{eq L}^{-1}$								$\mu\text{g Al L}^{-1}$	$\text{mg C L}^{-1}$	
I	7.17	333	92.1			200			462			
A	6.83	434	64.2	33.5	13.6	373	11.3	26.4	125	64	9	2.0
B	7.54	898	125	46.5	15.9	433	14.1	65.3	573	19	8	2.1

### Vegetation

At this mountain site there are very few Masson pines. The forest was dominated by other Pinus species, but sufficient ramets of Masson pine were available for photosynthetic activity measurements. All these trees were located close to arable land. The specimens were probably planted and looked after by the local farmer, and thus given good growth conditions. This fits in with the measured  $F_v/F_M$  ratio (Table C.20), showing very high values in spite of the sub-optimal climate conditions at this altitude. The standard deviation, 0.005, is the lowest registered at all our sites. This may indicate that the six measured trees are exposed to the same low level of air pollutants. Measurements of Chinese fir showed a relatively low photosynthetic activity (mean  $F_v/F_M$  ratio 0.79). Similar values are only found in other mountain areas like Leishan and Dinghushan. The trees at Simian Shan are less defoliated than at Nanshan and Tie Shan Ping. However, Simian Shan, where only a few Masson pine trees occur, is not

an appropriate reference site for the affected Masson pine forests near Chongqing. Needle samples were therefore not taken for chemical analysis.

### Invertebrates and diatoms

Diatom samples were taken from the same streams as the water samples. The sample from site A had a relatively rich and diverse diatom flora with *Achnanthes minutissima*, *Sellaphora pupula*, *Gomphonema spp.*, *Navicula spp.*, *Pinnularia spp.*, and *Cocconeis spp.* suggesting circumneutral water. The site B sample had a range of taxa characteristic of circumneutral to alkaline waters including large *Cymbellas*, *Achnanthes spp.*, *Gomphonema spp.* and *Fragilaria spp.* We observed no invertebrates in the two streams, neither through visual inspection of the substrate nor by later examination of the diatom samples. This is surprising particularly for the stream between two small artificial dams. These dams are stocked with carp which shows that the overall water quality should be suitable for fish. The cause of absence of stream invertebrates could in this case be drought and for the other stream, which drains the cultivated valley, perhaps local use of agrochemicals. This case still remains somewhat obscure because, as shown to us by the local people, the nearby main river was fished in and had several species.

## 3.2 Guiyang area

The Guiyang city covers an area of 8000 km<sup>2</sup>. The city is located about 1000 m a.s.l. and is surrounded by mountains. The temperature is 4 - 8°C in winter and 24-28°C in summer. Frequent inversion periods occur during the winter (80% of the time), and 27% of the year there is almost no wind. The annual precipitation amount is 1175 mm and the major part falls during summer.

The energy source is primarily coal, with a sulfur content of 3 - 5%. Many countermeasures towards particle and SO<sub>2</sub> emission have been conducted or are planned. In 1980, 95% of the energy was based on coal, while in 1997 this was reduced to 70-80%. Before the reductions the urban SO<sub>2</sub> emissions in Guiyang were 270 000 tons yr<sup>-1</sup>. According to Guiyang EPB, the emission of SO<sub>2</sub> in the rural area of Guiyang is today reduced by about 60 000 tons yr<sup>-1</sup>. In addition the total emission of solid particles (TSP) has been reduced by 30 000 tons yr<sup>-1</sup>. Accordingly, while the annual average SO<sub>2</sub> concentration in the city was 460 µg m<sup>-3</sup> in 1990, it was reduced to about 300 µg m<sup>-3</sup> in 1996. Thus, the concentration in Guiyang is still about 6 times greater than recommended by the National standard, Class II, and by WHO.

The pH of the precipitation was reported to have increased from around 4 to around 5 during the same period (1990-1996). This increase is unexpectedly large, especially because emissions of alkaline dust particles have decreased concurrently with SO<sub>2</sub> emissions. Continued observations of precipitation chemistry are necessary to study the trends. In 1990 pH < 5.6 occurred in almost 60% of the rain events.

Guiyang city is the major source for air pollution in this province. Accordingly, countermeasures in this city are essential for the future emissions of this province. The authorities are particularly concerned about corrosion, health problems and tree health (e.g. early shedding of leaves in the autumn and yellowing of conifer needles) due to air pollution.

Besides having meetings with local scientists and the local EPB, we also visited different demonstration projects for air emission reductions at three factories in Guiyang city. We visited a tire-factory, which by using a lime (CaCO<sub>3</sub>) injecting technique, had managed to reduce their SO<sub>2</sub>

emission by 70 - 80%. Today the annual emission of SO<sub>2</sub> from this factory is about 3 000 tons. The lime injection technique produces about 50 000 tons of relatively polluted CaSO<sub>4</sub> waste, a product primarily used in bitumen surfaces. Further, we visited a factory producing special steel for airplanes, which had a cyclone boiler flue gas scrubber using limewater for removal of SO<sub>2</sub> and particles. This was only a small demonstration unit. The removal of S by this technique was about 85%. Finally we visited Guiyang Steel Mills which started in 1958 and has 9 300 employees at present. They produce annually about 300 000 tons raw steel, and 250 000 tons more refined steel. The annual emissions are 6 500 tons (SO<sub>2</sub>), 1 350 tons dust, and 1 200 tons industrial dust. About 65 000 tons of industrial wastes are stored in land deposits. The factory is also discharging 23 mill tons of waste water annually. The factory is therefore an important emitter of both SO<sub>2</sub> and dust to the Guiyang city. However, the process is old, and the old boilers will be closed in the nearest future and/or retrofitted with gas fired boilers. In a few years time, all the ovens will be run on low sulfur coal gas instead of raw coal with a S-content of about 5%. Thus in 1998, the SO<sub>2</sub> emissions are expected to be about 1000 tons yr<sup>-1</sup>. Large efforts are being directed towards solving the dust problems, and it was claimed that they will be under control in 1999.

At present, there are three major countermeasures for reducing the emissions of S in Guiyang:

- 1) Limestone injection technique
- 2) Production of desulfurized coal briquettes or patent fuel.
- 3) More LPG (Liquid Petroleum Gas) products, e.g. the private stoves are today using more gas for cooking instead of raw coal.

### 3.2.1 Liu Chong Guang

#### Site description

The catchment is located about 10 km north-east of Guiyang center (26°34'N, 106°38'E). The catchment is about 7 ha, and the elevation ranges from 1320 to 1400 m a.s.l.. It is drained by two small streams which below their confluence run into a small artificial dam. Another stream draining the neighboring catchment enters the outlet stream from the dam. The annual discharge has been estimated at 630 mm which means that the evapotranspiration is roughly 45%. SO<sub>2</sub> concentrations in air (approximately 1 month's average) varied between 10 and 100 µg m<sup>-3</sup> which can be characterized as medium to high pollution level in China.

The predominant vegetation is Chinese fir, Masson pine and wild rose bushes. The catchment was deforested in 1958 and hit by fire soon after, but replanted after the fire. Thus, the catchment is in a dynamic succession stage, clearly experienced by the vegetation changes that have occurred since last time the Norwegian scientists (Rolf Vogt and Thorjörn Larssen, UiO) visited the catchment in 1995. During previous cooperation between Guizhou Institute for Environmental Sciences (GIES) and Dept. of Chemistry, UiO, equipment for sampling precipitation, throughfall and soil water has been installed at 7 plots within the catchment. A lot of scientific activity has been conducted or is still going on in this catchment. Previous results from this site are presented in Larssen et al. (1997) and their major findings are given below.

Soils are acidic yellow mountain soils with a relatively high clay content (Table C.15). The soils on the steep hill-slopes in the catchment are clearly affected by solifluction processes and by landslides. At several places we observed layers of colluvium (a layer of erosion products forming an inhomogeneous mixture of material from A, B and C horizons) on top of buried A horizons (the original top soil; see Table C.1). The colluvium layer can be in various stages of soil development. Because erosion

contributes to the supply of fresh mineral material to the soil surface this process may help keeping up the acid neutralization capacity in the root zone. There is a discontinuous O horizon in the catchment ranging from 0 to 15 cm in depth. Generally there is an A horizon underneath (ranging from 1 to 10 cm). In some cases the O horizon occurs directly on colluvium. In most places there is a transition zone (from 4 to 25 cm thick) between A and B horizon, indicated as an AB horizon, which still has a relatively high organic matter content.

When we visited the catchment, we observed several landslides which had occurred during a very high flow in early spring 1996. Parts of the streambed were covered by deposits from the slides, and some trees had fallen over the stream. To date parts of the slides have been eroded by the stream water. Prior to our visit in the catchment, there had been several days with rain. The streams were at high flow and the water in the dam had a grayish color, probably caused by soil colloids.

### **Air pollution**

The median precipitation pH was 4.40. Sulfate was the dominant anion in the precipitation and accounted for between 80 and 90 % of the total anionic charge on an annual basis (Table 3.5). The dominating cation in the precipitation was calcium, usually accounting for between 45 and 60 % of the total positive charge. The concentrations are generally similar to those found in Chongqing in the neighboring province, Sichuan. Large differences in concentration in throughfall at the different plots were observed, probably mainly due to differences in vegetation density. A clear increase in base cation concentrations and a decrease in H<sup>+</sup> concentrations in the throughfall compared to precipitation strongly indicate that dry deposition of alkaline dust is important for the total deposition chemistry.

Larssen et al. (1997) estimated a likely average total deposition of sulfur to be between 8.5 to 10 g S m<sup>-2</sup> yr<sup>-1</sup>, which is about 10 times higher than at Birkenes, southernmost Norway, an area strongly affected by acidification. Average concentration of SO<sub>2</sub> in air in 1993/94 was reported to be 44 µg m<sup>-3</sup> in the Liu Chong Guang catchment.

**Table 3.5.** Median and quartile concentrations (µeq L<sup>-1</sup>) of major ions in the precipitation in the catchment (Larssen et al., 1997).

	pH	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	F <sup>-</sup>	Cl <sup>-</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	NH <sub>4</sub> <sup>+</sup>
Median	4.40	228	19	5	6	133	30	29
Lower quartile	4.19	147	11	0	0	79	16	6
Upper quartile	4.74	334	35	10	14	199	53	79

### **Soil and soil water**

Most data from the Liu Chong Guang catchment, except for the soil transect and plot H, have been published in Larssen et al. (1997). Precipitation-, throughfall-, soil water- and stream water samples were analyzed at GIES in Guiyang. Some samples have in addition been analyzed at Department of Chemistry, UiO. Soils from 8 soil plots were analyzed at UiO. Soils from plot H and a transect have been analyzed for C/N ratios at NISK (Table C.2) and particle size distribution and mineralogy at Department of Geology, UiO.


Soil physical characteristics at Liu Chong Guang are quite variable. At plot C bulk density is rather low with values of about  $0.80 \text{ kg dm}^{-3}$  at least to 55 cm depth. Soil porosity is nearly constant with depth (about 64%). At plot H the soil is considerably more dense ( $1.05 \text{ kg dm}^{-3}$  in the A and  $1.32 \text{ kg dm}^{-3}$  in the AB) and less porous (porosity of 59% and 48% in the A and AB horizon, respectively). The differences in the soil's density and porosity at plots C and H are also reflected in the water retention characteristics, particularly in the sub-soil. For example, between 20 and 25 cm depth soil moisture content decreases from 44% at field capacity to 12% at wilting point on plot C, whereas these values are 37% (field capacity) and 18% (wilting point) at plot H (Table C.5). We estimate that the soil root zone (0 to 60 cm depth) has a storage capacity for plant available water amounting to about 200 mm (plot C) and 120 mm (plot H). This indicates that drought stress may occur in some parts of the catchment, but not in others.

Aluminum is the major exchangeable cation in the soil. In the mineral soil, the median aluminum saturation (AIS) is 83.7%. Hence the base saturation is generally quite low (Table C.1). The median effective cation exchange capacity ( $\text{CEC}_E$ ) for the mineral soil is high ( $86.2 \text{ meq/kg}$ ). In the upper horizon, rich in organic matter, the  $\text{CEC}_E$  is especially high, up to  $223 \text{ meq/kg}$ . Generally  $\text{CEC}_E$  decreases with depth together with the organic matter content. The median pH in the soils ( $\text{pH}_{\text{H}_2\text{O}}$ ) is 4.09. As shown in Table C.13, the sulfate adsorption is low despite the relatively high clay and sesquioxide content (Liao et al., 1994). The soil mineralogy is rather homogeneous and similar to what was found for the Tie Shan Ping site. The soils are dominated by quartz and clay minerals (median percentages 78 and 14 respectively). The remaining percentage was made up by varying amounts of plagioclase and K-feldspar. The main clay mineral is kaolinite and there is some vermiculite, partly hydroxy-interlayered. Also various amounts of residual illite and chlorite were found. C/N ratios decrease from about 18 in the O horizon to values between 11 and 18 in the A horizon. In the AB and B horizons the C/N ratio is between 6 and 14. As in the soils investigated at other sites on our trip in China, the C/N ratios are low, suggesting that nitrogen is available in considerable amounts. The total store of nitrogen in the upper 59 cm of the soil at plot H was estimated  $7\,400 \text{ kg ha}^{-1}$  (Table C.4).

As in throughfall, there are large differences in ion concentrations in soil water within the catchment (Figure 3.2, Table C.10). The median total ion concentration varies with a factor of five. This large spatial variation in the catchment makes generalizations on the soil water chemistry difficult. At all plots, the pH decreases from throughfall to the upper soil horizon. This is probably partly due to high concentrations of natural organic acids in the upper soil layer and partly due to ion exchange with soil organic matter and to base cation uptake and release of protons by plant roots. pH in soil water tends to increase with depth. Sulfate is the dominating anion also in the soil water. In some horizons nitrate is of significance. The source of the nitrate is not certain; the direct input is low, but there is considerable deposition of  $\text{NH}_4^+$  which may be oxidized in the upper soil layer. Considering the low C/N ratios, nitrate production in the soil either by decay of organic matter or by nitrogen fixing bacteria, is also possible. Mineralization of organic matter as an effect of lysimeter installation cannot be excluded, even though there is no clear time trend in the data. The median total fluoride concentrations vary from  $1 \mu\text{eq L}^{-1}$  to about  $60 \mu\text{eq L}^{-1}$ , suggesting that minerals in the soil are likely to be an important source. Calcium accounts for roughly half of the cationic charge and magnesium gives a significant contribution. Generally hydrated  $\text{Al}^{3+}$  was the dominating Al species in soil water accounting for more than 50% of the  $\text{Al}_a$  fraction.

As found for the Tie Shan Ping catchment, the  $\text{Al}^{3+}$  concentrations do not appear to be controlled by the solubility of a gibbsite mineral ( $\text{Al}(\text{OH})_3$ ). Instead we observe the typical increasing undersaturation with decreasing pH. A somewhat better agreement was found for a jurbanite ( $\text{AlSO}_4\text{OH}$ ) equilibrium, although this does not necessarily imply the occurrence of a mineral of this type.

The concentrations of aluminum in many soil water samples are high, and likely to increase with increased acid deposition. However, in many cases high concentrations of base cations may dampen biological effects. The median molar ratio  $Al_i/BC^{2+}$  for the different plots in the catchment, exceeds unity in a few horizons. The temporal variation is large. One should also note the large differences even within this small catchment, which illustrate the difficulties in making critical load maps for an area. Laboratory studies (Liao et al., 1997a,b) indicate that the soil may tolerate the present deposition for decades before the  $Al_i/BC^{2+}$  ratio increases substantially.



**Figure 3.2.** Soil water chemistry in the various plots at the Liu Chong Guang catchment. Left and right side bars represent cations and anions respectively (from Larssen et al., 1997).

Rough critical load mapping has been implemented in the RAINS-Asia program on a raster of  $1^\circ \times 1^\circ$  grid cells (Hettelingh et al., 1995). In the Guiyang area a critical load of  $0.32\text{--}0.80 \text{ g S m}^{-2} \text{ yr}^{-1}$  was estimated (with 75% of the ecosystems protected). Since the studied catchment is not likely to be particularly resistant to acidification, the values seem low considering that the  $\text{Al}_i/\text{BC}^{2+}$  ratios are generally  $< 1$  and that only minor effects on the vegetation have been reported (see below) with a present deposition estimate in the catchment of  $8.5 \text{ to } 10 \text{ g S m}^{-2} \text{ yr}^{-1}$ . There may be several explanations for this. The critical load calculation does not say how fast damage will occur. Hettelingh et al. (1995) stated that the deposition of base cations is a major uncertainty; the value used is very low compared to the present deposition in the catchment.

### Surface water

The streams in the Liu Chong Guang catchment are relatively acidic, although pH is considerably higher than typically found in soil water. The samples collected in April 1997 showed ANC about  $-50 \mu\text{eq L}^{-1}$  or less. The predominant acidifying compound is sulfuric acid, mirrored by the high sulfate concentrations ( $700\text{--}1400 \mu\text{eq L}^{-1}$ ) in the streams (Table 3.6). This concentration is almost 10 times higher than in the Birkenes catchment in southernmost Norway. However, the acid neutralizing capacity in the catchment is much lower at the Norwegian site, so that the ANC in surface water is only slightly higher at Birkenes. The nitrate concentration, which is high in some soil water samples, is fairly low. The  $\text{Al}_i$  concentration, although high especially in the April samples, is lower than in soil water. It is likely that acid neutralizing and aluminum removing processes in deep soil layers (where we have no lysimeters) and the sedimentary sandstone bedrock, reduce the acidity of stream water compared to soil water. Even though the concentrations of aluminum in the streams are similar to those found in other parts of the world, where damage to organisms has been observed in surface water (Rosseland et al., 1990), the high concentrations of base cations may dampen harmful effects of aluminum.

The highest concentrations of several heavy metals and trace elements (in the 1997 survey) were recorded at this site. This was the case for Cu, Zn, Cr, Ni, Co, Sr, U, Be, Li, and Y. Also the concentration of Pb was high (Table C.7). High concentrations of these metals are probably a combination of atmospheric inputs and rather high mobility of metals because the Liu Chong Guang streams are relatively acidic.

**Table 3.6.** Average concentrations of major chemical compounds in two small streams draining the Liu Chong Guang catchment. Data denoted Ia and IIa are unpublished average values from 1992 for the western and eastern stream, respectively (Xiong and coworkers at GIES), while Ib and IIb are from the same streams collected during our survey April 28, 1997. During our survey we also sampled 20-30 m downstream the confluence of the two streams when the discharges were  $\approx 40\text{--}50 \text{ L sec}^{-1}$  (III-1) and  $5\text{--}10 \text{ L sec}^{-1}$  (III-2).

Site	pH	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{Na}^+$	$\text{K}^+$	$\mu\text{eq L}^{-1}$			ANC	$\text{Al}_a$	$\text{Al}_i$	TOC
						$\text{SO}_4^{2-}$	$\text{Cl}^-$	$\text{NO}_3^-$				
											$\mu\text{g Al L}^{-1}$	$\text{mg C L}^{-1}$
Ia	5.01	362	245	32.5	28.6	670	13.2	3.2	-18	368	306	0.57
Ib	4.68	459	253	26.5	34.0	781	19.7	19.3	-48	600	582	0.81
IIa	5.21	405	291	31.6	30.3	732	17.0	5.1	4	243	214	0.48
IIb	4.71	768	577	26.5	39.1	1415	28.2	51.4	-85	1010	973	1.80
III-1	4.61	454	338	28.7	32.7	885	19.7	27.1	-79	868	837	0.95
III-2	4.73	433	335	31.2	32.0	854	19.7	15.0	-57	512	497	0.52



We also measured pH in the stream in the neighboring catchment (see site description). The pH was 7.36. This demonstrates the large differences in geology that may exist within a relatively small area, and the influence it has on streamwater chemistry.

### Vegetation

The Liu Chong Guang catchment shows similar damage to Masson pine as the Tie Shan Ping catchment in Sichuan. On the ridges, more exposed to air pollutants and on shallow soils, the defoliation of Masson pine was more pronounced than on less exposed sites on thicker soils, in topographic depressions and in the valley bottom. Therefore, we measured photosynthetic activity of species on both types of localities (Table C.20). On the ridges the mean  $F_v/F_M$  ratio was lower than in the depressions (0.79 vs. 0.80), although one of the measurements on the ridge showed a very high value, 0.84, which is the highest value recorded. The standard deviation of these measurements is similar to those at sites close to the cities (0.02), but higher than for most of the measurements done at remote sites. Corresponding measurements of Chinese fir show high values, indicating that this species is not affected by air pollutants at current levels. Most of the specimens were in excellent condition, although some seemed to be affected by soil erosion, which causes changes in light and temperature conditions and damages to the root systems.

As at most of the other investigated sites in China, the needles of Chinese fir and Masson pine suggest sub-optimal availability of phosphorus. In general the needles of Chinese fir have higher contents of base cations (Ca, Mg, K) than the needles of Masson pine. In contrast, the content of aluminum is highest in Masson pine. Molar Ca/Al ratios in Masson pine needles are close to 12.5, which may indicate aluminum damage. Molar Ca/Al ratios in fir needles range from 41 to 141, which is well above the levels where aluminum damage is likely to occur (Table C.3).

### Invertebrates and diatoms

Diatoms were collected from site I (western stream), site II (eastern stream), site III (below the confluence of the streams), site IV (the neutral tributary with pH 7.36 and conductivity  $254 \mu\text{S cm}^{-1}$  at  $12.8 \text{ }^\circ\text{C}$  entering the stream below the dam), and from site V (50-100 m below the confluence of streams III and IV, pH 7.16, conductivity  $157.7 \mu\text{S cm}^{-1}$  at  $13.6 \text{ }^\circ\text{C}$ ). The site I and II samples contained much fine silt, and diatoms were very sparse. Only a few *Eunotias* were observed, suggesting quite acid waters. The site III sample was very similar to samples from sites I and II in overall appearance with much fine silt. However, a greater range of diatom taxa was observed with *Pinnularia spp.*, *Eunotia spp.* and *Achnanthes spp.* being the most abundant. The sample from site IV also contained much fine silt and few diatoms. The dominance by *Achnanthes spp.* and small *Navicula spp.* suggests circumneutral waters. Site V is one of the richest sites in the series. Diatoms are abundant and diverse with taxa indicative of circumneutral waters and of a higher nutrient status. Taxa with affinities close to *Gomphonema parvulum*, *Achnanthes minutissima* and *Nitzschia palea* were the most common.

Invertebrate dipnet samples exist from site V and site III (low flow), and from site IV where some stones were inspected. In the acid confluent stream (site III: pH 4.6-4.7) we found no animals in the dipnet sample, but by examining the diatoms samples we found a few (6) small stoneflies (Plecoptera), 2 midge larvae (Chironomidae) and 1 as yet unidentified larvae. From site IV, where the substrate is affected by siltation, we collected a few invertebrates, i.e. two very small pulmonate snails (Gastropoda), some (10) stoneflies (Plecoptera) and one unidentified insect pupae. At site V on the other hand, we collected a variety of invertebrates belonging to several taxonomic groups: worms

(Oligochaeta) (4 specimens), pulmonate snails (Gastropoda) (2), mayfly larvae (Ephemeroptera) (1), stonefly larvae (Plecoptera) (16), caddislarvae (Trichoptera) (10), midges (Chironomidae) (24), seed or mussel shrimps (Ostracoda) (8) and two unidentified insect larvae (possibly Diptera). The clear differences in stream fauna between the acid stream sites and the more neutral are most probably related to water quality, particularly since snails are very acid sensitive and frequently used as indicators for acidification. But in addition to the considerable spatial variation in water quality observed in Liu Chong Guang, catchment processes like erosion and siltation also affect stream fauna and flora and should be noted.

### 3.2.2 Leishan

#### Site description

The Leishan or Leigong Mountains are located about 140 km to the east of Guiyang in the Guizhou Province. The sampling site (26°10'N, 108°10'E) is at 1600 m a.s.l. about 40 km south-east of the city Kaili where a large coal-fired power plant is under construction. In a catchment located 100-200 meters upstream of a plant bottling natural water, samples were collected on both sides of the small steep valley (slope facing west: plot 1; slope facing east: plot 2). On a ridge some way down the road another sampling site was selected in order to include a site with Masson pine (plot 3).

There are 10 weather stations in these mountains, 5 of which have data from the last 30 years. Air temperature, amount of precipitation, wind speed, wind direction, cloud cover and extreme weather conditions are normally recorded. Highest amounts of rain occur on the southern side of the mountains. A weather station located immediately downstream of the catchment was run from June 1, 1985 to December 31, 1988. Based on these monitoring data, annual precipitation in the catchment area varied from 1200 to 1600 mm, and about 80% of the rain was found to fall during April-September. At this weather station up to 315 days with fog have been recorded during a year (in 1987). Only two mountain areas in China have more foggy days than the Leishan Mountains. The number of foggy days decreases significantly further down in the valleys. Annually the area has 1200-1400 sun hours. Data from the different weather stations showed large temperature and precipitation gradients with respect to altitude in these mountains. The amount of precipitation increases with height, and is about 1700 mm in the upper areas (1800 - 2100 m a.s.l.). Snow may occur during winter, primarily at altitudes > 1400 m a.s.l. Last February, a snow fall caused severe damage to many trees in the area. Wind speed is 4 m sec<sup>-1</sup> 80-90% of the time at 1600 m a.s.l., but on the mountain tops wind speeds up to 30 m sec<sup>-1</sup> have been recorded. The temperature decreases with 8-10°C from the valley bottom to the mountain tops. At altitudes > 1500 m a.s.l. the air temperature is seldom > 20°C. Relative humidity is often > 80%.

Soil development is different for the three sampled plots. This difference is most pronounced with respect to the thickness of the transitional AB horizon, ranging from 10 cm in the mixed stand (Masson pine, Chinese fir, broadleaf; plot 3) to 20 cm under Chinese fir (plot 1) and 30 cm under broadleaf (plot 2). The profiles on the slopes (plots 1 and 2) had a 2-3 cm organic litter layer on thick A and AB horizons. The yellow soil in plot 3 had only a 1 cm thick fermentation (F) layer on thin A and AB horizons. The soil at plot 2 was covered by colluvium deposits of shale derived soil (about 1 cm thick) from the mountain side above.

## **Air pollution**

As the area is long away from large emission sources, we may assume that the air is relatively clean, and that the sulfate concentration in precipitation is similar to, or lower than, the concentrations reported for Simian Shan ( $\leq 2 \text{ mg S L}^{-1}$ ). However, the concentration levels of light hydrocarbons show that the area is influenced by pollution sources on the regional scale.

## **Soil**

The content of organic matter in the A and AB horizons was relatively high (Table C.1). The highest organic matter content was found in the A horizon of the broadleaf plot. Also the mineral sub-soils at these sites have the highest content of soil organic matter (see Table C.1). Soil mineralogy was determined in plot 1 and plot 3 (Table C.15). Except for a somewhat lower K-feldspar content, the composition is quite similar to that found in soils originating from granitic bedrock. This indicates a generally low degree of chemical weathering and that the indigenous sandstone bedrock is unusually rich in the more easily weatherable plagioclase and feldspar minerals. The quartz content was lower in plot 1 (<60%) than in plot 3. Furthermore, the clay fraction in plot 1 contained no secondary kaolinite minerals in contrast to plot 3. This may indicate an even lower degree of chemical weathering of the soil profile in plot 1 than in plot 3.

The A-horizon soil in the mixed forest plot (3) was a sandy loam (33.3% less than  $63 \mu\text{m}$ ), while the B-horizon soil in the Chinese fir plot (1) contained less silt and clay, thus making it a sand (89.6% greater than  $63 \mu\text{m}$ ). Stones were observed to be common throughout all the profiles at plots 1 and 2. Furthermore the stone content decreased down into the plot 2 profile, indicating that other pedogenetic processes than chemical weathering are dominating also here.

The base saturation (BS) generally decreases with depth. In the A horizon BS is well over 50%, and in the AB and B horizon it decreases to values between 20% and 40%. The soils in the small valley (Chinese fir and broadleaf stands) are less acidic and have higher BS (see Table C.1) than plot 3 (on the ridge). In the mixed stand BS decreases from 17% in the A to 6% in the B horizon. Nutrient cycling is likely to contribute to the observed decrease in BS with depth. However, other processes may also play a role. Considering the mineralogy and the observed distribution of stones, the high BS in plots 1 and 2 may be caused by the steep slopes causing enhanced physical erosion. The strong physical erosion exposes fresh primary minerals (richer in basic constituents) to chemical weathering. Furthermore, the B-horizons were rather compact, probably allowing little water to penetrate; lateral water flow in the A horizon may therefore be an important hydrological pathway. This flowpath may cause a transport of basic constituents down-slope within the upper horizons instead of down in the profile. Due to the distance to anthropogenic emission sources, including farming activities, base cation deposition is not likely to be the cause of the observed decrease in BS with depth.

The high BS at plots 1 and 2 indicates that these sites are rather resistant to soil acidification. However, the large amounts of stones may imply that the actual reservoirs of exchangeable cations are fairly limited and the sites therefore more sensitive than intuitively estimated from the BS.

The  $\text{CEC}_E$  ranges from the highest to lowest values among the sampled sites. The high values were found in soils with high organic matter content and there is a good correlation between organic matter content (i.e. tot C %) and  $\text{CEC}_E$  ( $r^2 = 0.81$ ) at these sites and horizons.

Plot 3 has high  $\text{CEC}_E$  even in the B-horizon (Table C.1). This is partly due to relative high organic content and partly to the clay minerals. At this plot there is much less supply of fresh soil material and

the profile has the lowest base saturation levels among these soils. The soil water at plot 3 will be rather sensitive to acid deposition due to low BS.

C/N ratios in the soils of the Chinese fir and broadleaf stands are low, particularly in the A horizon (about 11). In the mixed stand the C/N ratios are slightly higher, but still low compared to values common in forest ecosystems of the temperate and boreal zone (Table C.1).

### Surface water

The electric conductivity in the two streams investigated was extremely low, i.e. 11-12  $\mu\text{S cm}^{-1}$ . This is presumably related to low atmospheric deposition and lack of significant amounts of carbonates in the catchment. Even so, both streams have relatively high ANC and pH. Compared to the other sites visited, sodium ( $\text{Na}^+$ ) and not calcium was the predominant base cation in the streams (Table 3.7), which indicates a sodium rich soil. We also checked the conductivity in the main river about 20 km downstream from the main sampling site (14-15  $\mu\text{S cm}^{-1}$ ) and in Leishan city (35  $\mu\text{S cm}^{-1}$ ). This confirms a very electrolyte poor water in large areas of the Leishan Mountain. Accordingly, the water bodies in the Leishan Mountain area could be relatively sensitive to acid rain. At present impacts from acid rain are minor.

The very high concentrations of tin (Sn) in stream I (25.4  $\mu\text{g Sn L}^{-1}$ ) is difficult to explain (Table C.7). This stream should be resampled to confirm this result, especially since a mineral water company bottling natural water was located nearby. Tin is a very toxic element.

**Table 3.7.** Average concentrations of major chemical compounds in two small streams draining a catchment in the Leishan Mountains, 1600 m a.s.l.. Site I is the stream draining the catchment containing soil plots 1 and 2.

Site	pH	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{Na}^+$	$\text{K}^+$	$\mu\text{eq L}^{-1}$			ANC	$\mu\text{g Al L}^{-1}$		TOC $\text{mg C L}^{-1}$
						$\text{SO}_4^{2-}$	$\text{Cl}^-$	$\text{NO}_3^-$		$\text{Al}_a$	$\text{Al}_i$	
I	6.75	25.5	14.8	60.9	3.07	25.0	5.64	14.6	59	5	0	0.20
II	6.71	26.0	15.6	61.3	2.81	29.2	5.64	22.1	49	5	0	0.28

### Vegetation

Leishan Mountains contain many different vegetation units, which cover a number of habitats for different vascular plants and bryophytes. Therefore, the biodiversity of plant species is very high, compared to the other visited areas. Some of the species were damaged by chilling or spring frost. The last winter was very hard in this area, compared to normal conditions, so some trees, shrubs and herbs showed visible defects like malformation and discoloration. Defoliation was observed on Masson pine, and a number of needles showed necrosis. The unripe tips of many new shoots have been killed, while well developed needles were brown from the tip. Such damage was, to a lesser degree, also seen on Chinese fir. Compared to the Simian Shan catchment the PEA measurements show surprisingly low values: The mean  $F_V/F_M$  ratio for healthy ramets of Masson pine and Chinese fir was 0.80 and 0.79 respectively, and for both species the standard deviation showed very high values (0.03 and 0.02). This was expected due to the obvious effect of frost and chilling. Still it is interesting to observe that the  $F_V/F_M$  ratio is higher in this mountain area with harsher climatic condition than in areas close to the cities (Table C.20).

### **Invertebrates and diatoms**

Diatoms and stream invertebrates were collected from stream I and II. These streams were fast flowing with small pockets of water between boulders and stones. The streambed consisted of gravel, coarse sand and some organic debris. The diatom sample from site I contained a relatively diverse flora, dominated by *Diatoma sp.*, *Cocconeis cf. placentula*, *Nitzschia sp.*, *Gomphonema spp.*, *Synedra ulna* and a *Surinella sp.* Diatoms from site II (the left brook facing upstream) appeared very similar to the sample from site I. These taxa are typical of circumneutral (pH 6-7) water.

These sites had also a quite diverse stream fauna with at least 9 and 8 taxonomic groups respectively and with relatively high numbers of individuals in the samples (184 and 126). At these sites we found flatworms (Turbellaria), small aquatic worms (Oligochaeta), mayfly larvae (Ephemeroptera), stonefly (Plecoptera) larvae, larval and adult water beetles (Coleoptera), caddisfly larvae (Trichoptera), midge larvae (Chironomidae), blackfly larvae (Simuliidae), some primary terrestrial insects and some specimens which currently are unidentified. No snails were found, a fact which perhaps can be related to the low concentration of base cations, Ca and Mg (see above). Although we lack information about the species composition of these samples and their respective acid tolerance, we believe that these streams have intact biological communities and may serve as reference site for comparable sites in more impacted areas.

### **3.3 Guangzhou area**

Of the 29 ecosystem monitoring stations in China run by the Chinese Academy of Science (CAS), three are located in the Guangdong province; at Dinghushan, Heshan and Chonghua.

Guangzhou city (Canton) is situated on the Pearl river delta, about 120 km north-west of Hong Kong. For 1988 Qi and Wang (undated) gave the following concentrations (in  $\mu\text{eq/L}$ ) in precipitation:  $\text{Ca}^{2+}$ : 175,  $\text{NH}_4^+$ : 141,  $\text{NO}_3^-$ : 33,  $\text{SO}_4^{2-}$ : 255 and  $\text{H}^+$ : 41. It was stated that the annual average  $\text{SO}_2$  concentration had decreased from  $90 \mu\text{g m}^{-3}$  in 1981 to about  $50 \mu\text{g m}^{-3}$  at present. In contrast, the  $\text{NO}_x$  concentration had increased from  $50 \mu\text{g m}^{-3}$  in 1981 to about  $100 \mu\text{g m}^{-3}$  mainly due to the increasing number of motor vehicles.

About 80 km north of Guangzhou city lies the Conghua city. It is located 230 km south of the city Shaoguan, having the highest emissions of air pollutants in the Guangdong Province. In Conghua, acid rain has been documented back to 1985/86. The Guangzhou Environment Monitoring Center has an acid rain monitoring site on the roof of their building in Conghua. In 1991 pH in rain was  $< 5.6$ , 61% of the year, while in 1996 only 30% of the time. While weighted average pH was 4.1 in 1991, it was 5.0 in 1996. The concentration of  $\text{SO}_4^{2-}$  in precipitation has been reduced from  $7.1 \text{ mg L}^{-1}$  down to  $3.9 \text{ mg L}^{-1}$  in 1995. The precipitation sampler was just a simple container, which also collected dry deposition and allowed water to evaporate. Air concentrations of  $\text{NO}_x$  and  $\text{SO}_2$  are  $< 60 \mu\text{g m}^{-3}$  which are lower than the Class II national standard.  $\text{NO}_x$  emissions are likely to increase significantly in the coming decades as in Guangzhou city. The reduction in S-concentration, and the resulting increase in rain pH, are mainly caused by the building of high stacks in the area, and the reduced emissions in Hong Kong the last years. There are also data on damage to materials caused by air pollution in the area.

### 3.3.1 Dinghushan

#### Site description

About 80 km west of Guangzhou city nearby the city of Zhaoqing, the Dinghushan mountains rise above the plains. Annual mean temperature is 20.9° C, while mean annual precipitation and evapotranspiration are 1956 mm and 1115 mm, respectively. Most of the precipitation falls from April to September, when mean monthly rainfall normally exceeds 200 mm. Mean annual relative humidity is 81.5%.

Dinghushan Biosphere Reserve (DHSBR) covers major parts of the Dinghushan mountain area. The area is very hilly, and a few lakes are also present in the area. Total area of the reserve is 1155 ha and about 79% is forested (Kong et al., 1993). DHSBR is located at 23°10' N and 112°32' E, near the Tropic of Cancer. While 2/3 of the area of the world near the Tropic of Cancer consists of deserts and semideserts, the Dinghushan area is covered by tropical-subtropical forests. This is mainly due to the unique geographic location and climatic conditions. Many forest types with high biodiversity and vast genetic-stock resources are present in the area. Compared to surrounding degraded forests which have been subjected to long-term human disturbance, DHSBR contains a rare, more than 400 years old, primary forest. The geology in Dinghushan consists of sandstone, sandy shale, shale and quartz sandstone. Research has been performed in DHSBR since the 1950's. However, systematic and multidisciplinary research was initiated only after 1978, the year that Dinghushan Forest Ecosystem Research Station (DHSFERS) of CAS was established here. There are several ongoing international projects at DHSFERS.

The soils are in general shallow yellow soil, between 30 - 65 cm deep, with only a thin (1 cm) litter (L) layer and no organic (O) horizon. The soils in the sampled areas consist mainly of lateritic red-earth.

Three sites were visited in the Dinghushan mountain:

A) One soil profile was collected below a very old tree, approximately in the middle of a more than 400 years old monsoon evergreen broadleaf forest (MEBF). This forest is located above the Qingyun Temple, built towards the end of the Ming Dynasty. The catchment (about 370 m a.s.l.) is drained by a small stream.

B) The site is located on a steep hillside (about 200 m a.s.l.) facing an artificial hydroelectric reservoir (named Grass pond). The site was managed (terraced) some 40 years ago giving rise to a 30 cm ploughed A horizon (Ap) top soil. The vegetation is primarily pine broadleaf mixed forest (PBMF) and secondary monsoon evergreen broadleaf forest (MSEB).

C) A site called Tang e Ling which primarily is a 40 years old pine forest (PF). Typical soil depth is 30 cm. The ground vegetation is removed by local farmers every year. There is also a problem connected with increasing number of tombs in the area.

In addition, a water sample from the Lake Tianhu (Heaven or Sky lake in English), a headwater situated at 358 m a.s.l., was collected from a pipeline close to where the soils of site B were sampled. The sample was taken 5 minutes after the water was turned on. Samples were also collected for local scientists (CAS) for intercalibration purposes.

### Air pollution

Wet-deposition chemistry data exist but we have not received this information. However, according to Qi and Wang (undated), the precipitation pH in this area was about 4.7 in 1988.

### Soil

The effective cation exchange capacity ( $CEC_E$ ) is high in these soils relative to the organic matter content, that range from the lowest measured value (0.37%) in the B horizon to moderate values (up to 5.33%) in the A-horizon. In all three forest stands at Dinghushan the soils are strongly acidic with base saturation below 10%, even in the A horizons (Table C.1). The exchange sites are therefore dominated by aluminum. Despite the high  $CEC_E$ , the low BS causes these soils to have poor capacity to buffer any acid deposition by base cation release. Instead, leaching of aluminum is to be expected. This was confirmed by labile aluminum values close to  $1 \text{ mg L}^{-1}$  in the stream (see below). The C and N contents are low, as is the C/N ratio, indicating well decomposed organic matter, due to the warm subtropical conditions.

These soils are old and are comprised of almost entirely secondary clay minerals. The high clay content (not analyzed) explains the relatively high  $CEC_E$ . The age of the soils is probably the main reason for the lack of base cations on the ion exchanger.

The highest organic matter content and  $CEC_E$ -values in the A-horizons were, as could be expected, found in the monsoon evergreen broadleaf forest (site A). The lowest  $CEC_E$  was found in the mixed forest plot (site B). This plot has been managed into terraces about 40 years ago. The low  $CEC_E$  may therefore partly be due to the man-made inclusion of deeper soils containing less organic matter and clay. In addition cultivation *per se* may cause oxidation and decreased inputs of organic matter. These impacts may also explain the increase in BS from the A to the B horizon. The highest BS was found in the A horizon in plot C on the border of the natural park reserve, neighboring extensive areas of farmland and a small city. It is therefore likely that a considerable dust deposition may contribute to this high value.

### Surface water

The pH in the stream draining the monsoon evergreen broadleaf forest (MSEB, site I) was very low, 4.27 (Table 3.8). In contrast to the other sites in this survey, the stream water contains a relatively high concentration of chloride. The site is fairly close to the ocean and the ratio between Na and Cl concentrations indicates a contribution of NaCl from sea spray aerosols. The TOC concentrations are too low to have any significant effect on water pH. Since the pH in rain water is relatively high in the area, the acid runoff is probably mainly a result of a natural acidification processes, but S- and/or N-deposition may contribute. Kong et al. (1993) found a low pH i.e. 4.22 in a stream feeding into the Tianhu lake. All the other 9 surface water sites presented in Kong et al. (1993) had pH between 5.0 and 6.6.

The sampling site I is located very close to the Qingyun Temple, a site often clouded by smoke from joss sticks and candles burnt for offerings. Totally 700 000 to 800 000 tourists visit this site every year. This may cause a local air pollution problem, which might be a reason for a high concentration of lead (Pb) in this stream (Table C.7). The stream also had an unexpectedly high concentration of organic nitrogen (Org-N). Since the total concentration of carbon in the stream was very low (Table C.6), more samples are needed to check the validity of this unexpected observation.

**Table 3.8.** Average concentrations of some chemical compounds in a small stream (Site I) draining a monsoon evergreen broadleaf forest (MSEB), and from tap-water from the Tianhu Lake (Site IV).

Site	pH	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	μeq L <sup>-1</sup>			ANC	Al <sub>a</sub>	Al <sub>i</sub>	TOC
						SO <sub>4</sub> <sup>2-</sup>	Cl <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>		μg Al L <sup>-1</sup>	mg C L <sup>-1</sup>	
I	4.27	33.4	48.5	46.5	7.67	75.0	67.7	35.3	-42	939	919	0.74
IV	6.54	99.8	40.3	32.2	6.39	119	39.5	48.9	-28	16	11	0.60

The other water sample was collected from a pipeline 1-2 kilometers from the Tianhu Lake (site IV), and might therefore not reflect the real water conditions in the lake. This water was primarily taken for intercalibration purposes. Another sample was therefore collected for analysis by Chinese scientists. The water quality in the lake was relatively good, with high pH, but a very extreme concentration of zinc (1 904 μg Zn L<sup>-1</sup>) was recorded. Obviously this was a result of dissolution of this element from the water pipeline. If accounting for the high Zn value, the ANC of the water will be about 20 μeq L<sup>-1</sup> instead of -28 μeq L<sup>-1</sup>. In addition to the very extreme concentration of Zn, the water also contained the highest concentration of cadmium (Cd) and antimony (Sb) of all sites included in our survey (Table C.7). It is important to note that the water sample was collected after letting the tap run for 5 minutes. Thus, the high concentrations of Zn, Cd and Sb are potential dangerous to people drinking this water.

### Vegetation

At Dinghushan, the vegetation was investigated at two locations: Station B, close to the main road within the DHSBR and Station C, a few km away from DHSBR (Tang e Ling). The mean  $F_v/F_M$  ratio measured for both Masson pine and Chinese fir was lower in the site close to the road (Table C.20). For Masson pine the ratio was high compared to values observed in the areas close to Guangzhou city (0.79 and 0.81).

The content of phosphorus in the needles of Chinese fir and Masson pine is low, similar to what is found at the other sites investigated (Table C.3). Despite the low base saturation in the soil the content of base cations (Ca, Mg, K) in the needles does not indicate a lack of these nutrients. Also the Ca/Al ratio in the pine needles is relatively high, indicating that aluminum damage is unlikely. As observed at the other sites, the content of aluminum is considerably higher in the pine than in the fir needles.

### Invertebrates and diatoms

We took diatom and invertebrate samples from site I. The water, which runs over the cliffs and with some cobbles in the crevices, provides few microhabitats for stream invertebrates. The water is very acid and low in dissolved salts. The diatom sample from this site is of very low diversity, almost completely dominated by small *Eunotia* taxa, confirming an extremely acidic water.

The invertebrate sample consisted of worms (Oligochaeta)(1), water mites (Hydracarina)(1), adult beetles (Coleoptera)(2), Caddisfly larvae (Trichoptera)(3), midge larvae (Chironomidae)(12) and one unidentified specimen. The specimens were all very small in size. The general impression is that of a poor stream fauna. At this point it is impossible to fully assess which main factors are responsible for the apparently poor fauna at this site. Data on freshwater fauna are largely lacking. Some data on freshwater fauna are presented in Kong et al., 1993.



### 3.3.2 Heshan

#### Site description

Heshan is also one of 29 ecosystem monitoring stations in China run by CAS, and is primarily a forest research station. The area is divided into 8 subcatchments with different types of vegetation. A local poultry farm located in this area emits large quantities of ammonia (NH<sub>3</sub>). We performed our measurements and sampling in subcatchment 2. The site was located at 80-100 m a.s.l. (22°41'N, 112°54'E). The subcatchment was classified as a mixed pine deciduous forest (PDMF), however with few coniferous trees. Soil samples were collected in a bamboo forest with some other broadleaf bushes and some Masson pines. It was located on the left side in the lowest part of the catchment. Soils at the sampling site lack an O-horizon and have a relatively thin organic-rich A horizon on top of a shallow B-horizon (10 cm thick.). The site was cultivated and most of the soils were terraced. The samples were collected from an area that appeared undisturbed.

This catchment had a drained swamp in the valley bottom and no other streams than the drainage water from the bog/marsh. The water flow was monitored by a V-notch weir. Runoff was minimal during our visit.

We got no data on freshwater biology, perhaps because some or most of the drainage streams only carry water during the monsoon.

We received many publications from research conducted at this station. Most of them deal with input/output studies of chemical macro constituents, some inventories of soil chemistry (no soil water samples), stream water chemistry, soil organisms (major taxonomic groups only), and some biomass estimates.

#### Air pollution

Table 3.9 shows that the precipitation in the area is heavily polluted mainly with ammonium and sulfate. The very high amounts of ammonium are clearly related to the local poultry farm, which emits significant amounts of NH<sub>3</sub>.

**Table 3.9.** Average chemical composition ( $\mu\text{eq L}^{-1}$ ) of rain (R), throughfall (TF), stem flow (SF), percolating water (PW) and surface water (SW) in an *Acacia mangium* forest in Heshan during 1993 (Fang et al., 1995).

	pH	Ca <sup>2+</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	NH <sub>4</sub> <sup>+</sup>	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	PO <sub>4</sub> <sup>3-</sup>
R	4.44	51.4	21.4	10.2	191	440	32.9	7.2
TF	4.87	209	88.0	97.4	284	1092	57.1	5.8
SF	3.86	273	153	125	344	2794	73.6	7.6
PW	5.15	147	57.6	29.2	351	420	11.4	7.5
SW	6.64	34.9	82.3	50.4	227	389	13.6	4.4

## Soil

Soils at the Heshan site have intermediate  $CEC_E$  and BS values in the A-horizon; in the B-horizons the BS is low (Table C.1). BS in the A horizon is about 13% which is slightly higher than the values observed at Dingushan. Soil pH in forest soils is low with values ranging from 3.7 in the forest floor to 4.1 at 50-80 cm depth. Furthermore, increase in the  $Si/(Al+Fe)$  ratio with depth indicates relocation of Al and Fe within the soil profile (Li et al., 1995). C/N ratios of soil organic matter are low and similar to those found at Dingushan. Our values are in good agreement with those of Li et al. (1995) who reported that the ratio decreases from 13 in the forest floor to 9 at 50-80 cm depth. The nitrogen store in the upper 15 cm of the soil at Heshan (including A and B horizon) is about  $2700 \text{ kg ha}^{-1}$  (Table C.4).

The soil texture was mainly clay loam and loamy clay. The bulk density of the soil at Heshan is rather high and the porosity low (Table C.4). In this respect the soil at Heshan is similar to the one at Tie Shan Ping. Soil moisture content in the A horizon at Heshan decreases from 37% at field capacity to 25% at wilting point (Table C.5). Water retention is even stronger in the B horizon where the soil moisture content decreases from 36% at field capacity to 32% at wilting point, i.e. even at high suction little of the water can be removed from the soil. Water available for plant in the upper 60 cm of this soil is estimated at only 35 mm, which makes this soil extremely drought sensitive.

## Surface water

Runoff in the stream draining this catchment was minimal during our sampling and the turbidity (concentrations of particles) was very high compared to that in other surface waters in this survey. Sodium is the predominant cation in the stream, and much higher than chloride, indicating a Na-rich soil (Table 3.10).

**Table 3.10.** Average concentrations of some chemical compounds in a small stream (Site I) draining catchment 2 in Heshan Forest Research area.

Site	pH	$Ca^{2+}$	$Mg^{2+}$	$Na^+$	$K^+$	$SO_4^{2-}$	$Cl^-$	$NO_3^-$	ANC	$Al_a$	$Al_i$	TOC
		$\mu\text{eq L}^{-1}$								$\mu\text{g Al L}^{-1}$	$\text{mg C L}^{-1}$	
I	6.91	38.4	51.8	101	39.9	39.6	28.2	0.36	163	15	3	1.20

**Table 3.11.** Yearly inputs by rain (R), outputs by surface water (SW) and percentage leakage  $[(R/SW)*100]$  of major chemical compounds in an *Acacia mangium* forest in Heshan during 1993 (Fang et al., 1995).

	Unit	$Ca^{2+}$	$Mg^{2+}$	$K^+$	$NH_4^+$	$SO_4^{2-}$	$NO_3^-$	$PO_4^{3-}$	Sum
R	$\text{kg ha}^{-1}$	8.793	2.363	4.751	47.57	75.25	4.88	1.029	144.6
SW	$\text{kg ha}^{-1}$	0.178	0.267	0.629	1.031	1.174	0.053	0.015	3.423
Leakage	%	2.0	11.3	13.2	2.3	1.6	1.1	1.5	2.4

Since the ANC of water is high, the catchment should be well protected against acid precipitation. However, the acid neutralization processes may be much less efficient during high flow periods. Despite the very high inputs of  $\text{NH}_4^+$  in the area, leakage of both  $\text{NH}_4^+$  and  $\text{NO}_3^-$  is very low. Generally percent leakage of all major ions investigated is extremely low in this catchment (Table 3.11). Further studies on this phenomenon seem warranted. Of trace elements, the concentration of rubidium (Rb) in the stream was very high ( $8.16 \mu\text{g L}^{-1}$ ). Similar values were observed in another stream in the same region (Baiyun Shan; Table C.7). Since these stream waters also had the highest turbidity (concentration of particles), it is reasonable to assume that Rb primarily is associated with clay particles.

### **Vegetation**

The area around Heshan biological station presents a wide range of vegetation types, containing a number of different habitats from more or less natural to heavy manipulated habitats. Studies of photosynthesis activity was performed in areas with as little as possible human influence. For both pine and fir the ratios were high. The measured value of one of the Masson pines was as high as 0.83, the highest value measured by us. This site seems to have favorable climatic conditions in addition to good nutrient conditions and limited influence of air pollution. It can be observed that the standard deviations for both species are very low (0.009), also reflecting the favorable conditions (Table C.20).

Also at Heshan the content of phosphorus in the needles of Chinese fir and Masson pine are rather low (Table C.3). As observed at the other sites, the content of aluminum is considerably higher in the pine than in the fir needles. The Ca/Al ratios in the needles of Masson pine are among the lowest observed in this study and suggest a high risk for aluminum damage. Unfortunately, a direct comparison of the low Ca/Al ratio in the pine needles with soil acidity characteristics at this site cannot be made, because we did not sample the soils in the vicinity of the investigated pine and fir trees.

### **Invertebrates and diatoms**

The small stream draining catchment 2 had no stone substrate suitable for sampling stream fauna and diatoms. No samples of freshwater biota were therefore collected. No other data on freshwater biology exist probably due to lack of permanent water flow.

## **3.3.3 Guangzhou Botanical Garden**

### **Site description**

South China Botanical Garden is situated in the northeastern suburb of Guangzhou. It was established in 1956, and is now the largest botanical garden in the subtropics of China, with total area of 300 ha. The garden is divided into two major parts, an exhibition and an experimental section. Its main objective is research on introduction and acclimatization of plants. There are registered about 5000 plant species within the garden, both domestic and aliens. The horticultural impact is of course prominent, but there are corridors and ridges in the landscape and some more or less inaccessible areas for the public that partly show unaffected vegetation structure and composition. Small groves of coniferous and deciduous trees are similar to corresponding areas at other sites. A unit belonging to the Chinese Academy of Sciences (CAS) is situated within the garden, in the Long Dong area. The academy has a staff of well qualified scientists in taxonomy, plant physiology and environmental ecology with access to well equipped laboratory facilities. Only some vegetation measurements were

conducted at this site. However, a number of published papers are based on scientific activities from "natural" and "manipulated" habitats within the park, most of them in Chinese. Scientists at CAS have an extensive collaboration with colleagues in the US and Japan. Many papers based on dose-response studies have been published in international journals.

### **Vegetation**

Within the South China Botanical Garden the health condition of Masson pine varies a lot. On some of the ridges defoliation is common. The distance to Guangzhou is short and some areas are directly exposed to pollution from the city, so it is not surprising that symptoms of severe injuries are observed. The mean  $F_V/F_M$  ratios for healthy ramets of Masson pine and Chinese fir were respectively 0.80 and 0.82, and for both of the species the standard deviation approaches the mean values for our measurements in the Guangdong province (Table C.20). There were no severe injuries seen on Chinese fir within the site. Non of the selected specimens seem to be affected by direct exposure to air pollutants from Guangzhou. The same is true for a number of herbs within the park. Many of the species were surprisingly healthy and not affected by stress. The only species that appears to be sensitive is Masson pine.

## **3.3.4 Baiyun Shan**

### **Site description**

The site is located 189 m a.s.l. (23°03'N, 113°19'E) at the base of the Baiyun Shan mountain, 7 km from the downtown of Guangzhou. It is located quite close to the Guangzhou airport and there is a water reservoir higher up in the catchment. The sampled stream probably dries out temporarily, but this stream was chosen because a presumably more permanent inlet stream to the reservoir was strongly influenced by human activities (recreation, agriculture and dwellings).

Annual average temperature is 21.8 °C and precipitation is about 1700 mm. The soil is laterite, developed from granite and sandstone. Organic matter in the soil varies between 2.5 - 4.2%, and pH in soil solution is < 4.8. A relatively thin A horizon overlies an AB and B horizons low in organic matter content.

### **Air pollution**

We have no air chemistry data from this site, but as mentioned above, it is located close to the airport, 7 km from downtown Guangzhou. Hence, it presumably has rather high concentrations of air pollutants.

### **Soil**

The  $CEC_E$  ranges from close to median values (47.2 meq  $kg^{-1}$ ) for the sampled sites in the A-horizon to less than the 20 percentile value in the AB- and B-horizons (23.4 and 22.5 meq  $kg^{-1}$ , respectively). Similarly, the aluminum saturation (AIS) on the ion exchanger ranges from the median value (89.6%) in the A-horizon to the highest values found (95.7%) in the B-horizon. The highest recorded AIS and relatively low  $CEC_E$  render this site one of the most sensitive to soil water acidification among the sites included in this survey. As observed elsewhere, C/N ratios in soils are low. The proximity to the large

emissions in Guangzhou and Hong Kong gives therefore rise for concern regarding the soil water quality in the future (e.g. the  $\text{Al}^{3+}/\text{BC}^{2+}$  molar ratio may become high).

### Surface water

The stream water seems to be well buffered against acidification (ANC:  $282 \mu\text{eq L}^{-1}$ ). Since the soil profile sampled showed very low base saturation, the water must be in contact with other soils and/or bedrock which are richer in base cations. As in Heshan and a tributary to Liu Xi river,  $\text{Na}^+$  is the predominant cation, and this streamwater is the only waterbody within this survey where  $\text{Cl}^-$  exhibits the highest concentration of the strong acid anions (Table 3.12). The reason may be that this site is located closer to the ocean than all the other sites in this survey.

**Table 3.12.** Average concentrations of some chemical compounds in a small stream in Baiyun Shan.

Site	pH	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{Na}^+$	$\text{K}^+$	$\mu\text{eq L}^{-1}$			ANC	$\text{Al}_a$	$\text{Al}_i$	TOC
						$\text{SO}_4^{2-}$	$\text{Cl}^-$	$\text{NO}_3^-$				
											$\mu\text{g Al L}^{-1}$	$\text{mg C L}^{-1}$
I	7.10	142	92.1	149	51.2	56.2	59.2	36.4	282	11	4	1.30

As mentioned earlier, the concentration of Rb in this stream was the highest recorded in this survey (Table C.7). This stream also showed the highest concentration of Fe, despite high pH and low TOC. Most likely, these high concentrations are a consequence of the high turbidity (Table C.6). Accordingly, both Rb and Fe are primarily associated with clay particles. This may also be the explanation why the highest concentration of arsenic (As) was recorded in the same stream. To what extent high concentration of As might be due to other factors, e.g. the short distance to the Guangzhou airport, is difficult to assess at present.

### Vegetation

Data from Baiyun Shan show particularly low values for the chlorophyll fluorescence measurements on the Masson pine (Table C.20). The median  $F_v/F_m$  ratio of 0.78 and a minimum value of 0.69 are approaching the record low values from the Nanshan catchment in Chongqing. The standard deviation is very high compared to all the other sites, indicating sub-optimal growth condition.

The needles of China fir have relatively high contents of both calcium and aluminum. Yet the Ca/Al ratio in the needles is rather high. Element contents in the needles of Masson pine are quite similar to the levels observed at the other sites. The Ca/Al ratios in the pine needles at Baiyun Shan are all below 12.5. As observed at all other sites, the phosphorus contents in the needles of pine and fir indicate a sub-optimal availability of this nutrient (Table C.3).

### Invertebrates and diatoms

No diatoms or invertebrates were collected in the stream sampled for water chemistry because it probably dries out during dry spells. The more permanent inlet stream to the reservoir was also rejected due to heavy human influence in the catchment. The reservoir holds fish but more detailed information is lacking.

### 3.3.5 Liu Xi River

#### Site description

Nearby Conghua we can find one acid rain monitoring site and one vegetation monitoring site 10 km apart. Besides measuring air temperature, evapotranspiration and tree growth, some soil chemistry data exist (Pb, Cu, Zn, Ni, Cr, Cd, Hg, As, Mo, P and B). The soil in this area is red-soil on granite. There should also be some surface water and other biological data, but we have so far not received such information.

These sites were very difficult to reach because of very bad roads and lack of time. Therefore, we found another site within this area, located some distance from the city Conghua (23°28'N, 113°10'E). The site was close to the main drinking water reservoir of Guangzhou city, the Liu Xi reservoir. The stream in the chosen catchment enters the Liu Xi River a few hundred meters downstream from the outlet of the reservoir/artificial lake. The water was very clear, running directly on the granite bedrock in a very steep catchment. Just above the road, terraces are planted with tea shrubs. Soil samples were collected above this area, on the right hand side of the stream looking upwards. The sampled site had a mixed forest dominated by Masson pine and some bamboo. The soil is sandy and contains many stones. There is a thick layer of litter (grass, leaves) above a thick soil layer of A and AB horizons with many roots. The B horizon contained many large rocks.

#### Air pollution

We did not receive air quality data from this area, but according to Qi and Wang (undated) the pH in precipitation is about 4.7 in the region.

#### Soil

According to the particle size distribution, the soil material is ranging from loamy sand in the A-horizon to sand in the B-horizon, in this respect similar to the soils of Simian Shan. The  $CEC_E$  values are all less than the 15 percentile of our data (Table C.1). This low exchange capacity is likely to be due to the lack of finer material in the soil. Base saturation of the soil ranges from 31%, which is the highest value found among the sampled sites, in the A horizon, to 14% in the AB-horizon. The low  $CEC_E$  and high BS render these soils sensitive to soil acidification, although the effects of a minor acidification may not be serious. Similarly to the soil at Baiyun, the carbon content in the soil at the Liu Xi River site is a few percent in the A and less than 1% in the AB and B horizon. C/N ratios in soil organic matter are low in all soil horizons.

#### Surface water

The electrical conductivity of the stream water was extremely low,  $10.8 \mu\text{S cm}^{-1}$ , but pH was relatively high pH (6.42).  $\text{Na}^+$  is the predominant cation and  $\text{SO}_4^{2-}$  is predominant anion (Table 3.13) Despite high pH, the ANC is not very high in this stream. Accordingly, the streamwater may be sensitive to acidification in spite of fairly high base saturation in the sampled soils. Since granite predominates in these areas, this region is sensitive to increased inputs of acid rain. An important drinking water reservoir for the Guangzhou city is located nearby. It is therefore important to follow the trends in water chemistry by a monitoring program with focus on acidification effects. Chemical/biological monitoring has been carried out for this reservoir, but the parameters analyzed

are not applicable for documentation of water acidification. The stream had the highest concentration of uranium (U) within this survey (Table C.7). However, the concentration is not very high, and not unexpected because granites often contain a certain amount of this element. The highest concentration of molybdenum was also measured in this stream, but also this concentration was relatively low.

**Table 3.13.** Average concentrations of some chemical compounds in a small tributary entering into the Liu Xi River.

Site	pH	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	μeq L <sup>-1</sup>				Al <sub>a</sub>	Al <sub>i</sub>	TOC
						SO <sub>4</sub> <sup>2-</sup>	Cl <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	ANC			
I	6.42	19.5	6.58	49.6	12.8	31.2	14.1	17.1	26	24	4	1.10

### Vegetation

Data from the Liu Xi River site show, just like data from Guangzhou Botanical Garden, particularly low values for the chlorophyll fluorescence measurements on the Masson pine. The median  $F_v/F_M$  ratios of 0.77 and a minimum value of 0.72 indicate sub-optimal growth conditions (Table C.20). The standard deviation is high, but not as high as in Guangzhou Botanical Garden. Data from the measurements on Chinese fir show the opposite: Here the  $F_v/F_M$  ratio is slightly higher than the mean value for the Guangdong province.

Both fir and pine needles at this site have relatively high Ca/Al ratios (Table C.3). Even for pine this ratio is considerably higher than 12.5. These relatively high ratios in the needles correlate well with the relatively high base saturation levels found in the soil.

### Invertebrates and diatoms

The tributary to the Liu Xi River was sampled for diatoms and stream invertebrates. The stream runs in a steep sloping terrain with bare rock and huge boulders forming small pools with whitewaters in-between. The riverbed consists of open rock and large stones and cobbles and no gravel or sand was observed. Despite extensive dipnetting, no invertebrates were found. This can be related to the inhospitable character of this habitat. It is not known if this is a permanent stream carrying water in the dry season. The diatom sample contains a slightly acidophilous (pH 6-7) flora with *Achnanthes minutissima*, *Gomphonema* and a range of *Eunotia* taxa.

### 3.4 Volatile organic compounds in air

#### Introduction

Sampling for volatile organic compound analyses was conducted during the PIAC trip. The analyses and interpretations are made by Arne Semb, Christian Dye and Norbert Schmidbauer (all at NILU).

Emissions of sulfur dioxide and airborne particles have so far been the central air pollution issues in China, while emissions of nitrogen oxides and the photochemical formation of ozone have received little attention so far. This is understandable, since the emissions of nitrogen oxides are still relatively small in China compared to Europe and North America, but the number of motor vehicles is increasing. The building of new coal-fired power plants will also increase the emissions of nitrogen oxides in the next decades. The photochemical production of ozone is initiated by the reaction of hydrocarbons with OH radicals or ozone, and requires nitrogen oxides as catalysts. Hydrocarbons are released from evaporation of solvents and gasoline, and from incomplete combustion. Incomplete combustion occurs in motor vehicles, but also in small domestic fires, burning of litter and firewood, and coal combustion at low temperatures.

While natural hydrocarbons are mostly alkanes, combustion processes tend to remove hydrogen so that alkenes and alkynes are formed. Acetylene is a good indicator of hydrocarbons from combustion processes. Even benzene and other aromatic hydrocarbons may be formed in combustion processes, by condensation of 3 alkyne molecules. However, benzene and other aromatic hydrocarbons are also found in coal tar and are produced in petroleum refineries, as major constituents both in solvents and in motor gasoline.

Measurements of volatile hydrocarbons are useful for characterizing air quality and, particularly, as background information for modeling of photochemical oxidant formation. For this latter purpose, emission surveys of nitrogen oxide emissions are also needed.


Grab samples may conveniently be taken in steel flasks for subsequent analysis and determination of individual hydrocarbons up to C8 by use of a simple membrane pump. Carbonyls, which are the first stable intermediate reaction products, are sampled using 2,4-dinitrophenylhydrazine, and the resulting hydrazones are determined by high performance liquid chromatography (HPLC). Both these techniques have been used successfully in the EMEP program in Europe to provide documentation for the concentration level and emissions of hydrocarbons, and the chemical reactions which determine the ozone formation. The sampling could easily be carried out under field conditions during the PIAC study.

#### Results and discussion

The results of the chemical analyses are given in Appendix (Table C.18 and C.19) and Figure 3.3 and 3.4. Because of high field blanks, the measured concentrations of acetone in some samples are less reliable than the other aldehydes and ketones. The other components are generally well above their detection limits.

For comparison purposes, measurement results at some European background sites are also given. These are Birkenes in South Norway, Waldhof in Germany, and Kosetice in the Czech Republic. Hydrocarbons and carbonyl compounds have been measured at these sites since 1992, and show a well-defined seasonal trend with the highest concentrations of hydrocarbons in the winter, when the





**Figure 3.3.** Measured concentrations of hydrocarbons at sites in China (April/May 1997) compared with monthly mean concentrations for 3 EMEP sites in June 1996.

concentration of hydroxyl radicals is low, and progressively lower concentrations during the spring and summer. Aldehydes and ketones show an opposite trend, with higher concentrations in summer. Since South China is much further south, with correspondingly higher UV radiation levels, the concentrations at European sites in June are used for comparison with the Chinese values for both hydrocarbons and carbonyl compounds.

It is seen that the hydrocarbon concentration levels are generally higher in the flask samples from China, even for the very remote sampling sites Simian Shan and Leishan. This is particularly the case for the relatively reactive unsaturated compounds such as ethene, propene and the butenes. Substantially higher concentrations are seen in the samples from and near the cities. The concentrations of acetylene are particularly high, indicating large emissions from combustion sources, as are also the concentrations of benzene, toluene and xylenes. These hydrocarbons are also indicators of combustion, particularly exhaust emissions from motor vehicles. Since hydrocarbons are broken down primarily by hydroxyl radicals and ozone, their atmospheric residence times vary with both the ozone concentrations and the UV radiation level, as well as with the reaction rate of the compounds with each of these oxidizing species.

The typical emission ratio of acetylene to xylenes is less than 30% (Leggett, 1996) which corresponds to a molar ratio of less than 1:1. This can be used to estimate the "age" of the hydrocarbon mixture. Evidently, the samples from Chongqing and Guiyang-Guizhou are more aged than the samples from the Guangzhou area, which appear to be more influenced by sources within less than a few hours atmospheric transport distance, i.e. a few hundred kilometers.

The concentrations of aldehydes and ketones were low at Simian Shan and in the first sample from Guizhou Botanical Garden, but comparable with the European concentration levels at Leishan, and with the second sample from Guiyang. Much higher concentrations were found at Dinghushan and at Conghua, while the sample from Heshan had low concentrations. The aldehyde concentration levels

are clearly influenced by the weather conditions. While the sky was never quite clear during sampling, the highest concentrations are all from days with some sunshine and the low concentrations are from days with more dense cloud cover. The concentration levels of aldehydes and reactive hydrocarbons indicate that the photochemical activity in this part of China is high, with a high potential for ozone formation.

### 3.5 Intercalibration of water analyses

During our visit to the Dinghushan, two water samples were collected from tap water, taken from a pipeline 1-2 kilometers from the water source, the Tianhu Lake. The bottles collected for South China Institute of Botany SCIB were collected immediately after the bottle for NIVA was collected. Thus, they are not fully comparable. However, the analytical results (Table 3.14) clearly demonstrate the need for intercalibration between different laboratories.

Results from an intercalibration between CIESM and UiO on soil water from the Tie Shan Ping area, indicate fairly good agreement for the chemical parameters having the highest concentrations (Figure 3.4). The remaining parameters showed similar or poorer agreement. This documents a pertinent need for further intercalibration studies. Intercalibration will be a central issue in a future main collaborative project on acid rain monitoring and research in China. This is also a central point in the final report of the Fourth Expert Meeting on Acid Deposition Monitoring Network in East Asia (EMAD, 1997).

**Table 3.14.** Water chemical data of tap water from the Tianhu Lake analyzed by induced coupled plasma-mass spectroscopy (ICP-MS) at NIVA and SCIB.

Element	unit	NIVA	SCIB	Element	unit	NIVA	SCIB
<b>Pb</b>	$\mu\text{g L}^{-1}$	0.67	0.14	<b>Al</b>	$\mu\text{g L}^{-1}$	69.6	12.6
<b>Cd</b>	$\mu\text{g L}^{-1}$	0.37	0.38	<b>Sb</b>	$\mu\text{g L}^{-1}$	0.338	0.330
<b>Cu</b>	$\mu\text{g L}^{-1}$	0.70	12.3	<b>Bi</b>	$\mu\text{g L}^{-1}$	<0.02	<0.02
<b>Zn</b>	$\mu\text{g L}^{-1}$	1904	1700	<b>Tl</b>	$\mu\text{g L}^{-1}$	0.012	0.018
<b>Cr</b>	$\mu\text{g L}^{-1}$	<0.1	0.11	<b>U</b>	$\mu\text{g L}^{-1}$	<0.004	0.001
<b>Ni</b>	$\mu\text{g L}^{-1}$	0.60	0.35	<b>Be</b>	$\mu\text{g L}^{-1}$	<0.01000	0.0050
<b>Co</b>	$\mu\text{g L}^{-1}$	0.30	0.26	<b>Li</b>	$\mu\text{g L}^{-1}$	0.817	0.851
<b>Fe</b>	$\mu\text{g L}^{-1}$	89.1	27.5	<b>Rb</b>	$\mu\text{g L}^{-1}$	1.16	1.56
<b>Mn</b>	$\mu\text{g L}^{-1}$	62.1	95.2	<b>Mg</b>	$\mu\text{g L}^{-1}$	524	644
<b>V</b>	$\mu\text{g L}^{-1}$	<0.3	0.09	<b>Ca</b>	$\mu\text{g L}^{-1}$	2072	2490
<b>Ti</b>	$\mu\text{g L}^{-1}$	7.40	0.47	<b>Mo</b>	$\mu\text{g L}^{-1}$	<0.04	0.008
<b>As</b>	$\mu\text{g L}^{-1}$	0.36	0.24	<b>Y</b>	$\mu\text{g L}^{-1}$	0.01	0.003
<b>Ba</b>	$\mu\text{g L}^{-1}$	7.50	14.7	<b>Sn</b>	$\mu\text{g L}^{-1}$	<0.040	0.028
<b>Sr</b>	$\mu\text{g L}^{-1}$	4.50	8.30	<b>Ga</b>	$\mu\text{g L}^{-1}$	0.249	0.008




Figure 3.4. pH, Al, Ca<sup>2+</sup> and SO<sub>4</sub><sup>2-</sup> in soil water from Tie Shan Ping analyzed at UiO and CIESM.

## 3.6 Data Synthesis

### 3.6.1 Deposition and precipitation chemistry


The deposition of sulfate by precipitation in the visited areas is very high, compared to deposition levels in Europe and North America. There are also very important differences with respect to the average chemical composition of precipitation samples, shown in Figure 3.5. The concentration of base cations, particularly calcium, is very high in China compared to Europe. This is caused by anthropogenic as well as by natural sources. Fly ash from coal-burning is alkaline, and many process industries such as cement manufacturing, lime production, and iron and steel production emit dust which contain calcium carbonate and other acid-neutralizing material. It has been shown that the base cation concentrations in precipitation have been strongly reduced both in Europe and North America (Hedin et al., 1994), presumably as a result of more efficient control of anthropogenic emissions. An analysis by Semb et al. (1995) showed that anthropogenic emissions are still the dominating source for base cations in air and precipitation in Northern Europe, and Saharan dust was only of importance in Southern Europe.

Windblown dust from the arid areas in Central Asia and Northern China contributes strongly to the deposition of alkaline material and base cations in China, so that most of the areas in Northern and Eastern China do not experience acid rain. Chang et al. (1996) have presented model estimates for calcium deposition from this source. In south-western China, where precipitation with  $\text{pH} < 4.5$  does occur, the monthly deposition of calcium from this source is estimated to  $< 10 \text{ mg m}^{-2}$ . However, anthropogenic emissions are considerable, as evidenced by the very high concentrations of airborne particles in the cities of Chongqing and Guiyang. Control of the emissions of particulate matter is relatively inexpensive and highly desirable for improving human health and the quality of living in urban and industrial areas. It is quite likely that new measures will reduce the concentrations of base cations in precipitation in the future also in China.

Quite high concentrations of fluoride ion are also reported for Chinese precipitation samples. Fluoride in precipitation is not routinely monitored in Europe, but concentration levels in Southern Norway are at least 1-2 orders of magnitude lower than in China (Sæther et al., 1993). These data indicate high levels of hydrogenfluoride in air, which may have important phytotoxic implications. Fluorides in air may originate from coal combustion, possibly also from production of brick and tiles.

The concentration level of nitrate ions in precipitation is still relatively low in China. However, emissions of nitrogen oxides have increased substantially during this decade, due both to increased road transport and electric power production, and this trend will continue. In rural areas the concentrations of ammonia are generally high.

Data on airborne concentrations are more scarce, and it is therefore difficult to estimate dry deposition. Since the distances to the emission sources are comparatively short, dry deposition of sulfur dioxide is expected to contribute relatively more to the total deposition than sulfate in precipitation. Experiments with collection of throughfall precipitation seem to confirm that this may be the case, but there is lack of more thorough observations of throughfall and comparison with airborne concentrations and dry deposition estimated by the inference method. Dry deposition of particles and sedimentation of particles during dry periods may also be of interest.



**Figure 3.5.** The bars show volume-weighted average ionic composition in precipitation ( $\mu\text{eq L}^{-1}$ ) for some sites in Norway, The Czech Republic, Poland and China. The left bar represents cations, the right bar anions; the numbers above the bars give the estimated total (wet + dry) deposition of sulfur in  $\text{g S m}^{-2}$ . HUMEX is a nearly pristine site on the Norwegian west coast. The major components in precipitation are from sea salts. Birkenes is in southernmost Norway which is the region of the country most affected by acid rain. Kosetice and Czerniawka are stations in the Czech Republic and southern Poland, respectively. These stations are not much affected by local emissions. However, Czerniawka is in the region in Europe with the highest S-deposition caused by long-range transported pollutants. Ratanica is also in southern Poland, but more affected by local pollution than Czerniawka. LCG stands for Liu Chong Guang (near Guiyang) and TSP stands for Tie Shan Ping (near Chongqing).

### 3.6.2 Soil and soil water

In Sichuan and Guizhou provinces forest soils are largely acidic yellow mountain soils on sandstone bedrock. In the Guangzhou area (Guangdong province) red lateritic soils on sandstone or granite bedrock are more common. Due to the warm climate there is generally no organic horizon. Southern China was not glaciated during the last ice age and the soil material may therefore date as far back as to the tertiary period. Due to long weathering time most of the soils have a relatively high clay content. The main clay mineral is kaolinite, though also various amounts of residual illite and chlorite are found. The C/N ratios are in general very low (Figure 3.6) Especially low values are found in the B horizon at Dinghushan and Baiyun Shan.



Soil base saturation (BS) generally decreases with depth (Figure 3.7). In the B horizon, BS is about half of what is found in the A horizon ( $\text{BS}_{\text{A-hor}} = 0.5\text{BS}_{\text{B-hor}}$ ;  $r^2 = 0.79$  for all sites). The highest BS values are found for Leishan (plots 1 and 2), Liu Xi River and some of the Liu Chong Guang plots. The high values at Leishan are probably due to fresh soil material produced by strong mechanical weathering. At other

sites the higher values in the A-horizon may indicate contribution of basic dust deposition (either of natural or man-made origin) in addition to natural cycling of nutrients.

The relatively low C/N ratios in the Chinese forest soils, combined with soil nitrogen stores that probably exceed 5000 kg ha<sup>-1</sup> (when considering complete profiles), suggest that nitrate leaching may occur. Obviously this strongly depends on the ability of the vegetation and soil biota to assimilate mineralized nitrogen. As has been shown in Europe and North America, destabilization of forest ecosystems due to various stress factors (e.g. drought, acidification, cutting and thinning) may increase nitrogen mineralization and decrease the demand for this nutrient. Consequently, considerable nitrate leaching may occur with possible eutrophication and health hazards. The relatively low C/N ratios in the Chinese soils suggest that forest destabilization may lead to even more pronounced nitrate leaching than reported for temperate and boreal forests in Europe and North America. We have only soil water data from two sites, Tie Shan Ping and Liu Chong Guang; both are strongly affected by acid deposition. The C/N ratios in soils at these sites are relatively high in the A horizon and close to average values in the B horizon (Figure 3.6). The Tie Shan Ping site experiences high concentrations of nitrate in soil water while the concentrations are more variable at the Liu Chong Guang catchment.


As mentioned previously, it has been suggested that molar ratios  $Al/BC^{2+}$  in soil water greater than 1 may cause forest damage (Sverdrup and de Vries, 1994). This assumption is generally used in calculations of critical loads. Although the basis for the limit is weak (Løkke et al., 1996) and it has not been tested on Chinese environments, the values give an indication of possible harmful effects. In Figure 3.8,  $Al/BC^{2+}$  ratios for Tie Shan Ping and Liu Chong Guang are compared to values found at a Polish site where forest damage from pollution is severe. At Liu Chong Guang some values exceed 1, but in general the Chinese values are less than 1, mainly due to high calcium concentrations.

Large variations in soil properties and atmospheric deposition within short distances make assessments of critical loads very difficult. Acidification of soils is clearly not only affected by the S- and N-deposition; the deposition of base cations is equally important. Except at plots 1 and 2 at Leishan, base saturation is low and the aluminum saturation is high (see Fig. 3.7 and Table C.1). This implies that there is a danger for increased acidity and aluminum concentrations in soil water with increasing S- and/or N-deposition. These increases may be transferred to streamwater. However, streamwater is made up of different types of soil water and the sensitivity towards acidification depends strongly on water pathways. From our surveys and the scientific literature it seems that in most places there is sufficient neutralizing capacity, usually in deeper soils and bedrock, to make water bodies relatively resistant towards acidification. Increased concentrations of sulfate will then be accompanied mainly by increased concentrations of calcium and magnesium, generally with no harmful effects. Exceptions are found, usually in small headwater streams.



**Figure 3.6.** *The C/N ratios in soils.*

### A-horizon



### B-horizon





Figure 3.7. Base saturation in soils





**Figure 3.8.** Molar ratios  $Al/BC^{2+}$  in soils. Czerniawka is a site in southern Poland strongly affected by acid deposition, cf. Figure 3.5.

### 3.6.3 Vegetation

One of the tasks of the PIAC project was to assess potential ecological effects on terrestrial plants exposed to different loads of air pollution in selected sites in China. A number of vascular plants (trees, ferns and herbs) were studied to estimate response to photosynthetic activity. Two important and widespread coniferous tree species were selected as representatives for respectively sensitive and non-sensitive species: Masson pine (*Pinus massoniana*) and Chinese fir (*Cunninghamia lanceolata*). For almost all plant species, photosynthetic activity, measured as the  $F_v/F_M$  ratio of chlorophyll fluorescence, seems to increase from Northwest (Sichuan province) to Southeast (Guangdong province). Most of the species show higher photosynthetic activity in the urban areas compared to areas out in the district and up in the mountains, except for the air pollution sensitive Masson pine, which shows the opposite trend (Table C.20). The empirical data show a significant correlation between low  $F_v/F_M$  ratio of chlorophyll fluorescence, and high standard deviations. In general the standard deviation is higher for Masson pine than for Chinese fir and all other species measured. This may indicate that measurements of sample variance and dispersion are suitable for detecting biotic responses to different stressors, i.e. air pollutants and acidified soils.

The stand near Conghua (Liu Xi River) is the only one in our investigation where the molar Ca/Al ratios in pine needles are well above 12.5, the threshold level below which it has been suggested that there is a 50% risk for aluminum damage (Cronan and Grigal, 1995). However, we do not know if this threshold value applies to Masson pine. In all other Masson pine stands, molar Ca/Al ratios in needles are close to or even below this threshold value of 12.5. In the less sensitive Chinese fir there is relatively little accumulation of aluminum in needles and the molar Ca/Al ratios are high, suggesting that this tree has effective means of avoiding uptake of toxic aluminum. Both Chinese fir and Masson pine needles are low in phosphorus at all sites. Possibly phosphorus stores in the soils are low due to centuries-long intensive use of the forests (collection of litter, herbs, mushrooms, cutting of wood). In addition, mobilization of aluminum in the soil due to acidification may have made soil phosphorus less soluble.

The C/N ratio in soils may be used as an indicator of the availability of nitrogen for plants. Generally a lower C/N ratio indicates a higher nitrogen availability. The C/N ratios observed in the investigated soils

in China are extremely low compared to the values commonly found in forest soils of the temperate and boreal regions. Considering all sites, the nitrogen contents in the needles of Chinese fir and Masson pine range from 1.0% to 1.6%. These levels suggest a reasonably good supply with nitrogen. Our data on soils and vegetation suggest therefore that phosphorus rather than nitrogen is growth limiting for both Chinese fir and Masson pine.

Vegetation damage has mainly been found for Masson pine. The generally low Ca/Al ratios in needles may indicate that soil conditions are of some importance and there may be a lack of available phosphorus. At the most affected sites, the SO<sub>2</sub> concentrations are so high that direct effects may be expected. In the future, ozone is likely to become an increasingly important factor in connection with vegetation damage. The main dangers to ecosystems in China due to acid precipitation and its precursors seem therefore to be increased aluminum concentrations in soil water with accompanying effects on the vegetation and direct effects of harmful gases, in particular SO<sub>2</sub> and ozone.

### 3.6.4 Surface water

The acid neutralizing capacities (ANC) of surface waters were highly variable (Figure 3.9), reflected by large variations in water pH. The streams in Tie Shan Ping and in Liu Chong Guang were all significantly affected by acid precipitation, primarily by sulfuric acid. They do not represent the acidification status in larger areas of these provinces. An illustration of this is the stream very near Liu Chong Guang with high pH and ANC. The very acid surface water draining a Monsoon Evergreen Broadleaf Forest in Dinghushan in the Guangdong province is probably a combination of acid precipitation and specific catchment properties in this subtropic forest. Interesting areas investigated were the Leishan Mountains and Liu Xi River area. The ionic concentrations in these waters were extremely low, but because these areas are almost unaffected by acid rain, at present, the surface water is not acidified. However, important aspects of the air pollution abatement strategy in China are to reduce emissions of particles (which may be rich in base cations) and to build higher stacks for the main air pollution sources. Therefore, the acid rain problem may increase in areas further away from the emission sources. Even though available results show that water acidification is not likely to become a serious regional problem in China, one should be aware of possible future acidification in some very sensitive areas like Leishan Mountains and the Liu Xi River area. Only large regional surveys can document the extent of surface water acidification in China.

As illustrated in Figure 3.10, a significant increase in the concentration of Al<sub>i</sub> occurs when pH drops below 5.5. It is well documented in Europe and USA that this Al fraction includes the main toxic compounds in acidified surface waters. Also the concentrations of many heavy metals tend to be high at low pH, as illustrated for lead (Pb) in Figure 3.11. For heavy metals the concentrations are often a combination of increase in atmospheric inputs as a direct consequence of high air pollution, and increased mobilization of metals as the acid inputs contribute to soil and surface water acidification.

The surface water analyzes also revealed extremely high concentration of zinc (Zn) in the tap water from the Lake Tianhu in the Dinghushan area, and very high concentration of tin (Sn) in one stream in the Leishan Mountains. Since Lake Tianhu is used as drinking water for the local people and a factory bottling fresh water is located near the Simian Shan stream, these sites should be carefully followed up very soon.

The parallel analyses of surface water from the Lake Tianhu conducted at NIVA and SCIB, and of soil water from Tie Shan Ping at UiO and CIESM, illustrate the need for further intercalibration of methods.




Figure 3.9. The acid neutralizing capacity (ANC) in the different surface waters analyzed.




Figure 3.10. The relationship between pH and Al<sub>i</sub> in the surface waters analyzed.




Figure 3.11. The relationship between pH and lead (Pb) in the surface waters analyzed.


### 3.6.5 Aquatic invertebrates

Terrestrial and particularly freshwater ecosystems in Europe and North America are currently being impacted by anthropogenic acidification. One task of the PIAC project was to assess potential ecological effects on soil invertebrates and freshwater fish and invertebrates at selected sites in China with different atmospheric pollution. Measurements of physiological and biochemical indicators of the level of stress in individual specimens, as proposed in the project description, have not been performed. The study sites were small first order streams devoid of fish, crayfish, snails and other larger forms of stream invertebrates. In addition test organisms from adjacent catchment soils (earthworms, snails and woodlice) were extremely scarce and difficult to collect in adequate sizes and numbers suitable for such measurements. Instead, it was decided to perform qualitative sampling of smaller invertebrates and diatom algae in the permanent streams. Both groups are extensively used for monitoring and as indicators of surface water acidification through shifts in community composition. The samples have been analyzed for major taxonomic groups and in some cases down to the generic and species levels (diatoms). The invertebrates are analyzed by NINA and the diatoms by Environmental Change Research Center (ECRC), UK. These two data sets have been used to assess the more general relationship between stream fauna and diatom flora composition and water chemistry.


It has not been possible at this stage to determine which species are present in our samples. This requires more in depth analysis and contacts with Chinese taxonomic expertise. However, we may offer the following preliminary conclusions (see also Figure 3.12):

1. The stream fauna samples from the acid streams appear to have fewer taxonomic groups and specimens than the circumneutral streams. Absence of invertebrates in some non-acid streams is presumably due to factors associated with local land use and/or poor habitat quality.


A



B



C



**Figure 3.12.** Data from invertebrates collected at PIAC sites in China. **A:** The relationship between the number of taxa identified in individual samples vs. stream pH; **B:** : The relationship between the number of collected specimens in individual samples vs. stream pH; **C:** The relationship between number of registered taxa and the number of specimens occurring in individual samples.

2. The preliminary analysis of the diatom samples shows that there is an excellent correspondence between the approximate acidity of the water inferred from the diatoms and the measured pH. This indicates that biotic changes in some of the investigated freshwater communities can be attributed to the current level of water acidity.

3. If resources become available to support further work to refine the identification of the organisms and to increase the number of samples, we can develop powerful, quantitative relationships between freshwater communities and water chemistry that can be used for sophisticated environmental analysis and monitoring.

## 4. Other information related to acid rain in China

### 4.1 The Chinese-Norwegian seminar

A most valuable start of our work in China was an invitation to a joint seminar on acid rain. It was arranged by NEPA on April 15. This Chinese/Norwegian seminar represented the first bilateral arrangement of NEPA on acid rain, and the Norwegian team was the first group of scientists to be invited by NEPA to work in this field. The seminar gave an opportunity to share information on the state of knowledge on acid rain research in both countries, and to discuss the work program of the Norwegian team. The contacts made at the seminar helped to strengthen the basis for future cooperation between Norwegian and Chinese scientists.

The seminar demonstrated that Chinese authorities are giving increasing attention to acid rain problems in China and that the scientific understanding of the impacts of acid rain is growing.

On April 14, the Norwegian research group visited CRAES and met with senior managers and researchers of the academy. CRAES is a multidisciplinary research institution under NEPA. It was established in 1979 to conduct applied research on a wide range of environmental issues.

### 4.2 The RAINS-Asia model

The Asian version of atmospheric dispersion and deposition module of RAINS-Asia (**R**egional **A**ir **p**ollution **I**nformation and **S**imulation model) has been reviewed by D. M. Whelpdale and H. Ueda (in Dovland, 1995). The model builds on experience from Europe and North America, and has been modified to correspond to Asian conditions. The strength of this type of model is the mass balance, the deposition of sulfur is closely related to emissions, and the atmospheric lifetimes will typically correspond to average transport distances of a few hundred kilometers.

The major uncertainties involved in the model are spatial resolution, correct atmospheric chemistry with respect to the transformation of sulfur dioxide to sulfate aerosol, and deposition processes. Spatial resolution is particularly important in areas which are affected by emission sources within the nearest 100 kilometers. This is certainly the case for the areas most heavily impacted by acid precipitation in China. In these areas, model predictions in the RAINS-Asia grid can only give an order-of-magnitude prediction of the annual deposition to a small watershed or research plot.

The transformation of sulfur dioxide to sulfate aerosol follows two pathways. One is by the gas-phase reaction with hydroxyl radicals (Calvert and Stockwell, 1984), the other involves absorption of sulfur dioxide into cloud droplets or deliquescent aerosols and oxidation by hydrogen peroxide or ozone. Ammonia may significantly increase the conversion rate in the latter case if hydrogen peroxide is not available. In the RAINS-Asia model, higher OH levels have been assumed than in the original model because of more UV radiation, resulting in more rapid transformation to sulfate aerosol than in Europe.

Limited measurements are available for validation of the models. Sulfur dioxide has been measured by passive samplers, and precipitation samples from a number of monitoring sites have been analyzed (Wang et al., 1997b). However, since many of the sampling sites for precipitation were located in cities and suburbs, it is difficult to obtain a coherent picture of the deposition pattern. Dry deposition estimates by the inferential method (e.g. Hicks et al., 1987) also require knowledge of the vegetation cover and photosynthetic activity in relation to ambient concentration levels.

In the report from the Hiroshima meeting (EMAD, 1997), the Chinese representative stated that China did not have a proper monitoring network to satisfy the needs of the Acid Deposition Monitoring Network in East Asia. This is in accordance with our impressions, with some modifications. During our visit, we have seen that scientists and monitoring agencies have considerable experience and knowledge, and that the quality of the instruments and equipment varies. The measurements have either been short-term research activities with external funding, or are part of municipal monitoring with sampling in the cities. Measurements at remote sites are scarce, because of the costs involved in shipment of samples, lack of electricity and difficulties in finding qualified attendants in the rural districts. Some programs have been conducted in remote areas and in natural reserves which have been developed for tourist activities and which are also used as field stations for scientific research. Examples of such sites are Emei Mountains and Fanjin Mountains, which both are high-elevation sites. Even if the height of the mixing layer is higher in South China than in Europe, measurements at sites which are above 1500 m a.s.l. will generally give concentrations of pollutants in air and precipitation which are smaller than at lower altitudes. This also applies to the mountain of Leishan in Guizhou, which was visited by our team.


Critical loads have been computed and mapped in China (Hettelingh et al., 1995). Critical loads are part of the impact module of the RAINS-Asia model. A critical load map has been derived based on 31 vegetation types, taking into account information on soil types and climatic and meteorological conditions which are specific to the Asian region. The ecosystems incorporated ranged from tropical rain forests to near desert conditions and from mangroves to areas of taiga with associated soil types. Variations in land use have been accounted for. Therefore, extensive validation procedures are required to improve the assessment of the risk of damage on more local scales. To further improve the model, better input data are required. In the PIAC project, we have contributed with data that may be used for this purpose. Further work is also required to include other causes of damage in addition to acid deposition. SO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub> in air have been shown to cause direct damage to natural ecosystems and crops, as well as to materials and health, in large urban areas.

### **4.3 Management and scientific relationships**

There are several scientific institutions and environmental agencies/commissions working with pollution problems in China. The central units collaborating with the Norwegian PIAC-team are presented in Figure 4.1.

The PIAC-team was invited to visit China by the National Environmental Protection Agency (NEPA) of China. NEPA was formally established as a directorate as late as in 1988 and has its head office in Beijing with about 300 employees. NEPA is the key governmental body in China with branches at province level and bureaus in the larger cities, among others the cities we visited. The provincial and local units are called EPB's and are directly under the local government. There are 2 300 local EPB's in China, with 88 000 people employed. NEPA and EPB also have Environmental Monitoring Centers (NEMC or EMC), but they are not directly under NEPA (Figure 4.1), but relatively independent units, which is also the case for the Chinese Research Academy of Environmental Sciences (CRAES).





**Figure 4.1.** Organization map of central scientific institutions and environmental agencies in China. *NEPA: National Environmental Protection Agency; BCD: Bilateral Co-operation Division, Dept. of International Co-operation; APCD: Atmospheric Pollution Control Division, Dept. of Pollution Control; STD: Science and Technology Division, Dept. of Science, Technology and Standards; CRAES: China Research Academy of Environmental Sciences; NEMC: National Environmental Monitoring Center with a network division; EPB: Environmental Protection Bureau; EMC: Environmental Monitoring Center; CAS: Chinese Academy of Sciences; RCEES: Research Center of Eco-Environmental Sciences; NRC: Nature Resources Center; SCIB: South China Institute of Botany.*

Among NEPA's responsibilities are drafting environmental laws, rules, regulations and guidelines. NEPA prepares national environmental plans and is responsible for monitoring and control of emissions. Furthermore, NEPA supervises environmental impact assessments of large projects and has responsibility for environmental public relations and education. NEPA is not linked to a separate ministry, but is under the Environmental Protection Commission (EPC), which incorporates leaders from the state ministries and central governmental organs. EPC meets quarterly to assess the national environmental policy. NEPA is the secretariat of EPC.

In November 1995 a collaboration agreement between The Norwegian Ministry of Environment and NEPA was established. The first working program should start, hopefully in 1998, and will consist of 8-10 projects, with a total economic frame of 5 mill NOK pr. year.

Professor Tang Dagang was our contact person in CRAES and was responsible for all arrangements regarding our travel and work in China. Tang is director of the Atmospheric Institute at CRAES. This institute has a professional staff of more than 40 and consists of six divisions: Atmospheric Physics; Atmospheric Chemistry; Global Climate Change; Wind Tunnel; Aerosols; and Air Pollution Control.

The Chinese Academy of Sciences (CAS) is organized in central and local units. The universities are under the CAS-organization. Some places the local CAS might be directly under the local government but totally independent of NEPA/EPB. A large number scientists with great experience in environmental studies are within the CAS-system.

Based on our experience, the collaboration between local, provincial and central CAS units and corresponding EPB's needs to be significantly improved. Accordingly it is an important challenge in a future collaborative monitoring and research project to improve these relationships. The positive contacts we established within both the CAS and NEPA/EPB family may put us in a very important in-between position in order to improve this climate. It is highly important that the information of data flows smoothly between the local scientists running the local monitoring sites, and the local, provincial and central Environmental Agencies, which also need the same data in their policy planning. To obtain this, it is important that the scientists generating the data also get the possibility to publish their results. If not, the data flow from scientists to policy makers will be far more difficult.

Regarding future projects, the economic contribution from the Chinese counterparts has to come from the NEPA-system, since CAS has no money for such initiatives. On the other hand, having research cooperation with China without involving leading scientists from CAS will mean a great loss of scientific knowledge. Thus, a joint Chinese-Norwegian project on acid rain and its effects in China has to involve both CAS and NEPA.

#### **4.4 Acid rain research activity in China**

CRAES plays an important role in the national acid rain projects under implementation in China. These projects are included in the National Ninth Five-Year Plan for Environmental Protection and the Long-Term Targets for the Year 2010 (NEPA, 1997). According to the plan the amount of sulfur dioxide emissions will be reduced from 22.3 mill. tons in 1995 to 22.0 mill. tons in the year 2000, a reduction of 1.5%. Of a total of 1 591 environmental projects to be implemented in the five year period, 109 are related to acid rain pollution control. 36 of these are located in Central China, 28 in South, 36 in Southwest and 9 in East China. In addition, there will be 209 projects on control of air pollution in major cities.

Total investments in prevention and control of air pollution amount to 208 billion Yuan and represents 46% of all environmental protection investments under the five year plan 1995-2000.

The basic principle is that *total amount control* shall be enforced for emission of sulfur dioxide. The key objective is to restrict the discharge amount of sulfur dioxide back to the level at the end of the Eighth Five-Year Plan period (i.e. 1995). After a national overall balance is worked out, the permissible discharge amount of pollutants during the plan period will be distributed among the provinces, autonomous regions and municipalities, directly under the control of Central government. The provincial-level authorities are responsible for the implementation of the plan. Annual plans will be worked out and annual inspections and examination will be conducted. The progress will be made public on a regular basis.

Two key principles are important when implementing the environmental projects; i.e. the polluter pays and that input of funds shall mainly come from enterprises and local authorities.

The desired results of the reduction strategy in major cities within the acid rain- and the SO<sub>2</sub> pollution control regions are as follows: 1) Flue gas desulfurization in power plants generating 9.3 million kW; 2) A coal washing and blending capacity of 24 million tons a year; 3) A gas supply capacity of 1.41 million cubic meter a day; 4) An additional central heating area of 240 million square meter; 5) Smoke abatement and dust removal equipment in boilers generating 13.000 tons of steam shall be renovated. This will lead to an annual reduction of emissions in sulfur by 1.8 million tons and in soot by 1.5 million tons.

## 4.5 Conclusions from the Forth Expert Meeting on Acid Deposition in East Asia.

The conclusions from the PIAC-project are similar to the conclusions drawn in the final report of the Fourth Expert Meeting on Acid Deposition Monitoring Network in East Asia (EMAD, 1997) in Hiroshima, February 1997. This meeting concluded that acid rain in China is a large and difficult task and it is necessary to further optimize the acid rain monitoring network. They further concluded that at present, in terms of acid rain monitoring, there is a gap between the practical situation and the State's needs, and that monitoring technology in China is inferior compared to that in many developed countries. This was indicated by the following points mentioned in the report from the meeting:

- Better technological guidelines for acid rain monitoring are needed and more comprehensive information should be obtained with reasonable financial input.
- Too few monitoring locations for acid rain relative to the large regional variance in acid rain. There is especially a lack of data from background stations. As a result, information is not sufficient to evaluate the air quality in large regions.
- Too few parameters are monitored. All major anions and cations have to be analyzed.
- Quite weak quality control of acid rain, and the applied technology is not up to date.
- Unsatisfactory financial input and instruments for acid rain monitoring. Except for very few cities, the sampling for acid rain monitoring is manual.

The Chinese Government is attaching greater importance to environmental protection and more financial input is expected in the near future. The general target for modernized acid rain monitoring in China is to monitor dry fall out and wet precipitation by using advanced and uniform technologies and sampling means. This will enable scientists to collect comparable data. On this basis accurate time trends may be obtained.

With regard to the East-Asia Acid Rain Monitoring Network, NEPA supports this initiative and is willing to participate and to take a leading role. NEPA suggests that the establishment of the Network and its activities should follow the principle of *conducting step by step* and *gradually developing*. Specific suggestions from NEPA include:

- The cities proposed to be in the Network should be representative and be identified according to their specific situation. The monitoring station should have sufficient expertise.
- The national centers should be chosen among the participators of the Network. The national centers are responsible for technical management and data integration. At present, the activities within the Network should focus on information exchange.
- In the future, uniform guidelines for sampling, sampling frequency and analyses should be required to produce comparable data, provided that sufficient funds are guaranteed.

Some information about essential ongoing research and countermeasures towards acid rain in China is presented in Chapter 4.4.

## 5. Main conclusions and recommendations

### 5.1 Main findings


This report is based on a study tour by a team of Norwegian scientists and Dr. Tang Dagang, Director of Atmospheric Environment Institute at the China Research Academy of Environmental Science (CRAES), Beijing. A map showing the locations visited is presented in Figure 5.1. The group visited a number of sites in China affected by, or expected to be sensitive to, acid precipitation. Such areas are mostly in the southern part of the country. In most other regions soils and water are not likely to be affected by acid deposition although direct effects of pollutants on flora and fauna may occur. During the trip, field measurements of surface water pH and net photosynthesis (mainly of Masson pine and Chinese fir) were carried out. Samples of needles, soils and water were collected and later analyzed. Our results are supplemented with information obtained from Chinese scientists and literature studies as well as some results from ongoing Chinese-Norwegian cooperation projects.

Many studies have documented high concentrations of sulfur dioxide (SO<sub>2</sub>) and airborne particles in air as well as low pH in precipitation in Chinese cities and nearby areas. This is largely because of the widespread use of coal with high ash and sulfur contents, inefficient combustion and low release heights of the combustion products. The acidity of the precipitation and the ecological effects are largely governed by emissions of SO<sub>2</sub>, basic particles and (in rural areas) of ammonia (NH<sub>3</sub>).

Efforts being made to improve the air quality include introduction of cleaner fuels, such as LPG (Liquid Petroleum Gas) for cooking purposes, mixing of limestone with coal in coal briquettes, and control measures in polluting industries. This has already improved the air quality in some cities with respect to SO<sub>2</sub> and particles. Increased demand for electricity will lead to increased emissions of SO<sub>2</sub> from power plants unless strict mitigation measures are introduced in new as well as old plants. Yet the intention of the Chinese National Ninth Five-Year Plan for Environmental Protection and the Long-Term Targets for the Year 2010 is to reduce the SO<sub>2</sub> emissions from 22.3 million tons in 1995 to 22.0 million tons in the year 2000, a reduction of 1.5 per cent. During the Ninth Five-Year Plan, 109 projects related to acid rain pollution control will be implemented, most of these are located in central or southern China. In addition, there will be 209 projects on control of air pollution in major cities. Total investments in prevention and control of air pollution amount to 208 000 million Yuan, representing 46% of all environmental protection investments under the five year plan 1995-2000.

In contrast to the reduction plans for SO<sub>2</sub> and particles, emissions of nitrogen oxides (NO<sub>x</sub>) and the photochemical formation of ozone (O<sub>3</sub>) have so far received little attention in China. The emissions of NO<sub>x</sub> are still relatively low in China compared to Europe and North America, but the number of motor vehicles is dramatically increasing. The building of new coal-fired power plants will also increase the emissions of NO<sub>x</sub> in the next decades. The photochemical production of O<sub>3</sub> is initiated by the reaction of hydrocarbons with OH radicals or O<sub>3</sub>, and requires nitrogen oxides as catalysts. The concentration levels of reactive hydrocarbons and carbonyl compounds have been determined at sites in the Chongqing region and in the Guizhou and Guangdong provinces during the PIAC journey. The measurements show that volatile organic carbon (VOC) concentration levels are high, so that emissions of NO<sub>x</sub> are probably the controlling factor of O<sub>3</sub> formation. Photochemical reactivity is high, as demonstrated by the high concentrations of carbonyl compounds. This indicates high OH concentration levels leading to more rapid conversion of SO<sub>2</sub> to sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) and sulfate (SO<sub>4</sub><sup>2-</sup>), compared to European conditions. The relative high concentration levels of the less reactive hydrocarbons, such as acetylene and benzene, show that even the remote sites are affected by the high regional emissions.

## The PIAC project



**Figure 5.1.** Map of China showing the main sites visited in the Sichuan, Guizhou and Guangdong provinces.

Our data as well as already published material document that air pollution is a severe problem in China and ecological degradation already occurs as a consequence of air pollution and acidification. The deposition of acidifying compounds in the most polluted areas is much higher than in Central and Eastern Europe, regions considered to have very severe pollution problems. It is clear that some plant species including coniferous trees are affected in the most air-polluted areas. This is confirmed by our measurements of photosynthesis. Although the Al/Ca ratios are high and phosphorous (P) content low in needles of Masson pine in polluted areas, it is not possible to assess at present if the damage is due to aluminum in soil water, to reduced availability of phosphorus or to direct effects of SO<sub>2</sub> and/or O<sub>3</sub>. Other factors, e.g. nutrient depletion due to intensive use of forests, may also have negative impacts on plants. The C/N ratios in needles are low, indicating that nitrogen is not a limiting factor. We emphasize that the data base is too small and the quality of data too variable to assess the extent of damage due to air pollution on Chinese ecosystems.

**Table 5.1.** Estimates of atmospheric inputs, catchment sensitivity to acidification and present acidification status on different levels in the ecosystems investigated. H: high; M: Medium, L: Low

Site	Air		Catchment sensitivity to acidification			Net effects of atmospheric inputs on ecosystems at present				
	Acid <sup>1</sup> input	Alkaline input	Soil water <sup>2</sup>	Vegetation	Surface water	Soil water <sup>2</sup>	Vegetation	Surface water	Diatom algae	Aquatic invertebrates
Nanshan Tie Shan	H	H	H	H		H	H	no water	no water	no water
Ping Simian Shan	H	M	M	H	M	H	M/H	H	H	H
Liu Chong Guang <sup>3</sup>	M	L	H	H	L	M	L	L	L	not found
Leishan	H	M/H	M	H	M	H	M/H	H	H	H
Dinghushan <sup>3</sup>	L	L	L/M	M	H	L	L	L	L	L
Heshan Baiyun Shan	M	L/M	H	M	H	M	L/M	M	M	M
Liu Xi River	M	L/M	H	H	L	M	L	L	not sampled	not sampled
	M	M/H	H	H	L	M	H	L	not sampled	not sampled
	L/M	L	M	M	H	L/M	M	L	L	not found

<sup>1</sup> The input of acidifying substances is considered, essentially (sulfate + nitrate - calcium) in precipitation + dry deposition, essentially SO<sub>2</sub>.

<sup>2</sup> We consider changes with respect to soil water rather than soil properties. Most of the soils have low base saturation (BS), implying that only severe acid deposition is able to further reduced BS. However, increased acid input will affect the composition of soil water, in particular increase the aluminum concentrations. In terms of changes in BS (i.e. soil acidification), soils with intermediate BS (20 - 40 %) and low CEC<sub>E</sub> are most sensitive, but minor reductions in BS are not likely to have serious ecological effects.

<sup>3</sup> Liu Chong Guang refers to the monitored catchment, while a neighboring catchment showed much higher buffering capacity, based on surface water measurements. This indicates large local differences in geological properties. Estimates in Dinghushan is also very uncertain, because soil and surface water indicate a relatively sensitive system with acid, Al-rich water, even though the inputs of acidic compounds are rather low, based on the data we have.

In the areas investigated in this study, the top- and sub-soils are generally quite acid with low base saturation. Deeper soils and bedrock seem in most cases to have greater neutralizing capacity judging from water chemistry. The ratios of organic carbon/organic nitrogen are low in accordance with the ratios found in needles. The main problems, at present and in the future, are related to soils and soil waters with possible consequences for the vegetation. Soil waters have been studied only at two heavily polluted sites. The aluminum/(calcium + magnesium) molar ratios are in some soil water samples greater or close to 1, a limit often used in critical load estimates. Surface water acidification is not likely to become a severe regional problem. However, there is very limited information on variations in streamwater acidity with flow. In some areas the percolating water may not get into sufficient contact with neutralizing material during storm flow events. Furthermore, our survey documented relatively large rural mountainous areas where the surface waters had very low acid buffering capacity. With the strategy of building higher stacks as well as eliminating the particle emissions, one should be aware of possible future acidification in some very sensitive areas which at present do not receive high amounts of acid rain. Increased emissions of nitrogen compounds may also cause a gradual eutrophication of surface water. Thus, monitoring of surface water should be incorporated in future acid rain monitoring programs in China.

There are very few field studies on soil and surface water organisms documenting ecological effects of high acidity in China. Some of our observations resemble those of acidified systems in Norway, but there are several important differences. Assessments of current status and future biotic changes based on scientific studies in sensitive soils and freshwaters are needed.

Attempts to assess the inputs of acid and alkaline compounds (A), catchment sensitivity towards acidification (B) and net effects of A and B at the different ecosystems investigated are presented in Table 5.1. The net effects are an "at present estimate". More detailed discussions of our findings are given in the Chapter 3.6 - Data Synthesis. The PIAC-project has also made a status report about acidification in China, based on international and Chinese literature. This report, which will be published shortly in *Environmental Science*, is also presented in Appendix A.

## **5.2 A framework for further cooperation**

One major goal in the PIAC project was to propose a frame for a monitoring and research program on acid rain in affected areas in China, i.e. the Chongqing, Guiyang and Guangzhou areas. The framework is presented in Appendix D, and is already presented to our Chinese counterparts, in order to have them formulate a request for cooperation. We expect to receive their reactions early 1998.

According to the framework presented, the objective of the main project should be:

*To obtain a reliable picture of the extent and severity of effects of acid precipitation on Chinese ecosystems and to give predictions of their future development under different scenarios.*

In order to meet this objective, the following has to be done:

1. Establish a network of Chinese institutions in cooperation with institutions from Norway and other countries, in order to develop best possible basis for research and monitoring of acid precipitation and the effects.
2. Establish a monitoring program for determining the extent, severity and trends of acid precipitation.

3. Improve the knowledge of processes determining the impacts of acid deposition on ecosystems.
4. Use the obtained knowledge to improve acidification models that are suitable for predictions of future effects (e.g. the RAINS-Asia model).
5. Cooperate with decision makers in applying the knowledge to make cost-efficient countermeasures.

To be as useful as possible for decision makers, one should combine determination of physical, chemical and biological effects *with estimates of economic values* of effects on the ecosystems.

Our survey and contact with Chinese scientists and people in local, provincial and central Environmental Agencies reveal large variations in knowledge within acidification research. We also found infrastructure problems and large differences with respect to scientific equipment, spare parts and the scientific quality of monitoring and research. During the establishment and implementation of a monitoring program in the different areas particular consideration will be given to these differences. Intercalibration of techniques should be an important component of the project.

Many conclusions from the PIAC-project are similar to those drawn in the final report from the Fourth Expert Meeting on Acid Deposition Monitoring Network in East Asia (EMAD, 1997). This meeting concluded that acid rain is an important environmental problem in China and that it is necessary to improve the acid rain monitoring network with respect to data quantity and quality. The monitoring technology in China is not state of the art and needs further improvements. Lack of data from background stations was also criticized. Our survey documented that relatively large rural regions with low acid neutralizing capacity exist both in the Guizhou and Guangdong provinces. According to our data, these acid sensitive areas are little affected by air pollution at present. Thus, these sites may serve as background monitoring stations. To what extent these areas will be affected by acid rain in the years to come is difficult to assess at present, but they are located relatively close to large urban areas. Accordingly, a monitoring program should be initiated relatively soon in these areas, prior to the anticipated increase in air pollution. The PIAC-survey has also resulted in enough knowledge to locate other suitable monitoring/research sites, besides those already running. The selected locations should include sites receiving both high and low atmospheric inputs of acidifying air pollution as well as high and low resistance towards inputs of these pollutants. NEPA suggests that the establishment of the Network and its activities should follow the principle of "conducting step by step" and "gradually developing", and has several special suggestions for a proper strategy, see Chapter 4.5.

We recommend a careful monitoring of soil, soil water, vegetation and soil invertebrates, as well as streamwater and aquatic invertebrates where permanent surface water is present. Monitoring at other ecosystem levels than air quality and deposition is also in accordance with the Chinese policy. EMAD (1997) stated that "improvements of air quality are highly important not only for the human health, but also for the eco-environment" in China. Our program proposal also incorporates research activities of importance for dose-response assessments at different ecosystem levels. Such documentation is absolutely necessary for emission reduction plans or other countermeasures to be effectuated. The proposed framework for a monitoring and research program in China is presented in Appendix D, ("Elements in a framework for further cooperation"). Data from corresponding monitoring programs in Europe have been used in RAINS-Europe to successfully assess abatement strategies of sulfur emissions in this region. High quality data, that may be obtained through a comprehensive research and monitoring program, are of great importance, not only for China but for the whole Southeastern Asia. Based on such data it should be possible to make relevant estimates for critical loads on different ecosystems in China. At present the scientific basis for the critical loads is generally weak,



and for Asian ecosystems the situation is even worse due to lack of data. A thorough evaluation of the concept seems necessary.

The Chinese Government is paying more and more attention to environmental protection and increased financial input is expected in the near future. A collaborative Chinese-Norwegian monitoring and research program on acid rain, supported by NORAD and/or the WB, may help the Chinese Government to fulfill their ambitious intentions. The Norwegian PIAC-institutions will contribute with their knowledge and experience from corresponding programs in Europe in order to implement a high quality program in accordance with the strategy made by NEPA.

We expect our Chinese counterparts, coordinated by NEPA/CRAES, to develop their proposal for a monitoring, research and training program to be sent to NORAD and the World Bank, hopefully early in 1998. This proposal will be a collaborative product between local and central Chinese scientific institutions and NEPA/EPB's. We will recommend a proposal to NORAD for economic support to the establishment and daily running of selected monitoring sites.

We will further recommend another proposal to WB-Asten for economic support for the implementation of regional surveys. Data from the regional survey are highly needed for improving and validating the RAINS-Asia model. It should be an important issue for the WB to help improve the RAINS-Asia model so it may serve as a proper tool for assessing abatement strategies for emissions of air pollutants within China, and provide a basis for negotiations between countries in the area on this issue. The proposal for economic support to a monitoring/research program most likely will have a time frame of 5 years.

## 6. References

- Aunan, K., Seip, H.M. Aaheim, H.A., 1995. A model framework for ranking of measures to reduce air pollution with a focus on damage assessment. In: I. Hargittai and T. Vidóczy (eds.) *Combustion Efficiency and Air Quality*, Plenum Press, 269-283.
- Bian, Y.-M., Yu, S.-W., 1992. Forest decline on Nanshan, China. *For. Ecol. Managem.*, 51, 53-59.
- Blank, L.W., 1985. A new type of forest decline in Germany. *Nature*, 314, 311-314.
- Blank, L.W., Roberts, T.M., Skeffington, R.A., 1988. New perspective on forest decline. *Nature*, 336, 27-30.
- Björkman, O. and Demming, B. 1987. Photon yield of O<sub>2</sub> evolution and chlorophyll fluorescence characteristics at 77K among vascular plants of diverse origin. *Planta*, 170, 489-504.
- Byrne, J., Shen, B., Li, X., 1996. The challenge of sustainability. Balancing China's energy, economic and environmental goals. *Energy Policy*, 24, 455-462.
- Calvert, J. G., and Stockwell, W.R., 1984. Mechanism and rates of the gas-phase oxidations of sulfur oxides and nitrogen oxides in the atmosphere. In: Calvert, J.G.(ed.): SO<sub>2</sub>, NO and NO<sub>2</sub> Oxidation Mechanisms: Atmospheric Considerations. *Acid Precipitation Series*, Vol. 3. Boston, Butterworth, 1-62.
- Carr, M.R., 1990. Practical notes for using the computer programmes prepared for the training workshop on statistical treatment and interpretation of marine community data. - Plymouth Marine Laboratory, Plymouth.
- Chang Young-Soo, Arndt, R.L. and Carmichael, G.R., 1996. Mineral base-cation deposition in Asia. *Atmos. Environm.*, 30, 2417-2427.
- Clarke, B., 1991. Lecture notes for training workshop on multivariate analysis of benthic community data. - Univ. Oslo, Oslo.
- Cronan, C. S. and Grigal, D. F. , 1995. Use of calcium aluminum ratios as indicators of stress in forest ecosystems. *J. Environm. Qual.*, 24, 209-226.
- Derwent, R.G., 1993. Hydrocarbons in the Atmosphere: their sources, distributions and fates. *Conference Proc. Volatile Organic Compounds in the Environment*, London, 27-28 October 1993. Indoor Air International, London, 19-25.
- Dovland, H., 1995. Report on the International Peer Review of RAINS-ASIA (H. Dovland, ed.). Kjeller, Norwegian Institute for Air Research (NILU OR 49/95).
- EMAD, 1997. Acid deposition. Monitoring Network in East Asia. Final Report of the Fourth Expert Meeting on Acid Deposition Monitoring Network in East Asia. Hiroshima, 4.-6. February 1997. Japan Environment Agency, Tokyo, 185-188.

- Fang, W., Ding, M., Lu, D., Li, Z., Cai, X., Zhou, G., Yu Z., 1995. Hydrological dynamics and nutrient migration with precipitation of artificial *Acacia mangium* forest in low subtropical downland. *Acta Ecol. Sinica*, 15, 115-123.
- Galloway, J.N., Zhao, D., Thomson, V.E., Chang, L., 1996. Nitrogen mobilization in the United States of America and the Peoples Republic of China. *Atmos. Environ.*, 30, 1551-1561.
- Hanssen, J.E., Schaug, J., Semb, A., 1996 Manual for sampling and chemical analysis. EMEP/CCC - Report 1/95, revised 1996. Norwegian Institute for Air Research, Kjeller Norway (O-7726).
- Hedin, L.O., Granat, L., Likens, G.E., Buishand, T.A., Galloway, J.N., Butler, T.J., Rohde, H., 1994. Steep declines in atmospheric base cations in regions of Europe and North America. *Nature*, 367, 351-354.
- Hettelingh J-P., Sverdrup H., Zhao D., 1995. Deriving Critical Loads for Asia. *Water Air Soil Pollut.*, 85, 2565-2570.
- Hicks, B.B., Baldocchi, D.D., Meyers, T.P., Hosker, R.P., Matt, D.R., 1987. A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. *Water Air Soil Pollut.*, 36, 311-330.
- Hill, M.O. 1979. DECORANA - A FORTRAN program for detrended correspondence analysis and reciprocal averaging. Cornell Univ., Ithaca, NY.
- Kong, G., Liang, C., Wu H., Huang Z., 1993. Dinghushan Biosphere Reserve. Ecological research history and perspective. Science Press, Beijing, 39 pp.
- Larssen, T., Xiong, J., Vogt, R.D., Seip, H.M., Liao, B., Zhao, D., 1997. Studies of soils, soil water and stream water at a small catchment near Guiyang, China. *Water Air Soil Pollut.* (in press).
- Leggett, S., 1996. Forecast distributions of species and their atmospheric reactivities for the U.K. VOC emission inventory. *Atmos. Environ.*, 30, 215-226.
- Lei, H.C., Tanner, P.A., Huang, M.Y., Shen, Z.L., Wu, Y.X., 1997. The acidification process under the cloud in southwest China: observation results and simulation. *Atmos. Environ.*, 31, 851-861.
- Li Z., Fang, W., Lu D., 1995. Physical and chemical properties of soils in Heshan hilly land. *Acta Ecol. Sinica*, 15, Supp. A, 93-102. (In Chinese with English abstract).
- Liao, B., Larssen, T., Seip, H.M., Vogt, R.D., 1994. Anion adsorption and aluminum release from Chinese soils treated with different concentrations of  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ . *J. Ecol. Chem.*, 3, 281-301.
- Liao, B., Seip, H.M., Larssen, T., 1997a. Response of two Chinese forest soils to acidic inputs: Leaching experiment. *Geoderma*, 75, 53-73.
- Liao, B., Larssen, T., Seip, H.M., 1997b. Response of five Chinese soils to acidic inputs: batch experiment. *Geoderma* (submitted).

Løkke, H., Bak, J., Falkengren-Grerup, U., Finlay, R.D., Ilvesniemi, H., Nygaard, P.H., and Starr, M., 1996. Critical loads of acidic deposition for forest soils: is the current approach adequate? *Ambio*, 25: 510-516.

Moldan, G. and Schnoor, J.L., 1992. Czechoslovakia, examining a critically ill environment. *Environ. Sci. Technol.*, 26, 14-21.

NEPA, 1997. The National Ninth Five-year Plan for Environmental Protection and the Long-term targets for the year 2010. The National Environmental Protection Agency; Beijing, China.

Økland, R.H. & Eilertsen, O., 1996. Dynamics of understorey vegetation in an old-growth boreal coniferous forest, 1988-1993. *J. Vegetation Sci.* 7, 747-762.

Pylvänäinen, M., 1993. Manual for integrated monitoring. Program Phase 1993-1996. Environ. Data Centre, National Board of Waters and the Environment. Helsinki.

Qi, L. and Wang, W., (undated). The character of precipitation at Guangdong and Guangxi provinces in China. Report/Manuscript. Chinese Research Academy of Sciences, Beijing, China.

Rosseland, B.O., Eldhuset, T.D., Staurnes, M., 1990. Environmental effects of aluminum. *Environ. Geochem. Health*, 12, No 1/2, 17-27.

Sæther, O.M., Andreassen, B. Th. and Semb, A., 1995. Amounts and sources of fluoride in precipitation over southern Norway. *Atmos. Environm.*, 29, 1785-1794.

Seip, H.M., Zhao, D., Xiong, J., Zhao, D., Larssen, T., Liao, B., Vogt, R.D., 1995. Acidic deposition and its effects in southwestern China. *Water Air Soil Pollut.*, 85, 2301-2306.

Semb, A., Hanssen, J.E., Francois, F., Maenhaut, W. and Pacyna, J.M., 1995. Long range transport and deposition of mineral matter as a source for base cations. *Water, Air, Soil Pollut.*, 85, 1933-1940.

Skjelkvåle, B.L., Henriksen, A., Vadset, M., Røyset, O. (1996) Trace elements in Norwegian lakes. Preliminary results for 473 lakes (In Norwegian). Norwegian Institute for Water Research, Oslo, SNR 3457-97.

Solberg, S., Dye, C., Schmidbauer, N., Herzog, A., Gehrig, R., 1996. Carbonyls and Nonmethane Hydrocarbons at Rural European Sites from the Mediterranean to the Arctic. *J. Atmosph. Chem.*, 25, 33-66.

Solberg, S., Dye, C., Reimann, S., Schmidbauer, N., 1997. VOC measurements 1996. Norwegian Institute for Air Research Kjeller, EMEP/CCC-Report 7/97.

Sverdrup, H., and de Vries, W., 1994. Calculating critical loads for acidity with the simple mass balance method. *Water Air Soil Pollut.*, 72, 143-162.

ter Braak, C.J.F., 1986. Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology*, 67, 1167-1179.

ter Braak, C.J.F., 1987a. The analysis of vegetation-environment relationships by canonical correspondence analysis. *Vegetatio*, 69, 69-77.

ter Braak, C.J.F., 1987b. CANOCO - a FORTRAN program for canonical community ordination by (partial) (detrended) (canonical) correspondence analysis, principal components analysis and redundancy analysis (version 2.1). TNO Inst. of Appl. Computer Science, Wageningen.

ter Braak, C.J.F., 1990. Update notes: CANOCO version 3.10. Agricult. Math. Group, Wageningen.

Vogt, R.D., Godzik, S., Kotowski M., Niklinska, M., Pawlowski, L., Seip, H.M., Sienkiewicz, J., Skotte, G., Staszewski, T., Szarek, G., Tyszka, J., Aagaard, P., 1994. Soil, soil water and streamwater chemistry in some Polish sites with varying acid deposition. *J. Ecol. Chem.*, 3, 325-356.

Wang R., Yang, J., Tang, H., 1996. Impacts of Acid Deposition on Ecosystems in China. Research report, Research Center for Eco-Environmental Sciences, Academia Sinica, P.O. Box 934, Beijing 100083, China.

Wang W., Chen F., Zhang, Y., Hong S., Tang, D., Zhang, W., 1997a. The current state of acid rain in China. Report. Chinese Research Academy of Sciences, Beijing, China.

Wang W., Ding, G., Wang, T., 1997b. Geographical distribution of rain acidity over China. *Water Air Soil Pollut.* (submitted).

Xue, H.B., and Schnoor, J.L., 1994. Acid deposition and lake chemistry in Southwest China. *Water Air Soil Pollut.*, 75, 61-78.

Zhao, D. and Sun B., 1986. Air pollution and acid rain in China. *Ambio*, 15, 2-5.

Zhao, D. and Wang A., 1994. Estimation of anthropogenic ammonia emissions in Asia. *Atmos. Environm.*, 28, 689-694.

Zhao D., Xiong J., Xu Y., Chan, W.H., 1988. Acid rain in southwestern China. *Atmos. Environm.*, 22, 349-358.

Zhao D., Seip, H.M, Zhao Di., Zhang D., 1994. Pattern and cause of acidic deposition in the Chongqing region, Sichuan Province, China. *Water Air Soil Pollut.*, 77, 27-48.

## **7. Appendix**

# **APPENDIX A**

# Acid rain and its effects in China

Background report from the PIAC project.

Thorjørn Larssen (ed.)<sup>Δ</sup>\*, Hans Martin Seip\*, Arne Semb\*\*, Jan Mulder#, Ivar P. Muniz<sup>###</sup>, Rolf D. Vogt\*, Espen Lydersen<sup>‡</sup>, Valter Angell<sup>°</sup>, Tang Dagang<sup>≡</sup> and Odd Eilertsen<sup>###</sup>.

---

<sup>Δ</sup> E-mail: thorjorn.larssen@kjemi.uio.no

\* Department of Chemistry, University of Oslo, P.O.Box 1033, Blindern, 0315 Oslo, Norway

\*\* Norwegian Institute for Air Research, P.O.Box 100, 2007 Kjeller, Norway

# Norwegian Forest Research Institute, Høyskoleveien 12, 1432 Ås, Norway

### Norwegian Institute for Nature Research, P.O.Box 736, Sentrum, 0105 Oslo, Norway

‡ Norwegian Institute for Water Research, P.O.Box 173, Kjelsås, 0411 Oslo, Norway

° Norwegian Institute of International Affairs, P.O.Box 8159, Dep. 0033 Oslo, Norway

≡ Atmospheric Environment Institute, China Reaserach Academy of Environmental Science (CRAES), Beiyuan, Beijing, China 100012.



## Preface

This report is prepared for the Environmental and Natural Resource Division, Asia Technical Department, the World Bank as a part of the project *Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China* (PIAC). The report is based on information available in printed form, basically from international and Chinese scientific journals. Further information will be given and the present situation described in more detail in the forthcoming reports from PIAC.

Thorjørn Larssen

Oslo, June 27, 1997

### Chemical symbols and abbreviations

SO <sub>2</sub>	Sulfur dioxide	TgSyr <sup>-1</sup>	Terra (10 <sup>12</sup> ) gram (mill. tons) sulfur per year
NO <sub>x</sub>	Nitrogen oxides (NO and NO <sub>2</sub> )	TgNyr <sup>-1</sup>	Terra (10 <sup>12</sup> ) gram (mill. tons) nitrogen per year
NH <sub>3</sub>	Ammonia	mg	milli (10 <sup>-3</sup> ) gram
NH <sub>4</sub> <sup>+</sup>	Ammonium	µg	micro gram (10 <sup>-6</sup> gram)
HF	Hydrogen fluoride	µeq	micro (10 <sup>-6</sup> ) equivalent (= µmol charge)
Ca <sup>2+</sup>	Calcium	ANC	Acid neutralizing capacity
SO <sub>4</sub> <sup>2-</sup>	Sulfate	GNP	Gross national product
Al	Aluminum	GDP	Gross domestic product
Cd	Cadmium		
CaSO <sub>4</sub>	Calcium sulfate (gypsum)		
CaCO <sub>3</sub>	Calcium carbonate (lime)		

## **Contents**

<b>1. Introduction</b>	<b>4</b>
1.1. History of the acid rain problem in China	4
1.2. Why deal with acid rain in China?	4
1.3. Purpose of this report	4
<b>2. Background information</b>	<b>5</b>
2.1. Emissions	5
2.2. Geographical distribution	7
2.3. Atmospheric dispersion and deposition	7
2.3.1. Atmospheric concentrations of SO <sub>2</sub>	7
2.3.2. Measurements and monitoring of precipitation chemistry	7
2.3.3. Dust and particles in the atmosphere	11
2.3.4. Evaluation of the monitoring	12
2.4. Studies of effects on ecosystems	13
2.4.1. Forest and trees	13
2.4.2. Other wild vegetation	14
2.4.3. Crops	14
2.4.4. Soils and soil water	15
2.4.5. Surface water	16
2.4.6. Integrated studies	17
2.4.7. Critical loads	17
2.5. Effects on health	18
2.6. Effects on materials	19
2.7. Economic losses caused by acid rain and related pollutants	20
<b>3. Emission control and ongoing research activities</b>	<b>20</b>
3.1. Information from the State Council of the People's Republic of China	20
3.2. Law on the control of atmospheric pollution	21
3.3. Ongoing projects	22
<b>4. Conclusions</b>	<b>22</b>

## 1. Introduction

### 1.1. History of the acid rain problem in China

Acid rain was recognized as a potential environmental problem in China in the late 1970s and early 1980s (Zhao and Sun, 1986; Zhao et al., 1988; Wang et al., 1997a; Wang et al., 1996). The First National Symposium on Acid Rain was convened in November 1981. In 1982 the National Environmental Protection Agency (NEPA) organized and sponsored a National Survey of Acid Rain, in addition to local research projects in several provinces (Zhao et al., 1988). Based on the findings of the first survey, the Second National Survey on Acid Rain was initiated in 1985 and lasted for two years; the third national acid rain research project lasted from 1986 to 1990 (the 7th five-year plan) and the fourth national project from 1991 to 1995 (the 8th five-year plan) (Wang et al., 1997a). The two first projects focused mainly on emission of SO<sub>2</sub>, distribution and deposition of acid rain, while the two next projects also involved studies on effects.

### 1.2. Why deal with acid rain in China?

The Chinese energy consumption increased 5.3% annually over the period 1980-1991 (Byrne et al., 1996). Coal accounted for more than 3/4 of the commercial energy production and it is likely that coal will still be the major energy carrier in the next decades. The SO<sub>2</sub> emissions have shown a lower growth rate due to cleaner technology on new power plants and boilers, but the SO<sub>2</sub> emissions are still increasing. All reasonable scenarios tell that the Chinese SO<sub>2</sub> emissions will continue to increase, and also the NO<sub>x</sub> emission due to the fast increasing number of motor vehicles in the cities. Hence increased acidification in China is likely (e.g. Rodhe et al., 1992; Seip et al., 1995). Preliminary calculations, e.g. by the RAINS-Asia model suggest that the critical loads of forest ecosystems are seriously exceeded in many areas (Hettelingh et al., 1995). However, the critical loads are uncertain and largely untested, as discussed later.

The combination of high emissions of acid gases and acid sensitive ecosystems requires a better understanding of the relationship between emissions and environmental effects than what is available today.

### 1.3. Purpose of this report

The purpose of this report is to describe the current knowledge related to acid rain in China. The background for preparing the report is the involvement of several Norwegian institutions in acid rain research in China through the PIAC-project (*Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China*). The focus in the report is on emissions important for formation or neutralization of acid rain, deposition and effects of acid rain. We concentrate on ecosystem effects rather than effects on health and materials because the latter two are largely local effects, particularly health effects which are primarily caused by the acid rain precursor SO<sub>2</sub> (and particles) rather than the acidity of rain.

## 2. Background information

### 2.1. Emissions

Estimated total SO<sub>2</sub> emissions in China vary from 16 to 22 million tons per year (see e.g. Wang et al., 1996; Cao, 1989; Arndt et al., 1997; Akimoto and Narita, 1994). The official figure for 1995 is ~19 million tons SO<sub>2</sub>, while the RAINS-Asia model used 22 million tons (Wang et al., 1996). Akimoto and Narita (1994) discuss the emission figures and explain much of the differences with different average sulfur content of coal used in the calculations. Wang et al. (1996) explain the differences with different emission sectors included in the estimates, e.g. small domestic sources are probably underestimated in the official figures. The annual total SO<sub>2</sub> emission has been increasing the last decades (Figure 1) and is likely to increase further in the near future. Coal is the major energy source in China, accounting for about 75% (Wang et al., 1996). The average sulfur content of the coal consumed is 1.2%. In Guizhou and Sichuan provinces the averages were S-content of coal is 3.2% and 2.8%, respectively.

The nitrogen emissions in China are dominated by NH<sub>3</sub> from fertilizer and domestic animal waste (Zhao and Wang, 1994; Galloway et al., 1996). Commercial fertilizers account for about 80% of China's total 20 TgNyr<sup>-1</sup> mobilization to the atmosphere. The nitrogen mobilization is expected to increase significantly in the coming decades due to increased use of fertilizer as well as enhanced fossil fuel combustion (Galloway et al., 1996).




Figure 1 Sulfur emission in China from 1982 to 1995 (Statistics China, 1983-1996)




Figure 2 Map of the people's republic of China

## 2.2. Geographical distribution

According to Zhao and Sun (1986), two core areas for acid rain were identified in the beginning of the 1980s. This was the Chongqing-Guiyang core zone and the Nanchang core zone. In the beginning of the 1990s two more core zones were observed: one in the southeast coastal area (Fuzhou-Xiamen-Shanghai) and one in the coastal north area surrounding Qingdao in Shandong Province (Wang et al., 1996).

## 2.3. Atmospheric dispersion and deposition

### 2.3.1. Atmospheric concentrations of SO<sub>2</sub>

Monitoring of SO<sub>2</sub> in 88 Chinese cities in 1994 showed that annual average concentrations varied from 2 to 472 µg/m<sup>3</sup> (Wang et al., 1996). Average concentrations were 89 µg/m<sup>3</sup> in northern cities and 83 µg/m<sup>3</sup> in southern cities. 48 of the 88 cities exceeded the Chinese National Air Quality Standard Class II (60 µg/m<sup>3</sup> annual avg.) for SO<sub>2</sub> (Wang et al., 1996). High SO<sub>2</sub> concentrations are observed in both southern and northern China, while acid rain is mainly observed in the south. SO<sub>2</sub> concentrations in Chinese cities, from different sources, are given in Table 1.

### 2.3.2. Measurements and monitoring of precipitation chemistry

In the 1980s, most of the acid rain research was focused on the situation in the Sichuan, Guizhou, Guangxi and Guangdong provinces (Zhao and Sun, 1986; Wang et al., 1997a). The Research Center for Eco-Environmental Sciences (Chinese Academy of Sciences) in Beijing focused on Sichuan and Guizhou provinces, while the Chinese Research Academy of Environmental Sciences (CRAES) focused on Guangxi and Guangdong provinces. Detailed information on precipitation chemistry in Sichuan and Guiyang is available from the early 1980s (e.g. Zhao et al., 1988; Zhao and Zhang, 1990; Zhao and Xiong, 1988; Zhang et al., 1996; Xue and Schnoor, 1994). The general picture was that pH was low and sulfate, calcium and ammonium concentrations high in the city centers of Chongqing and Guiyang. The total ion concentration decreased out from the urban area, but with only a slight increase in pH (Table 2). This can be explained by the high emission from small, domestic sources inside the city centers, with large emissions of both SO<sub>2</sub> and dust.

From 1986 to 1990 there was a monitoring network containing 70 monitoring stations in Guangdong and Guangxi provinces (Qi and Wang, 1990). Generally the chemical composition of precipitation in these provinces is quite similar to that in Guizhou and Sichuan, with high concentrations of sulfate, calcium and ammonium. However, the concentrations were generally somewhat lower in Guangdong and Guangxi than Guizhou and Sichuan.

In the beginning of the 90s, the National Acid Deposition Monitoring Network (NADMN) was started, with the purpose of increasing the knowledge about acid deposition in the other provinces than the southern and southwestern. There are all together 261 monitoring sites in this network throughout China, jointly managed by the National Meteorological Agency, the Water Conservation Agency and NEPA. This network mainly covers the southeastern provinces. The sites are divided into three categories (Wang et al., 1997a):

- General or routine sites
- Additional sites (additional to the sites managed at local level)
- Special chemical study sites

Precipitation sampling and analyses are standardized nationally. The samples are analyzed locally, but quality assessments have been conducted regularly by CRAES (Wang et al., 1997b). The volume-weighted annual average concentrations of ions in precipitation in the 8 provinces dominating this monitoring network are shown in Table 2. At all sites sulfate is the dominating anion and calcium and ammonium the major cations. The pH is relatively low, with weighted averages less than 4.5 in several provinces. The acid rain situation in Chongqing and Guiyang is particularly serious, with a volume-weighted average pH of about 4.1 in the urban areas. However, it is likely that the situation has been improved recently in Guiyang, due to countermeasures induced by the local government.

The information on acid rain in rural areas is relatively scarce since few monitoring data are available. However, some model studies indicate the level of air pollution in rural areas and the degree of long range transport. Meng et al. (1996; 1997) combined use of a simplified three dimensional Eulerian model and a Lagrangian trajectory model to study the transport and deposition of sulfur in the Minnan area in Fujian province. They concluded that long range transport (i.e. from sources outside the Fujian province), accounted for almost 60% of the total sulfur deposition.

The chemical composition of cloudwater was studied at the top of Emei mountain in Sichuan during a sampling period in 1980; the average pH was 4.64, with a minimum of 3.77 (Wang et al., 1991).

In northern parts of China, the pH of precipitation is generally above 6, due to the high levels of neutralizing soil dust in the atmosphere (e.g. Zhao et al.; 1988, Wang et al., 1997b). Hence acid rain is not an important problem in this area, but of course direct effects from acid gases (and particles) may be of great importance to both health and vegetation.

The composition of the rain samples from China differs from the composition of precipitation in Europe mainly in that the concentrations of calcium relative to sulfate are very high, and the concentrations of nitrate relative to the other components are low. In relation to soil acidification and long-term effects, it is the balance of acidifying versus acid-neutralizing components which is of significance. Usually the acid input equals the sulfate and part of the nitrogen input, while alkaline fly ash and soil dust represent the acid-neutralizing part. pH may be an important parameter in relation to short-term effects on materials, and to some extent on vegetation, but is misleading in many other connections.

Calcium in precipitation is very sensitive to the location of the collection site and the sampling procedures. In cities and industrial areas with emissions of dust and particles, settling of coarse particles may be considerable. Resuspension of dust and soil from the surface near the sampler may also be a problem. Wet-only samplers have been widely recommended in order to avoid effects of locally generated dust at sampling sites. However, alkaline dust particles from industrial processes may also be transported over long distances in amounts which are significant in relation to acid rain. In China, natural alkaline dust from the arid areas in Central Asia and northern China influences the precipitation chemistry over very large areas. It is largely this effect which limits the area of acid precipitation to the southern and southwestern part of China. Dust and particles in the atmosphere are discussed further below.

Table 1 Concentrations of SO<sub>2</sub> and total suspended particles (TSP) in air at different sites in China.

Place	SO <sub>2</sub> conc. (µg/m <sup>3</sup> )	TSP (µg/m <sup>3</sup> )	avg. period	year measured	no. of measure spots included	reference
Northern	110	740	24 hr.	1985	31	Cao, 1989
Southern	100	444	24 hr.	1985	33	Cao, 1989
Beijing	115	380	1 year	1991	1	Wells et al., 1994
Chengdu., urban	85	658	annual	'85-'89	1	Lei et al., 1997
Chongq., Jiulongpo district	280		annual	1990	?	Ma, 1990
Chongq., Ba county	240		annual	1990	?	Ma, 1990
Chongq., Dadukou district	320		annual	1990	?	Ma, 1990
Chongq., Jiangbei district	400		annual	'88-'89	?	Ma, 1990
Chongq., Nanan district	310		annual	1990	?	Ma, 1990
Chongq., Nanshan park	50		1 month	Jan. '89	1	Ma, 1990
Chongq., Nanshan, Huangshan	130		1 month	Jan. '89	1	Ma, 1990
Chongq., Nanshan, Tieliao	450		1 month	Jan. '89	1	Ma, 1990
Chongq., Nanshan, Wenfongtai	133		24 hr.	Sept '86	1	Bian and Yu, 1992
Chongq., Nanshan, Wenfongtai	254		24 hr.	July '88	1	Bian and Yu, 1992
Chongq., Nanshan, Zhenwushan	400		1 month	Jan. '89	1	Ma, 1990
Chongq., Shizhong district	540		annual	1990	?	Ma, 1990
Chongq., Simian shan	5	100	annual	'85-'89	1	Lei et al., 1997
Chongq., Simian shan	8		annual	'89-'90	1	Ma, 1990
Chongqing	422		annual	1990	?	Zhao et al., 1995
Chongqing, Nanshan park	40		24 hr.	Sept '86	1	Bian and Yu, 1992
Chongqing, Nanshan park	100		24 hr.	July '88	1	Bian and Yu, 1992
Chongqing, Nanshan, Zhenwushan	126		24 hr.	Sept '86	1	Bian and Yu, 1992
Chongqing, Nanshan, Zhenwushan	214		24 hr.	July '88	1	Bian and Yu, 1992
Chongqing, suburb 1	138	500	annual	'85-'89	1	Lei et al., 1997
Chongqing, suburb 2	27	300	annual	'85-'89	1	Lei et al., 1997
Chongqing, urban	402	690	annual	'85-'89	1	Lei et al., 1997
Guangzhou	90		annual	1988	?	Cao et al., 1991
Guangzhou	47	180	annual	1991	1	Wells et al., 1994
Guiyang	504	496	winter season	1990	6	UNDP, 1991
Guiyang	356		annual	1990	?	Zhao et al., 1995
Guiyang catchment	44		1 month	1993	1	Larssen et al., 1997
Guiyang, suburb 1	40	500	annual	'85-'89	1	Lei et al., 1997
Guiyang, suburb 2	48	300	annual	'85-'89	1	Lei et al., 1997
Guiyang, urban	157		1 month	1993	1	Larssen et al., 1997
Guiyang, urban	423	970	annual	'85-'89	1	Lei et al., 1997
Huhot	119	375	winter season	1990	5	UNDP, 1991
Qingdao	470	335	winter season	1990	4	UNDP, 1991
Qingdao	233		annual	1990	?	Zhao et al., 1995
Shanghai	85	220	1 year	1991	1	Wells et al., 1994
Shenyang	152	380	1 year	1991	1	Wells et al., 1994
Xian	47	510	1 year	1991	1	Wells et al., 1994



Table 2 Volume weighted annual average concentrations of ions in precipitation in China (concentrations in  $\mu\text{eq/l}$ ).

Period	Fujian <sup>a</sup>		Jiangxi <sup>a</sup>		Hunan <sup>a</sup>		Zhejiang <sup>a</sup>		Hubei <sup>a</sup>		Anhui <sup>a</sup>		Jiangsu <sup>a</sup>		Shandong <sup>a</sup>		Chongqing <sup>b</sup>		Chongqing <sup>c</sup>		Chongqing <sup>b</sup>		Chongqing <sup>c</sup>		Guiyang <sup>b</sup>		Guiyang <sup>b</sup>		Guiyang <sup>d</sup>			
	Average	1992-93	Average	1992-93	Average	1992-93	Average	1992-93	Average	1992-93	Average	1992-93	Average	1992-93	Average	1992-93	Urban	1982/84	Urban	1987-89	Rural	1982/84	Urban	1982/84	Rural	1987-89	Urban	1982/84	Suburban	1992-95	Suburban	1982/84
rainfall (mm)	1379	1555	1274	1550	1108	1550	1108	1550	1108	1550	1020	1212	597	597	700	700	4.11	4.44	4.11	4.33	4.44	4.44	4.07	4.42	4.33	4.07	4.42	4.43	4.43	4.58	4.58	
pH	4.48	4.49	4.78	4.69	4.47	4.69	4.47	4.69	4.47	4.69	5.37	5.48	6.10	6.10	4.14	4.14	4.14	4.44	4.14	4.33	4.44	4.07	4.42	4.33	4.07	4.42	4.43	4.43	4.58	4.58		
H <sup>+</sup>	33.2	32.4	16.6	20.5	34.0	20.5	34.0	20.5	34.0	4.3	3.3	3.3	0.8	0.8	72.4	77	77	36.3	77	47	36.3	84.5	37.9	49.2	47	84.5	37.9	37.2	37.2	26.3	26.3	
NH <sub>4</sub> <sup>+</sup>	70.1	51.3	81.7	68.5	100.5	68.5	100.5	68.5	100.5	58.5	60.2	60.2	52.4	52.4	106	123	123	64.1	123	116	64.1	78.9	49.2	23.9	116	78.9	23.9	23.9	50.6	50.6		
Ca <sup>2+</sup>	52.3	64.5	63.0	49.3	64.8	49.3	64.8	49.3	64.8	65.2	116.7	116.7	167.6	167.6	110	125	125	42.0	125	74	42.0	231.2	198	125.3	74	231.2	125.3	125.3	87.7	87.7		
Mg <sup>2+</sup>	7.3	11.5	10.1	9.6	9.5	9.6	9.5	9.6	9.5	8.6	14.0	14.0	52.9	52.9	48.3	31	31	18.3	31	15	18.3	56.5	44.6	26.6	15	56.5	26.6	26.6	29.4	29.4		
Na <sup>+</sup>	14.8	22.5	8.9	24.5	8.9	24.5	8.9	24.5	8.9	18.6	29.6	29.6	41.4	41.4	51.4	17	17	45.4	17	22	45.4	10.1	11.2	6.7	22	10.1	6.7	6.7	5.9	5.9		
K <sup>+</sup>	5.1	10.42	7.8	8.4	7.3	8.4	7.3	8.4	7.3	10.8	8.9	8.9	28.0	28.0	7.4	17	17	23.4	17	17	23.4	26.4	10.5	6.0	17	26.4	6.0	6.0	7.0	7.0		
SO <sub>4</sub> <sup>2-</sup>	104	100	128	110	129	110	129	110	129	107	166	166	161	161	307	299	299	165	299	200	165	411	281	188	200	411	188	188	167	167		
NO <sub>3</sub> <sup>-</sup>	14.0	19.6	18.1	18.0	22.4	18.0	22.4	18.0	22.4	21.4	20.9	20.9	22.6	22.6	31.6	23	23	18.0	23	20	18.0	21.0	25.3	15.2	20	21.0	15.2	15.2	15.9	15.9		
Cl <sup>-</sup>	19.3	14.9	16.3	20.3	16.4	20.3	16.4	20.3	16.4	24.4	28.0	28.0	91.4	91.4	15.0	30	30	23.9	30	21	23.9	8.2	11.8	10.1	21	8.2	10.1	10.1	21.1	21.1		
F <sup>-</sup>	11.0	16.4	11.7	8.3	3.2	8.3	3.2	8.3	3.2	1.0	10.7	10.7	21.5	21.5	35	35	35	33	35	33	33	4.8	4.8	4.8	33	4.8	4.8	4.8	4.8	4.8		
Σ+	182	192	188	180	225	180	225	180	225	166	232	232	343	343	395	390	390	229	390	291	229	488	351	226	291	488	226	226	207	207		
Σ-	149	151	174	156	171	156	171	156	171	154	226	226	297	297	354	387	387	207	387	274	207	440	318	218	274	440	218	218	204	204		
(Σ+ - Σ-)100	10.2	12.0	3.8	7.3	13.5	7.3	13.5	7.3	13.5	3.9	1.5	1.5	7.2	7.2	5.6	0.4	0.4	5.2	0.4	3.0	5.2	5.11	4.97	1.7	3.0	5.11	1.7	1.7	0.71	0.71		
(Σ+ + Σ-)																																

<sup>a</sup>From Wang et al. (1997a); <sup>b</sup>From Zhao et al. (1988); <sup>c</sup>From Zhao et al. (1994); <sup>d</sup>From Larssen et al. (1997)

Possible effects of the large emissions of reduced nitrogen compounds ( $\text{NH}_3$ ,  $\text{NH}_4^+$ ) have not been studied in detail in China. However, these emissions is of importance, both because it may neutralize acidity of the rain, and because it may cause acidification of soils after deposited.

The concentration of fluoride in precipitation appears also to be high in China. This may be linked to combustion of coal with high fluoride contents, but also to production of brick and tiles from clay with high fluoride contents. Emissions occur as hydrogen fluoride (HF), which is strongly toxic to plants.

### 2.3.3. *Dust and particles in the atmosphere*

Wind-blown soil dust is an important feature in the Chinese atmosphere (Chang et al., 1996). In addition there are high concentrations of anthropogenically derived aerosol particles. Zhao et al. (1988) suggested, based on the work of Wang et al. (1981), that the ratio of coal burning dust to wind blown soil dust is about 2:1 in southern China, and between 2:3 and 3:2 in northern China. Zhao et al. (1994) estimated 40% of the dust in Chongqing to originate from coal burning, however, the authors pointed out that the method was uncertain. Wang and Wang (1996) suggested that generally 30-70% of the particulate matter in urban areas originate from soils.

Generally the airborne anthropogenically derived particles are smaller than the soil dust particles (Ning et al., 1996; Yin et al., 1996). Ning et al. (1996) compared the particle aerosol size in a rural area (outside Shenyang) with that in urban areas in cities in northern China (Beijing, Shenyang, Lanzhou and Taiyuan). At the rural site, most of the particles had an aerodynamic diameter larger than  $10\mu\text{m}$ . In the cities the smaller fractions dominate. They also found the pH and  $\text{Ca}^{2+}$  content of the particles to decrease and  $\text{SO}_4^{2-}$  and  $\text{NH}_4^+$  content to increase with decreasing particle size. In northern China the content of base cations in airborne soil particles is considerably larger than in southern China. According to Wang and Wang (1996) the contents of calcium and sodium are respectively about 3% and 1.5% in the north and about 0.1% and 0.5% in the south.

Yin et al. (1996) compared the chemical composition of the aerosol particles in Beijing, Chongqing and Guiyang. As can be seen from Table 3, the pH of the particles dissolved in water was much lower in Chongqing and Guiyang than in Beijing, while the base cation concentration, particularly  $\text{Ca}^{2+}$ , was higher in Beijing. The sulfur content was high all three places, though there are differences in the temporal variation. These results clearly show that the atmospheric particles in northern China have a large ability to neutralize acid rain, compared to the situation in the southwest. To what extent aerosols in the southwest have an alkalizing effect on the precipitation is not clear, although, Liu and Huan (1993) showed that in Chongqing the aerosols have some neutralizing effect on the precipitation. However, it is not clear in what forms  $\text{Ca}^{2+}$  is bound in the particles, e.g. as non-neutralizing  $\text{CaSO}_4$  or neutralizing  $\text{CaO}$  and  $\text{CaCO}_3$  (Chang et al., 1996).

In addition to the possible neutralizing effect of the aerosol particles, the particle surfaces also play an important role in sulfate and nitrate formation (Chang et al., 1996).

Model studies by Chang et al. (1996) suggested that about half of the total  $\text{Ca}^{2+}$  deposition in China is due to wet deposition.

Table 3: Mass percentage concentration and pH in aerosol particles dissolved in water (from Yin et al., 1996).

Element	Chongqing	Guiyang	Beijing
S <sub>(March)</sub>	4.2	2.8	3.2
S <sub>(Sept.)</sub>	2.2	0.8	4.5
K	0.7	0.3	1
Ca	1.3	1	4
pH	4.1	4.3	6.8

Lei et al. (1997) discussed the chemical composition of precipitation samples compared to cloud-water samples collected simultaneously from aircraft in 1985-1989. The concentrations of ammonium, calcium, hydrogen and sulfate in rainwater samples collected at urban sites in southwest China are all lower than the corresponding concentrations in simultaneously collected cloudwater samples. This is attributed to washout by precipitation over urban areas. This was also found for suburban precipitation samples, but the effect was less pronounced at rural sites. In particular, the concentration of calcium in rural rainwater samples was less than in urban and suburban samples, and were comparable to the concentrations in cloudwater samples. Simulation of the washout process indicated that the sulfate concentration in urban rainwater samples is limited by the hydrogen peroxide concentration rather than the sulfur dioxide concentration. In a chamber study in Guangzhou, Lin et al. (1992b) found that the transformation rate of SO<sub>2</sub> was strongly dependent on the number of condensation nuclei. Shao et al. (1995) combined statistical factor analyses with accelerator mass spectrometry to investigate source identification of aerosols in Beijing, Shandong and Hunan. Several authors have discussed the long-range transport of dust from the northern Chinese deserts to Korea, Japan and the South China Sea, which in parts of the year may be considerable (e.g. Chang et al., 1996; Hashimoto et al., 1994; Suzuki et al., 1993).

In the suburban area of Guiyang, the dry deposition and the wet deposition of sulfur were estimated to be about the same, approximately 4.3 gS/m<sup>2</sup> (Larssen et al., 1997).

#### 2.3.4. Evaluation of the monitoring

The acid rain monitoring in China was evaluated by an expert group at the *Fourth Expert Meeting on Acid Deposition and Monitoring Network in East Asia* in February, 1997 (EMAD, 1997). The expert group concluded that it is necessary to further optimize the acid rain monitoring network in China and that there currently are some gaps between the State's needs and the actual situation. As weakest points in the present monitoring programme network they identified the use of non-state-of-the-art technology, the regional representativity of the selected sites, the lack of data from background stations, measurement of too few parameters and the quality assurance. They concluded that the financial support to acid rain monitoring is too low and should be increased.

## 2.4. Studies of effects on ecosystems

### 2.4.1. Forest and trees

Damages to forests may be due to direct effects of the acid rain precursors  $\text{SO}_2$  and  $\text{NO}_x$ , or to indirect effects involving soil acidification and mobilization of toxic metals as aluminum. Events with extremely acid rain may also cause direct damages of leaf surfaces. Most research on forest damage in China has been related to direct effects from  $\text{SO}_2$ , acid mist or extremely acid rain events.

Forest decline in large areas has up to now not been reported in China (Bian and Yu, 1992). However, forest decline in smaller areas, particularly near cities and industrial areas is observed. In the early 1980s serious forest damage has been observed on Nanshan mountain, just outside Chongqing city. More than 50% of 1800 ha. masson pine (*Pinus massoniana*) stand died (Wang et al., 1996). Several researchers have discussed the reason for the forest decline (e.g. Bian and Yu, 1992; Totsuka et al., 1994; Liu and Du, 1991). Important factors are considered to be high concentrations of  $\text{SO}_2$  and hydrogen fluoride (HF), acid rain and attacks by insects and fungi. Several different symptoms were observed on trees, as tip necrosis of needles, reduced needle length, premature abscission, crown thinning, branch dieback and reduced radial growth (Bian and Yu, 1992). Bian and Yu (1992) investigated three sites with different pollution loading at Nanshan. They found good correlations between the air concentration of  $\text{SO}_2$ , the sulfur content of the needles and the extent of damage. However, even better correlations were found between fluorine content in needles and damage. Unfortunately analyses of HF in air was not available. Differences in soil properties at sites with healthy and damaged pine were not observed, hence Bian and Yu (1992) concluded that direct effects from the gases and not acidification of soils were important for the observed damage. Other scientists believe that soil acidification has been of major importance (Ma, 1990). The final dieback of the trees is believed to be caused by insect pests (Bian and Yu, 1992). The annual forest biomass production (of masson pine) was found to be 3-4 times lower at Nanshan than at three less polluted areas (Wang et al., 1996).

Regarding the discussions about the causes of the Nanshan forest decline, Totsuka et al. (1994) compared the conditions of masson pine and camphor tree at one heavily polluted and one lightly polluted area outside Chongqing. They also conducted laboratory experiments to clarify interactive effects of  $\text{SO}_2$  and soil acidification on growth. They found a slower growth with high  $\text{SO}_2$  concentration and acid soil, but did not try to generalize the observations e.g. in terms of dose-response relationships.

According to Wang et al. (1996) 40% of *A. f. xabri (mast.) craib* died at Emei mountain in Sichuan. However, this observation has not been confirmed by other sources. Cao et al. (1988) and Wang et al. (1988) discussed the relationship between acid precipitation and decline of fir at high elevations at Emei mountain and conclude that acid rain and acid fog may result in both direct and indirect effects.

A forest damage assessment study was carried out in Liuzhou, Guangxi province (Wang et al., 1996). Of 436 tree species investigated, 84 were affected; 30 of these seriously. *Pinus massoniana* and *Cinnamomum burmannii* were the two most heavily injured tree species. Wang et al. (1996) do not discuss the relationship between the observed damages and the pollution types and -levels.

According to Wang et al. (1996) 32% of the forested area (280 000 ha) in Sichuan province (including Chongqing) is influenced by air pollutants and acid rain. In Guizhou province 15,000 ha

are influenced. Loss of masson pine was estimated from results showing differences in growth in clean and polluted areas in 11 Chinese provinces. The annual growth rate in acidified compared to non-acidified areas was found to be 40% - 50% (Wang et al., 1996).

Cao et al. (1991) and Shu et al. (1993) presented a multiple regression equation for yield loss of coniferous trees as a function of rain pH, SO<sub>2</sub> concentration and soil depth. They did not report details about model development and did not discuss uncertainties which obviously are large.

#### 2.4.2. *Other wild vegetation*

Very little information exists about possible effects of air pollution and acid rain in China on other wild vegetation than the most common trees. However, this kind of vegetation is important in China, for example is a large variety of wild herbs and firms collected for food, medicin and fodder.

We are also not aware of any comprehensive studies of lichens or other specially pollution sensitive species.

#### 2.4.3. *Crops*

Effects of SO<sub>2</sub> and HF on crops have been studied in China since the 1970s (Cao, 1989). However, few studies have been carried out on the effects of other gaseous pollutants as ozone and NO<sub>x</sub>. Cao (1989) reported short-term exposure concentrations of SO<sub>2</sub> and HF causing 5% injury to three sensitivity groups of crops. The sensitive species are reported to have a 5% injury at 880-1430 µg/m<sup>3</sup> SO<sub>2</sub> and 12-48 µg/m<sup>3</sup> HF during 8 hour exposure (Cao, 1989). How the injury was measured was not reported. Dose-response curves for yield loss caused by SO<sub>2</sub> were given for short-term exposure, the investigated plants have decreasing sensitivity as follows:

Cabbage > Pinto Bean > Potato > Wheat > Soybean > Corn > Rice

For long-term exposure of SO<sub>2</sub>, a yield loss of 5% is reported for 30-50 µg/m<sup>3</sup> SO<sub>2</sub> for sensitive species (potato, bean, Chinese cabbage). In the most polluted Chinese cities a yield loss of 5-25% should be expected for the most common crops (Cao, 1989). Chang and Hu (1996) reported that the average yield reduction for vegetables in Chongqing is 24.4%. In an other study Cao et al. (1989) studied effects of both SO<sub>2</sub> and acid rain in open top chambers. Rice was among the most resistant species both to SO<sub>2</sub>, acid rain and the combination of the two. "Most vegetables" are classified as sensitive.

A ranking of the sensitivity of acid rain for the most common crops in China is given by Wang et al. (1996):

rape > wheat > corn > barley > soybean > rice > tobacco > jute

Of the most common vegetables tomato, aubergine, and cucumber are among the most sensitive, while cabbage, spinach and carrot are among the least sensitive for acid rain (Wang et al., 1996). The sensitivity ranking from Cao (1989) and from Wang et al. (1996) are rather different. It is not clear if the differences are due to differences between SO<sub>2</sub> and acid rain effects, or if they are due to experimental uncertainties.

Cao et al. (1990) studied acute injury of acid solutions to 105 species of garden and flower plants. Plants were either sprayed with, or shots soaked in, solutions with pH ranging from 1.5 to 3.5. Most species showed acute injury when pH was below 3.0; pH above 3.5 did usually not cause acute injury.

In all the above mentioned studies, the effects have not been related to soil properties, which are of major importance for the growth.

#### 2.4.4. *Soils and soil water*

Direct effects of air pollutants on vegetation are most commonly used to explain damages in China. However, changes in soils caused by acidification are a likely long-term effect of acid rain. Dai et al. (1997) and Pan (1992) compared soil analyses conducted 30 years ago with recent results from several sites in southern China. They found the soil pH to have decreased considerably; between 0.1 and 1.0 pH units. These results clearly suggest that soil acidification has occurred and may become a serious problem in China in the future. This is in agreement with modeling results (Zhao and Seip, 1991). However, experimental and modeling studies involve relatively large uncertainties. Furthermore acid deposition is not the only possible cause of soil acidification; for example changes in vegetation type, ecosystem succession etc. may have similar effect (e.g. logging of broad-leaved vegetation followed by planting of coniferous trees, as was the case in large areas of China in the late 50s and early 60s).

Liao et al. (1997a; 1997b) conducted laboratory experiments in which forest soils from 5 sites in southern China (Chongqing, Guiyang, Fujian, Hunan and Nanchang) were extracted with different salt- and acid solutions. The experiments showed that soil acidification is likely to occur in the most sensitive soils, if the acid deposition continues at current level or increases. The soils from the Fujian and Nanchang were found to be the most sensitive of these soils. However, one must be careful in generalizing to larger geographical scales because of the heterogeneity of soils even within small geographical areas (Larssen et al., 1997). To what extent effects on vegetation via soil acidification may occur in the future will depend strongly on the development of both the sulfur and the base cation deposition. Lin et al. (1992a) studied the buffering capacity of soils from several places in China. They concluded that soil from Nanshan (outside of Chongqing) had a low buffering capacity, soils from Guiyang and Hunan relatively high, and soils from Hebei and Xian the highest buffer capacity. Such soil experiments give valuable information about the particular soil sampled.

Sulfate absorption in soils is a process which may retard the acidification process. Liao et al. (1994) studied this process in soils from Guiyang and Nanchang in laboratory experiments and found that the sulfate adsorption was relatively low at sulfate concentrations corresponding to ambient sulfate deposition. Larssen et al. (1997) estimated the sulfate retention in a catchment outside Guiyang to be 30-40% of the input.

Results from studies of soil water have been published from sites in Chongqing (Xu et al., 1994a) and Guiyang (Larssen et al., 1997). In Chongqing, high concentrations of sulfate (80-335 mg/l) and calcium (10-51 mg/l) were found, but low concentrations of aluminum (38-150 µg/l) (Xu et al., 1994a). In the Guiyang study concentrations of all these three components were high (SO<sub>4</sub><sup>2-</sup>: 18-111 mg/l; Ca<sup>2+</sup>: 3-25 mg/l; Al<sup>3+</sup>: 0.4-13 mg/l) (Larssen et al., 1997). Precipitation, soil and soil water chemistry in the Guiyang catchment were compared with several catchments in Poland and Norway,

showing that concentrations of sulfate, aluminum and calcium were especially high in Guiyang (Larssen et al., 1996). Singer (1993; 1988) has conducted detailed studies of the weathering patterns of soils in Guangxi; the results may be useful in acidification modeling at a later stage.

The number of soil micro-organism was found to be considerably lower in soils at Nanshan compared to the less acidified area at Simianshan (100 km. south of Chongqing); Wang et al. (1996) put these observations in relation to the increased number of insect attacks at Nanshan. This may affect the decomposition rate of organic matter, the fixation rate of nitrogen etc., and may cause imbalance in the soil nutrient state (Wang et al., 1996). Liao et al. (1991) compared the toxicity of simulated acid rain to soil microbes in Liuzhou and Nanning and concluded that acid rain affects soil microbes in Liuzhou, but not yet in Nanning. Because soil micro-organisms are difficult both to quantify and qualify these studies have large uncertainties.

#### 2.4.5. *Surface water*

Surface water acidification in China is considered to be a minor problem. Xue and Schnoor (1994) published results from a survey of 16 streams and lakes in southwestern China: All investigated waters had a pH above 6.5 and base cation concentration above 300  $\mu\text{eq/l}$ , due to considerable acid neutralization in the soils and high deposition of dust. At high elevation Xue and Schnoor (1994) found some waters that may be sensitive to acidification ( $\text{ANC} < 150 \mu\text{eq/l}$ ), but not acidified at present. Williams et al. (1995) studied the ion concentration in the Ürümqi river in northwest China and concluded that this river is not sensitive to acidification. Larssen et al. (1997) reported rather low pH in a small stream outside Guiyang. However, the water is rapidly neutralized downstream, probably due to mixing with drainage water from more alkaline soils. Model studies by Zhao and Seip (1991) using the MAGIC model (Cosby et al., 1985) also indicated that acidification of surface water is not likely to become a serious problem.

There are very few studies on the ecological effects of high acidity on Chinese freshwater ecosystems. Xia et al. (1994) studied effects on freshwater biota in four small ponds and two lakes in the Chongqing area. They compared acid, fish-less ponds (pH 4.2 and 4.7) with high concentration of aluminum (2.3 and 1.8 mg/l) and sulfate (20.0 and 20.5 mg/l) with less acid ponds (pH 5.1 and 5.5) and neutral control lakes (pH 7.0 and 8.5). The two moderately acid ponds and the control lakes had lower concentration of aluminum (0.24 to 0.32 mg/l) but the sulfate levels were high (16.8 to 52.0 mg/l). Furthermore the most acid ponds had higher transparency, less species of phytoplankton, lower algal cell-density, biomass and chlorophyll-a concentrations. Experimental evidence suggested that low pH and low phosphorous levels were limiting factors for these primary producers. The numbers and mean densities of zooplankton were also much reduced at low pH and tests indicated critical pH's for zooplankton in the range 5.0 - 5.5. Zoobenthos density and biomass, however, were highest in the most acid ponds and not correlated to the measured water quality parameters. According to the authors this could be attributed to the absence of fish. They also pointed out that the reason for the high acidity was not clear.

Even though available observations show that large scale water acidification is unlikely, one should be aware of possible future acidification in some sensitive areas if the present low loadings are changed to higher inputs of acid rain.

#### 2.4.6. *Integrated studies*

In order to understand the acidification processes in the Chinese yellow soil (Haplic Alisol in the FAO classification system) in more detail, a small catchment outside Guiyang was equipped for sampling of SO<sub>2</sub> gas, precipitation, throughfall, soil water and surface water in 1992. The results from the first years of sampling are given in Larssen et al. (1997). The relatively low pH in the precipitation (average: 4.4) causes high concentration of inorganic aluminum in soil water. However, possible toxic effects of aluminum are probably counteracted by high concentration of base cations, in particular calcium. The study also showed that the commonly used mechanisms for describing aluminum chemistry (gibbsite equilibrium and Gaines-Thomas ion exchange) were not able to describe the field data in a satisfactory way. Surface water acidification occurred in the lower order small streams in the upper part of the catchment, but is neutralized further downstream (Larssen et al., 1997). They concluded that the present high base cation deposition dampens possible effects of the high sulfur deposition.

#### 2.4.7. *Critical loads*

Critical load is often defined as “the maximum input of acid deposition to an ecosystem which will not cause long-term damage to ecosystem structure and function”. The concept of critical loads is widely used in Europe and estimation of critical loads has also been done in China (Wang et al., 1996; Hettelingh et al., 1995; Zhao et al., 1995). The areas most sensitive to acid deposition in China are in the south. Wang et al. (1996) pointed out that the Nanchang area at present has a high loading of sulfur and low critical loads and hence is an area where damages may be expected to occur in coming decades. Other areas, where the critical load for mixed forest is seriously exceeded, are the Chongqing-Guiyang area, the Shanghai region and smaller spots in the northern part of the country (Wang et al., 1996). According to the RAINS-Asia calculations of Wang et al. (1996), the critical loads for irrigated farmlands are also exceeded several places.

Zhao et al. (1995) compared different critical load calculations for different soils in China. They used the steady state mass balance (SSMB) method (Sverdrup and de Vries, 1994), the Profile (Warfinge and Sverdrup, 1992) and MAGIC (Cosby et al., 1985) models as well as values obtained in the project RAINS-Asia. The results are shown in Table 4. Zhao et al. (1995) note that the critical load estimates are uncertain and are based on many assumptions; which also can be seen in the differences in the figures from the different methods.

Zhao et al. (1995) also made a case study for Guizhou province. Based on the same critical load approximations and maps as used in RAINS-Asia large parts of the province were found to have a critical load of 200-500 eq/ha/yr (approximately 0.3-0.8 gSm<sup>-2</sup>yr<sup>-1</sup>).

Critical loads are based on the assumption that the molar ratio of Ca<sup>2+</sup> / Al<sup>3+</sup> in soil water is crucial for effects of acidification on trees. Gao and Cao (1989) conducted laboratory experiments in which aluminum toxicity to masson pine seedlings was related to ionic strength, pH and Ca<sup>2+</sup> / Al<sup>3+</sup> ratio. The results indicated that the aluminum toxicity increased with decreasing pH and decreasing ion strength and Ca<sup>2+</sup>-concentration. In another study aluminum toxicity to masson pine seedlings was related to biophysiological parameters and growth; the growth was inhibited markedly at an aluminum concentration of 15 mg/l (Cao et al., 1992).



Table 4 Critical loads for Chinese soils using different methods (from Zhao et al., 1995). Unit:  $\text{gSm}^{-2}\text{yr}^{-1}$ .

	SSMB	Profile	MAGIC	RAINS-Asia
Yellow soil	0.72	1.12	0.6	0.5-1.0
Red soil	0.69	0.87		0.2-0.5
Lateritic red soil	0.66	0.69		0.2-0.5
Latosol	1.04	0.44		0.2-0.5

One should be aware that the basis for critical loads is weak (e.g. Løkke et al., 1996). The present critical load estimates assume ecosystem effects to be directly related to the  $\text{Ca}^{2+} / \text{Al}^{3+}$  ratio and is based on a too simple description of the soil chemistry. The critical load of sulfur will also depend strongly on deposition of base cations. In addition the critical load calculations for China are only preliminary and based on few observations.

Traditional use of forests in China has been gathering of edible plants as well as collection of litter and understory vegetation for use both as domestic fuel and for fertilizing of cultivated land, usually after composting. This practice reduces the fertility of the forests because of the removal of plant nutrients which would otherwise be returned to the soil with the litterfall. This problem has been addressed by scientists in the Dinghushan Biosphere Reserve, who have conducted mass balance studies of nutrients in litterfall and material removed by local residents (Brown et al., 1995; Mo et al., 1995). The removal of nutrients, particularly potassium and calcium by harvesting and litter-collection, should also be considered in connection with estimation of critical loads.

### 2.5. Effects on health

In discussing health effects the focus is not on acid rain per se, but on precursors ( $\text{SO}_2$ ,  $\text{NO}_x$ ) or pollutants which to a large degree originate from the same sources (e.g. particles).

WHO (1995) has given air quality guidelines for Europe for some pollutants and for different averaging times. The maximum allowable annual average for  $\text{SO}_2$  is  $50 \mu\text{g}/\text{m}^3$  and for  $\text{NO}_2$  40 -  $50 \mu\text{g}/\text{m}^3$ . For particles WHO has decided not to give guidelines since there is no evident threshold for effects, while US EPA recently proposed that daily average concentration of particles with diameter less than  $2.5 \mu\text{m}$  ( $\text{PM}_{2.5}$ ) should not exceed  $50 \mu\text{g}/\text{m}^3$  (Reichardt, 1996).

Comparing with Table 1, it is clear that the  $\text{SO}_2$  guideline is exceeded in many places in China, the exceedance may be up to 8 times. In comparison,  $\text{NO}_x$  poses a minor problem at present. The concentrations of airborne particles in many Chinese cities/towns are very large and serious health effects are accordingly expected.

A large number of epidemiological studies in Europe and the USA have shown a significant correlation between level of air pollution and mortality. The correlation seems to be stronger between particles and mortality than between  $\text{SO}_2$  and mortality (Aunan, 1996). In contrast, Wells et al. (1994) found the clearest relationship between  $\text{SO}_2$  and mortality in Beijing and Shenyang.

Concerning morbidity, there seems to be a clear effect of air pollution on respiratory diseases. It is highly probable that air pollution also may increase the frequency of lung cancer, but quantitative relationships are difficult to obtain.

Studies in Western Europe and the USA generally support a linear relationship between SO<sub>2</sub> (or particles) and health effects. There are indications that these functions overstate the response in areas with high pollution levels (WHO, 1995); i.e. the dose-response function levels off as the pollution increases. Studies in Beijing and Shenyang seem to support this conclusion (Wells et al., 1994; Xu et al., 1994b).

In a World Bank study in Chongqing, air pollution was found to be significantly associated with daily mortality from cardiovascular diseases and increased frequency of hospital visits (Xu, 1996). The prevalence of respiratory system illness in the downtown area of Chongqing is very high; 34.3% according to Chang and Hu (1996). Pope and Xu (1993) found a significant, and nearly additive, effect of passive cigarette smoke and coal heating on respiratory symptoms in a study in the Anhui Province. Indoor pollution may be particularly important as a cause of lung cancer in many places in China (Xu et al., 1989).

In addition to effects mentioned above, acidification of soils and waters may lead to mobilization of toxic metals (e.g. Al, Cd), which may be accumulated in vegetables. High dietary intake of fluorides produces mottled teeth and skeletal disorders in both animals and man.

In summary: there are many studies on health effects of air pollutants: but, most of them have been carried out in Western Europe and the USA. The results cannot directly be applied to Chinese conditions.

## 2.6. *Effects on materials*

Atmospheric corrosion of materials is a cumulative, irreversible process taking place also in the absence of pollutants. The reactivity to various air pollutants varies greatly for different materials and pollutants. Direct uptake of SO<sub>2</sub> is the main cause of material damage, but for porous building materials acid rain may also be of importance. High concentrations of sulfur dioxide cause increased corrosion of metals, particularly steel and galvanized steel, and deterioration of calcareous building materials, including concrete and rendering. High ambient temperatures and humidity increase the corrosion and deterioration rates.

More knowledge about deterioration processes and better methods for assessing stock at risk have increased the possibilities for better damage assessment of building materials in recent years (see e.g. Kucera and Fitz, 1995). Wang et al. (1995) developed dose-response functions for material damage based on experiments at different levels of SO<sub>2</sub> and pH in China. Chang and Hu (1996) reported that life-times of outdoor painting in Chongqing are only 1/2 to 1/3 of that found under background conditions. Effects on materials have also been studied in Guangzhou.

Costs associated with material damage may be estimated from assessments of exposed material, and the costs of early replacements or necessary repairs or protective measures, e.g. painting. In many European cities, reduced sulfur dioxide concentration levels have resulted in substantial reductions in

building maintenance costs, and savings in this sector are often comparable to the costs of reducing emissions. (e.g. Kucera et al., 1993). It is likely that this is also the case in many Chinese cities.

### *2.7. Economic losses caused by acid rain and related pollutants*

Cao (1989) estimated that 2.66 million ha. crops are affected by SO<sub>2</sub> and about the half by hydrogen fluoride. causing an economic loss of 550 mill. USD annually. Cao et al. (1990) estimated the yield loss of grain and vegetables to be 2530 and 536 thousand tons in Guangxi and Guangdong provinces. Based on a multiple regression model they further estimated loss of timber to about 4 million tons in the same provinces. Shu et al. (1993) estimated the annual cost of forest damage by acid rain in Guangxi to 650 mill. yuan (~80 mill. USD). Obviously these data have extremely large uncertainties.

Ou et al. (1996) estimated the economic loss due to acid rain damages to crops and materials in the Xiamen area to 53 million RMB yuan (about 6 mill. USD), which equals about 1% of the GNP. The uncertainties are not discussed. Chang and Hu (1996) report that the annual damage from air pollution in Chongqing in 1993 was 1765 million yuan which is 4.4% of GNP. They included damages to health, agriculture, forestry, materials (including additional cleaning costs) as well as increased transportation costs due to reduced visibility.

## **3. Emission control and ongoing research activities**

### *3.1. Information from the State Council of the People's Republic of China*

*On the occasion of the annual World Environment Day, June 5, 1996, the Information Office of the State Council of the People's Republic of China prepared a status report on environmental protection in China. The information in this chapter is from this report, reflecting the Chinese government view on what has been done on environmental issues in China.*

In the 1980s China enacted and implemented a series of principles, policies, laws and measures for environmental protection. China will gradually change strategy from end-of-pipe pollution control to pollution control during the whole industrial process. The Chinese government has drawn up three major policies for environmental protection:

- Putting prevention first and combining prevention with control.
- Making the polluter responsible for treating the emission
- Intensifying environmental management

In addition, policies for comprehensive utilization of resources, technical improvement on industrial emitters, improvement of the urban environment and environmental protection technology and industries. One of the results of these policies is an increase in energy efficiency annually with 5.8% during the 8th 5-year plan (1991-95) in terms of coal consumed per yuan in GDP. Further, China has made substantial scientific and technological achievements in some research areas, such as [...] background value and environmental capacity of the nation's major soils, acid deposition and its impacts and controls [...]. By the end of 1995 some 390 scientific research bodies engaged in environmental protection had been established nationwide, including a comprehensive research

system composed of the Chinese Academy of Sciences. A large number of environment related publications are available: The China Environment News, a national-level professional newspaper on environmental protection was established in 1983, the China Environmental Press (established in 1980) had by 1995 published 5 million copies of books of 860 environment-related titles. China Environment Yearbook has been published annually since 1990, and in English since 1994. In addition several hundred professional periodicals are published.

### *3.2. Law on the control of atmospheric pollution*

“The Law of the People’s Republic of China on the Prevention and Control of Atmospheric Pollution” was adopted on September 5, 1987; entered into force on June 1, 1988 and amended on August 29, 1995.

Important points include:

- NEPA shall set national standards for atmospheric environmental quality and discharge. The national standards may be made more stringent by provincial/autonomous region/municipal government. (Article 6 and 7)
- Anyone who sends a pollutant into the air must follow the local discharge standard. (Article 7)
- Every new construction/expansion/reconstruction must comply with the states regulations on environmental protection. No construction project which fails this may be put into production or use. (Article 10)
- Every unit exceeding the discharge standard must take effective measures and pay a fee. (Article 12)
- Every enterprise shall adopt in priority production techniques which have a high energy utilization efficiency and low discharge of pollutants. (Article 15)
- The central and local governments shall adopt measures to improve urban fuel structure, develop urban gas supply and popularize the production of shaped coal. (Article 22)
- Any newly-constructed coal mines which dig high sulfur and high ash content must install corresponding coal washing and dressing facilities. Already constructed coal mines shall install such equipment within a predescribed period of time. (Article 24)
- The state may designate areas where acid rain or serious SO<sub>2</sub> pollution occur or may occur as acid rain and SO<sub>2</sub> control areas. Thermal power plants and large and medium-sized enterprises within these areas must take measures to remove SO<sub>2</sub> and dust and gradually also NO<sub>x</sub>. (Article 27)
- Prevention and control measures shall be taken to reduce emissions from motor vehicles and ships. Vehicles which emit pollution exceeding the prescribed standards must not be manufactured, marketed or imported. (Article 37)
- The environmental protection department or any other supervision and control department may give a warn or impose a fine on the violator of this law. (Article 39)
- Enterprises failing an emission reduction within the time limit required shall pay an excess pollutant fee, may be fined, ordered to suspend its operation or closed down.

### 3.3. Ongoing projects

In addition to RAINS-Asia, a large number of research projects related to air pollution are going on in China at present. Air pollution in cities and related health effects are of major concern, but several projects are also dealing with problems related to acid rain. We will here mention a few large projects related to acid rain problems.

United Nations development Program (UNDP) has recently (December, 1996) initiated a project titled *Capacity Development for Acid Rain and SO<sub>2</sub> pollution Control in Guiyang* (UNDP, 1996). The main objective is to reduce adverse social, environmental and economic consequences of air pollution and acid rain in the Guiyang area through controlling sulfur dioxide emissions. The project will, based on available research results on acid rain and its effects in southwest China, improve the EPB (Environmental Protection Bureau) officials' and municipal leaders' understanding of the impact of regional acid rain by providing a set of acid rain assessment and prediction methods and models. Where required existing information will be supplemented by additional field observations and studies (UNDP, 1996). Involved institutions are CRAES and Guiyang EPB. The Acid Rain Control Center, located within the Guiyang EPB, will be responsible for data collection etc.

The Norwegian Consortium for Energy and Environment (NORCE) has a NORAD-funded project in Guangzhou called *Air Quality Management and Planning System of Guangzhou*. The project will mainly deal with local air pollution sources and local effects from the pollutants. However, to some extent the project will work with subjects related to acid rain. In addition, UNDP has a new started projects also dealing with local air pollution, especially from motor vehicles, in Guangzhou.

## 4. Conclusions

China's economy is fast developing and the energy demand in the coming decades is likely to increase substantially. There is little doubt that coal will account for most of the increase since other energy carriers will generally be more expensive in China. Hence the sulfur emissions will continue to increase in the coming decades, even if several countermeasures are taken to reduce the emission per energy unit produced. In this situation it is important to present to the authorities assessments of the effects of acid rain as complete and detailed as possible.

This report shows that there has been carried out considerable research on acid rain and related issues in China. However, it is also clear that there still is a strong need for increased knowledge within the fields of acid rain and its effects. Major tasks for acid rain research in China, all central for choosing the best countermeasures, include:

- The relationship between anthropogenic emissions, natural emissions (particularly dust) and the components in the precipitation should be better understood. One important aspect is to expand the focus from only pH and sulfate to also include the other major components, as calcium, magnesium, ammonium and nitrate. Another important aspect is the transportation to, and deposition in, rural areas. With an increasing tendency to move industry out of the cities and building tall stacks, acid deposition will increase in rural areas.
- Long-term changes in the soil chemistry is a serious threat to the ecosystems. The soil acidification processes are poorly understood, which in turn means that estimates of future exceedances of critical loads are very uncertain. A combination of regional surveys, integrated monitoring sites

(including deposition, soils and soil water) laboratory experiments with soils and modelling is necessary to make better predictions of future soil acidification.

- Effort should be made to substitute the primitive steady-state models used in critical load estimates at present with a process-based dynamic model. Model calibration and testing is important to make a useful model; this requires field data for comparison. Such a model would, if incorporated in regional integrated assessment models, as RAINS-Asia, improve the effect part considerably.
- Little is known about effects of long range transported air pollutants on forest and agro-ecosystems. A systematic inventory of effects on forest ecosystems using standardized techniques should be carried out at the national level.
- The critical load concept should be improved and modified to better fit Chinese ecosystems. A few well-designed and controlled experiments with a selection of vegetation species should be carried out, in addition to careful long-term monitoring of changes on several ecosystem levels.
- Water acidification has only been observed in very limited areas. Although it is not likely to become a major problem, additional surveys may be useful. In some areas careful monitoring is recommended.

There is great interest among Chinese and Norwegian scientists in cooperation on these issues. The Chinese authorities have also expressed interest for such cooperation and are clearly aware of acid rain as an important environmental problem. This can for example be seen in the interest for the RAINS-Asia model. However, there is need for improvements since the RAINS-Asia model has its obvious limitations at present, partly due to lack of observations for model calibration and testing.

#### References:

- Akimoto, H. and Narita, H., 1994. Distribution of SO<sub>2</sub>, NO<sub>x</sub> and CO<sub>2</sub> emissions from fuel combustion and industrial activities in Asia with 1° X 1° resolution. *Atmos. Environ.* 28:213-225.
- Arndt, R.L., Carmichael, G.R., Streets, D.G. and Bhatti, N., 1997. Sulfur dioxide emissions and sectorial contributions to sulfur deposition in Asia. *Atmos. Environ.* 31:1553-1572.
- Aunan, K., 1996. Exposure-response functions for health effects of air pollutants based on epidemiological findings. *Risk Analysis* 16: 693-709.
- Bian, Y.-M. and Yu, S.-W., 1992. Forest decline in Nanshan, China. *For. Ecol. Manage.*, 51:53-59.
- Brown, S.; Lenart, M., Mo J., and Kong G., 1995. Structure and Organic Matter Dynamics of a Human-Impacted Pine Forest in a MAB Reserve of Subtropical China. *Biotropica*, 27, 276-289.
- Byrne, J., Shen, B. and Li, X., 1996. The challenge of sustainability. Balancing China's energy, economic and environmental goals. *Energy Policy* 24:455-462.
- Cao Hongfa et al., 1991 Study on economic loss of crops and forest productivity caused by acid deposition in Guangdong province and Guangxi autonomous region, China. p. 110-114.
- Cao Hongfa, 1989. Air pollution and its effects on plants in China. *J. Applied Ecol.* 26:763-773.
- Cao Hongfa, Gao Jixi and Shu Jianmin, 1992. Study on the response of *Pinus massoniana* seedling to aluminium. *Acta Ecologica Sinica*. 12:239-246. (In Chinese with English abstract).
- Cao Hongfa, Gao Xuping and Shu Jianming, 1990. Response of 105 species of plants to simulated sulfur acid rain. *China Environ. Sci.* 1:26-32.
- Cao Hongfa, Gao Yingxing, Shu Jianming, Liu Yangyun, Chen Yanzhi, Wang Wei and Sun Wenshun, 1988. Preliminary study on the relationship between acid precipitation and fir decline of high elevation in the Emei Mountain. *Research of Environ. Sci.* 1:10-11. (In Chinese with English abstract).
- Chang, Y.-S., Arndt, R.L. and Carmichael, G.R., 1996. Mineral base-cation deposition in Asia. *Atmos. Environ.* 30:2417-2427.
- Chang Yongguan and Hu Tao, 1996. The economic cost of damage from atmospheric pollution in Chongqing. Manuscript, Chongqing Environmental Protection Bureau.
- Cosby B.J., Hornberger, G.M., Galloway, J.N. and Wright, R.F., 1985. Modeling the effects of acid deposition assessment of a lumped parameter model of soil water and stream water chemistry. *Water Resources Res.* 21:51-63.

- Dai, Z., Liu, Y., Wang, X., Zhao, D., 1997. Decrease in soil pH in southern China within about 30 years. Submitted Water Air Soil Pollut.
- EMAD (Expert Meeting on Acid Deposition), 1997. Final report from the Fourth Expert Meeting on Acid Deposition and Monitoring Network in East Asia, 4-6 February, 1997. Environ. Agency of Japan, Hiroshima, Japan.
- Galloway, J.N., Zhao, D., Thomson, V.E. and Chang, L., 1996. Nitrogen mobilization in the United States of America and the Peoples Republic of China. *Atmos. Environ.*, 30:1551-1561.
- Gao Jixi and Cao Hongfa, 1989. Effects of ionic strength, pH and Ca/Al ratio on aluminum toxicity of masson pine seedlings. *Acta Scientiae Circumstantiae*. 11:194-198. (In Chinese with English abstract).
- Hashimoto, Y., Sekine, Y., Kim, H.K., Chen, Z.L. and Yang, Z.M., 1994. Atmospheric fingerprints of East Asia, 1986-1991. An urgent record of aerosol analysis by the JACK network. *Environ.* 28:1437-1445.
- Hettelingh J-P., Sverdrup H. and Zhao D., 1995. Deriving Critical Loads for Asia. *Water, Air and Soil Pollution*, 85: 2565-2570.
- Kucera, V. and Fitz, S., 1995. Direct and indirect air pollution effects on materials including cultural monuments. *Water Air Soil Pollut.*, 85:153-165.
- Kucera, V., Henriksen, J., Knotkova, D., Sjöström, C., 1993. Model for calculation of corrosion cost caused by air pollution and its application in three cities. In: Costa, J.M. and Mercer, A.D.(eds.): progress in the understanding and Prevention of Corrosion., Volume I, pp 24. London, Institute of Materials.
- Larssen, T., Vogt, R.D., Seip, H.M., 1996. A comparison of soil- and water chemistry in a catchment in China with sites in Poland and Norway. In Pawlowski, L. (ed.) *Chemistry for the Protection of the Environment II*. Plenum press. pp. 421-434.
- Larssen, T., Xiong, J., Vogt, R.D., Seip, H.M., Liao, B. and Zhao, D., 1997. Studies of soils, soil water and stream water at a small catchment near Guiyang, China. *Water Air Soil Pollut.* (in press)
- Lei, H.-C., Tanner, P.A., Huang, M.-Y., Shen, Z.-L. and Wu, Y.X., 1997. The acidification process under the cloud in Southwest China Observation results and simulation. *Atmos. Environ.*, 32:851-861.
- Liao Ruizhang, Cao Hongfa, Liu Liangui and Gao Jinlan, 1991. Study on the toxicity of acid rain to microbiota in soils. *J. Environ. Sci. (China)* 3:17-26.
- Liao, B., Larssen, T., Seip, H.M. and Vogt, R.D., 1994. Anion adsorption and aluminum release from Chinese soils treated with different concentrations of  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ . *J. Ecol. Chem.*, 3:281-301.
- Liao, B., Seip, H.M. and Larssen, T., 1997a. Response of two Chinese forest soils to acidic inputs: Leaching experiment. *Geoderma* 75:53-73.
- Liao, B., Seip, H.M. and Larssen, T., 1997b. Response of five Chinese forest soils to acidic inputs: Batch experiment. Manuscript submitted *Geoderma*.
- Lin Gouzhen, Liao Bohan and Ding Ru, 1992a. A study on the buffering capacity of the forest soil against acidic precipitation in several areas of China. *Chinese Geographical Sci.*, 2:126-132.
- Lin Ziyu et al., 1992b. Study of the transformation rate of  $\text{SO}_2$  and aerosol formation in the ambient air of Guangzhou city. *Research of Environ. Sci.* 5:23-24. (In Chinese with English abstract).
- Liu Houtian and Du Xiaoming, 1991. A preliminary study on the characteristics of fogwater in the masson pine forest in Chongqing, China. *J. Environ. Sci. (China)*. 3:11-16.
- Liu Shuairan and Huang Meiyuan, 1993. The effects of below-cloud aerosol on the acidification process of rain. *J. Atmos. Chem.* 17:157-178.
- Løkke, H., Bak, J., Falkengren-Grerup, U., Finlay, R.D., Ilvesniemi, H., Nygaard, P.H. and Starr, M., 1996. Critical loads of acidic deposition for forest soils. Is the current approach adequate? *Ambio*. 25:510-516.
- Ma Liang-Qing, 1990 Absorption capacity of forest to  $\text{SO}_2$  as well as injury of acid rain and atmospheric pollution to it in Chongqing. Report.
- Meng Fan, Wang Zhenkui, Ge Ming, 1996. Study on simulation of  $\text{SO}_x$  transport-deposition in Southeastern China. *Research of Environ. Sci.* 9(5):20-25. (In Chinese with English abstract)
- Meng Fan, Xu Jun, Zhang Jing and Wang Zhenkui, 1997. Sulfur transport/deposition modeling in southeastern China. Manuscript. Chinese Research Academy of Sciences, Beijing, China.
- Mo J., Brown, S., Lenart, M., and Kong G.(1995). Nutrient Dynamics of a Human-Impacted Pine Forest in a MAB Reserve of Subtropical China. *Biotropica*, 27, 290-304.
- Ning, D.-T., Zhong, L.-X. and Chung, Y.-S., 1996. Aerosol size distribution and elemental composition in urban areas of northern China. *Atmos. Environ.* 30:2355-2362.
- Ou Shouming, Pan Liqing and Zhuang Mazhan, 1996. Estimation of the effects and economic loss caused by acid deposition in Xiamen area. *Research of Environ. Sci.* 9(5):57-59. (In Chinese with English abstract)
- Pan, G.X., 1992. Acidification of Soils in Mount Lushan over the Last 35 Years. *Pedosphere*, 2:179-182.
- Pope III, C.A. and Xu, X., 1993. Passive cigarette smoke, coal heating and respiratory symptoms of nonsmoking women in China. *Environm. Health Perspectives* 101: 314-316.

- Qi Liwen and Wang Wenxing, 1990. The character of precipitation at Guangdong and Guangxi provinces in China. Report/Manuscript. Chinese Research Academy of Environmental Sciences, Beijing, China.
- Reichhardt, T., 1996. New US air pollution regulations are set for a bumpy ride. *Nature*, 384, 392.
- Rodhe, H., Galloway, J. and Zhao, D., 1992. Acidification in Southeast Asia - Prospects for the Coming Decades. *Ambio* 21:148-150.
- Seip, H.M., Zhao, D., Xiong, J., Zhao, D., Larssen, T., Liao, B. and Vogt, R.D., 1995. Acidic deposition and its effects in southwestern China. *Water Air Soil Pollut.* 85:2301-2306.
- Shao Min, Tang Xiaoyan, Li Jinglong, Li Kun and Yuan Sixum, 1995. The application of accelerator mass spectrometry (AMS) in the study of source identification of aerosols in China. *Pure & Appl. Chem.* 67:1457-1459.
- Shu Jianmin, Cao Hongfa, Gao Yingxing, Liu Liangui, Gao Jixi, Yong Yanjun, 1993. Ecological criteria for masson pine effected by acidic precipitation and evaluation of economic loss. *China Environ. Sci.* 13(4):279-283.
- Singer, A., 1988. Properties and Genesis of Some Soils of Guangxi Province, China. *Geoderma*, 43:117-130.
- Singer, A., 1993. Weathering patterns in representative soils of Guangxi province, south-east China, as indicated by detailed clay mineralogy. *J. Soil Sci.* 44:173-188.
- Statistics China, 1983-1996. China Environmental Statistics Annual Report, Beijing, China.
- Suzuki, K., Maeda, K., Sasa, Y., Okada, A., Sakamoto and Ozawa, 1993. Application of PIXE to source identification of Kosa aerosol: analysis of desert soils in China. *Nucl. Instr. and Meth.* B75:317-320.
- Sverdrup, H. and de Vries, W., 1994. Calculating critical loads for acidity with the simple mass balance method. *Water Air Soil pollut.*, 72:143-162.
- Totsuka, T., Liang Yang, Fuzhu Zhang and Zhongwei Feng, 1994. Effects of acid precipitation on growth of masson pine trees in Chongqing area, China. In: Proceedings of China-Japan joint symposium: Impacts of Salination and Acidification on Terrestrial Ecosystems and Their Rehabilitation in East Asia. pp.142-145.
- UNDP, 1991. Control of Atmospheric Pollution from Coal Combustion. United Nations Development Programme. Project document CPR/91/213/B/01/99.
- UNDP, 1996. Capacity Development for Acid Rain and SO<sub>2</sub> Pollution Control in Guiyang. United Nations Development Programme. Project document CPR/96/304/A/01/99.
- Wang Rusong, Yang, J. and Tang, H., 1996. Impacts of Acid Deposition on Ecosystems in China. Research report, Research Center for Eco-Environmental Sciences, Academia Sinica, P.O.Box 934, Beijing 100083, China.
- Wang Wei, Wang Wenxing, Chen Yanzhi, Qi Liwen, Cai Yiqi, Ye Qin and Huang Xingyuan, 1991. Chemical properties of cloud water in Jinding, Ermei mountain. *China Environ. Sci.*, 11(4):245-246. (In Chinese with English abstract).
- Wang Wei, Zao De Shan, Cheng Yanzi, Tang Dagang, Jiang Zebguan, Han Yingjian and Ning Jie, 1988. Research of interaction between aerosol and precipitation. *Research of Environ. Sci.* 1(1):43-44. (In Chinese with English abstract).
- Wang Wenxing and Tao Wang, 1996. On acid rain formation in China. *Atmos. Environ.* 30:4091-4093.
- Wang Wenxing, Fu Chen, Yutian Zhang, Shaoxian Hong, Dagang Tang and Wanhua Zhang, 1997a. The current state of acid rain in China. Report. Chinese Research Academy of Sciences, Beijing, China.
- Wang Wenxing, Guoan Ding and Tao Wang, 1997b. Geographical distribution of rain acidity over China. Submitted to *Water Air Soil Pollut.*
- Wang Wenxing, Hong Shaoxian and Zhang Wanhua, 1995. Development of material damage function for acid deposition. *Acta Scientiae Circumstantiae.* 15(1):30-31. (In Chinese with English abstract)
- Wang, A., Huang, Y., Ma, C., Wang, Q., Yang, S., Liu, H., Li, M. and Liu, J., 1981. Chemical characteristics of airborne particules in the Beijing area. *Acta Scientiae Circumstantiae.* 1:220-232. (In Chinese)
- Warfinge, P. and Sverdrup, H., 1992. Calculating critical loads of acid deposition with PROFILE - a steady state soil chemistry model. *Water Air Soil Pollut.*, 63:119-143.
- Wells, G.J., Xiping, X., Johnson, T.M., 1994. Valuing the health effects of air pollution: Application to industrial energy efficiency projects in China. Report no. 8 of China: Issues and Options in Greenhouse Gas Emission Control. The World Bank.
- WHO, World Health Organization, 1995. Update and revision of the Air Quality Guidelines for Europe. Meeting of the Working Group "Classical" Air Pollutants, Bilthoven, The Netherlands, 11-14 October 1994.
- Williams, M.W., Yang, D., Liu, F., Turk, J. and Melack, J.M., 1995. Controls on the major ion chemistry of the Ürümqi river, Tian Shan, Peoples Republic of China. *J. Hydrol.* 172:209-229.
- Xia, Yicheng, Sakamoto, M., Qijun Kuang, Dehui Zhuang, Baoyuan Liu, Zhihong Lei, Xiamin Liang and Xiaoqing Xu, 1994. Effects of acidification on freshwater ecosystem in southwest China. In: Proceedings of Cina-Japan joint symposium: Impacts of Salination and Acidification on Terrestrial Ecosystems and Their Rehabilitation in East Asia. pp.152-158.
- Xu Guangshan, Zhouping Sun and M. Okazaki, 1994a. Effects of acid precipitation on some soils under different vegetation in Chongqing, China. In: Proceedings of China-Japan joint symposium: Impacts of Salination and Acidification on Terrestrial Ecosystems and Their Rehabilitation in East Asia. pp.173-178.
- Xu Xiping, 1996. Air Pollution and its Health Effects in Chongqing, China. Presentation at World Bank.



- Xu, X., Dockery, D. and Chen, Y., 1994b. Air pollution and daily mortality in residential areas of Beijing, China. *Archives Environm. Health*, 49:216-222.
- Xu, Z-Y., Blot, W.J., Xiao, H.-P., Yi-Ping Feng, A.W., Stone, B.J., Sun, J., Ershow, A.G., Henderson, B.E. and Fraumeni, J.F., 1989. Smoking, air pollution, and high rates of lung cancer in Shenyang, China. *J. National Cancer Institute*, 81:1900-1806.
- Xue, H.B., and Schnoor, J.L., 1994. Acid deposition and lake chemistry in Southwest China. *Water Air Soil Pollut.*, 75:61-78.
- Yin, Sha, Wang Anpu, Yang Shulan, Liu Pingsheng, 1996. Correlation of acid rain with the distributions of acid and alkaline elements in aerosols. *Nucl. Instr. and Meth. B109/110*:551-554.
- Zhang, F., Zhang, J., Zhang, H., Ogura, N. and Ushikubo, A., 1996. Chemical composition of precipitation in a forest area of Chongqing, Southwest China. *Water Air Soil Pollut.* 90:407-415.
- Zhao Dawei, Seip, H.M, Zhao Dianwu, and Zhang Dongbao., 1994. Pattern and cause of acidic deposition in the Chongqing region, Sichuan Province, China. *Water Air Soil Pollut.*, 77:27-48.
- Zhao Dianwu and Seip, H.M., 1991. Assessing effects of acid deposition in southwestern China using the MAGIC model. *Water Air Soil Pollut.*, 60:83-97.
- Zhao Dianwu and Sun Bozhen, 1986. Air pollution and acid rain in China. *Ambio* 15:2-5.
- Zhao Dianwu and Wang Anpu, 1994. Estimation of anthropogenic ammonia emissions in Asia. *Atmos. Environ.*, 28:689-694.
- Zhao Dianwu and Xiong Jiling, 1988. Acidification in Southwestern China. In: Rodhe, H. and Herrera, R.(eds.) *Acidification in tropical countries*, SCOPE, John Wiley & Sons.pp. 317-346.
- Zhao Dianwu and Zhang Xiaoshan, 1990. Energy related environmental issues in China: Air pollution and acid rain. Report, Res. Center for Eco-Environ. Sci., Beijing. Presented at Fukuoka Int. Symp. '90: Global environ, & Energy issues, 1990, Japan.
- Zhao Dianwu, Chen Chuyin, Xiong Jiling, Zhang Xiaoshan, Dai Zhaohua, Mao Jietai, Seip, H.M. and Vogt, R., 1995/6. Acid reign 2010 in China?.
- Zhao Dianwu, Mao, j., Xiong, J., Zhuang, X., Yang, J., 1995. Critical load of sulfur deposition for ecosystem and its application in China. *J.Environ. Sci.*, 7:325-337.
- Zhao Dianwu, Xiong Jiling, Xu Yu and Chan, W.H., 1988. Acid rain in southwestern China. *Atmos. Environ.* 22:349-358.

# **APPENDIX B**

**Table B.1** Sampling program during the China visit. **1:** Sampling volatile organic air compounds (VOC), **3:** Stream invertebrate sampling; **4:** Stream diatoms sampling; **5:** Measurements of chlorophyll fluorescence of coniferous trees and high frequent vascular plants; **6:** Sampling of coniferous tree needles; **7:** Soil sampling for texture/density analysis; **8:** Soil sampling for soil chemistry; **9:** Soil water sampling for main chemistry; **10:** measured pH, Conductivity and water temperature at site; **11:** Collected samples for main water chemistry and trace metals; **12:** Extra samples for intercalibration purpose.

Area, site name	Sites	Date	Location	NILU	NINA	NISK	UjO	NIVA
Nanshan west	I	17/04/97	29°32'N, 106°39'E	1		8	8	
Nanshan east	II	"			5, 6	8	8	
Tie Shan Ping	I	18/04/97	29°38'N, 106°41'E	1	3,4,5,6	6,7,8	8,9	10
	A	"			3,4			10,11
	B	"			3,4			10,11
Simian Shan, Piao Chang Qoy	A	19/04/97	28°40'N, 106°22'E	1	3,4,5,6	6,7,8	8	10,11
	B	20/04/97			3,4			10,11
Liu Chong Guang	II-a	23/04/97	26°34'N, 106°38'E	1	3,4,5,6	6,7,8	8,9	10,11
	II-b	"						10,11
	III-1	23/04/97						10,11
	III-2	28/04/97					8,9	10
	IV	28/04/97						10
	V	23/04/97						10
Leishan	I	26/04/97	26°10'N, 108°10'E	1	3,4,5,6	?6,8	8	10,11
	II	"			3,4			10,11
	IV	"			5,6	6,8	8	
On a ridge along the road	V	"						10
Bridge, 20 km downstream	VII	"						10
Leishan City, main river								
Dinghushan, Qingyun Temple	I	02/05/97	23°09'N, 112°30'E	1	3,4	8	8	10,11
Grass Pond	II	"			5,6	6,8	8	
Tang e Ling	III	02/05/97			5,6	6,8	8	
Tianhu Lake	IV	03/05/97			5,6	6,8	8	10,11, 12
Heshan Forest, Catchment 2.	I	"	22°41'N, 112°54'E	1	3,4,5,6	6,8	8	10,11
Guangzhou Botanical Garden	I	04/05/96			5,6			
Baiyun Shan	I	05/05/97	23°03'N, 113°19'E	1	5,6	6,8	8	10,11
Liu Xi river	I	06/05-97	23°28'N, 113°10'E	1	5,6	6,8	8	10,11

# **APPENDIX C**

**Table C.1.** Site and soil information about the sampled forest stands. ( LCG = Liu Chong Guang near Guiyang.) Data include the soil content of organic carbon (C) and of nitrogen (N) in % of the fine earth fraction ( $< 2\mu\text{m}$ ), the effective cation exchange capacity ( $\text{CEC}_E$ ) in  $\text{meq kg}^{-1}$  of fine earth and the proportion of the exchange sites occupied by base cations (BS, i.e. the sum of calcium, magnesium, potassium and sodium) and aluminum (AIS).

Site	Vegetation	Soil horizon	Depth cm	tot. C %	tot. N %	C/N	$\text{CEC}_E$ $\text{meq kg}^{-1}$	BS %	AIS %
Nanshan east	Masson pine	A	0 - 3	4.07	0.27	15.0	47.1	9.9	89.0
		B	3 - 6	1.02	0.10	10.5	43.5	9.5	89.4
Nanshan west	Masson pine	A	0 - 4	1.33	0.11	11.9	40.9	18.6	80.0
		B	4 - 8	0.58	0.05	11.1	34.4	11.7	87.2
Tie Shan Ping 2	Masson pine	A	0 - 3	7.44	0.46	16.1			
		B1	3 - 13	1.35	0.12	11.0			
		B2	13 - 28	0.74	0.09	8.1			
Simian Shan	Chinese fir	A	3 - 7	14.06	0.83	17.0	101.6	8.5	90.4
		B1	7 - 13	4.81	0.31	15.6	53.2	7.6	91.9
		B2	13 - 18	2.87	0.19	15.2	33.2	6.4	93.1
LCG A	Masson pine Chinese fir	AO	0-4	14.89			138.0	28.4	66.5
		AB	4-11	1.98			106.4	18.6	77.3
		B2	>11	0.68			94.1	12.5	85.0
LCG B	Broadleaves	OA	0-4	24.06			223.2	42.9	39.3
		AB	4-12	1.90			84.1	21.0	75.5
		B	12-40	3.63			63.8	22.0	75.5
LCG C	Masson pine	A1	0-15	3.38			81.4	9.0	87.6
		A2	15-30	3.61			57.6	4.2	93.4
		BC1	30-60	0.67			26.2	3.7	93.9
		BC2	>60	0.68			24.5	4.1	93.3
LCG D	Masson pine	A	0-4	4.57			99.0	20.6	77.1
		B1	4-22	1.29			88.3	15.4	82.3
		B2	22-38	0.62			83.8	17.4	80.6
		C	38-60	1.10			81.5	40.3	57.8
LCG E	Masson pine	OA	0-2	9.23			137.0	59.5	38.1
		B	2-30	1.09			57.8	6.7	91.9
		C	30-50	0.24			72.4	6.1	92.3
LCG F	Broadleaves	OA	0-2	8.36			188.4	25.0	71.1
		B	2-22	0.88			141.9	13.6	84.1
LCG G	Masson pine Chinese fir	A	0-5	7.70			124.7	9.7	83.2
		B2	5-37	1.03			90.7	5.9	92.5
		C	37-60	0.45			90.8	5.9	92.5
LCG H	Masson pine	A	4 - 8	3.61	0.28	12.9			
		AB	11 - 15	2.31	0.20	11.5			
		B1	30 - 34	0.80	0.08	10.4			
		B2	55 - 59	0.59	0.08	7.9			
Leishan 1	Chinese Fir	A	3 - 7	12.67	1.08	11.7	71.8	63.1	36.6
		AB	7 - 27	6.79	0.64	10.6	44.1	29.8	69.7
		B	> 27	1.81	0.27	6.6	10.9	24.4	74.9
Leishan 2	Broadleaves	A	4 - 14	14.73	1.37	10.7	124.1	75.9	23.8
		AB	14 - 44	6.39	0.63	10.1	37.6	23.7	75.9
		B	> 44	4.09	0.45	9.1	13.0	36.8	62.5

**Table C.1. Continues**

Site	Vegetation	Soil	Depth	tot. C %	tot. N %	C/N	CEC meq kg <sup>-1</sup>	BS %	AIS %
Leishan 3	Masson pine,	A	1 - 5	8.16	0.62	13.2	75.6	17.2	82.0
	Chinese fir,	AB	5 - 15	7.99	0.60	13.3	69.4	9.1	90.3
	Broadleaves	B	15 - 45	3.77	0.36	10.6	46.9	6.8	92.7
Dingushan A	Monsoon	A	1 - 4	5.33	0.44	12.1	78.9	6.7	92.4
	Evergreen	AB	4 - 14	2.07	0.22	9.3	54.6	4.2	95.0
		B	14 - 40	0.51	0.10	5.1	33.6	4.1	94.9
Dingushan B	Mixed forest	Ap	10 - 30	1.42	0.15	9.5	38.3	5.3	93.7
		B	45 - 65	0.37	0.08	4.8	26.4	6.2	92.8
Dingushan C	Masson pine	A	0 - 5	3.87	0.33	11.8	56.8	9.5	89.4
		B	5 - 30	1.09	0.14	8.0	37.0	4.6	94.5
Heshan	Bamboo	A	0 - 5	2.75	0.23	11.8	41.6	12.9	86.3
		B	5 - 15	0.75	0.09	8.5	38.8	4.2	95.1
Baiyun Shan	Masson pine	A	1 - 4	2.35	0.17	13.9	47.2	9.0	89.6
		AB	4 - 27	0.80	0.08	9.5	23.4	4.9	94.0
		B	30 - 45	0.40	0.07	5.9	22.5	3.4	95.7
Liu Xi River	Bamboo,	A	1 - 5	1.75	0.15	11.5	17.4	31.1	67.9
	Masson pine	AB	5 - 25	0.93	0.08	12.0	14.4	13.8	85.2
		B	40 - 50	0.41	0.03	12.5	11.0	15.8	82.9

**Table C.2.** Soil samples from a 275 meter long transect across the Liu Chong Guang site (Guiyang) with limited analyses. The site denomination gives the distance in meters along the transect from east to west.

Site	Vegetation	Soil horizon	Depth cm	tot. C %	tot. N %	C/N
0	Masson pine	A	7 - 8	7.24	0.41	17.5
		C	10 - 20	0.89	0.08	10.7
25	Masson pine, Chinese fir	A	6 - 7	8.58	0.47	18.2
		B	7 - 17	1.69	0.13	13.0
50	Masson pine	H layer	8 - 11	26.40	1.47	18.0
		Colluvium	11 - 23	3.19	0.23	14.1
		A (buried)	23 - 31	3.18	0.18	17.7
		B	>31	1.21	0.10	11.9
75	Masson pine, Maple (?)	H layer	8 - 11	43.44	2.43	17.9
		AB	12 - 17	7.79	0.56	14.0
		B	>17	1.87	0.16	12.0
100	Masson pine	A	15 - 21	3.85	0.24	16.1
		B1	21 - 31	1.44	0.13	10.9
125	Masson pine	H layer	4 - 5	22.99	1.46	15.7
		Colluvium	5 - 7	3.57	0.25	14.4
		A (buried)	7 - 12	3.59	0.22	16.1
150	Broadleaves	A	0 - 10	4.02	0.25	15.9
		AB	12 - 23	2.07	0.17	12.0
		B	>23	0.48	0.08	6.4
175	Masson pine, Broadleaves	A	2 - 5	2.83	0.25	11.2
		AB	5 - 30	1.62	0.15	11.1
		B	>30	0.54	0.09	6.4
200	Masson pine	H layer	13 - 15	23.69	1.32	17.9
		AB	16 - 31	2.79	0.21	13.1
		B	>31	0.49	0.08	6.4
225	Broadleaves	A	0 - 2	7.14	0.59	12.1
		AB	2 - 24	1.31	0.14	9.1
		B	>24	0.86	0.11	8.1
250	Broadleaves	A	4 - 11	13.02	0.85	15.3
		AB	12 - 28	2.82	0.19	14.6
		B	>28	0.93	0.10	9.2
275	Masson pine	F & H layer	2 - 6	35.18	1.96	18.0
		A	6 - 9	5.12	0.31	16.7
		AB	9 - 19	1.84	0.13	14.2
		B	> 19	0.70	0.08	8.5

**Table C.3.** The content of selected elements in current (1997) and one year old needles (1996) of Chinese fir and Masson pine. Values are mmol kg<sup>-1</sup> of dry weight.

Site	Vegetation		Al	Ca	K	Mg	N	P	Ca/Al
	Specie	Year							
Tie Shan Ping		1996	5.55	280.4	176.7	50.8	754	24.2	50.5
Tie Shan Ping		1997	5.51	295.3	176.0	54.2	1084	35.2	53.6
Liu Chong Guang		1996	6.18	253.7	264.5	41.3	1076	29.1	41.1
Liu Chong Guang		1996	3.14	403.9	214.7	85.3	972	39.5	128.6
Liu Chong Guang		1997	2.00	281.9	237.8	80.3	1080	43.1	141.0
Liu Chong Guang		1997	3.50	185.6	337.7	54.1	1118	47.6	53.0
Dinghushan	Chinese	1996	4.59	263.9	175.9	88.4	1078	23.9	57.5
Dinghushan	Fir	1997	3.70	254.1	220.0	107.2	1226	28.1	68.7
Heshan		1996	4.24	537.6	173.9	56.0	905	27.2	126.8
Heshan		1997	4.05	375.7	176.5	53.8	1072	31.1	92.8
Baiyun Shan		1996	6.30	559.2	248.8	72.3	998	31.6	88.8
Baiyun Shan		1997	6.41	525.7	214.2	83.9	1073	37.1	82.0
Liu Xi River		1996	2.60	391.3	275.6	59.9	998	24.7	150.5
Liu Xi River		1997	1.91	332.6	264.7	66.3	1062	33.8	174.1
Tie Shan Ping		1996	11.08	164.1	135.6	58.3	1125	32.0	14.8
Tie Shan Ping		1996	6.48	78.4	186.9	41.9	1060	64	12.2
Liu Chong Guang			15.85	201.5	183.7	52.5	1008	33.2	12.7
Liu Chong Guang		1996	12.66	139.3	129.3	54.9	1206	36.3	11.0
Liu Chong Guang		1997	12.63	177.0	216.7	62.3	959	39.2	14.0
Liu Chong Guang		1996	10.71	250.1	83.7	30.9	968	23.6	23.4
Liu Chong Guang		1997	6.87	79.7	146.0	24.5	951	47.7	11.6
Dinghushan	Masson	1996	9.68	311.7	130.7	52.8	1033	26.0	32.2
Dinghushan	Pine	1997	8.49	112.4	233.8	50.2	971	51.8	13.2
Heshan		1997	8.84	78.4	191.3	49.9	858	46.2	8.9
Heshan		1996	14.29	123.5	94.1	42.2	884	21.3	8.6
Baiyun Shan		1996	12.55	100.2	90.6	41.3	1180	23.9	8.0
Baiyun Shan		1996	11.19	114.8	103.7	45.7	1050	16.0	10.3
Baiyun Shan		1997	5.71	63.1	288.7	44.4	950	31.9	11.1
Liu Xi River		1996	7.33	137.0	85.2	53.7	924	20.4	18.7
Liu Xi River		1997	2.93	92.4	169.8	42.6	813	29.9	31.5



**Table C.4.** The stores of nitrogen in four selected forest soils. NB Two of the soils were sampled to 15 and 18 cm depth only. «dens» indicates soil density. Liu Chong Guang site is named Guiyang.

Horizon	Tie Shan Ping 2			Simian Shan			Guiyang H			Heshan		
	depth cm	Masson pine		depth cm	Mixed		depth cm	Masson pine		depth cm	Bamboo	
		dens. kg dm <sup>-3</sup>	N store kg ha <sup>-1</sup>		dens. kg dm <sup>-3</sup>	N store kg ha <sup>-1</sup>		dens. kg dm <sup>-3</sup>	N store kg ha <sup>-1</sup>		dens. kg dm <sup>-3</sup>	N store kg ha <sup>-1</sup>
A	0 - 3	1.25	<b>1729</b>	3 - 7	0.57	<b>1881</b>	4 - 8	1.05	<b>1176</b>	0 - 5	1.25	<b>1463</b>
AB							8 - 15	1.32	<b>1857</b>			
B1	3 - 13	1.25	<b>1538</b>	7 - 13	0.57	<b>1057</b>	15 - 34	1.32	<b>1931</b>	5 - 15	1.42	<b>1250</b>
B2	13 - 28	1.34	<b>1829</b>	13 - 18	0.92	<b>869</b>	34 - 59	1.32	<b>2475</b>			
TOTAL			<b>5095</b>			<b>3807</b>			<b>7439</b>			<b>2712</b>

**Table C.5.** Volumetric soil moisture content (in %) as a function of pF (logarithm of the water potential in mbar). pF = 2.0 represents field capacity and pF = 4.2 represents wilting point. Plant available water is water that can be extracted from the soil between pF 2.0 and pF 4.2. «Field» indicates the soil moisture content at the time of sampling. Liu Chong Guang site is named Guiyang.

pF	Tie Shan Ping 2		Simian Shan		Guiyang Plot C			Guiyang Plot H		Heshan	
	5-10 cm	15-20 cm	5-10 cm	15-20 cm	5-10 cm	20-25 cm	50-55 cm	A	AB	0 - 5 cm	20-25 cm
0.0	49.7	47.3	69.3	58.4	64.3	62.1	63.4	59.1	47.6	52.6	47.1
1.3	38.7	38.9	44.9	44.6	48.4	49.1	53.0	48.2	38.8	40.2	38.7
2.0	36.4	36.3	39.9	38.6	43.2	43.6	47.6	45.2	36.5	36.8	36.1
3.0	34.4	34.5	37.4	35.7	39.8	40.4	45.2	43.2	34.2	34.6	34.1
4.2	22.3	25.3	9.1	11.8	10.6	11.5	13.2	13.1	18.4	25.4	31.5
Field	35.9	35.8	39.9	38.3	42.5	43.7	48.2	45.9	36.6	34.5	34.1

**Table C.6.** Concentration of major chemical compounds in Chinese streams in April/May 1997.  
 Units: Temperature : °C; Conductivity (Cond):  $\mu\text{s cm}^{-1}$ ;  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Al}^{\text{nt}}$ ,  $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$ ,  $\text{Cl}^-$ ,  $\text{NO}_3^-$ , F, alkalinity (Alk), sum cations ( $\Sigma\text{Cat}$ ), sum anions ( $\Sigma\text{An}$ ), charge balance (CB), acid neutralizing capacity (ANC):  $\mu\text{eq L}^{-1}$ ;  $\text{Al}_\text{f}$ ,  $\text{Al}_\text{o}$ ,  $\text{Al}_\text{i}$ , organic nitrogen (org-N):  $\mu\text{g L}^{-1}$ ; Total organic carbon (TOC) and  $\text{SiO}_2$ :  $\text{mg L}^{-1}$ ; Turbidity: FTU; Ionic strength (IS):  $\mu\text{molar}$ .

Site	Chongqing area				Guiyang area					Guangzhou area				
	Tie Shan Ping		Simian Shan		Liu Chong Guang			Leishan		Dingushan		Heshan	Baiyun Shan	Liu Xi
	A	B	A	B	II	I	III	I	II	I	IV			
Temp	17.2	16.4	13.0	12.6	12.8	12.6	12.8	11.5	12.1	21.9	22.9	23.2	22.7	23.2
Cond	65.4	79.4	63.5	113	175	108	119	11.8	11.9	55.9	30.1	25.0	48.5	11.7
Turb	0.32	0.13	0.85	2.00	0.35	0.24	0.54	2.60	3.70	0.25	0.35	8.40	4.10	0.29
pH	5.34	4.89	6.83	7.54	4.71	4.68	4.61	6.75	6.71	4.27	6.54	6.91	7.10	6.42
$\text{Ca}^{2+}$	298	341	434	898	768	459	454	25.5	26.0	33.4	99.8	38.4	142	19.5
$\text{Mg}^{2+}$	128	136	64.2	125	577	253	338	14.8	15.6	48.5	40.3	51.8	92.1	6.58
$\text{Na}^+$	43.9	54.8	33.5	46.5	26.5	26.5	28.7	60.9	61.3	46.5	32.2	101	149	49.6
$\text{K}^+$	16.4	23.8	13.6	15.9	39.1	34.0	32.7	3.07	2.81	7.67	6.39	39.9	51.2	12.8
$\text{Al}^{\text{nt}}$	7.4	14.5	0.3	0.0	72.2	46.2	68.7	0.1	0.1	86.2	0.1	0.0	0.0	0.6
$\text{NH}_4^+$	1.07	3.36	0.43	0.36	0.36	0.36	0.36	0.36	0.36	0.36	3.36	0.36	1.07	0.29
$\text{Al}_\text{f}$	127	204	64	19	1010	600	868	5	5	939	16	15	11	24
$\text{Al}_\text{o}$	6	8	55	11	37	18	31	5	5	20	5	12	7	20
$\text{Al}_\text{i}$	121	196	9	8	973	582	837	0	0	919	11	3	4	4
$\text{SO}_4^{2-}$	408	475	373	433	1415	781	885	25.0	29.2	75.0	119	39.6	56.2	31.2
$\text{Cl}^-$	59.2	79.0	11.3	14.1	28.2	19.7	19.7	5.64	5.64	67.7	39.5	28.2	59.2	14.1
Alk	9.95	3.26	122	552	3.88	0.27	2.93	62.6	53.6	0.00	39.7	172	296	23.8
F	1.05	3.16	0.53	0.53	17.37	7.90	9.47	0.53	0.53	5.26	1.58	0.53	5.79	6.84
$\text{NO}_3^-$	40.7	57.5	36.4	65.3	51.4	19.3	27.1	14.6	22.1	35.3	48.9	0.36	36.4	17.1
$\text{SiO}_2$	4.9	5.4	4.8	5.0	7.1	6.6	6.9	9.7	9.0	6.2	3.1	13.7	18.3	10.0
TOC	0.54	0.64	2.00	2.10	1.80	0.81	0.95	0.20	0.28	0.74	0.60	1.20	1.30	1.10
SCat	500	586	546	1086	1503	840	947	105	106	276	182	232	435	89.7
SAn	519	618	542	1065	1517	828	944	108	111	183	248	240	454	93.1
CB	-19	-32	4	21	-14	12	3	-3	-5	93	-66	-8	-19	-3.4
IS	926	1083	950	1610	2921	1605	1822	118	128	377	338	248	487	123
ANC	-21	-56	125	573	-85	-48	-79	59	49	-42	-28	163	282	26
Org-N	55	28	104	90	55	45	55	70	50	1970	48	122	100	81

**Table C.7.** Concentration ( $\mu\text{g L}^{-1}$ ) of heavy- and trace-metals in Chinese streams in April/May 1997.  
 < d.l.: less than detection limit. Actual detection limits ( $\mu\text{g L}^{-1}$ ): Cd: < 0.02; Cu: < 0.2; Cr: < 0.01; Ni: < 0.01; Co: < 0.02; Fe: < 15; V: < 0.3; As: < 0.1; Bi: < 0.02; Tl: < 0.006; U: < 0.01; Mo: < 0.04; Sn: < 0.04..

Site	Chongqing area				Guiyang area					Guangzhou area				
	Tie Shan Ping		Simian Shan		Liu Chong Guang			Leishan		Dingushan	Heshan	Baiyun Shan	Liu Xi	
	A	B	A	B	II	I	III	I	II	I	IV			
<b>Pb</b>	0.12	0.13	0.09	0.14	0.58	0.53	0.88	0.15	0.19	0.92	0.67	0.77	0.14	0.09
<b>Cd</b>	0.21	0.15	0.09	< d.l.	0.24	0.26	0.29	< d.l.	< d.l.	0.13	0.37	< d.l.	< d.l.	< d.l.
<b>Cu</b>	0.60	0.60	0.50	0.40	3.40	1.40	1.70	0.40	< d.l.	1.30	0.70	0.60	< d.l.	< d.l.
<b>Zn</b>	11.1	12.1	2.20	0.50	53.4	31.7	36.7	1.10	0.30	13.6	1904	3.30	1.30	0.60
<b>Cr</b>	0.10	< d.l.	0.10	< d.l.	0.50	< d.l.	0.20	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.
<b>Ni</b>	6.60	3.10	0.70	0.70	37.1	21.4	25.7	< d.l.	< d.l.	2.60	0.60	0.50	0.20	< d.l.
<b>Co</b>	0.50	3.80	0.20	0.10	25.7	12.6	15.3	0.10	0.10	1.90	0.30	0.20	0.50	< d.l.
<b>Fe</b>	165	221	248	450	349	218	223	31.5	41.9	< d.l.	89.1	588	639	< d.l.
<b>Mn</b>	295	632	14.0	3.90	215	149	148	3.70	6.50	81.8	62.1	21.4	35.4	2.40
<b>V</b>	< d.l.	< d.l.	< d.l.	0.80	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	0.40	< d.l.	< d.l.
<b>Ti</b>	28.4	32.2	39.6	80.3	71.8	44.8	44.1	2.20	2.30	2.80	7.40	3.40	10.1	1.50
<b>As</b>	0.31	0.28	< d.l.	0.11	< d.l.	< d.l.	< d.l.	< d.l.	0.18	0.29	0.36	2.18	4.07	0.23
<b>Ba</b>	47.1	42.0	29.3	41.3	116	142	151	1.00	1.00	16.1	7.50	13.4	23.4	2.80
<b>Sr</b>	45.8	53.2	24.0	125	143	49.1	69.0	4.30	5.20	3.30	4.50	4.60	17.8	2.40
<b>Sb</b>	0.02	0.03	0.06	0.07	0.05	0.03	0.02	0.02	0.05	0.03	0.34	0.03	0.02	0.03
<b>Bi</b>	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.	< d.l.
<b>Tl</b>	0.01	0.02	< d.l.	< d.l.	0.01	0.01	0.02	< d.l.	0.01	0.01	0.01	0.03	0.05	0.03
<b>U</b>	0.02	0.02	0.01	0.03	0.19	0.04	0.05	< d.l.	0.02	0.05	< d.l.	0.08	0.03	0.20
<b>Be</b>	0.44	0.30	0.03	< d.l.	2.48	1.32	1.43	< d.l.	< d.l.	0.13	< d.l.	0.05	0.02	0.22
<b>Li</b>	1.67	0.97	0.15	0.09	8.81	4.91	6.28	0.09	0.08	0.68	0.82	2.10	1.83	0.39
<b>Rb</b>	1.88	3.10	0.81	0.52	3.00	2.80	2.77	0.16	0.19	0.64	1.16	8.16	8.20	4.52
<b>Mo</b>	< d.l.	< d.l.	< d.l.	0.04	< d.l.	< d.l.	< d.l.	< d.l.	0.04	< d.l.	< d.l.	0.18	0.08	0.34
<b>Y</b>	0.58	0.44	0.12	0.06	2.38	1.13	1.85	0.08	0.07	0.93	0.01	0.13	0.14	0.31
<b>Sn</b>	< d.l.	< d.l.	0.05	< d.l.	0.15	< d.l.	< d.l.	25.4	0.54	0.20	< d.l.	0.09	< d.l.	< d.l.

**Table C.8.** Median, maximum and minimum values in the Chinese water bodies sampled during April/May 1997 and in 473 Norwegian Lakes (Skjelkvåle et al., 1997). Concentrations in  $\mu\text{g L}^{-1}$

		<b>Pb</b>	<b>Cd</b>	<b>Cu</b>	<b>Zn</b>	<b>Cr</b>	<b>Ni</b>	<b>Co</b>	<b>Fe</b>	<b>Mn</b>	<b>V</b>	<b>Ti</b>	<b>As</b>
China	Med	0.17	0.11	0.60	7.20	0.10	0.70	0.40	219	48.8	<0.3	19.23	0.21
	Max	0.92	0.37	3.40	1904	0.50	37.1	25.7	639	632	0.80	80	4.07
	Min	0.09	0.09	0.40	0.30	0.10	0.20	0.10	31.5	2.40	0.40	1.50	0.11
Norway	Med	0.18	0.02	0.4	1.70	<0.1	0.33	0.05	60.7	3.43	<0.3	4.86	
	Max	3.62	0.26	37.7	139	4.85	4.82	3.15	7680	327	2.43	1220	
	Min	0.03	0.02	0.2	0.30	0.10	0.10	0.02	15.0	0.20	0.30	0.40	

		<b>Ba</b>	<b>Sr</b>	<b>Sb</b>	<b>Bi</b>	<b>Tl</b>	<b>U</b>	<b>Be</b>	<b>Li</b>	<b>Rb</b>	<b>Mo</b>	<b>Y</b>	<b>Sn</b>
China	Med	26.4	20.9	0.03	<0.02	0.01	0.03	0.09	0.89	2.33	<0.04	0.23	0.04
	Max	151	143	0.34	0.00	0.05	0.20	2.48	8.81	8.20	0.34	2.38	25.4
	Min	1.00	2.40	0.02	0.00	0.01	0.01	0.02	0.08	0.16	0.04	0.01	0.05
Norway	Med	3.08	5.91	0.03	<0.02	<0.006	0.04	<0.01	0.17	0.49	<0.04	0.09	0.061
	Max	147	3861	0.36	3.62	0.05	2.22	1.34	134	71.1	6.95	2.7	3.52
	Min	0.11	0.32	<0.01	<0.02	<0.006	<0.004	<0.01	<0.01	0.052	<0.04	<0.003	<0.04

**Table C.9.** Soil water chemistry on the Tie Shan Ping catchment. P and H denote plot designation and soil horizon, respectively. N gives number of samples. The data were provided by Dr. Zhao and co-workers at CIESM.

P	H	N	pH	Al <sub>a</sub>	Al <sub>o</sub>	Al <sub>i</sub>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	NH <sub>4</sub> <sup>+</sup>	SO <sub>2</sub> <sup>-4</sup>	NO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	F <sup>-</sup>	Fe <sup>2+</sup>	Mn <sup>2+</sup>
				μM						μeq L <sup>-1</sup>							
1	A	8	5.25	14	1	13	314	199	51	12	15	395	123	90	8	1	5
2	A	6	3.81	235	26	209	824	285	82	190	52	2032	430	211	29	4	28
3	A	8	4.01	178	11	167	354	169	44	91	81	906	578	73		3	17
4	A	9	4.16	184	21	163	1342	487	195	79	161	1633	936	205	59	13	74
5	A	9	4.09	244	13	228	535	259	76	67	45	1323	425	135	60	9	50
6	A	2	3.91	942	30	912	5418	619	202	124	147	5172	2424	423	168	-	-
7	A	2	4.18	241	24	217	531	476	47	68	59	1720	643	106	40	-	-
1	B	9	4.97	1	1	16	269	209	47	9	5	382	126	77	7	2	7
2	B	6	3.97	236	13	224	674	201	87	107	92	1622	312	120	34	1	19
3	B	9	4.08	207	17	190	341	161	48	64	48	917	533	75	21	6	13
4	B	9	4.42	103	8	95	632	251	69	23	25	680	475	126	25	10	31
5	B	7	4.20	178	7	190	547	243	73	71	37	1064	244	151	30	9	30
6	B	1	4.76	171	-	-	1033	483	114	108	62	1201	1587	195	40	-	-
7	B	2	4.70	-	-	-	666	429	106	122	88	939	640	232	13	-	-

**Table C.10.** Median soil water chemistry in the Liu Chong Guang catchment. P and H denote plot designation and soil horizon, respectively. N gives number of samples. Data are from Larssen et al., 1997.

P	H	N	pH	$\mu\text{eq L}^{-1}$									
				Al <sup>3+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	NH <sub>4</sub> <sup>+</sup>	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	F <sup>-</sup>
B	OA	2	4.10	295	1,109	124.9	22.0	34.3	9.8	1,227	311.0	85.6	8.0
F	OA	5	4.65	417	1,218	222.1	15.7	113.0	5.0	1,133	403.2	403.6	10.3
A	A	3	4.37	684	759	220.5	22.1	82.3	10.0	1,707	3.3	104.3	61.8
C	A1	6	4.22	449	491	142.3	22.0	27.4	9.8	864	123.2	180.8	2.4
C	A2	8	4.43	259	160	73.1	7.3	11.4	9.8	407	109.0	11.4	2.9
D	A	6	4.55	122	812	230.0	21.6	71.5	2.5	1,028	54.4	28.9	4.2
E	A	3	4.45	342	544	150.2	28.4	25.2	2.8	923	1.0	19.2	4.1
G	A	3	4.30	526	600	156.1	17.4	62.0	3.9	1,398	1.0	157.7	6.0
A	AB	3	4.20	609	634	199.2	21.6	43.2	5.0	1,496	1.0	166.5	20.0
B	AB	6	4.95	54	832	108.3	19.1	50.0	9.8	760	3.3	97.2	2.9
A	B	1	4.27	1,069	1,233	86.9	25.7	28.9	9.8	2,074	73.9	248.5	20.2
B	B	7	4.30	396	852	121.7	21.5	48.1	5.0	958	53.9	110.0	4.6
C	B	9	4.50	280	213	52.7	10.4	7.9	3.9	492	72.9	18.6	1.7
D	B1	2	4.71	189	831	266.1	12.4	79.1	15.0	1,245	133.8	14.9	35.1
D	B2	8	4.91	63	961	266.0	24.7	39.1	7.1	1,088	10.2	49.5	2.6
E	B	2	4.31	456	537	82.1	20.1	7.6	15.0	1,149	4.2	2.9	16.1
F	B	9	4.75	203	933	267.1	14.4	76.5	10.0	1,051	482.8	64.9	15.9
G	B1	3	4.20	816	649	152.5	16.5	97.6	5.5	1,747	1.0	33.6	28.8
G	B2	2	4.03	912	534	142.4	11.0	56.0	6.7	1,770	1.8	21.2	28.9
C	C	10	4.45	296	185	61.9	12.3	5.4	9.8	364	222.5	35.4	2.6
D	C	7	5.20	26	878	283.8	34.9	14.1	9.8	971	3.4	61.3	1.0
E	C	9	4.66	215	599	79.8	21.6	3.4	4.4	764	1.0	50.8	2.0
F	C	5	4.50	265	754	269.6	10.0	98.1	10.0	963	464.8	67.1	15.5

**Table C.11.** Soil physical characteristics to soils from Liu Chong Guang. P and H denote plot designation and soil horizon, respectively. Data are from Larssen et al., 1997, except for plot H.

P	H	Depth cm	Density g cm <sup>-3</sup>	Sand _____	Silt %	Clay _____
A	AO	3		56.6	29.9	13.5
A	AB	9	0.99	36	33.9	30.1
A	B2	30	1.1	46.2	32.3	21.5
B	OA	1				
B	AB	6	1.17			
B	B	30	1.32			
C	A1	10	0.82			
C	A2	25				
C	BC1	55	0.84			
C	BC2	90				
D	A	3				
D	B1	15				
D	B2	30				
D	C	55				
E	OA	2				
E	B	25		49.6	28.2	22.2
E	C	50		40.6	32.7	26.8
F	OA	2				
F	B	15				
G	A	4				
G	B2	35				
G	C	53				
H	A			66.2	33.8	
H	AB			63.4	36.5	
H	B1			85.0	15.1	
H	B2			74.1	25.9	

**Table C.12.** Cation exchange capacities of soils from Liu Chong Guang. P and H denote plot designation and soil horizon, respectively. N gives number of samples. Data are from Larssen et al., 1997.

P	H	pH		CEC meq kg <sup>-1</sup>	BS	AIS	HS %	FeS
		H <sub>2</sub> O	BaCl <sub>2</sub>					
A	AO	3.85	3.16	138.0	28.4	66.5	4.18	0.78
A	AB	3.89	3.31	106.4	18.6	77.3	3.67	0.38
A	B2	3.87	3.54	94.1	12.5	85.0	2.13	0.35
B	OA	3.81	3.41	223.2	42.9	39.3	1.46	16.2
B	AB	3.95	3.46	84.1	21.0	75.5	2.49	0.92
B	B	4.44	3.74	63.8	22.0	75.5	2.07	0.31
C	A1	3.95	3.54	81.4	9.0	87.6	2.89	0.43
C	A2	4.09	3.72	57.6	4.2	93.4	2.00	0.31
C	BC1	4.36	3.86	26.2	3.7	93.9	1.79	0.57
C	BC2	4.38	4.00	24.5	4.1	93.3	1.91	0.61
D	A	3.89	3.54	99.0	20.6	77.1	2.17	0.03
D	B1	4.22	3.64	88.3	15.4	82.3	2.02	0.17
D	B2	4.43	3.68	83.8	17.4	80.6	1.77	0.13
D	C	4.66	3.81	81.5	40.3	57.8	1.78	0.01
E	OA		3.41	137.0	59.5	38.1	2.37	0.04
E	B	4.48	3.96	57.8	6.7	91.9	1.26	0.12
E	C	4.41	3.82	72.4	6.1	92.3	1.32	0.14
F	OA	3.72	3.20	188.4	25.0	71.1	3.21	0.62
F	B	3.96	3.46	141.9	13.6	84.1	2.04	0.17
G	A	3.61	3.18	124.7	9.7	83.2	4.63	2.39
G	B2	4.21	3.75	90.7	5.9	92.5	1.42	0.12
G	C	4.30	3.80	90.8	5.9	92.5	1.49	0.06



**Table C.13.** Sulfate adsorption capacities to soils from Liu Chong Guang. P and H denote plot designation and soil horizon, respectively. Data are from Liao et al., 1994.

P	H	Dissolvable (native) SO <sub>4</sub> <sup>2-</sup>	SO <sub>4</sub> <sup>2-</sup> adsorption capacities (meq kg <sup>-1</sup> )	
		meq kg <sup>-1</sup>	at 2 meq L <sup>-1</sup>	at 5 meq L <sup>-1</sup>
A	AO	6.8	1.52	2.30
A	AB	4.7	1.45	2.20
A	B2	6.5	1.73	3.46
E	B	2.88	2.33	4.00
E	C	3.26	2.88	5.64

**Table C.14.** Aluminum pools in soils from Liu Chong Guang. P and H denote plot designation and soil horizon, respectively. Data are from Liao et al., 1997b.

P	H	Al pyrophosphate	Al dithionite	Al oxalate	Al CuCl <sub>2</sub>	Al NaOH
		meq kg <sup>-1</sup>				
A	AO	332.4	400.4	340.4	198.1	1061.3
A	AB	224.4	303.3	173.3	116.4	1205.3
A	B2	256.3	317.7	260.2	107.3	1197.2
C	A1	342.6	461.5	347.7	146.1	1638.2
C	A2	433.2	516.3	432.8	160.8	1787.7
C	BC1	446.8	527.3	497.8	86.0	3102.7
C	BC2	332.8	430.2	410.0	78.7	3129.1
E	OA	175.1	330.1	234.8	127.2	1453.1
E	B	263.8	438.1	241.7	108.9	2975.2
E	C	218.0	343.5	204.8	100.3	3332.1

**Table C.15.** Particle size distribution and mineralogy of soils from Liu Chong Guang. Sand is defined as the size fraction between 2000  $\mu\text{m}$  and 63  $\mu\text{m}$  and silt + clay as the fraction passing through a sieve with grating 63  $\mu\text{m}$ . P and H denote plot designation and soil horizon, respectively.

Site	P	H	Depth cm	Particle size distribution in the fine fraction		Total clay	Minerology		
				Sand %	Silt+ clay %		Quartz	Plagioclase %	K-feldspar
Tie Shan Ping	2	A	0 - 3	55.5	44.5	12	76	6	6
Tie Shan Ping	2	B1	3 - 13	61.7	38.3	13	74	8	5
Tie Shan Ping	2	B2	13 - 28	77.3	22.7	16	72	4	8
Simian Shan		A	3 - 7	82.2	17.8	8	83		9
Simian Shan		B1	7 - 13	79.7	20.3	10	82		8
Simian Shan		B2	13 - 18	88.3	11.7	8	86		6
Liu Chong Guang	H	A	4 - 8	65.7	34.3	12	80		8
Liu Chong Guang	H	AB	11 - 15	63.1	36.9	14	80		6
Liu Chong Guang	H	B1	30 - 34	84.5	15.5	15	72	4	9
Liu Chong Guang	H	B2	55 - 59	73.5	26.5	18	74		8
Leishan	1	AB	7 - 27	-	-	19	59	15	7
Leishan	1	B	> 27	89.6	10.4	22	56	18	4
Leishan	3	A	1 - 5	66.2	33.8	22	73	10	5
Liu Xi		A	1 - 5	76.8	23.2	14	62	12	22
Liu Xi		AB	5 - 25	78.7	21.3	12	44	16	28
Liu Xi		B	40 - 50	86.8	13.2	11	51	14	24

**Table C.16.** Sampling sites, time of sampling and weather conditions for hydrocarbon and carbonyl air samples.

	Date	Hydrocarbon sample:	Carbonyl sample:	Weather conditions
Nanshan	17.04.97	1530		Overcast
Simian Shan	19.04.97	1600	1605-1735	Overcast, low cloud base
Liu Chong Guang	23.04.97	1315	1055-13-45	Overcast
Leishan	26.04.97	1245	1330-1410	Overcast, thin cloud cover
Leishan (2)	26.04.97	1450		Overcast, thin cloud cover
Liu Chong Guang (2)	28.04.97	1105	0945-1255	Cloudy, some sun
Dinghushan	02.05.97	1435	1443-1543	Sunny, haze
Heshan	03.05.97	1400	1407-1547	Overcast, some drizzle
Baiyun Shan	05.05.97	0950		Overcast
Liu Xi	06.05.97	1205	1212-1312	Sunny, warm

**Table C.17.** Typical atmospheric residence times in days, according to Derwent (1993).

Compound	Days	Compound	Days
Ethane	116	Benzene	25.3
Ethene	3.6	Toluene	5.2
Acetylene	34.5	o-Xylene	2.3
Propane	27	m-Xylene	1.3
Propene	1.2	p-Xylene	2.2
n-Butane	12.2	Formaldehyde	3.2
1-Butene	1.0	Acetaldehyde	2
2-Butene	0.6	Propanone	137
n-Hexane	10		

**Table C.18.** Measured concentrations of hydrocarbons in air at the indicated sites in China during April and May, 1997. Average concentrations at 3 European sites during June 1996 (from Solberg et al., 1996) are shown for comparison. Unit: ppb (v).

Date	970417	970419	970423	970426	970426	970428	970502
Time	1530	1600	1315	1245	1450	1105	1435
Site name	Nanshan	Simian Shan	Liu Chong Guang	Leishan	Leishan	Liu Chong Guang	Dinghushan
Ethane	7.18	3.37	3.22	2.5	2.56	3.83	2.33
Ethene	5.33	2.28	1.28	1.18	1.33	3.61	1.47
Propane	1.67	0.75	0.65	0.46	0.48	0.75	0.57
Propene	0.61	0.31	0.26	0.26	0.31	0.73	0.42
Acetylene	5.23	2.8	2.25	1.63	1.72	5.27	2.07
I-Butane	0.35	0.29	0.11	0.07	0.07	0.28	0.25
n-Butane	0.58	0.43	0.18	0.11	0.12	0.54	0.36
Sum Butenes	0.36	0.44	0.22	0.18	0.2	0.58	0.37
i-Pentane	0.46	0.22	0.1	0.05	0.05	0.87	0.42
n-Pentane	0.22	0.1	0.06	0.03	0.04	0.39	0.14
1,3-Butadiene	0.03	0.01	0.01	0.01	0.01	0.08	0.06
Sum Pentenes	0.17	0.18	0.07	0.08	0.05	0.59	0.23
Sum Unresolved C6	1.83	0.19	0.23	0.02	0.12	1.25	1.49
n-Heptane	0.21	0.05	0.02	0.01	0.01	0.22	0.05
Benzene	2.53	0.72	0.63	0.4	0.42	2.37	0.61
Toluene	1.08	0.23	0.19	0.11	0.14	1.95	0.44
Ethylbenzene	0.21	0.05	0.03	0.02	0.02	0.21	0.05
m+p Xylenes	0.38	0.09	0.04	0.02	0.02	0.57	0.09
o-Xylene	0.17	0.05	0.07	0.01	0.01	0.24	0.08

Date	970503	970505	970506	9606	9606	9606
Time	1400	950	1205	1200-1400	1200-1400	1200-1400
Site name	Heshan	Baiyun Shan	Liu Xi	Birkenes	Waldhof	Kosetice
Ethane	1.36	1.2	1.98	1.49	1.58	1.42
Ethene	0.92	2.23	1.89	0.286	0.195	0.303
Propane	0.18	1.69	0.66	0.405	0.499	0.402
Propene	0.28	0.63	0.61	0.117	0.071	0.09
Acetylene	0.8	3.23	1.83	0.336	0.379	0.53
I-Butane	0.05	0.61	0.21	0.163	0.245	0.239
n-Butane	0.11	0.99	0.42	0.349	0.13	0.184
Sum Butenes	0.22	0.73	0.56			
i-Pentane	0.08	0.68	0.36	0.097	0.098	0.29
n-Pentane	0.09	0.28	0.19	0.251	0.146	0.408
1,3-Butadiene	0.02	0.09	0.07			
Sum Pentenes	-0.01	0.56	0.32			
Sum Unresolved C6	1.16	1.31	1.08			
n-Heptane	0.1	0.25	0.11			
Benzene	0.36	1.55	0.43	0.114	0.134	0.168
Toluene	5.43	3.29	1.05	0.162	0.162	0.233
Ethylbenzene	0.04	0.36	0.2			
m+p Xylenes	0.06	0.7	0.47			
o-Xylene	0.02	0.26	0.17			

**Table C.19.** Measured concentrations of carbonyl compounds in ppb(v) at the indicated sites in China in April and May 1997. Comparable average concentrations for 3 European sites in June 1996 (Solberg et al., 1997) are shown for comparison.


Date	970419	970423	970426	970428	970502	970503	9705066	9606	9606	9606
Time	1605-1735	1055-1345	1330-1410	0945-1255	1443-1543	1407-1547	1212-1312	1200-1400	1200-1400	1200-1400
Sampling site	Simian Shan	Liu Chong Guang	Leishan	Liu Chong Guang	Dinghushan	Heshan	Liu Xi	Birkenes	Waldhof	Kosetice
Methanal	0.55	0.79	1.13	0.69	3.92	1.49	3.49	0.76	1.38	1.28
Ethanal	0.43	0.92	0.98	1.42	2.13	1.17	8.21	0.44	0.63	1.04
Propanone	n.d.	n.d.	4.40	4.01	6.40	n.d.	4.81	n.a.	2.03	3.11
Propenal	n.d.	0.11	0.30	n.d.	0.11	0.26	n.d.	n.a.	n.a.	n.a.
Propanal	n.d.	0.28	0.20	0.40	0.47	0.25	0.41	0.08	0.11	0.21
3-Butene-2-one	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.a.	n.a.	n.a.
Butanone	n.d.	0.12	n.d.	0.66	1.10	n.d.	0.29	0.16	0.29	0.41
2-Methylpropenal	n.d.	n.d.	0.10	n.d.	n.d.	n.d.	n.d.	n.a.	n.a.	n.a.
Butanal+isobutanal	0.29	0.38	0.20	0.44	1.33	0.73	0.86	n.a.	n.a.	n.a.
Benzencarbaldehyde	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.a.	n.a.	n.a.
Pentanal	0.09	n.d.	n.d.	0.31	0.39	n.d.	n.d.	n.a.	n.a.	n.a.
Ethandial	0.09	n.d.	n.d.	0.17	0.18	0.19	n.d.	n.a.	n.a.	n.a.
Hexanal	0.15	0.08	0.28	0.08	0.46	n.d.	0.64	n.a.	n.a.	n.a.
2-Oxopropanal	0.11	n.d.	n.d.	0.17	0.16	0.11	n.d.	n.a.	n.a.	n.a.

**Table C.20.** Data for chlorophyll fluorescence measurements (by PEA) from the visited sites. **A:** Masson pine; **B:** Chinese fir.

**A**

*Pinus Massoniana*


	Mean	Min.	Max.	Med.	SD
<b>Sichuan province, Chongqing area</b>	0,7934				0,0770
Nashan, East side	0,7575	0,6920	0,8020	0,7700	0,0428
Tie Shan Ping	0,8022	0,7660	0,8400	0,8015	0,0251
Simian Shan	0,8207	0,8130	0,8270	0,8210	0,0052
<b>Guizhou province, Guiyang area</b>	0,7982				0,0056
Liu Chong Guang, Low	0,8047	0,7720	0,8430	0,8035	0,0247
Liu Chong Guang, High	0,7928	0,7650	0,8260	0,7880	0,0203
Leigong Shan	0,7973	0,7490	0,8320	0,8030	0,0329
<b>Guangdong province, Guangzhou area</b>	0,8034				0,0560
Dinghushan, High	0,8103	0,7680	0,8300	0,8195	0,0236
Dinghushan, Low	0,8202	0,8140	0,8290	0,8180	0,0063
Heshan station	0,8247	0,8060	0,8320	0,8280	0,0010
Guangzhou Bot. Garden	0,7995	0,7640	0,8300	0,7985	0,0235
Baiyun Shan	0,7870	0,6920	0,8310	0,8065	0,0544
Liu Xi River	0,7785	0,7200	0,8120	0,7865	0,0329



**B**

*Cunninghamia lanceolata*

	Mean	Min.	Max.	Med.	SD
<b>Sichuan province, Chongqing area</b>	0,7994				0,0097
Nashan, East side	0,8043	0,7930	0,8160	0,8040	0,0076
Tie Shan Ping	0,7973	0,7840	0,8050	0,8005	0,0082
Simian Shan	0,7965	0,7800	0,8040	0,7995	0,0088
<b>Guizhou province, Guiyang area</b>	0,8049				0,0372
Liu Chong Guang, Low	0,8157	0,8090	0,8260	0,8140	0,0063
Liu Chong Guang, High	0,8137	0,8050	0,8230	0,8130	0,0073
Leigong Shan	0,7853	0,7630	0,8100	0,7845	0,0201
<b>Guangdong province, Guangzhou area</b>	0,8078				0,0304
Dinghushan, High	0,7923	0,7720	0,8060	0,7950	0,0122
Dinghushan, Low					
Heshan station	0,8063	0,7930	0,8160	0,8085	0,0090
Guangzhou Bot. Garden	0,8172	0,8010	0,8320	0,8165	0,0124
Baiyun Shan					
Liu Xi River	0,8155	0,8080	0,8230	0,8140	0,0055



# **APPENDIX D**



## **D. Proposed framework for future research, monitoring and training program**

One of the main goals for the PIAC-project was to identify a framework for future cooperation between Norway and China on acid rain. In this appendix, the PIAC team have identified what they believe should constitute the key elements in this framework. This framework is intended as a point of departure for a dialogue between Chinese and Norwegian partners in order to develop a main project on acidification research and monitoring in China, including monitoring on air, precipitation, soil, soil water, vegetation (including forest), surface water and terrestrial and aquatic biota.

In our opinion the main goal of the project should be:

*To assist Chinese researchers and institutions to obtain a reliable picture of the extent and severity of effects of acid precipitation on Chinese ecosystems.*

In order to help Chinese researchers and institutions reach this goal, the following key elements have been identified:

1. Establish a network of Chinese institutions in cooperation with institutions from Norway and other countries, in order to develop best possible basis for research and monitoring of acid precipitation and the effects.
2. Establish a monitoring program for determining the extent, severity and trends of acid precipitation.
3. Improve the knowledge of processes determining the impacts of acid deposition on ecosystems by drawing experience from other countries.
4. Use the obtained knowledge to improve acidification models that are suitable for predictions of future effects (e.g. the RAINS-Asia model).
5. Cooperate with decision makers in applying the knowledge to make cost-efficient countermeasures.

In order to provide a best possible understanding of acid rain issues for decision makers, one should combine determination of physical, chemical and biological effects *with estimates of economic values* of effects on the ecosystems.

To develop a successful program on acid rain abatement, both interdisciplinary and international efforts are necessary. A corner stone in the main project on acid rain will therefore be to establish an effective network between central and local Environmental Authorities (The NEPA-system), scientific institutions within the CAS-system, and Norwegian institutions. Some of these relationships are already established. The action plan for the main project should lay the foundations for a high quality monitoring, research and training program. If we fulfill our intentions, this program will hopefully be highly useful for both Chinese and Norwegian scientists and policy makers, and may act as a template for other research/monitoring programs in China. Data from the project will also be very useful for validation and improvement of the RAINS-Asia model. In addition to the monitoring, research and training program, intercalibration of methods and analyses will be a central issue.

The framework we are presenting is focusing on the scientific elements of the program. We expect our Chinese partners to identify in detail the need of training that is necessary for a successful implementation of the program. Also, in the request for assistance to be made by our Chinese partners, a detailed specification of the technical equipment needed has to be made.

## **D.1 Selection of sites**

We propose to incorporate the Sichuan, Guizhou and Guangdong provinces in the monitoring, research and training project, with preferably at least two sites in each province. Priority is given to initiate and implement collaboration between Chinese institutions at local and central levels, to exchange knowledge (e.g. on analytical methods), to identify training needs, and to work out the logistics to assure functional equipment (e.g. supply of spare parts, chemicals, sampling equipment). Based on what we learned by visiting actual sites and other knowledge and information, we feel it is possible at this stage to identify a number of suitable monitoring and research sites. The need of incorporating or establishing more background stations, as also emphasized in the final report of the Fourth Expert Meeting on Acid Deposition Monitoring Network in East Asia (EMAD, 1997), will be followed up.

We will also highly recommend a large regional survey to be included in the program, in order to reveal the regional impacts of acidification. A regional survey will provide important insights to policy makers as well as valuable for the calibration of the RAINS-Asia model.

## **D.2 Selection of Chinese partners**

The PIAC-team have established close contacts with the following Chinese institutions, which accordingly also are among the likely institutions to participate in a future collaborative monitoring/research and training program on acid rain in China:

### **Beijing:**

- National Environmental Protection Agency (NEPA)
- China Research Academy of Environmental Science (CRAES)
- Chinese Academy of Sciences (CAS) and connected research centers.

### **Chongqing area:**

- Chongqing Environmental Protection Bureau
- Chongqing Inst. of Environmental Science and Monitoring
- Forestry Bureau of Chongqing

### **Guiyang area:**

- Guizhou Provincial Environmental Protection Bureau
- The Acid Rain Control Center of Guiyang, Guiyang Environmental Protection Bureau
- Guizhou Academy of Sciences, Guiyang
- Guizhou Inst. of Environmental Sciences, Guiyang
- Guizhou Botanical Garden, Guiyang

### **Guangzhou area:**

- Guangzhou Environmental Monitoring Center
- Guangzhou Municipal Environmental Protection Bureau
- South China Inst. of Botany, Guangzhou

### **D.3 Administrative structure**

We propose that key persons from the participating institutions should establish a Management group or Steering Committee for the project. We will also propose to establish 4 scientific research groups within each province:

- Meteorology and precipitation chemistry
- Soil, soil water and surface water
- Forest health
- Other terrestrial and aquatic flora and fauna

Each group should contain Chinese scientists at local/provincial level, in close collaboration with the Norwegian experts. Each of these 4 research groups should have one group leader with responsibility for the implementation of their monitoring/research and training parts. These group leaders should be in close contact with the leader of the project in the province. Close cooperation between the groups is essential. Coordination may be promoted by establishing a common *group for data analysis and modeling*.

Annually, there should be a status meeting with participation of the Environmental Agencies in China and Norway and the researchers involved in the project. If necessary, a status report should be issued after each annual meeting.

### **D.4 Monitoring program**

In Norway we have derived benefits from studying effects of acidification at three levels:

- a) A few sites with highly frequent sampling programs for detailed process studies. Both highly affected areas and control areas with minor anthropogenic effects are incorporated. Many research projects with international cooperation have been conducted at such sites in Norway.
- b) A few sites for temporal trend analyses. The sampling frequency should vary for different systems, e.g. be quite frequent for surface water and much less frequent for soils. These sites might very well be the same as in a).
- c) Large surveys with 5 or 10 years' intervals over a large geographical area (country or province) to get an idea of the effects at this scale. This information is of great importance for policy makers.

Accordingly, we will recommend these three levels as a basis for establishing a monitoring program for acid rain in China. As a supplement, laboratory studies are recommended.

## **D.4.1 Sampling**

### **Air and precipitation**

The monitoring program may serve two purposes: estimation of the atmospheric input to the selected ecosystem(s), and establishment of a deposition field by interpolation of data from a network of stations. In addition, data from the measurement sites will be useful for developing models describing the relationships between atmospheric emissions, airborne concentrations and deposition.

The basic measurement program should conform to international standards with respect to sampling, analytical methods and quality assurance. We suggest sampling of precipitation on a weekly basis with a wet-only sampler, and determination of sulfur dioxide and sulfate in particles by sampling with a filterpack method, in accordance with EMEP recommendations (cf. Table D.4.). This requires that electricity is available, and can interfere with the selection of an ideal sampling site. If this becomes a serious problem, it may be considered to use a NILU bulk sampler for precipitation sampling, and passive samplers to determine sulfur dioxide. A limited number of parallel samples should be analyzed at NILU for quality control purposes.

In addition to this standard program, more short-term measurement activities should be considered in order to characterize the air chemistry and the deposition at the sites. This could include the determination of base cations in airborne particles (sampled in two size fractions), airborne concentrations of ammonia, ammonium, nitrate and gaseous nitric acid, hydrogen fluoride and other components of special interest.

Throughfall water should be collected weekly to fortnightly with collectors similar to the ones used to collect bulk precipitation.

### **Soil and soil water**

Data on soil and soil water are essential to understand acidification mechanisms and therefore for modeling of effects. Soil water from the dominating soil types within the intensive studied sites should be sampled by means of porous suction cup lysimeters installed in the genetic soil horizons. Samples should routinely be collected monthly, but more frequent sampling during some hydrologic episodes is recommended. Soil samples will be collected adjacent to the lysimeter installations. For details see Table D.4.

### **Forest ecosystems**

Monitoring programs of forest ecosystems should focus on stands with Masson pine and Chinese fir along environmental gradients. Inventories of tree vitality and understory vegetation might make up the standard of reference and habitat stratification. These data might be used in different GIS applications on fine scale level (see section 5.4.4). The aims are to quantify the year-to-year composition changes, and to elaborate a basis for long-term vegetation trends. This includes spatial-temporal studies of changes in biodiversity and interpretation of species dynamics along identified major environmental gradients (cf. Økland & Eilertsen 1996). Species richness and changes in richness might be studied within different structural and taxonomic groups, between different forest types and at comparative sites with different loads of pollutants such as SO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub>.

Statistically significant vegetation changes might be tested by direct gradient analysis, i.e. constrained ordination procedures like Canonical Correspondence Analysis, CCA (ter Braak 1986; 1987a), with time or a selected exploratory parameter as the constraining variable. This could be demonstrated for

year-to-year intervals or for long-term periods. Data should also be prepared for other multivariate statistics, i.e. ordination methods like Correspondence Analysis, CA (Hill 1979; ter Braak 1987b; 1990) or Multidimensional Scaling, MDS (Carr 1990; Clarke 1991), where vegetation changes along identified gradients are measured as plot displacement along interpreted ordination axes.

Intensive monitoring in forest ecosystems includes the estimation of throughfall fluxes (water dripping from the tree canopy), as well as fluxes and concentrations of the major elements in soil water. The monitoring of forest ecosystems also includes estimation of nutrient fluxes associated with litterfall and net nutrient uptake by trees. Litterfall should be collected monthly using litterfall samplers (about 1 m<sup>2</sup> in size per collector). Net nutrient uptake is estimated on an annual basis from the chemical composition of the trees and the annual increment. The chemical composition of current and previous-years needles, crown density and crown color should be estimated annually.

### **Water and surface water chemistry**

If runoff is monitored, we will recommend a V-notch weir with a continuous water level recorder. The stream water may be collected using the "grab" spot method, using well cleaned high density polypropylene bottles. Samples should be collected weekly. Stream water might be more frequently sampled during some hydrologic episodes. The samples should be stored at 4°C. See Table D.4. for details.

### **Regional survey**

In addition to the sampling program above, we also recommend to implement regional surveys, preferably in the beginning of the project period, in the participating provinces. The surveys should include: sampling of soil, vegetation (forest, ground vegetation), surface water, freshwater fauna and flora (e.g. invertebrates and diatoms), soil invertebrates and microorganisms. In addition one should obtain information about past and present fish status as well as relevant atmospheric data in the provinces.

Regarding forest ecosystems, the regional surveys should focus on tree health parameters (annual observations of fixed plots in selected stands) as proposed by ICP Forest (Pylvänäinen, 1993) in relation to site index, stand age, climate, pollution stress (e.g. acid deposition) and soil characteristics. Critical load maps should be made in collaboration with RAINS-Asia.

### **D.4.2 Analytical program**

Table D.4. presents the analytical physico-chemical methods used in the monitoring programs in Norway based on well described analytical methods. Depending on the economical frames of the project and the equipment already present at the different Chinese institutions involved in this project, different analytical methods might be used at different institutions. Intercalibration between Chinese institutions and the Norwegian institutes is therefore highly important to produce high quality and comparable data in the project. Intercalibration should be implemented at least once during the project period, but likely we have to repeat intercalibrations for several parameters.

In the regional survey, the analyses will be the same as present in Table D.4. for soil, vegetation and surface water.

**Table D.4. Frequency of precipitation and surface water sampling and description of the applied physical, chemical methods.**

Sampling frequency	Parameter/Method	Comments
	<b>1) Air analysis</b>	
Weekly	<b>Dry-deposition</b> Gas: SO <sub>2</sub> , NO <sub>2</sub> Gas/particles: ΣNH <sub>4</sub> /NH <sub>3</sub> , ΣNO <sub>3</sub> /HNO <sub>3</sub> Particles: SO <sub>4</sub> , Ca, Mg, Na, K, Cl	SO <sub>2</sub> : hydrogen-peroxide extraction of filter. Sulfate formed analyzed by ion chromatography (IC); SO <sub>4</sub> : water extract of filter and analyzed by IC. NO <sub>2</sub> : TGS-colorimetric method (NS 4855); ΣNH <sub>4</sub> <sup>+</sup> /NH <sub>3</sub> : colorimetry (CM); ΣNO <sub>3</sub> <sup>-</sup> /HNO <sub>3</sub> : Water extraction of filter containing sodium-hydroxide, and analyzed with respect to nitrate by IC. Ca, Mg, K: Water extract of a sulfate filter and analyzed by Flame Atomic Absorption Spectroscopy (FAAS); Na: Flame Atomic Emission Spectroscopy (FAES); Cl: IC
Weekly	<b>Wet-deposition</b> Precipitation (mm), pH, Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> , K <sup>+</sup> , NH <sub>4</sub> <sup>+</sup> , SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> ; Trace element: As, Pb, Cd, Cr, Co, Zn, Ni, Cu	pH: potentiometry (PM); Cl <sup>-</sup> , SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> : IC; Na <sup>+</sup> : FAES; Ca <sup>2+</sup> , Mg <sup>2+</sup> , K <sup>+</sup> : FAAS NH <sub>4</sub> <sup>+</sup> : CM; Trace metals: Induced Coupled Plasma Mass Spectroscopy (ICP-MS)
	<b>2) Soil/soil water analysis</b>	
5 - 10 years intervals	<b>Soil</b> Grain size distribution, texture, moisture characteristics, mineralogy, mineral composition, pH <sub>H2O</sub> , CEC <sub>E</sub> , CEC <sub>P</sub> , %BS, %AIS, %HS, exchangeable cations, Al-pools, organic C, total N, extractable P, SO <sub>4</sub> adsorption	Organic matter content: Loss on ignition; Grain size distribution: sieve and sedimentation; Mineral composition: X ray diffraction/scanning electron microscopy (XRD/SEM); pH <sub>H2O</sub> : pH in water extracts of the soil; CEC <sub>E</sub> , %BS, %AIS, %HS: Soil are extracted with BaCl <sub>2</sub> solution and the extractant analyzed for pH, Ca, Mg; Na, K, and Al by FAAS; CEC <sub>P</sub> , exchangeable base cations and acidity: extraction using extraction with NH <sub>4</sub> NO <sub>3</sub> . Al-pools: soil extracts of pyrophosphate, dithionite, oxalate, CuCl <sub>2</sub> and NaOH and the extractants analyzed for Al by FAAS. Organic C and total N by element analysis; P by extraction with HNO <sub>3</sub> /HClO <sub>4</sub> (microwave), SO <sub>4</sub> adsorption by extraction with PO <sub>4</sub> .
Monthly	<b>Soil water</b> Cond., Temp, H <sup>+</sup> (pH), Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> , K <sup>+</sup> , NH <sub>4</sub> <sup>+</sup> , SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> , TOC/DOC, Aluminum (Ala, Alo), total F, H <sub>4</sub> SiO <sub>4</sub>	Conductivity: electrometry; pH: PM; Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> , K <sup>+</sup> : FAAS; NH <sub>4</sub> <sup>+</sup> : Cation IC; SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> : Anion IC; TOC/DOC: Catalytic high temperature combustion to CO <sub>2</sub> and measured spectrometrically by infra-red (IR) gas analyser; Aluminum (Ala, Alo): Complexation to 8-HQ and extracted to MIBK before determination by CM on non-cation exchanged and cation exchanged samples respectively; total F: Ion selective electrodes (IS), H <sub>4</sub> SiO <sub>4</sub> : CM
	<b>3) Vegetation/forest</b>	
Yearly	Analysis of trees and understorey vegetation. Analysis of needles, branches and stem wood; analysis of tree litter. Contents of all major nutrients and several micro elements.	Measurements of chlorophyll fluorescence by PEA (Plant Efficiency Analyser) Extraction with HNO <sub>3</sub> /HClO <sub>4</sub> in microwave. N content using Kjeldal. Other elements using ICP

**Table D.4. Continuous**

Sampling frequency	Parameter/Method	Comments
	4) Surface water	
Weekly	Cond., Temp, Turbidity, pH, Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> , K <sup>+</sup> , NH <sub>4</sub> <sup>+</sup> , SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , Cl <sup>-</sup> , alkalinity, TOC/DOC, Tot-N, aluminum (RAL, ILAL), F <sup>-</sup> , H <sub>4</sub> SiO <sub>4</sub>	Cond: electrometry; pH: PM; Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> , K <sup>+</sup> : FAAS; NH <sub>4</sub> <sup>+</sup> : CM; SO <sub>4</sub> <sup>2-</sup> , Cl <sup>-</sup> : IC; NO <sub>3</sub> <sup>-</sup> : CM; Alkalinity: potentiometric titration with hydrochloric acid to pH 4.5; TOC: Organic carbon oxidized to CO <sub>2</sub> by peroxydisulfate and UV-radiation in strongly acidic environment and measured spectrometrically by infra-red (IR) gas analyzer. Tot-N: oxidizing all N to NO <sub>3</sub> and analyzed by CM as for NO <sub>3</sub> ; RAL, ILAL: CM (PCV-method); F <sup>-</sup> : ionic selective electrode (ISE); H <sub>4</sub> SiO <sub>4</sub> : CM.; Trace metals: Induced Coupled Plasma Mass Spectroscopy (ICP-MS)

### D.4.3 Comments on biological studies

#### Plants and vegetation

Air pollutants such as SO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub> are often dispersed over large areas or regions. Since there are no pollution-free «control» areas, the regional distribution of air pollutants makes their impact on plants difficult to assess. Therefore, the sampling design for plant studies and vegetation monitoring need to be carefully assessed by all participants before it is implemented in the final program. Strategies for data treatment and analysis should follow standardized procedures as described in section 5.4.1 (Forest Ecosystems). If necessary, an education and calibration program should be developed for all participants.

In addition, environmental differences between areas and sites, such as changes in soil types and contrasts in day length, temperature and water availability, make it difficult to compare air pollution impact data from different sites. Therefore, the scientists should consider additional studies of effects on plants and vegetation under controlled conditions. A possible approach is to interconnect results from fieldwork studies in «natural» sites with data from effect studies on plants and vegetation in open-top fumigation chambers. Plants are typically raised in these chambers for a full growing season and then assessed for injuries from different pollutants such as SO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub>. Botanists at CRAES in Beijing and at SCIB in Guangzhou have scientific experience from studies in open-top chambers.

#### Soil and surface water biota

The sampling and analysis strategy for monitoring and research on soil and surface water biota needs more careful evaluation based on discussion with potential Chinese counterparts and preferably ecological data from regional studies and some research data before this can be fully integrated in the monitoring program. However, regional surveys should include qualitative sampling of freshwater fauna and flora e.g. invertebrates and diatoms using standard methods and collection of local information about past and present fish status, species composition and, if possible, inventories of soil invertebrates and samples of soil microorganisms at some sites.

Besides continuing the Chinese comparative studies on soil micro-organisms at the selected sites with different levels of acid loading and sensitivity, and parallel toxicity studies of acid rain effects on soil microbes, we propose new studies of soil invertebrate fauna particularly macro-invertebrates (like

snails and worms) with regard to fauna composition and abundance. At a later stage we recommend biomarker studies to measure the general level of environmental stress in such organisms.

The program on freshwater biota should include microalgae/diatoms, macroalgae and macrophytes (higher plants), invertebrates and fish in stream- and lake/pond habitats, applying both field observations and experiments under field and laboratory conditions to establish biological dose-response functions suitable for environmental modeling and management strategies. The overall approach should be:

- (1) Initiate qualitative sampling of freshwater plants and animals for the determination of which species are present/absent at selected sites in a regional survey
- (2) Initiate quantitative sampling of freshwater plants and animals for the description of abundance and community composition at selected sites
- (3) Identify a set of key indicator organisms and/or biotic processes suitable for a longer term monitoring and research program and use this information to establish biological dose-response functions as suitable tools for environmental modeling and management strategies
- (4) to initiate more detailed studies of ecophysiological effects at the level of individuals and ecological studies of processes and interactions between species, communities and different level of organization, e.g. within the food chain.

Data from this program should be linked to physico-chemical data, in order to relate the level of ecological change to changes in the pollution level.

The inventory studies would require annual or biannual sampling of biota and identification of which species are present. The more detailed studies should involve more frequent sampling (daily - weekly) and collection of quantitative data in order to describe communities and the interrelationships between various functional groups, e.g. primary producers and consumers at higher levels in the food chain. Experimental work would require even more intensive sampling frequency and refined analytical protocols.

#### **D.4.4. Use of Geographical Information Systems (GIS)**

The project proposed by the Norwegian PIAC group addresses monitoring of pollution effects in China. The proposal includes a wide range of potential monitoring variables, covering physical, chemical as well as biotic parameters in air, freshwater and terrestrial ecosystems. These varied parameters originate and have effects on a broad range of spatial scales, from small catchments to larger geographic regions. They are also affected by a large number of surrounding factors associated with both the physical landscape, the vegetation and various societal forces which all vary at different spatial scales. It is apparent that appropriate integration and synthesis of this thematically and spatially diverse set of parameters represent a general challenge within the project. One particular aspect of this challenge is to integrate across spatial scales and to handle scale shifts from the local observational scale to the regional modeling scale of, e.g., RAINS-Asia.

We suggest to apply Geographical Information Systems (GIS) for this purpose. It seems convenient to employ two spatial scales. The finer scale should be at the level of local valleys and catchments, i.e., an area of up to about 5 km. This scale will allow reasonably detailed modeling of local observation patterns in relation to actual factors of influence. The larger spatial scale should be at the regional level, i.e., an area of 100–500 km. This will be particularly well suited for integration of observations



from the proposed regional surveys. This will also allow an integration of local observations to the approximate cell size of continental scale models like RAINS-Asia (i.e., about 150 x 150 km).

The availability of data, particularly appropriate geo-referenced digital data, will represent a critical constraint for the extent and successful execution of the analyses.

We propose to employ a variety of PC-based GIS programs. The various programs will be adopted to fit the appropriate modeling task at hand in any given case. However, as an integrative platform we propose to use ArcView which is effective for integrated presentations and some relevant modeling. It is also widely distributed with well-known data exchange facilities. Using a PC basis implies constraints on the amount of data which can be handled. However, the relatively low cost and availability of PCs and relevant programs should make it more readily available among cooperating Chinese counterparts.

## **D.5 Need of local people, training and relevant equipment**

Access to local infrastructure and relevant equipment is important for implementing the program. All logistics, including transport, housing, field assistance, power supply, laboratory facilities, supplies of minor field equipment and storage of material etc., should preferably be organized locally.

Norwegian and/or other foreign experts should be allowed to spend enough time in China to ensure transfer of expertise, support and conduct training and supervise when needed. As mentioned earlier, the training need of Chinese scientists and experts has to be identified by themselves on the basis of the scientific and technical aspects of the project. It is evident that training will constitute a central element in this project.

## **D.6 Time table**

The objective of the project is to help our Chinese counterparts to be able to run a long term acid rain monitoring program themselves. Given the challenges at hand and their present need for foreign scientific, technical and financial support, we suggest that the monitoring program should be run for 5 years, by support from NORAD and/or WB-ASTEN.

# Acid Rain Research Reports

- 1/1982 Henriksen, A. 1982. Changes in base cation concentrations due to freshwater acidification. 50 pp. Out of print.
- 2/1982 Henriksen, A. and Andersen, S. 1982. Forsuringssituasjonen i Oslomarkas vann. 45 pp. Out of print.
- 3/1982 Henriksen, A. 1982. Preacidification pH-values in Norwegian rivers and lakes. 24 pp. Out of print.
- 4/1983 Wright, R.F. 1983. Predicting acidification of North American lakes. 165 pp.
- 5/1983 Schoen, R., Wright, R.F. and Krieter, M. 1983. Regional survey of freshwater acidification in West Germany (FRG). 15 pp.
- 6/1984 Wright, R.F. 1984. Changes in the chemistry of Lake Hovvatn, Norway, following liming and reacidification. 68 pp.
- 7/1985 Wright, R.F. 1985. RAIN project. Annual report for 1984. 39 pp.
- 8/1985 Lotse, E. and Otabbong, E. 1985. Physiochemical properties of soils at Risdalsheia and Sogndal: RAIN project. 48 pp.
- 9/1986 Wright, R.F. and Gjessing, E. 1986. RAIN project. Annual report for 1985. 33 pp.
- 10/1986 Wright, R.F., Gjessing, E., Semb, A. and Sletaune, B. 1986. RAIN project. Data report 1983-85. 62 pp.
- 11/1986 Henriksen, A., Røgeberg, E.J.S., Andersen, S. and Veidel, A. 1986. MOBILLAB-NIVA, a complete station for monitoring water quality. 44 pp.
- 12/1987 Røgeberg, E.J.S. 1987. A coulometric Gran titration method for the determination of strong and weak acids in freshwater. 28 pp.
- 14/1988 Hauhs, M. 1988. Water and ion movement through a minicatchment at Risdalsheia, Norway. RAIN project. 74 pp.
- 15/1988 Gjessing, E., Grande, M. and Røgeberg, E.J.S. 1988. Natural Organic Acids. Their Role in Freshwater Acidification and Aluminium Speciation. 28 pp.
- 16/1988 Wright, R.F. 1988. RAIN project. Annual report for 1987. 77 pp.
- 17/1988 Wathne, B.M. and Røgeberg, E.J.S. 1988. Buffering effects of river substrates under acidic conditions. 19 pp.
- 18/1989 Lotse, E.G. 1989. Soil Chemistry 1983-86 at the RAIN Project Catchments. 66 pp.
- 19/1989 Reuss, J.O. 1989. Interpretation of Soil Data from the RAIN Project. 81 pp.
- 20/1990 Skjelkvåle, B.L. and Wright, R.F. 1990. Overview of areas sensitive to acidification: Europe. 20 pp.
- 21/1990 Hindar, A. 1990. Chemistry and fish status of 67 acidified lakes at the coast of Aust-Agder, Southern Norway, in relation to postglacial marine deposits. 47 pp.
- 22/1990 Henriksen, A., Lien, L. and T. Traaen, T.S. 1990. Critical Loads for Surface Waters. Chemical Criteria for Inputs of Strong Acids. 45 pp.
- 23/1990 Wright, R.F. 1990. Reversibility of acidification: soils and surface waters. 39 pp.
- 24/1991 Wright, R.F. RAIN project. Report for the years 1988, 1989 and 1990. 156 pp.
- 25/1991 Wright, R.F., Holmberg, M., Posch, M. and Warfvinge, P. 1991. Dynamic models for predicting soil and water acidification: Application to three catchments in Fenno-Scandia. 40 pp.
- 26/1992 Wohlfeil, I.C. and Müller, D.I. 1992. RAIN-project: Vegetation mapping at Risdalsheia 1991. 42 pp.
- 27/1992 Rosseland, B.O. and Kroglund, F. 1992. Reversibility of acidification; Fish responses in experiments at Risdalsheia. 50 pp.
- 28/1992 Wathne, B.M., Norton, S.A. and Henriksen, A. 1992. Buffering capacities and effects of river substrates during acidic episodes. 24 pp.
- 29/1992 Henriksen, A., Mill, W.A., Kot, M., Rzychon, D. and Wathne, B.M. 1992. Critical loads of acidity to surface waters: A case study from Polish Tatra Mountains. 34 pp.
- 30/1992 Kroglund, F., Dalziel, T., Rosseland, B.O., Lien, L., Lydersen, E. and Bulger, A. 1992. Restoring Endangered Fish In Stressed Habitats. ReFISH project 1988-1991. 43 pp.
- 31/1992 Rosseland, B.O., Brandrud, T.E., Raddum, G.G. 1992. Effects of Aluminium in Acidified Aquatic Ecosystems. 34 pp.
- 32/1993 Henriksen, A., Forsius, M., Kämäri, J., Posch, M. and Wilander, A. 1993. Exceedance of Critical Loads for Lakes in Finland, Norway and Sweden: Reduction Requirements for Nitrogen and Sulfur Deposition. 46 pp.
- 33/1993 Wathne, B.M., Mill, W.A., Kot, M., Rzychon, D., Henriksen, A., and Tørseth, K. 1993. Critical loads of acidity to lakes in the Polish Tatra Mountains. A study of the yearly variations in the precipitation and water chemistry. 37 pp.
- 34-A/94 Lydersen, E. 1994. Long term monitored catchments in Norway - a hydrologic and chemical evaluation -, 306 pp. + Appendix.
- 34-B/94 Lydersen, E. 1994. Long term monitored catchments in Norway - a hydrologic and chemical evaluation -, Summary Report, 66 pp.
- 35/1994 Wright, R.F. 1994. Modelling long-term hydrochemical response at ENCORE catchments in the UK and Norway. 42 pp.
- 36/1994 Wright, R.F. 1994. RAIN PROJECT. Risdalsheia data report for June 1990 - May 1994. 165 pp.
- 37/1995 Závodský, D., Wathne, B.M., Henriksen, A., Sverdrup, H. et al. 1995. Mapping critical levels/ loads for the Slovak Republic. 74 pp.
- 38/1995 Wright, R.F. 1995. NITREX: Phase 2 Final Report 56 pp.

- 39/1995 Wright, R.F. 1995. NIVA contributions to ACID REIGN 95? Conference in Gothenburg, Sweden, 26-30 June 1995. 244 pp.
- 40/1996 Henriksen, A. et al. 1996. Regional Lake Surveys in Finland - Norway - Sweden - Northern Kola - Russian Karelia - Scotland - Wales 1995. 30 pp.
- 41/1996 Norton, S.A., Appleby, P.G., Dauvalter V. and Traaen T. 1996. Trace metal pollution in Eastern Finnmark, Norway and Kola Peninsula, Northwestern Russia as evidenced by studies of lake sediment. 18 pp.
- 42/1996 Wright, R.F. 1996. Nitrex: Phase 3 Final Report. 28 pp.
- 43/1996 Závodský, D., Wathne, B.M., Henriksen, A. and Sverdrup, H. 1996. Mapping critical levels/loads for the Slovak Republic. Final Report. 93 pp.
- 44/1997 Lydersen, E., Rzychon, D., Worsztynowicz, A., Tørseth, K. Hanssen, J.E., Mill, W.A., Kot, M., Henriksen, A., Wathne, B.M. 1997. Critical loads of acidity to lakes in the Polish Tatra Mountains- Final Report. 49 pp.
- 45/1997 Henriksen, A., Mannio, J., Wilander, A., Moiseenko, T., Traaen, T.S., Skjelkvåle, B.L.S., Fjeld, E. and Vuorenmaa, J. 1997. Regional Lake Surveys in The Barents region of Finland - Norway - Sweden and Russian Kola 1995 - Results.
- 46/1997 Kjønås, O.J., Wright, R.F., and Cosby, B.J. 1997. MERLIN model applied to NITREX Gårdsjön. 38 pp.
- 47/1997 Henriksen, A., Skjelkvåle, B.L., Mannio, J., Wilander, A., Jensen, J.P., Moiseenko, T., Harriman, R., Traaen, T.S., Fjeld, E., Vuorenmaa, J., Kortelainen, P. and Forsius, M. 1997. Results of national lake surveys 1995 in Finland, Norway, Sweden, Denmark, Russian Kola, Russian Karelia, Scotland and Wales. 43 pp.
- 48/1997 Lydersen, E., Angell, V., Eilertsen, O., Larssen, T., Mulder, J., Muniz, I.P., Seip, H.M., Semb, A., Vogt, R.D., Aagaard, P. 1997. Planning of an Integrated Acidification Study and Survey on Acid Rain Impacts in China. Final Report. 77 pp. + Appendix.

**Norwegian Institute for Water Research**

P.O. Box 173 Kjelsås    Telephone: + 47 22 18 51 00  
N-0411 Oslo            Telefax:     + 47 22 18 52 00

By ordering the report, please use  
serial number 3719-97.

ISBN 82-577-3287-7