

**Statlig program for forurensningsovervåking**  
**Langtidsovervåking av miljøkvaliteten langs Norges kyst**

Rapport: 944/2005  
TA-nummer: 2141/2005  
ISBN-nummer: 82-577-4822-6

Oppdragsgiver: Statens forurensningstilsyn  
Utførende institusjon: Norsk institutt for vannforskning

National Comments regarding  
the Norwegian data for 2004

**Rapport  
944/2005**

NIVA rapport nr. 5112--2005

## Norwegian Institute for Water Research

## REPORT

Main Office	Regional Office, Sørlandet	Regional Office, Østlandet	Regional Office, Vestlandet	Akvaplan-NIVA A/S
P.O. Box 173, Kjelsås N-0411 Oslo Norway Phone (47) 22 18 51 00 Telefax (47) 22 18 52 00	Televeien 1 N-4890 Grimstad Norway Phone (47) 37 29 50 55 Telefax (47) 37 04 45 13	Sandvikaveien 41 N-2312 Ottestad Norway Phone (47) 62 57 64 00 Telefax (47) 62 57 66 53	Nordnesboder 5 N-5008 Bergen Norway Phone (47) 55 30 22 50 Telefax (47) 55 30 22 51	Søndre Tollbugate 3 N-9000 Tromsø Norway Phone (47) 77 68 52 80 Telefax (47) 77 68 05 09

Title  Joint Assessment and Monitoring Programme (JAMP). National Comments regarding the Norwegian Data for 2004  (Norwegian State Pollution Monitoring Programme Report no. 944/2005. TA-no. 2141/2005)	Serial No. 5112-2005	Date 24.12.05
	Report No. Sub-No. O-80106 O-25106	Pages Price 235
Author(s)  Norman W. Green Anders Ruus Merete Schøyen Lise Tveiten Mats Walday	Topic group Marine ecology	Distribution
	Geographical area Oslofjord to Varangerfjord	Printed NIVA

Client(s)  Norwegian Pollution Control Authority (SFT)	Client ref.
--	-------------

## Abstract

This report is part of the Norwegian contribution to the SIME 2006 meeting administrated by OSPAR. JAMP 2004 included the monitoring of contaminants in sediment (17 stations), blue mussel (58), dogwhelk (20), cod (9) and flatfish (11) along the coast of Norway from Oslo to Bergen, Lofoten and Varangerfjord. The results indicated elevated levels of contaminants, i.e., poorer than Class I in SFT's classification system, or over provisional "high background", in the inner Oslofjord (PCBs, mercury and lead in cod; PCBs in mussels), and Sørkjosen and Hardangerfjord (cadmium, lead, mercury and DDT (ppDDE) in mussels, and mercury in cod). The results from the remaining stations showed low or moderate levels of contamination in 2004. Considering the whole monitoring period, significant upward trends were found for mercury in cod from the inner Oslofjord and a downward trend was found for lead in mussels from Sørkjosen/Hardangerfjord. The "Pollution" index was between "marked" and "severe," as it was in 2003, in the SFT system. The "Reference" index was between "slight" and "moderate" in the system, as before. Contamination of organotin in mussels and imposeshell in dogwhelk were still apparent. No significant improvement was registered in comparison to 1992 investigations. The results from studies using biological effects methods in cod (4 stations) are discussed.

4 keywords, Norwegian  1. Miljøgifter 2. Organismar 3. Marin 4. Norge	4 keywords, English  1. Contaminants 2. Organisms 3. Marine 4. Norway
--	--

Norman W. Green  
Project manager

Kristoffer Næs  
Research manager

Øyvind Sørensen  
Project director

ISBN 82-577-4822-6

NIVA report no. 5112--2005

OSPAR CONVENTION FOR THE PROTECTION OF THE MARINE ENVIRONMENT OF  
THE NORTHEAST ATLANTIC

WORKING GROUP ON CONCENTRATIONS, TRENDS AND EFFECTS OF SUBSTANCES  
IN THE MARINE ENVIRONMENT (SIME)

LONDON 21-23 FEBRUARY 2006

---

**O-80106 / O-25106**

**JOINT ASSESSMENT AND MONITORING PROGRAMME (JAMP)  
NATIONAL COMMENTS REGARDING  
THE NORWEGIAN DATA FOR 2004**

Oslo, 24. December 2005

Submitted by:

NORWAY  
Norman W. Green  
Norwegian Institute for Water Research (NIVA)  
PB 173 Kjelsås  
N-0411 Oslo

Submitted to:

Oslo and Paris Commissions (OSPAR)  
att. Richard Emmerson  
New Court  
48 Carey Street  
London WC2A 2JQ

Copy to:

Norwegian Pollution Control Authority (SFT)  
att. Per Erik Iversen  
PB 8100 Dep.  
N-0032 Oslo  
International Council for the Exploration of the Seas (ICES)  
att. Marilynn Sørensen  
Palægade 2-4  
DK-1261 Copenhagen



## ***Foreword***

*This report presents the Norwegian national comments on the 2004 investigations for the Joint Assessment and Monitoring Programme (JAMP). JAMP is administered by the Oslo and Paris Commissions (OSPAR) and their Environmental Assessment and Monitoring Committee (ASMO). JAMP receives guidance from the International Council for the Exploration of the Sea (ICES). ASMO has delegated implementation of part of the programme to the Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME). The Norwegian 2004 investigations are directed to particular JAMP issues relating to contaminants and implemented by SIME. JAMP replaced Joint Monitoring Programme (JMP) in 1995.*

*The Norwegian JAMP for 2004 was carried out by the Norwegian Institute for Water Research (NIVA) by contract from the Norwegian Pollution Control Authority (SFT), (NIVA contract O-80106, O-25106).*

*The Norwegian contribution to the JMP/JAMP was initiated by SFT in 1981 as part of the national monitoring programme. It now comprises three areas: the Oslofjord and adjacent areas (Hvaler-Singlefjord area and Langesundsfjord, 1981-), Sørfjord/Hardangerfjord (1983-84, 1987-) and Orkdalsfjord area (1984-89, 1991-93, 1995-96).*

*Since the North Sea Task Force Monitoring Master Plan was implemented in 1990, additional areas have also been monitored. These include: Arendal, Lista and Bømlo-Sotra areas. On the initiative of SFT and NIVA "reference" or merely diffusely contaminated areas from Bergen to Lofoten have been monitored since 1992 and from Lofoten to the Norwegian-Russian border from 1994.*

**Reader's guide.** The comments are presented in accordance with the agreed standardised format (ASMO 1997, Annex 12). Following the SIME meeting in London, 21-23 February 2006, the full report in PDF-format can be downloaded from either of two websites: the SFT's website and using SFT's TA-number at <http://www.sft.no/arbeidsomr/overvaking/rapporter/get.cfm?l=1&kat=8> or from NIVA's website at <http://www.niva.no/symfony/infoportal/portenglish.nsf> and doing a search on the "løpenr", which is the NIVA-report number for this report.

**Acknowledgments.** Thanks are due to many colleagues at NIVA, especially: Lise Tveiten, Merete Schøyen, Åse Kristine Rogne, Sigurd Øxnevad, Åse Bakketun, for field work, sample preparations and data entry; Alfild Kringstad, Merete Grung, Torgunn Sætre, and their colleagues for organic analyses; Bente Hiort Lauritzen and her colleagues for metal analyses; Randi Romstad and her colleagues for biological effects measurements, Gunnar Severinsen for data programme management and operation; and to the authors Anders Ruus (biological effects methods), Mats Walday (organotin), and Eva Hagebø and her colleagues (analytical quality assurance). Thanks go also to the numerous fishermen and their boat crews for which we have had the pleasure of working with.

*Oslo, 24 December 2005.*

*Norman W. Green  
Project co-ordinator*



# Contents

<b>1. GENERAL DETAILS</b>	<b>1</b>
<b>1.1. Executive Summary / Sammendrag</b>	<b>1</b>
<b>1.2. Introduction</b>	<b>3</b>
<b>1.3. Information on measurements</b>	<b>4</b>
1.3.1. Oslofjord area	5
1.3.2. Sørfjord and Hardangerfjord	10
1.3.3. Lista area	21
1.3.4. Bømlo-Sotra area	21
1.3.5. Orkdalsfjord area	21
1.3.6. Open coast areas from Bergen to Lofoten	23
1.3.7. Exposed area of Varangerfjord near the Russian border	23
1.3.8. Norwegian Pollution and Reference Indices (The Index Programme)	24
<b>1.4. Biological effects methods for cod</b>	<b>25</b>
1.4.1. OH-pyrene metabolites in bile	26
1.4.2. ALA-D in blood cells	28
1.4.3. EROD activity in liver	30
1.4.4. Concluding remarks	32
<b>1.5. Effects and concentrations of organotin</b>	<b>33</b>
1.5.1. Dogwhelk	33
1.5.2. Mussels	36
1.5.3. Concluding remark	36
<b>1.6. Overall conclusions</b>	<b>38</b>
<b>2. TECHNICAL DETAILS</b>	<b>41</b>
<b>2.1. Compliance with guidelines/procedures</b>	<b>41</b>
2.1.1. JAMP programme	41
2.1.2. Overconcentrations and classification of environmental quality	41
2.1.3. Comparison with previous data	44
2.1.4. The effect of depuration and freezing on mussels	45
<b>2.2. Information on Quality Assurance</b>	<b>45</b>
<b>2.3. Description of the Programme</b>	<b>46</b>
<b>3. REFERENCES</b>	<b>47</b>
Appendix A Quality assurance programme	51
Appendix B Abbreviations	57
Appendix C Analytical overview, sediment and biota	65
Appendix D Participation in intercalibration exercises	97
Appendix E Overview of localities and sample count for biota 1981-2004	103
Appendix F Overview of localities and sample count for sediment 1987-2004	115
Appendix G Map of stations	119
Appendix H Overview of materials and analyses 2004	135
Appendix I Temporal trend analyses of contaminants and biomarkers in biota 1981-2004	141
Appendix J Geographical distribution of contaminants and biomarkers in biota 1990-2004	173
Appendix K Geographical distribution of contaminants in surficial sediment 1987-2004	195
Appendix L Results from INDEX determinations 1995-2004	211



# 1. General Details

## 1.1. Executive Summary / Sammendrag

The Norwegian JAMP 2004 included the monitoring of micropollutants (contaminants) in sediment (17 stations), blue mussel (58), dogwhelks (20), cod (9) and flatfish (11) from the border of Sweden in the south along the coast of Norway to the Bergen area, Lofoten and the Varangerfjord bordering Russia. The mussel sites include supplementary stations for the Norwegian Index programme. The results showed several cases of levels of contaminants, higher than Class I in the Norwegian Pollution Control Authority's (SFT's) classification system or over provisional "high background"). The major cases were found in:

- Part of JAMP area 26: Oslofjord (up to Class III for PCBs and to a lesser extent lead and mercury), where cod liver from the inner Oslofjord was markedly polluted with PCB (Class III), and a significant upward trend was found for mercury in cod fillet from both "large" and "small" individuals and for cadmium in cod liver from the inner Oslofjord 1984-2004;
- JAMP areas 63 and 62: Sørfjord and Hardangerfjord (for mussels, up to Class IV for lead, Class III for cadmium and DDE, and for cod, Class II for mercury and ppDDE). A significant downward trend was found for lead in mussels at two stations in Hardangerfjord 1987-2004, and an upward trend was found for mercury in fillet in flounder. Surficial sediment is contaminated with mercury and PAH (Class V), TBT (Class IV) and to a lesser extent cadmium, lead, zinc and copper (Class II or III).

Part of JAMP area 26: Langesundsfjord has been an area of concern partly due to concentrations of HCB in mussels. However, since 2001 concentrations have been low or moderate (Class I and II), and a downward trend was found for the period 1990-2004, and there is less of a reason for concern when considering this contaminant.

Two environmental indices have been applied annually since 1995 to assess the levels of contamination in mussels from "polluted" and "reference" areas. The 2004 Pollution Index result was between the classes "markedly" and "severely" polluted in the Norwegian Pollution Control Authority's (SFT's) classification system, the same class as in 2003. The Reference Index was between the classes "slightly" and "moderately" polluted" (Class II), as in years prior to 2004.

The biological effects methods OH-pyrene (pyrene metabolite; marker for PAH exposure), δ-aminolevulinic acid dehydrase (ALA-D; marker for lead exposure), and the activity of cytochrome P4501A (EROD; marker for planar hydrocarbons, such as certain PCBs/PCNs, PAHs and dioxins) were determined in cod from three to four stations along the coast from the Oslofjord to Hardanger. With respect to OH-pyrene metabolites in 2004, the Oslofjord showed the most elevated levels of OH-pyrene. Furthermore, levels of OH-pyrene were higher at Lista, than in the Sørfjord, as observed prior to 2002. Results for ALA-D indicated exposure of lead to cod from the inner Oslofjord and inner Sørfjord. EROD activity in the inner Oslofjord was higher than in the less contaminated Sotra-Bømlo area, while EROD activity in the inner Sørfjord was lower. Previous years have also shown that EROD in fish from the Oslofjord and the Sørfjord are not consistently higher than at other presumed cleaner stations.

The presence of organotin (as TBT) in Norwegian waters was still a problem in 2004, most evident close to harbours, but also at stations remote from known point-sources. Concentrations of organotin in mussels and dogwhelk were elevated, and biological effects from TBT were found in dogwhelk from all of the investigated areas. However, TBT concentrations in mussels and dogwhelk were mostly lower than in 2003, but no significant trends were found, nor significant difference when compared to 1992 investigations. It is a cause for concern that the restrictions on the use of TBT in antifouling agents on vessels has not lead to a clear improvement in the investigated areas.

## Sammendrag

JAMP (Joint Assessment and Monitoring Programme) er et internasjonalt program for miljøovervåking av kystfarvann. Norge er et av tolv land som gjennom Oslo-Pariskonvensjonen (OSPAR) har forpliktet seg til å delta i dette felles overvåkingsprogrammet. Programmet i Norge startet i 1981 og hovedmålsettingen er å overvåke miljøgifter i påvirkede områder og ellers langs hele norskekysten. Resultatene fra de minst påvirkede områdene benyttes for å angi "bakgrunnsnivåer". Resultatene rapporteres årlig.

I 2004 omfattet JAMP undersøkelse av sediment (på 17 stasjoner), blåskjell (58, inkludert de til SFTs forurensningsindeks og til overvåking av TBT), purpurnegl (20), torsk (9) og flatfisk (11) fra svenskegrensen i syd til Bergen, Lofoten og Varangerfjorden mot den russiske grensen. Resultatene tydet på forhøyede konsentrasjoner av miljøgifter, dvs. mer enn Klasse I i SFTs klassifiseringssystem, eller over antatt "høyt bakgrunnsnivå". Disse tilfellene ble registrert i:

- Oslofjorden med opp til Kl.III for PCB og i mindre grad bly og kvikksølv, hvor torskelever fra indre Oslofjord var markert forurenset med PCB (Kl.III), og det ble funnet signifikant økende trender for kvikksølv i torskefilet fra både "store" og "små" individer og for kadmium i torskelever fra indre Oslofjord 1984-2004;
- Sørkjorden og Hardangerfjorden for blåskjell med opp til Kl.IV for bly, Kl.III for kadmium og DDE og for torsk Kl.II når det gjaldt kvikksølv og kadmium. Det ble funnet en signifikant avtagende trend for bly i blåskjell på to stasjoner i Hardangerfjorden 1987-2004, og en økende trend for kvikksølv i skrubbefilet. Overflatesediment er forurenset med kvikksølv og PAH (Kl.V), TBT (Kl.IV), og i mindre grad kadmium, bly, zinc og kobber (Kl.II eller Kl.III).

Langesundsfjorden har vært et område med bl.a. høye konsentrasjoner av HCB i blåskjell. Men konsentrasjonene siden 2001 har vært lav eller moderat (Kl.I eller II), og en avtagende trend ble funnet for perioden 1990-2004, slik at det er mindre grunn til bekymring når det gjelder dette stoffet.

SFTs blåskjell-forurensningsindeks og blåskjell-referanseindeks har blitt brukt årlig siden 1995 på en gruppe "forurensede" og "referanse"- fjordområder. Forurensningsindeksen for 2004 betegnet sin gruppe mellom "markert" og "sterk" forurenset, den samme klassen som i 2003. Referanseindeksen har klassifisert sin gruppe mellom "lite" og "moderat" forurenset i hele perioden..

Følgende biologiske effekt-parametre ble undersøkt i torsk fra tre-fire stasjoner langs kysten fra indre Oslofjord til Hardanger: OH-pyren (pyren-metabolitt; markør for PAH-eksponering), δ-aminolevulinsyre dehydrase (ALA-D; markør for bly-eksponering), og aktivitet av cytokrom P4501A (EROD; markør for plane hydrokarboner, slik som spesifikke PCB/PCN, PAH og dioksiner).

Oslofjorden viste de høyeste OH-pyren-nivåene. Videre var OH-pyren nivåene ved Lista høyere enn i Sørkjorden, slik det har vært observert før 2002. Resultatene for ALA-D indikerte bly-eksponering for torsk fra indre Oslofjord og indre Sørkjord. EROD aktivitet i indre Oslofjord var høyere enn i det mindre forurensede Sotra-Bømlo området, mens EROD aktivitet i indre Sørkjorden var lavere.

Tidligere år har også vist at EROD i fisk fra Oslofjorden og Sørkjorden ikke er konsistent høyere enn på andre, antatt mindre forurensede stasjoner.

Effekter av organotin (bl.a. TBT) kunne fortsatt registreres i 2004, tydeligst i havner eller i områder med mye skipstrafikk, men også på stasjoner som var antatt lite påvirket. Konsentrasjoner av TBT i blåskjell og purpurnegl var forhøyet, og virkning av TBT (imposex) ble registrert på samtlige stasjoner. Ingen tydelig utvikling i imposex over tid ble registrert, men konsentrasjoner i blåskjell og purpurnegl var lavere enn tidligere år. Noen tidstrend for perioden 1997-2004 kunne imidlertid ikke påvises. Forbud mot bruk av TBT som begroingshindrende middel på båter har ikke ført til klar forbedring i de undersøkte områdene.

## 1.2. Introduction

The Norwegian contribution to the “Joint Assessment and Monitoring Programme (JAMP) was initiated by the Norwegian Pollution Control Authority (SFT) and is integrated with SFT’s State Pollution Monitoring Programme. The procedures and practice of JAMP has also provided a basis for other investigations of interest to SFT but not necessarily requested by JAMP (e.g. SFT’s Index Programme (Pollution and Reference Indices), chapter 1.3.8).

Data are submitted to ICES under three categories: for Purpose A (health assessment) on a voluntary basis, Purpose C (spatial distribution) on a voluntary basis and Purpose D (temporal trend assessment) on a mandatory basis. Where practical, data collection was in accordance to agreed procedures (OSPAR 1990, 1997). Data were screened and submitted to ICES in accordance with procedures outlined by ICES (1996).

This report focuses on issues and situations in Norway concerning contaminants and considered of interest to the implementation of JAMP (Table 1). It should be noted that these issues are being revised (cf., MON 2001). The Norwegian programme for JAMP 2003 has been outlined previously (Green 2003).

**Table 1.** Extract from list of JAMP issues, subjects and descriptions to which the Norwegian investigations for 2003 can be addressed (cf. ASMO 1997, Annex 30).

Issue	Subject	Description
1.2	Hg, Cd, and Pb	What are the concentrations and fluxes in sediments and biota?
1.3	TBT	To what extent do biological effects occur in the vicinity of major shipping routes, offshore installations, marinas and shipyards?
1.7	PCBs	Do high concentrations pose a risk to the marine ecosystem?
1.8	PCBs	Do high concentrations of non-ortho and mono-ortho CBs in seafood pose a risk to human health?
1.10	PAHs	What are the concentrations in the maritime area?
1.11	PAHs	Do PAHs affect fish and shellfish?
1.12	Other synthetic compounds	How widespread are synthetic organic compounds within the maritime area?
1.15	Chlorinated dioxins and dibenzofurans	What concentrations occur and have the policy goals (for the relevant parts of the maritime area) been met?
1.17	Biological effects of pollutants	Where do pollutants cause deleterious biological effects?
5.3	Chemical used [mariculture]	In which areas do pesticides and antibiotics affect marine biota?
6.1	Ecosystem health	How can ecosystem health be assessed in order to determine the extent of human impact?

The chapter structure of this report for the first and second level is according to agreed format (ASMO 1997, Annex 12) which *inter alia* presents results before methodology.

### 1.3. Information on measurements

An overview of JAMP stations in Norway is shown in the tables in Appendix E and maps in Appendix G. The stations and sample counts relevant to the 2004 investigations are noted in the tables in Appendix E. Data reports have been published recently for sediment 1986-1997 (Green *et al.* 2002a) and biota 1981-2002 (Green *et al.* 2002b-d).

Sediment was samples at 17 stations, blue mussel at 58 stations (including supplementary stations for Index and TBT), dogwhelk at 20, cod at 9 and flatfish at 11 stations from the border to Sweden in the south to the border to Russia in the north. Generally, mussels are not abundant on the exposed coastline from Lista (south Norway) to the North of Norway. A number of samples were collected from dock areas, buoys or anchor lines.

This chapter focuses on the principle cases where *median* concentrations exceeded provisional "high background" ("normal"). The median concentration can be derived from the tables in Appendix I or figures in Appendix J and Appendix K, depending on the year and concentration basis in question. Where possible, these medians are classified according to the Norwegian Pollution Control Authority's (SFT's) **environmental quality classification system** (cf. Molvær *et al.* 1997). An extract of the system that is applied in this report is shown in Table 5 and Table 6 and includes unofficial conversion to other bases. The system does not cover some contaminants for some species-tissues, however provisional "high background" concentrations have been determined and these are listed in Table 7. "High background" concentrations set the upper limit for Class I in SFT's system. The factor by which concentrations exceeded "high background" is termed **overconcentration**. "High background" concentration corresponds to the upper limit to Class I, or slightly (also stated in classifications system as "insignificantly") polluted. Below the median concentrations are assessed according to the SFT system, but where this is not possible overconcentrations are used. The term "significant" refers to the results of a statistical analysis of linear trends shown in Appendix I. More details concerning these terms and methods can be found in chapter 2.1.2.

### 1.3.1. Oslofjord area

Mussels from the inner Oslofjord were moderately polluted with  $\Sigma$ PCB-7 (SFT's Class II, Figure 1A). Cod liver from the inner Oslofjord was markedly polluted with  $\Sigma$ PCB-7 (Class III, Figure 2A). The median concentration in cod liver was 2620  $\mu\text{g}/\text{kg}$  w.w., higher than in 2003. Cod liver from the outer Oslofjord was slightly polluted with regard to  $\Sigma$ PCB-7 (st.36B, Figure 2B).

In 1994, and renewed in 2004, the Norwegian Food Safety Authority (SNT) advised not to consume liver of cod from the inner Oslofjord (north of st.31A, see Map 1 in Appendix G) due to concerns about PCB contamination (cf. Table 3).

A significant linear *downward* trend was detected (see method description in chapter 2.1.3) for  $\Sigma$ PCB-7 in mussels from the inner Oslofjord (30A and 31A Figure 1A, B) for the period 1988 to 2004.

Power analyses (see chapter 2.1.3) indicated that a hypothetical trend of 10% change per year in  $\Sigma$ PCB-7 concentration in the blue mussel from the mid and inner Oslofjord would take 10 to 13 years to be detected with 90% significance (Appendix I).

The fillet of "small"<sup>1</sup> (29-44 cm) and "large" cod (45-70 cm) from the inner Oslofjord in 2004 were moderately polluted with mercury (Class II, Figure 3A, B). A significant *upward* trend was detected for the period 1984-2004 for both size groups, even though the concentrations in 2002-2004 were lower than the three previous years. No significant trend was found for period 1998-2004.

Considering the entire period, the power, indicated as number of years to detect a hypothetical 10% change per year for mercury in cod fillet from either station, was slightly better for "small" fish (10-11 years) than "large" fish (13 years) (cf. Appendix I). Concentrations of mercury were significantly higher in "large" cod compared to "small" cod.





Median concentration of cadmium and lead in cod liver from the inner Oslofjord (30B) 2004 was 0.10 and 0.17  $\text{mg}/\text{kg}$  w.w., respectively. For lead, this was less than a third of the concentration found in 2002, which was the second highest found during the entire period (1990-2004). "High background" for these metals is 0.1  $\text{mg}/\text{kg}$  w.w. A significant *upward* trend was found for cadmium in these cod. A significant *upward* trend was also found for cadmium in mussel from one station in the inner Oslofjord (st. 30A). These mussels were moderately polluted with respect to both cadmium and lead in 2004.

The SFT's environmental quality classification system does not include cadmium and lead in cod liver.





It should be noted that the Index programme indicated severe concentrations of TBT in mussels from a station located in the inner Oslofjord (see chapter 1.3.8).

---





<sup>1</sup> The size of "small" and "large" cod depends on the station-year catch, and hence may vary (see section 2.1.3. The range given is the lower and upper quartile of the median lengths of the "small" or "large" fish.

**A** $\Sigma\text{PCB-7}$ , *Mytilus edulis*, 30A Inner Oslofjord**B** $\Sigma\text{PCB-7}$ , *Mytilus edulis*, 31A Solbergstrand**C** $\Sigma\text{PCB-7}$ , *Mytilus edulis*, 35A Mølen**D** $\Sigma\text{PCB-7}$ , *Mytilus edulis*, 36A Outer Oslofjord (Færder)

**Figure 1.** Median  $\Sigma\text{PCB-7}$  (sum of PCB 28, 52, 101, 118, 138, 153 and 180) concentration in blue mussel (*Mytilus edulis*) from inner (st.30A) to outer (st.36A) Oslofjord. (cf. Appendix G and Appendix I, and key in Figure 21).

**A** $\Sigma\text{PCB-7}$ , *Gadus morhua*, liver, 30B Inner Oslofjord**B** $\Sigma\text{PCB-7}$ , *Gadus morhua*, liver, 36B Outer Oslofjord**C** $\Sigma\text{PCB-7}$ , *Gadus morhua*, filet, 30B Inner Oslofjord**D** $\Sigma\text{PCB-7}$ , *Gadus morhua*, filet, 36B Outer Oslofjord

**Figure 2.** Median  $\Sigma\text{PCB-7}$  (sum of PCB 28, 52, 101, 118, 138, 153 and 180) concentration in liver and fillet of cod (*Gadus morhua*) from the inner (st.30B) to outer (st.36B) Oslofjord. (cf. Appendix G and Appendix I, and key in Figure 21).

**A****B****C****D**


**Figure 3.** Median mercury (Hg) concentration in fillet of cod (*Gadus morhua*): for the inner Oslofjord (st.30B) “small” (**A**) and “large” (**B**) fish, and for the outer Oslofjord (st.36B) “small” (**C**) and “large” (**D**) fish. (cf. Appendix G and Appendix I, and key in Figure 21).

Mussels from Langesundsfjord (st. 71A) in 2004 were slightly polluted with HCB (Class II, Figure 4). The median concentration for 2004 was higher compared to 2003, which was the lowest at this station since 1983. Concentrations have varied greatly since 1983 but median value have decreased distinctly since 1989 (Figure 4) due to about 99% reduction in discharge of HCB and other organochlorines from a magnesium factory (cf. Knutzen *et al.* 2001).

The power of the monitoring programme was 15 years for the period 1990-2004 and more than 25 years for the entire period (cf. Appendix I). The 1983-2004 data series and the 1990-2004 data series both had significant *downward* trends.

Extremely high lindane concentrations (Class V) were found in blue mussel from st. 71A in 2002 (cf. Green *et al.* 2004a), but the concentrations were low (Class I) in both 2003 and 2004. The reason for high values in 2002 has not been determined.

It should be noted that dioxin is one of the contaminants monitored to establish the Pollution Index (see section 1.3.8). Dioxin toxicity equivalents based on the Nordic model (TCDDN) in mussels from 71A were marked (SFT Class III) and severe values were found at a nearby Index station (I712) (Figure 34).



**Figure 4.** Median HCB concentration in blue mussel (*Mytilus edulis*) from Langesundsfjord (west of Oslofjord). (cf. Appendix G and Appendix I, and key in Figure 21). Vertical line indicates when a magnesium factory reduced it's discharge by 99%. **NB: log-scale.**

### 1.3.2. Sørfjord and Hardangerfjord

The development of the contaminant conditions in these connected fjords and the main remedial actions that have been taken, have been outlined in the JAMP National comments for 1989 (Green 1991)) and in recent reports concerning Sørfjord in particular (Skei 2000, 2001, Skei & Knutzen 2000, Skei *et al.* 1998). The results from JAMP 2004 are coupled to other studies in this area (cf. Knutzen & Green 2001a, Ruus & Green 2002, 2003, 2004, 2005) and confirm that the Sørfjord, and in some cases also Hardangerfjord, continue to be contaminated especially with cadmium (Figure 5 and Figure 6), lead (Figure 7 and Figure 13), mercury (Figure 8, Figure 9 and Figure 13), ppDDE (Figure 10, Figure 11 and Figure 12). PAH (represented by benzo[*a*]pyrene (Figure 13)) and to a lesser extent PCB (Figure 12).

In 2002 the Norwegian Food Safety Authority (*Mattilsynet*, earlier referred to as SNT) extended their advice against the consumption of mussels to include all seafood in the Sørfjord including deep-water fish due to concerns about metal and PCB contamination (Table 3).

Results for mussels collected from the Sørfjord indicated that these were moderately (Class II) or markedly polluted (Class III) with cadmium in respect to SFT's classification system (Figure 5, Appendix I). Mussels as far as Krossanes at the mouth of Sørfjord (st.57A) and about 35 km from Odda, were moderately polluted with cadmium (Figure 5). In 2003 this limit was as far as Lille Telløy in Hardangerfjorden (100 km from Odda). A significant *downward* trend was found for cadmium at three stations in Sørfjord (st.52A, 56A and 57A) and two in Hardangerfjord (st.63A and 65A) (Appendix I). Also, the median lead concentration at one station nearest Odda (st.51A) was severe (Class IV), whereas the other three stations in the Sørfjord were moderately or markedly polluted with lead. A *downward* trend was found for lead at st. 63A and 65A, 1990-2004. Three stations in Sørfjord were moderately polluted with respect to mercury.

Cod fillet from "small" (33-39 cm) and "large" individuals (40-47 cm) from the inner Sørfjord (st.53B) were moderately polluted with mercury (Class II). Overconcentrations for mercury were found in fillet in flounder (4-5 times "background") and an *upward* trend was also found. Overconcentrations were found for cadmium in cod liver and flounder liver from inner Sørfjord (3-5 times). Overconcentrations of lead in flounder liver were also observed (2 times).

The power of the sampling strategies for mussels was relatively poor for samples collected from Odda; the innermost part of Sørfjord (st.51A or 52A). For example for lead in mussels, it is estimated that it would take 19-23 years to detect a hypothetical trend of 10% per year with 90% significance (Appendix I). This reflects the large variability found in the data series from this area. The variability is mostly due to the irregular/accidental input of contaminated discharges. The power improved with distance from Odda, and at Ranaskjær (st.63A, ca.50 km from Odda) it was only 13 years.

Mussels at one station (st.56A) near the outer Sørfjord were markedly polluted with ppDDE (Class III); with a median concentration of 49.3 µg/kg d.w. The lower limit to Class IV, severely polluted, is 50 µg/kg d.w. Mussels elsewhere in the Sørfjord were moderately polluted (Class II) (Figure 10 and Figure 11). Cod fillet and liver from the Sørfjord were only moderately polluted with ppDDE (Figure 12A, Appendix I).

The liver of cod from Hardangerfjord for 2004 were only slightly polluted (Class I) with respect to ΣPCB-7. Since JAMP monitoring started in the Sørfjord and Hardangerfjord the median values have varied between 100 and 2400 µg/kg w.w. (Appendix I). This indicated that cod is subject to a variable exposure from PCB, but the cause of this variation is not clear.





No trends were evident for ppDDE and ΣPCB-7 in mussels and cod. A *downward* trend was found for these substances in flounder fillet from inner Sørfjord.

Sediment has been sampled from seven stations from the Sørfjord-Hardangerfjord region in 2004, and also previously in 1990 and 1997 (Appendix F and Appendix K). The position for sampling near Strandebarm (st.67S) was moved in 2004 about 250 m to avoid submerged power lines. The water depth was 680 m, about 30 m deeper than previous sampling. The sampling near Krossanes at the mouth of Sørfjorden was mistakenly taken 600 m east of previous sampling. The water depth was 170 m and 135 m shallower than previously. Comparisons between stations were based on concentrations in surficial sediment; usually 0-1 cm, but 0-2 cm for analyses of organic contaminants prior to 2004.




Surficial sediment from the inner Sørfjorden (st.52S) was extremely polluted with mercury and PAH (Class V), severely polluted with cadmium and lead (Class IV), markedly polluted with zinc and TBT (Class III) and moderately polluted with copper (Class II). Concentrations of these contaminants generally decreased with distance from Odda, but still above Class I well into Hardangerfjorden (Figure 13).

Concentrations of PCBs and organic pesticides were low (Class I).





Sedimentation rate is about 1 mm per year (Green *et al.* 2002a), and there has not been sufficient time nor sampling to do an adequate temporal trend assessment for the period of investigation (1990-2004). Furthermore, the problem of such an assessment is compounded for the organic contaminants because of the above mentioned change in surface sample depth (0-1 vs. 0-2 cm). Taking these aspects into consideration, there is some evidence that concentrations of PAH have decreased from 1990 to 2004 (e.g. PAH compound benzo[*a*]pyrene in Figure 13). It should also be noted that where surficial sediment was markedly polluted with DDE before (Class III), it was only slightly polluted in 2004 (Class I).

**A**Cd, *Mytilus edulis*, 51A Byrknes**B**Cd, *Mytilus edulis*, 52A Eitrheimsneset**C**Cd, *Mytilus edulis*, 56A Kvalnes**D**Cd, *Mytilus edulis*, 57A Krossanes





**Figure 5.** Median cadmium (Cd) concentration in blue mussel (*Mytilus edulis*) from inner (st.51A) to outer (st.57A) Sørfjord. NB: (cf. Appendix G and Appendix I, and key in Figure 21). Note: for some years the upper confidence interval line is off-scale in figures A-C. Note: horizontal lines for Classes I and II are near x-axis.

**A**Cd, *Mytilus edulis*, 63A Ranaskjær**B**Cd, *Mytilus edulis*, 65A Vikingneset**C**Cd, *Mytilus edulis*, 69A Lille Terøy





**Figure 6.** Median cadmium (Cd) concentration in blue mussel (*Mytilus edulis*) from Hardangerfjord (st. 63A, 65A and 69A). (cf. Appendix G and Appendix I, and key in Figure 21). **Note: horizontal lines for Classes I and II are near x-axis.**

**A**Pb, *Mytilus edulis*, 51A Byrkenes**B**Pb, *Mytilus edulis*, 52A Eitrheimneset**C**Pb, *Mytilus edulis*, 56A Kvalnes**D**Pb, *Mytilus edulis*, 57A Krossanes





**Figure 7.** Median lead (Pb) concentration in blue mussel (*Mytilus edulis*) from inner (st.51A) to outer (st.57A) Sørkjord. (cf. Appendix G and Appendix I, and key in Figure 21). Note: horizontal lines for Classes I and II are near x-axis.

**A**Hg, *Mytilus edulis*, 51A Byrkenes**B**Hg, *Mytilus edulis*, 52A Eitrheimneset**C**Hg, *Mytilus edulis*, 56A Kvalnes**D**Hg, *Mytilus edulis*, 57A Krossanes




**Figure 8.** Median mercury (Hg) concentration in blue mussel (*Mytilus edulis*) from inner (st.51A) to outer (st.57A) Sørfjord. (cf. Appendix G and Appendix I, and key in Figure 21). Note: for some years the upper confidence interval line is off-scale in figure A. Note: horizontal lines for Classes I and II are near x-axis.

**A****B****C****D**





**Figure 9.** Median mercury (Hg) concentration in fillet of cod (*Gadus morhua*): from Sørfjord (st.53B) for “small” (**A**) and “large” (**B**) fish and Hardangerfjord (st.67B) for “small” (**C**) and “large” (**D**) fish (cf. Appendix G and Appendix I, and key in Figure 21). **Note:** for some years the upper confidence interval line is off-scale in Figure B.

**A**ppDDE, *Mytilus edulis*, 51A Byrknes**B**ppDDE, *Mytilus edulis*, 52A Kvalnes**C**ppDDE, *Mytilus edulis*, 56A Kvalnes**D**ppDDE, *Mytilus edulis*, 57A Krossanes


**Figure 10.** Median ppDDE concentration in blue mussel (*Mytilus edulis*) from inner (st.51A) to outer (st.57A) Sørkjord. (cf. Appendix G and Appendix I, and key in Figure 21). Note: horizontal line for Class I is near x-axis.

**A**ppDDE, *Mytilus edulis*, 63A Ranaskjær**B**ppDDE, *Mytilus edulis*, 65A Vikingneset**C**ppDDE, *Mytilus edulis*, 69A Lille Terøy

**Figure 11.** Median ppDDE concentrations in blue mussel (*Mytilus edulis*) from Hardangerfjord (st. 63A, 65A and 69A). (cf. Appendix G and Appendix I, and key in Figure 21). **Note:** horizontal line for Class I is near x-axis.

**A**ppDDE, *Gadus morhua*, liver, 53B Inner Sørkjorden**B**ppDDE, *Gadus morhua*, liver, 67B Strandebarm (Hardangerfjord)**C** $\Sigma$ PCB-7, *Gadus morhua*, liver, 53B Inner Sørkjorden**D** $\Sigma$ PCB-7, *Gadus morhua*, liver, 67B Strandebarm (Hardangerfjord)

**Figure 12.** Median ppDDE and  $\Sigma$ PCB-7 concentrations in liver of cod (*Gadus morhua*) from Sørkjorden (st.53B) and Hardangerfjord (st.67B) (cf. Appendix G and Appendix I, and key in Figure 21). Note that for 1989 the upper confidence interval line is off-scale in Figure A.



**Figure 13.** Median concentration of mercury (**A**), lead (**B**) and benzo[a]pyrene (**C**) in surficial sediment (0-1 cm (alt. 0-2 cm)) ordered by distance from near Odda (st.52S) to Bømlo area (22S), almost 200 km seaward. The Station order is 52S, 56S, 57S, 63A, 65A and 22S and the results are from 1990 (dotted line or open circles), 1997 (dashed line) and 2004 (solid line). (cf. Appendix K, and key in Figure 21). **Note: log scale.**

### **1.3.3. Lista area**

Median concentrations of contaminants in mussels, cod and dab were slight, with one exception (moderate (Class II) for mercury in "large" dab) and no upward trends were found (st.15, Appendix I and Appendix J).

### **1.3.4. Bømlo-Sotra area**




It was impractical to continue sampling for flatfish at st.22F Borøyfjorden. Thus, a new station in Åkrafjorden, 21F Kyrping, was initiated in 2000. This station is located about 82km south-east of 22F, but like 22F, 21F is located in a reference area.

Mussels, cod and flounder from this area (22A, 23B, 21F) generally were only slightly polluted (Class I) or showed no overcentrations with respect to metals or organochlorines (Appendix I and Appendix J). One exception was the unexpectedly the high concentration of ppDDE in the single bulk sample of flounder liver from Kyrping. This was the highest found in not only flounder but also plaice and cod in this area (both for lever and fillet) since 1990. The concentrations was three times background and higher than any found in the Sørhfjord or Hardangerfjord for this species-tissue. The median concentration in flounder fillet from Kyrping was also above background (1.2 times).

### **1.3.5. Orkdalsfjord area**

Mussels from this area were monitored for the period 1984-1996, and then not again until 2004 when bulk samples from four stations were investigated (Trossavika – st.84A, Flakk – 82A or Ingdalsbukt – 87A). Median concentrations found in 2004 can be classified as "slight" in SFT's system (Class I), as found before.

Sediment was sampled from four stations from the Orkdalsfjord region in 2004, and also previously in 1987 and 1992 (Appendix F and Appendix K). Surficial sediment from the inner Orkdalsfjorden (st.89S) was markedly polluted with cadmium and copper (Class III) and moderately polluted with lead and zinc (Class II) (Appendix K, Figure 14). The concentrations of ΣPCB-7, PAH and TBT were low (Class I). No temporal trends were evident for the period 1987-2004.

**A****B****C**

**Figure 14.** Median concentration of cadmium (**A**), copper (**B**) and lead (**C**) in surficial sediment (0-1 cm (alt. 0-2 cm)) ordered by distance from near Orkanger(st.89S) to Trondheimsfjorden (90S), about 20 km seaward. The Station order is 89S, 84S, and 90S and the results are from 1987 (dotted line), 1992 (dashed line) and 2004 (solid line). (cf. Appendix K, and key in Figure 21).

### **1.3.6. Open coast areas from Bergen to Lofoten**

This stretch of coastline covers 7° of latitude to 68°N (Appendix G). Fourteen mussel stations were investigated in 2004, thirteen of these not since the period 1990-1993. The longest times series was obtained from mussels collected at the island of Skrova (st. 98A). Samples were collected in 1992-1993, however, during the period 1994-1996 mussels were not found at this station, but nearby in the Skrova harbour (98X). Since 1997 a "new" 98A location was established at Husvågen, roughly 18 km north of Skrova, in a small fjord remote from any apparent point source of contamination, and hence considered comparable.

In 2004, the mussels were only slightly contaminated (SFT's Class I), which was generally the case for the period 1997-2003. "Large" cod were moderately contaminated with mercury (Class II) (Appendix I and Appendix J).

The median concentrations in mussels from Sætervik (st.93A) 2004 were moderate (Class II) for ΣPCB-7. PCB and pesticides were not measured at this station previously.

Sediment was sampled from Sotra (st.24S) to Lofoten (st.98S) in 2004; a total of six stations (24S, 27S, 93S, 95S, 99S and 98S). Most of these stations were also investigated previously in 1990-1992 (Appendix F and Appendix K). Surficial sediment was markedly polluted with TBT at Sotra (st.24S), east of Statlandet (27S) and northeast of Raudøya (st.93S) (Class III) and moderately polluted with lead, HCB, benzo[a]pyrene east of Statlandet (Class II) (Appendix K). No temporal trends were evident for the period 1990-2004.

### **1.3.7. Exposed area of Varangerfjord near the Russian border**

The remaining and northern area of JAMP in Norway stretches north of 68°N and east from a longitude of 17 to 29°E (Appendix G). In 2004 only two mussel stations, one cod station and one plaice station were investigated in the Varangerfjord (at approximately 70°N).

Slight overconcentrations (less than 2 times "high background") of cadmium were found in liver of plaice (st.10F) (Appendix I and Appendix J).

### 1.3.8. Norwegian Pollution and Reference Indices (The Index Programme)

The Norwegian Pollution Control Authority (SFT) has requested a specific and small group of indices to assess the quality of the environment with respect to contaminants - The Index Programme. One index is based on the levels and trends of contaminant concentrations in blue mussel collected annually from a selection of the more contaminated fjords in Norway (Appendix L). SFT has also requested the testing of this index against "reference" stations from selected areas and fjords.

The Index scale varies from 1, when all areas or fjords are slightly polluted (Class I in SFT's environmental quality classification system (Molvær *et al.* 1997)), to 5, in which at least one sample from each area or fjord could be classified as extremely polluted or Class V in SFT's system.

Nine fjord areas were used to calculate the Pollution Index. Taking the supplementary stations (Strømtangen, Flåøya, Moholmen, and Toraneskaien) and analyses (TBT and dioxin) into consideration the Index was 3.4 for 2004 compared to 3.6 for 2003 (cf. Appendix L). A value between 3 and 4 would be between "Markedly" and "Severely" polluted in the SFT classification system. Indices calculated with and without supplementary stations and analyses has been presented earlier (cf. Green *et al.* 2004a, b).

Six areas were included in Reference Index for 2004 compared to five for 1998-2003, and seven-eight fjords used in previous years. With the new calculation where supplementary analyses of TBT are included, the Reference Index was 1.3 for 2004, compared to 1.8 for 2003. Comparison between the old and new calculations has been done for 2002 and 2003 (cf. Green *et al.* 2004a, b). A value between 1 and 2 would be classified between "Slight" and "Moderate" in SFT's system. Five of the six fjords/areas included TBT analyses.

The use of the indices to assess the general level of pollution in contaminated or reference areas of coastal water for the period 1995 to 1999 has been reviewed (Green & Knutzen, 2001). The conclusions were mainly that the sample and analytical strategies lacked adequate coverage of the relevant contaminants and geographical areas. Furthermore, the report suggested supplementing the assessment of this type with relevant analyses of sediment. In 2002 the programme was improved by including more stations and parameters relevant to the blue mussel Pollution Index.

It is not the intent of the application of the indices to give a station by station account, however, time trend analyses for the entire period (1995-2004) has been calculated and show both significant upward and downward trends in mussels (cf. Appendix I). Some cases are worth noting:

- I024 Kirkø in the Hvaler area (Map 1, Appendix G), where a *downward* trend for ΣPCB-7 was found, and where moderate concentrations (Class II) were found previously,
- 30A Gressholmen and I307 Ramtonholmen in the inner Oslofjord (Map 1, Appendix G), where *upward* trends in cadmium were found and where the 2004 median at 30A was in Class II,
- 30A Gressholmen, where a *downward* trend in ΣPCB-7 was found and where the 2004 median was in Class II,
- I306 Håøya and I307 Ramtonholmen in the inner Oslofjord, where *upward* trends in benzo[a]pyrene were found and where the 2004 median at I307 was in Class II,
- 71A Bjørkøya in the Langesund area (Map 3, Appendix G), where a *downward* trend in HCB was found and where the 2004 median was still in Class II,
- I713 Strømtangen also in the Langesund area (Map 3, Appendix G), where an *upward* trend in ppDDE was found but where the 2004 median was still in Class I,
- I133 Odderø in the Kristiansand harbour (Map 4, Appendix G), where a *downward* trend for ΣPCB-7 was found, and where moderate concentrations (Class II) were found previously,
- 52A Eitrheimsneset in the inner Sørfjord (Map 6, Appendix G) where the 2004 median for cadmium was in Class III and a *downward* trend detected.

## 1.4. Biological effects methods for cod

The JAMP-programme for 2004 included five biological effects methods (BEM): OH-pyrene, ALA-D, EROD, CYP1A and TBT (Table 2). The first four are discussed in this chapter (Figure 15 to Figure 17) and TBT is discussed separately (cf., section 1.5). Results for OH-pyrene, ALA-D, EROD and metallothionein (MT) in cod and flatfish, 1997-2001, have been published earlier (Ruus *et al.* 2003). For the 2004 investigations OH-pyrene, ALA-D, EROD and CYP1A were measured in Atlantic cod in the inner Oslofjord (30B), Sørfjord (st.53B), and Sotra-Børnlo area (23B). OH-pyrene was also measured in cod outside Lista (15B). It has become clear that cod caught in the open coastal area outside Lista are more strongly affected by PAHs than cod at the other stations, despite the large water exchange in that area (Ruus *et al.* 2003). Furthermore, stations from the inner Oslofjord and Sørfjord are considered to be more contaminated with metals and organochlorines than the other stations.

**Table 2.** Summary of biological methods employed by the JAMP-2004.

Code	Name	tissue sampled	Specificity
OH-pyrene	Pyrene metabolite	fish bile	PAH
ALA-D	$\delta$ -aminolevulinic acid dehydrase inhibition	fish red blood cells	Pb
EROD	Cytochrome P4501A-activity (CYP1A/P4501A1, EROD)	fish liver	planar PCB/PCNs, PAHs, dioxins
CYP1A	Relative amount of cytochrome P450 1A-protein	fish liver	Supporting parameter for EROD
TBT	Imposex/Intersex	snail soft tissue	organotin

The reason to use biological effects methods within monitoring programmes is to evaluate whether marine organisms are affected by contaminant inputs. Such knowledge can not be derived from tissue levels of contaminants only. In addition to enable conclusions on the health of marine organisms, some biomarkers assist in the interpretation of contaminant bioaccumulation. The biological effects component of the Norwegian JAMP is possibly the most extensive of its type in Europe and includes imposex in gastropods as well as biomarkers in fish. The four chosen methods for fish were selected for specificity, for robustness and because they are among a limited set of methods proposed by international organisations, including OSPAR and ICES.

Under controlled conditions the measures derived from OH-pyrene, EROD and CYP1A increase with increased exposure to their respective inducing contaminants. The activity of ALA-D on the other hand is inhibited by contamination (i.e., lead), thus lower activity means higher exposure.

As in most previous years, 25 individual cod were sampled for biological effects measurements at each station. However, since 2002 only three stations (four for OH-pyrene) were sampled, instead of eight stations as in previous years. Furthermore, no samples for BEM were taken from flatfish. All fish were collected by local fishermen and kept alive until sampling by NIVA staff within 5 days. There is a continuous process to train and inform the fishermen that collect fish for JAMP to ensure the quality of the material.

### 1.4.1. OH-pyrene metabolites in bile





Detection methods for OH-pyrene have been changed (improved) two times since the initiation of these analyses in the JAMP programme. In 1998 the support/normalisation parameter biliverdin was changed to measurement of light absorbance at 380 nm. Furthermore, in 2000, the use of single-wavelength fluorescence for quantification of OH-pyrene was discontinued and the use of HPLC separation with fluorescence detection was implemented. Although there is a good correlation between results from the two methods they can not be compared directly. The single wavelength fluorescence method is naturally more unspecific and will include fluorescence from more components than the HPLC method, which has extremely high specificity towards individual metabolites. The interpretation of OH-pyrene data is therefore primarily focused on the differences between the stations within each year.

In 2004 the median concentration of OH-pyrene metabolites in cod from the inner Oslofjord (30B), was over twice that of cod from Lista (15B), and more than a factor of 10 higher than that of cod from Inner Sørfjorden (53B) and the “reference” station on the west coast (23B) (Figure 15, Appendix I).

For 1998, 1999, 2001 the median concentrations of OH-pyrene in cod from Lista (15B) were higher than at stations 30B, 53B and 23B (no samples from 15B in 2000). In 2003 concentrations were below those at st. 30B but above st. 23B and st. 53B. The variability at Lista was relatively high in 2003, indicated by the standard deviation:median ratio of >3, suggesting large differences among the 25 cod sampled. In 2004 the ratios varied from 1.4 at 53B to 2.3 at 30B, and OH-pyrene levels at Lista (15B) were again higher than in Inner Sørfjorden (although still lower than in the inner Oslofjord, 30B). There was a slight increase in the median OH-pyrene-level at Lista from 2002-2003 to 2004 (Figure 15). Lista is located in an area with a large discharge of PAH to water from an aluminium-smelter. The fish were collected on the open coast and the discharge from the smelter occurred in a small bay about 2-3 km away. No significant temporal trends were found at these three stations.

Bile metabolites of PAH can be detected within a short period (hours) following exposure, and holding conditions prior to sampling may affect results. However, measures were taken in 1998 and 1999 to minimise or remove the problem. Given the precautions taken, it is unlikely that the observed levels have been caused by storage of fish prior to sampling.

In 2004, as in 2000 and 2002-2003, the median concentration of OH-pyrene in cod from st. 30B (inner Oslofjord) was higher than the other three stations. When considering the whole period (1998-2004), the yearly median concentration at 30B was the highest or next highest compared to the other 2-3 stations Appendix I). Furthermore concentrations at 53B Sørfjorden were usually higher than 23B. This presumably reflect the general contamination of the two areas (inner Sørfjord and inner Oslofjord).

**A****B****C****D**

**Figure 15.** Concentration of OH-pyrene ( $\mu\text{g}/\text{kg}$  ABS 380nm) in bile from Atlantic cod collected at the indicated stations 2004. There was insufficient data to present a time series from st. 15B. (cf. Appendix G and Appendix I, and key in Figure 21). **NB: log-scale.**

### 1.4.2. ALA-D in blood cells


Most years the activity of ALA-D in cod was generally inhibited (indicating the influence of lead contamination) in the inner Oslofjord (30B) and inner Sørfjord (53B), compared to reference stations, i.e. outer Oslofjord (36B), Sotra-Bømlo area (23B), and Varangerfjord (10B). This was the case for 1997, 1998, and 2000-2004 (cf. Appendix I, results for stations 30B, 53B, and 23B in Figure 16). For all years 1997-2004 the activity of the enzyme at st.53B in Sørfjord was generally lower than st. 23B on the open coast, about 130 km away.

Since 2002, ALA-D was measured only in cod at stations 23B, 30B and 53B. As in previous years, the inhibition was largest in the Sørfjord (53B) and the inner Oslofjord (30B) in 2003. (Figure 16, Appendix I). This indicates pollution of metals at 53B and 30B. The median ALA-D activity 1997-2002 and 2004 was lowest at st. 53B (Appendix I, cf. Green *et al.* 2004a, b). A slight increase in median ALA-D activity could be seen from 2002 to 2003-2004 indicating less exposure. This was supported by measurements of lead concentrations in cod liver, where the median decreased about a third from 2002 to 2003 (Appendix I). Furthermore, the total discharge of lead into the inner Sørfjorden in 2004 was about 3.2 tonnes, a 37% increase from 2003, but 80% of the 2002 discharge (Ruus & Green 2004, 2005). No significant temporal trends in ALA-D activity were found at any of the three stations.

Although ALA-D inhibition is lead-specific, it is not possible to rule out interference by other metals or organic contaminants. Previous studies indicate that only zinc may ameliorate the effect of lead to some extent, but the effect is variable and weak. Other studies have also shown ALA-D to be a remarkably robust biomarker and factors such as sex, age or season do not appear to affect the response.


**A**

ALAD, *Gadus morhua*, 30B Inner Oslofjord




**B**

ALAD, *Gadus morhua*, 53B Inner Sørkjorden



**C**

ALAD, *Gadus morhua*, 23B Sotra-Bømlo area



**Figure 16.** Activity of  $\delta$ -aminolevulinic acid dehydrase (ALA-D, ng PBG/min/mg protein) in red blood cells from Atlantic cod collected at the indicated stations 2004. (cf. Appendix G and Appendix I, and key in Figure 21).

### 1.4.3. EROD activity in liver




#### *EROD activity*

High activity of hepatic cytochrome P4501A activity (EROD) normally occurs as a response to the contaminants indicated in Table 2. It was expected that higher activity would be found at the stations that were presumed to be most perturbated by planar PCBs, PCNs, PAHs or dioxins, which were st.30B (inner Oslofjord) and 53B/F (inner Sørfjord). However, median EROD activity at 53B was lower than at reference station 23B (Figure 17, Appendix I). Previous years have also shown that EROD in fish at stations 30B and 53B are not consistently higher than at other stations. No significant temporal trends were found at these three stations.

Extreme concentrations of PCBs were found in four individuals of cod from the inner Sørfjord in 2002 (Green *et al.* 2004a), which should induce a subfamily of cytochrome P450A1 proteins (CYP 1A) and thus increase EROD activity in these individuals. Two of these fish were those with the highest EROD-activities (although not much higher than some individuals with moderate PCB-concentrations). However, the other two extreme-PCB-concentration-fish had moderate hepatic EROD activities. It was concluded from the finding that undetermined confounding factors, which affects EROD activity, were likely present, and that moderate EROD activities do not disprove an environmental problem, at least with respect to the PCB congeners in question (Green *et al.* 2004a). There were no individuals with correspondingly high PCB-concentrations in 2004 (unpublished data).

No adjustment for water temperature has been made. Fish are sampled at the same time of year (September-November) when differences between the sexes should be at a minimum. Statistical analyses indicate no clear difference in activity between the sexes (Ruus *et al.* 2003). It has been shown that generally higher activity has been found at more contaminated stations (Ruus *et al.* 2003). However, the response is inconsistent (cf. Appendix I), perhaps due to sampling of populations with variable exposure history. Besides, there is evidence from other fish species that continuous exposure to e.g. PCBs may cause adaptation, i.e. decreased EROD response.

In 2003, a correlation was shown between the EROD activity and the amount of CYP1A protein measured (Green *et al.* 2004b).

**A**EROD, *Gadus morhua*, 30B Inner Oslofjord**B**EROD, *Gadus morhua*, 53B Inner Sørkjorden**C**EROD, *Gadus morhua*, 23B Sotra-Bømlo area

**Figure 17.** Activity of cytochrome P4501A (EROD, pmol/min/mg protein) in liver from Atlantic cod collected at the indicated stations 2004. (cf. Appendix G and Appendix I, and key in Figure 21).

#### **1.4.4. Concluding remarks**

The application of BEM methods within JAMP through the years 1997-2001 has indicated that the location Lista, st. 15B, previously regarded as only diffusely polluted, has an input of PAH which is sufficient to markedly affect fish in the area. For 2002-2003, however, the median concentrations of OH-pyrene in cod at 15B were below those at the inner Oslofjord, st. 30B, and inner Sørfjord, st. 53B. In 2004 the levels at Lista (15B) were again higher than in inner Sørfjorden (53B), but still lower than in the inner Oslofjord (30B). The median concentration of OH-pyrene in cod was lowest at the reference station in the Børnlo-Sotra area, st. 23B. Chronic exposure to PAHs may lead to liver lesions and reproductive disorders in fish, as shown through National Ocean and Atmospheric Administration's (NOAA (USA)) studies in Puget Sound. The highest levels of PAH metabolites observed in the bile of cod from st. 15B, and recently from the inner Oslofjord, are high compared to other studies, but it is not at present possible to infer population effects on cod in the area. It would be relevant to include DNA adduct analyses at some stage to clarify whether the cellular repair system of cod is sufficient to protect against damage from PAH radicals.

Results for the period 1997-2004 clearly indicated that there are lead effects, shown by decreased activity of the enzyme ALA-D in the two most strongly polluted areas, i.e. cod from the inner Oslofjord (30B) and cod from the inner Sørfjord (53B).

Median EROD activity at st. 53B was lower than at the less contaminated 23B station. Previous years have also shown that EROD in fish at stations 30B and 53B are not consistently higher than at other, presumed cleaner stations.

## 1.5. Effects and concentrations of organotin

Effects from organotin in dogwhelk (*Nucella lapillus*) were investigated at twenty stations. Organotin concentrations in dogwhelk and blue mussel (*Mytilus edulis*) were quantified at twenty and twenty-eight stations respectively. The stations are located along the coast of Norway and samples were collected September-October 2004 (Appendix E and maps in Appendix G).

TBT-induced development of male sex-characters in females, known as imposex (VDSI and RPSI), was analysed according to OSPAR-JAMP guidelines. Detailed information about the chemical analyses of the animals is given in Følsvik *et al.* (1999).




### 1.5.1. Dogwhelk

Twenty stations were investigated in 2004; most of these were on the west coast. Effects from organotin were observed at all stations. VDSI varied from 0.29 to 4.47. The three lowest VDSI (<2) were found at Brashavn (st. 11G), in the Varangerfjord, northern Norway, Landfast (st.94G) on the east side of the island of Vega and at Klakholm (st.97G) northwest of Bodø (Appendix J). The VDSI at the remaining stations were above 3.2; the highest at Grinden (st.27G) between Stadt and Ulsteinvik. A significant *upward* trend was found at the station in Lofoten (st. 98G) (Appendix I, Figure 18).




Concentrations of organotin from the nineteen stations measured were relatively low (<0.040 mg/kg w.w., <0.26 mg/kg d.w.). As in 2003 the highest organotin levels were found at Haugesund (Appendix I, Appendix J, Figure 19). Concentrations decreased compared to 2003, however, no statistically significant temporal trends for the period 1997-2004 were found.

The results for 1997-2004 indicates elevated concentrations and imposex-indexes not only near harbours (e.g., Haugesund), but also in presumably less polluted areas (e.g. Grinden) (Figure 19, Appendix I).

A comparison of imposex was made with samples from nine stations collected both in 1992 and 2004. The stations included four from the Stadtland area (st. 25G, 26G, 27G, 28G), two from Orkdalsfjord area (82G, 87G) and three along the coast from the island of Vega and north to Lofoten (94G, 95G, 97G). The 1992 samples were kept frozen until their VDSI was assessed together with the 2004 samples. The effect of freezing should not effect the index (Minchin & Davies 1999). A paired-t-test revealed no difference between the two years ( $p>0.475$ ).

**A**VDSI, *Nucella lapillus*, 36G Outer Oslofjord**B**VDSI, *Nucella lapillus*, 227G Haugesund**C**VDSI, *Nucella lapillus*, 98G Lofoten area (Husvagen)

**Figure 18.** Imposex (VDSI) in dogwhelk (*Nucella lapillus*) at 2 stations in southern Norway; Færder (36G) and Haugesund (227G) and one at Lofoten (98G). Data from 1991 (Harding *et al.* 1992) and 1993 (Walday *et al.* 1997). (cf. Appendix G and Appendix I, and key in Figure 21).

**A**TBT, *Nucella lapillus*, 36G Outer Oslofjord (Færder)**B**TBT, *Nucella lapillus*, 71G Langesund (Fugløyskjær)**C**TBT, *Nucella lapillus*, 227G Haugesund (Melandholmen)





**Figure 19.** Median concentration of TBT (on a formulation basis) in dogwhelks (*Nucella lapillus*) from outer Oslofjord (st.36G), Langesundsfjord (west of Oslofjord) (st.71G) and Haugesund (St.227G), mg/kg (mg TBT/kg) dry weight. NB: (cf. Appendix G and Appendix I, and key in Figure 21). Note: for some years the upper confidence interval line is off-scale in Figures B and C.

### **1.5.2. Mussels**

Blue mussel was severely contaminated with organotin at one station in the inner Oslofjord (Index st. 301); Class IV in SFTs environmental classification system (Appendix J), and another station nearby was markedly contaminated (Cl.III, Figure 20). Moderately (Cl.II) or markedly polluted (Cl.III) mussels were not only found in other harbour areas (e.g. the Frierfjord (712, 713) and Haugesund (227A)) but also in areas in western Norway that are presumably remote from point sources (e.g. 22A, 27A, 93A). Low concentrations were found at the northern stations (11X) and at Farsund (15A) as well as some stations in western Norway. Levels (median) ranged between 0.006 and 2.8 mg/kg d.w.. Compared to 2003, most concentrations were lower in 2004, however, significant temporal trends were not detected at any station for the 1997-2004 period investigated.

### **1.5.3. Concluding remark**

The presence of organotin (as TBT) in Norwegian waters still exceeded acceptable levels in 2004, not only in harbour areas but also some stations presumably remote from known point sources. Concentrations of organotin in mussels and dogwhelk were elevated, and biological effects from TBT were found in dogwhelk from all of the investigated areas. No significant trends were found. Furthermore, there was no significant difference between the VDSI in 1992 compared to 2004 at nine stations on the westcoast. It is a cause for concern that the ban on the use of TBT in antifouling on boats <25 m of length, in effect since 1.January 2003, has not lead to a clear improvement in the investigated areas.

**A**TBT, *Mytilus edulis*, 30A Inner Oslofjord**B**TBT, *Mytilus edulis*, 36A Outer Oslofjord (Færder)**C**TBT, *Mytilus edulis*, 71A Langesund**D**TBT, *Mytilus edulis*, 227X Haugesund (Hølevarode)

**Figure 20.** Median concentration of TBT (on a formulation basis) in blue mussel (*Mytilus edulis*) from inner (st.30A) and outer (st.36A) Oslofjord, Langesundsfjord (west of Oslofjord) (st.71A) and Haugesund (St.227X), mg/kg (mg TBT/kg) dry weight. (cf. Appendix G and Appendix I, and key in Figure 21). Note: for 1997 in Figure A the upper confidence interval line is off-scale. Note: horizontal line for Class I is near x-axis

## 1.6. Overall conclusions

In regards to JMP/JAMP Purpose A (health assessment), attention should be called to the list from Norwegian Food Safety Authority (*Mattilsynet*) which names the restrictions and recommendations concerning the sale and consumption in Norway for seafood taken from Norwegian fjord areas (Table 3). Furthermore, *Mattilsynet* has issued general advice to avoid consumption of seafood taken in or in close proximity to harbours (cf. Økland, 2005).

In regards to JMP/JAMP Purpose C (spatial distribution assessment), the concentrations found in 2004 are indicated in the bar graphs shown in Appendix J and Appendix K. Provisional "high background" levels were used to identify elevated concentrations. This assessment revealed no new areas of concern that are not currently under surveillance.

In regards to JMP/JAMP Purpose D (temporal trend assessment), and considering where statistically significant linear trends have been found, the following cases should be noted:

- ΣPCB-7 in mussels from the inner Oslofjord has decreased since 1988;
- Mercury in cod fillet from the inner Oslofjord has increased since 1984;
- Cadmium in cod liver and mussels (1 st.) from the inner Oslofjord has increased since 1984;
- HCB in mussel from Langesundsfjorden has decreased since 1983;
- Mercury in flounder fillet from the inner Sørfjorden has increased since 1988;
- Cadmium in mussels (3 st.) in the Hardangerfjord/Sørfjorden has decreased since 1987;
- Lead in mussels (2 st.) in the Hardangerfjorden has decreased since 1990 (1987);

Study of the power of temporal trend monitoring was useful in assessing existing sampling strategies, however, modifications might be needed to account for local conditions (see Appendix O in Green *et al.* 2000).

The 2004 investigation also includes results on Norwegian Pollution Control Authority Pollution Indices (Appendix L), and discussion of the results of biological effects methods including imposex and intersex (chapters 1.4 and 1.5).

**Table 3.** Summary of action taken by the Norwegian Food Safety Authority (*Mattilsynet*) concerning the consumption and sale of fish products along the Norwegian Coast (based on review by Økland 2005). Restrictions on sale vary and may concern the whole or part of fish product.

Area of concern (km <sup>2</sup> )	Main parameters of concern	Last year of issue/evaluation	Main fish/shellfish product of concerned	Recommendations or restrictions of concern:
Mid <sup>1)</sup> and Inner Oslofjord (498.9) (includes Drammensfj.)	PCB	2002	fish liver, eel	Consumption and sale
Tønsberg area (23.7) (includes (Vrengen))	PCB	2003	fish liver, eel, mussels	Consumption
Inner Sandefjordfjord (1.5)	PCB	1999	fish liver	Consumption and sale
Grenlandsfjords, Langesundsfjord (90.3)	Chl.org <sup>2)</sup> / Dioxins	2002	fish, shellfish	Consumption and sale
Kragerø (3.2)	PAH Dioxins	2002	eel, mussels	Consumption
Tvedstrand (2.3)	PCB	2002	fish liver	Consumption and sale
Arendal (8.0)	PCB	2002	fish liver	Consumption and sale
Inner Kristiansandsfjord (33.3)	Chl.org <sup>2)</sup> / Dioxins/PCB	2000	fish, shellfish	Consumption and sale
Farsund area (42.0)	PCB PAH	2000	fish liver, mussels	Consumption and sale
Fedafjord (11.2)	PAH	1995	mussels	Consumption and sale
Flekkefjord (4.2)	PCB	2000	fish liver	Consumption and sale
Stavanger (4.0)	PCB PAH	2001	fish liver, mussels	Consumption
Sandnes (1.7)	PAH	2001	Mussels	Consumption
Karmsund-Eidsbotn, Vedavågen (???)	PCB, PAH	2005	fish liver <sup>3)</sup> , shellfish	Consumption and sale
Saudafjord (24.1)	PAH	1992	fish liver, mussels	Consumption and sale
Sørkjosen (62.2)	Cd Pb Hg PCB	2005	fish, shellfish	Consumption and sale
Bergen area (169.9)	PCB	2002	fish, shellfish	Consumption and sale
Årdalsfjord (30.4)	PAH	2002	mussels	Consumption and sale
Alesund, Asefjorden (16.7)	HBCDD <sup>4)</sup>	2005	fish, shellfish	Consumption
Sunndalsfjord (100.1)	PAH	2005	fish liver, mussels	Consumption and sale
Hommelvik (2.6)	PAH	1985	mussels	Consumption and sale
Inner Trondheimfjorden (1.2)	PAH PCB	2002	fish liver, mussels	Consumption
Brønnøysund (7.0)	PAH	2003	mussels	Consumption
Vefsnfjord (76.4) <sup>5)</sup>				
Sandnessjøen (0.4)	PAH	2005	mussels	Consumption
Inner Ranfjord (16.6)	PAH	2005	mussels	Consumption and sale
Ramsund (5.4)	PCB	2002	fish, shellfish	Consumption and sale
Harstad (2.9)	PCB Pb Cd	2003	fish liver, mussels	Consumption and sale
Narvik (11.6)	PCB PAH	2005	fish, mussels	Consumption
Tromsø (17.7)	PAH	2003	Mussels	Consumption and sale
Hammerfest (4.1)	PAH	2003	mussels	Consumption and sale
Honningsvåg (3.3)	PAH	2000	mussels	Consumption and sale

<sup>1)</sup> Includes, Hvitsten, Moss, Horten og Holmenstrand

<sup>2)</sup> Organochlorine compounds

<sup>3)</sup> Concerns only Eidsbotn

<sup>4)</sup> A brominated flame retardant

<sup>5)</sup> Grounds for concern were cleared in 2005

The JAMP issues to which these investigations are relevant are shown in Table 4.

**Table 4.** JAMP issues relating to the Norwegian JAMP (cf., SIME 2002).

Subject	JAMP issue	Question	Recent Norwegian contribution
<b>Hg, Cd and Pb</b>	JAMP issue 1.2.	What are the concentrations and fluxes in sediments and biota?	1996-1997: Levels in sediment (cf., Green <i>et al.</i> 2000) 2003: Levels and trends in biota (annual investigations since 1981, Chapter 1.3) 2003: INDEX for blue mussel from selected stations (annual investigations since 1995, cf. Chapter 1.3.8)
<b>TBT</b>	JAMP issue 1.3.	To what extent do biological effects occur in the vicinity of major shipping routes offshore installations, marinas and shipyards	2003: Levels and trends in mussels and snails (annual investigations since 1997, cf. Chapter 1.5)
<b>PCBs</b>	JAMP issue 1.7.	Do high concentrations pose a risk to the marine ecosystem	[as for JAMP issue 1.2]
<b>PCBs</b>	JAMP issue 1.8.	Do high concentrations of non-ortho and mono-ortho CBs in seafood pose a risk to human health?	1995: INDEX for blue mussel from selected stations (cf. Green 1997) 1996: Levels in cod (cf. Green <i>et al.</i> 2000)
<b>PAHs</b>	JAMP issue 1.10.	What are the concentrations in the maritime <sup>1)</sup> area?	1992: Levels in shellfish (Green <i>et al.</i> 1995) 1992-1993: Levels in fish and mussels for selected stations (Knutzen & Green 1995) 1996-1997: Levels in sediment (cf., Green <i>et al.</i> 2000) 2003: INDEX for blue mussel from selected stations (annual investigations since 1995, Chapter 1.3.8)
<b>PAHs</b>	JAMP issue 1.11.	Do PAHs affect fish and shellfish?	1998: Biological effects methods in cod (cf. Chapter 1.4)
<b>Other synthetic organic compounds</b>	JAMP issue 1.12.	How widespread are synthetic organic compounds within the maritime <sup>1)</sup> area?	2003: Levels and trends in biota (annual investigations since 1983 of selected organochlorines, cf. Chapter 1.3) 1996: Introductory investigation of organochlorines in cod livers (cf. Green <i>et al.</i> 2000)
<b>Chlorinated dioxins and dibenzofurans</b>	JAMP issue 1.15. <sup>2)</sup>	What concentrations occur and have the policy goals (for the relevant parts of the maritime <sup>1)</sup> area) been met?	2003: INDEX for blue mussel from selected stations (cf. Appendix L) 1996: Introductory investigation of organochlorines in cod livers (cf. Green <i>et al.</i> 2000)
<b>Biological effects of pollutants</b>	JAMP issue 1.17.	Where do pollutants cause deleterious biological effects?	2003: Southern Coast, planar PCBs, metals, PAHs in cod (annual investigations since 1997, cf. Chapter 1.4)
<b>Chemicals used</b>	JAMP issue 5.3.	In which areas do pesticides and antibiotics affect marine biota?	2003: Levels and trends in biota (cf. Chapter 1.3)
<b>Ecosystem health</b>	JAMP issue 6.1. <sup>2)</sup>	How can ecosystem health be assessed in order to determine the extent of human impact?	Results for the other issues are also relevant here

<sup>1)</sup> Not defined in original text

<sup>2)</sup> See SIME 1997

## 2. Technical Details

### 2.1. Compliance with guidelines/procedures

#### 2.1.1. JAMP programme

Samples were collected and analysed, where practical, according to OSPAR guidelines (OSPAR 1990, 1997) and screened and submitted to ICES by agreed procedures (ICES 1996). The most important point of concern are those stations where insufficient number of fish were collected (cf. Appendix H).

#### 2.1.2. Overconcentrations and classification of environmental quality

This report focuses on the principle cases where *median* concentrations exceeded the upper limit to Class I in the Norwegian Pollution Control Authority's (SFT's) environmental quality classification system (cf. Molvær *et al.* 1997). The relevant extract from the system is shown in Table 5 and Table 6, and show five classes from Class I, "slightly polluted", to Class V, "extremely polluted".

However, the system does not cover all the contaminants in indicator species-tissues used in JAMP. To assess concentrations not included in the system provisional "high background" values were used (Table 7). The factor by which concentrations exceeded "high background" is termed **overconcentration**. It should be noted that there is in general a need for periodic review and supplement of this list of limits in the light of results from reference localities and introduction of new analytical methods, and/or units. Because of changes in the limits, assessments of overconcentrations for years prior to 1997 made in this report may not correspond to figures and assessments made in previous national comments. The median concentration can be found in the tables in Appendix I or figures in Appendix J.

A review by Knutzen and Green (2001b) of provisional "high background" concentrations based on recent JAMP-data generally confirmed that the reference concentrations (i.e., upper limit for Class I) in SFT's classification system, but recommended the following revisions (concentrations in µg/kg wet weight):

Cod liver - ΣDDT: Either increase limit from 200 to 300 or preferably replace ΣDDT with p,p-DDE and keep the limit at 200,

Cod liver - ΣHCH: Decrease limit from 50 to 30,

Cod liver - TEPCDD/PCDF: Decrease limit from 0.015 to 0.0,

Cod fillet - ΣPCB7: Decrease limit from 5 to 3,

Cod fillet - ΣHCH: Decrease limit from 0.5 to 0.3,

Blue mussel - ΣPCB7: Decrease limit from 4 to 3.

Furthermore, the review, supplemented by other studies (cf. Green & Knutzen 2003), also suggested the following decreases for Class I in fillet of flounder (µg/kg w.w.):

ΣPCB7: from 5 to 3,

ΣDDT: from 2 to 1 for p,p-DDE only.

The review did not recommend changes in the Class I limits for mercury in fish fillet (1 mg/kg w.w.) or mercury, cadmium, lead, zinc and copper in mussels (in the same order 0.2; 2; 3; 200 and 10 mg/kg d.w.). However, for chromium and nickel in mussels limits should be decreased from 3 to 2 and from 5 to 3 mg/kg d.w., respectively. Further, reference values for organochlorines were indicated for fillet and liver of fish species that are not included in the classification system (dab, plaice, lemon sole) and for lead and cadmium in liver of cod.

These recommendations for changes have been taken into account in this report. However, corresponding adjustment of Classes II-V has not been done, but should be considered once the above mentioned Class I revisions have been accepted by SFT. SFT is considering these recommendations in a current review of their classification system.

No attempt has been made to compensate for differences in size groups or number of individuals of mussels or fish. The exception was with mercury in fish fillet where six data sets in both cod and flatfish in this study showed significant differences between "small" and "large" fish (Appendix I). With respect to mussels, there is some evidence that concentrations do not vary significantly among the three size groups employed for this study (i.e. 2-3, 3-4 and 4-5 cm) (WGSAEM 1993).

With respect to Purpose A (health risk assessment), the Norwegian Food Safety Authority (SNT) is responsible for official commentary as to possible health risk due to consumption of seafood. Hence, the results of the JAMP pertaining to this purpose are presented only as a partial basis for evaluation.

**Table 5.** Extracts of the Norwegian Pollution Control Authority revised environmental classification system of contaminants in blue mussel and fish (from Molvær *et al.* 1997 and revised (shaded) Class I concentrations as suggested in this report).

Contaminant		Classification (upper limit for Classes I-IV)				
		Degree of pollution				
		I <i>Slight</i>	II <i>Moderate</i>	III <i>Marked</i>	IV <i>Severe</i>	V <i>Extreme</i>
<b>BLUE MUSSEL</b>						
<b>Lead</b>	mg/kg d.w.	3	15	40	100	>100
<b>Cadmium</b>	mg/kg d.w.	2	5	20	40	>40
<b>Copper</b>	mg/kg d.w.	10	30	100	200	>200
<b>Mercury</b>	mg/kg d.w.	0.2	0.5	1.5	4	>4
<b>Zinc</b>	mg/kg d.w.	200	400	1000	2500	>2500
<b>TBT<sup>1)</sup></b>	mg/kg d.w.	0.1	0.5	2	5	>5
<b>ΣPCB-7</b>	µg/kg w.w. d.w. <sup>2)</sup>	3 15	15 75	40 200	100 500	>100 >500
<b>ΣDDT</b>	µg/kg w.w. d.w. <sup>2)</sup>	2 10	5 25	10 50	30 150	>30 >150
<b>ΣHCH</b>	µg/kg w.w. d.w. <sup>2)</sup>	1 5	3 15	10 50	30 150	>30 >150
<b>HCB</b>	µg/kg w.w. d.w. <sup>2)</sup>	0.1 0.5	0.3 1.5	1 5	5 25	>5 >25
<b>ΣPAH</b>	µg/kg w.w. d.w. <sup>2)</sup>	50 250	200 1000	2000 10000	5000 25000	>5000 >25000
<b>ΣKPAH</b>	µg/kg w.w. d.w. <sup>2)</sup>	10 50	30 150	100 500	300 1500	>300 >1500
<b>B[a]P</b>	µg/kg w.w. d.w. <sup>2)</sup>	1 5	3 15	10 50	30 150	>30 >150
<b>TE<sub>PCDF/D</sub><sup>3)</sup></b>	µg/t <sup>4)</sup> w.w.	0.2	0.5	1.5	3	>3
<b>COD, fillet</b>						
<b>Mercury</b>	mg/kg w.w.	0.1	0.3	0.5	1	>1
<b>ΣPCB-7</b>	µg/kg w.w.	3	20	50	150	>150
<b>ΣDDT</b>	µg/kg w.w.	1	3	10	25	>25
<b>ΣHCH</b>	µg/kg w.w.	0.3	2	5	15	>15
<b>HCB</b>	µg/kg w.w.	0.2	0.5	2	5	>5
<b>COD, liver</b>						
<b>ΣPCB-7</b>	µg/kg w.w.	500	1500	4000	10000	>10000
<b>ΣDDT</b>	µg/kg w.w.	200	500	1500	3000	>3000
<b>ΣHCH</b>	µg/kg w.w.	30	200	500	1000	>1000
<b>HCB</b>	µg/kg w.w.	20	50	200	400	>400
<b>TE<sub>PCDF/D</sub><sup>2)</sup></b>	µg/t <sup>4)</sup> w.w.	10	40	100	300	>300

<sup>1)</sup> Tributyltin on a formula basis

<sup>2)</sup> Conversion assuming 20% dry weight

<sup>3)</sup> TCDDN (Appendix B)

<sup>4)</sup> µg/1000 kg (Appendix B)

**Table 6.** Extracts of the Norwegian Pollution Control Authority revised environmental classification system of contaminants in sediment (from Molvær *et al.* 1997).

Contaminant		Classification (upper limit for Classes I-IV)				
		Degree of pollution				
		I Slight	II Moderate	III Marked	IV Severe	V Extreme
<b>SEDIMENT</b>						
Lead	mg/kg d.w.	30	120	600	1500	>1500
Cadmium	mg/kg d.w.	0.25	1	5	10	>10
Copper	mg/kg d.w.	35	150	700	1500	>1500
Mercury	mg/kg d.w.	0.15	0.6	3	5	>5
Zinc	mg/kg d.w.	150	700	3000	10000	>10000
TBT <sup>1)</sup>	µg/kg d.w.	1	5	20	100	>100
ΣPCB-7	µg/kg d.w.	5	25	100	300	>300
ΣDDT	µg/kg w.w.	0.5	2.5	10	50	>50
HCB	µg/kg w.w.	0.5	2.5	10	50	>50
ΣPAH	µg/kg w.w.	300	2000	6000	20000	>20000
B[a]P	µg/kg w.w.	10	50	200	500	>500
TE <sub>PCDF/D</sub> <sup>2)</sup>	µg/t <sup>3)</sup> w.w.	0.01	0.03	0.1	0.5	>0.5


<sup>1)</sup> Tributyltin on a formula basis<sup>2)</sup> TCDDN (Appendix B)<sup>3)</sup> µg/1000 kg (Appendix B)**Table 7.** Provisional "high background levels" of selected contaminants, in **mg/kg dry weight** (blue mussel) and **mg/kg wet weight** (blue mussel and fish). The respective "high background" limits are from Knutzen & Skei (1990) with mostly minor adjustments (Knutzen & Green 1995; Molvær *et al.* 1997), except for dab where the suggested limit is based on JAMP-data (Knutzen & Green 1995). Especially uncertain values are marked with "?".

Cont.	Blue mussel <sup>1</sup>	Cod <sup>1</sup>		Flounder <sup>1</sup>		Dab <sup>1</sup>		Plaice <sup>1</sup>	
		liver	fillet	liver	fillet	liver	Fillet	liver	fillet
		mg/kg d.w.	mg/kg w.w.	mg/kg w.w.	mg/kg w.w.	mg/kg w.w.	mg/kg w.w.	mg/kg w.w.	mg/kg w.w.
Lead	3.0 <sup>2)</sup>	0.6 <sup>3)</sup>	0.1	0.3 ?		0.3 ?		0.2 ?	
Cadmium	2.0 <sup>2)</sup>	0.4 <sup>3)</sup>	0.1	0.3 ?		0.3 ?		0.2 ?	
Copper	10 <sup>2)</sup>	2 <sup>3)</sup>	20	10 ?		30 ?		10 ?	
Mercury	0.2 <sup>2)</sup>	0.04 <sup>3)</sup>	0.1 <sup>2)</sup>	0.1 ?		0.1		0.1 ?	
Zinc	200 <sup>2)</sup>	40 <sup>3)</sup>	30	50 ?		60 ?		50 ?	
ΣPCB-7 <sup>8)</sup>	0.015 <sup>3,9)</sup>	0.003 <sup>2,9)</sup>	0.50 <sup>2)</sup> 0.003 <sup>9)</sup>	0.1 0.003 <sup>9)</sup>		0.5 0.005 <sup>9)</sup>		0.05 ? 0.004 <sup>9)</sup>	
ppDDE	0.010 <sup>3)</sup>	0.002 <sup>6)</sup>	0.2 <sup>9)</sup>	0.03 0.001 <sup>9)</sup>		0.1 0.002 <sup>9)</sup>		0.01 ? <sup>6)</sup> 0.001 <sup>9)</sup>	
γHCH	0.005 <sup>3)</sup>	0.001 <sup>6)</sup>	0.03 <sup>9)</sup> 0.0003 <sup>9)</sup>	0.01 0.0003 <sup>9)</sup>		0.03 0.0005 <sup>9)</sup>		0.005 ? <sup>6)</sup> 0.0003 <sup>9)</sup>	
HCB	0.0005 <sup>3)</sup>	0.0001 <sup>2)</sup>	0.02 <sup>2)</sup>	0.005 0.0001 <sup>9)</sup>		0.01 0.0002 <sup>9)</sup>		0.005 ? 0.0002 <sup>9)</sup>	
TCDDN	0.000001 <sup>3)</sup>		0.00001 <sup>9)</sup>						
			0.0000002 <sup>2)</sup>						

<sup>1)</sup> Respectively: *Mytilus edulis*, *Gadus morhua*, *Platichthys flesus* and *Limanda limanda*.<sup>2)</sup> From the Norwegian Pollution Control Authority Environmental Class I ("good") (Molvær *et al.* 1997).<sup>3)</sup> Conversion assuming 20% dry weight.<sup>4)</sup> Approximately 25% of ΣPCB-7 (Knutzen & Green 1995)<sup>5)</sup> 1.5-2 times 75% quartile (cf. Annex B in Knutzen & Green 1995)<sup>6)</sup> Assumed equal to limit for ΣDDT or ΣHCH, respectively, from the Norwegian Pollution Control Authority Environmental Class I ("good") (Molvær *et al.* 1997). Hence, limits for ppDDE and γHCH are probably too high (lacking sufficient and reliable reference values)<sup>7)</sup> Mean plus 2 times standard deviation (cf. Annex B in Knutzen & Green 1995)<sup>8)</sup> Estimated as sum of 7 individual PCB compounds (CB-28, -52, -101, -118, -138, -153 and -180) and assumed to be ca. 50% and 70% of total PCB for blue mussel and cod/flatfish, respectively.<sup>9)</sup> With respect to revisions suggested by Knutzen & Green (2001b) and Green & Knutzen (2003), see text.

### 2.1.3. Comparison with previous data

A simple 3-model approach has been developed to study time trends for contaminants in biota based on *median* concentrations (ASMO 1994). A variation of this method was applied to mercury in fish fillet to distinguish trends in "small" and "large" individuals, the size of which may vary from year to year, station to station, depending on the catch. To determine the "small" fish, the sample is sorted by length and split into two groups of one or even numbers. The fish with median length in the smaller group is the "small" fish, and the median length in the larger group is the "large" fish. The concentration in these two size groups (one per group) determine the concentrations in the two groups. The method was first used on a large-scale basis by the Ad Hoc Working Group on Monitoring that met in Copenhagen 8-12. November 1993 (MON 1993). At this meeting it was agreed to apply the method on contaminants in fish muscle and liver on a wet weight basis and contaminants in soft tissue of mussels on a dry weight basis. The results for this assessment are presented earlier (cf. ASMO 1994). The method has been applied to Norwegian data and results are shown in Appendix H. The results can be presented as in Figure 21.



**Figure 21.** Example presentation and variation in contaminant concentration with time, indicating median concentrations, running mean of median values (Loess smoother), 95% confidence intervals. The horizontal lines indicate the lower boundaries to SFT classes of pollution: Class II (moderate=upper boundary to Class I (slight)), III (marked), IV (severe) and V (extreme), or alternatively the Class II boundary is replaced by the upper boundary to provisional "high background level" as in which case no class-boundaries are shown. (see text and refer to Table 7).

The method of calculating the smoother is in accordance to the methods employed at Ad Hoc Working Group on Monitoring that met in Copenhagen 23-27. February 1998 (MON 1998). A Loess smoother is based on a running seven-year interval, a non-parametric curve fitted to median log-concentrations (Nicholson *et al.* 1997). For statistical tests based on a fitted smoother to be valid the

contaminants indices should be independent to a constant level of variance and the residuals for the fitted model should be lognormally distributed (cf. Nicholson *et al.* 1998). No transformation was applied to the imposex (VSDI) data.

The National Comments since 1994 have included two additional analyses. The first is that the smoothed median for the last three sampling years is linearly projected for the next three years. This deviates from previous reports where the upper 95 confidence interval was used to assess the likelihood of overconcentrations (Nicholson, *et al.* 1994). The projected estimate is based on the results for the temporal trend analyses of at least 6 years of data.

The second is an estimate of the power of the temporal trend series expressed as the number of years to detect a 10% change per year with a 90% power (cf. Nicholson *et al.* 1997). The fewer the years the easier it is to detect a trend. The power is based on the percentage relative standard deviation (RLSD) estimated using the robust method described by ASMO (1994) and Nicholson *et al.* (1998). The estimate was made for series with at least 3 years of data and covers the *entire* period monitored. This fixed means of treating all the datasets may give misleading results especially where non-linear temporal changes are known to occur, such as for HCB in blue mussel from Langesundsfjord (Figure 4).

The statistical analysis was carried out on temporal trend data series for cadmium, mercury, lead, ΣPCB-7 (sum of congeners: 28, 53, 101, 118, 138, 153, 180), ppDDE (ICES code DDEPP), HCB, non-dicyclic PAHs, sum carcinogenic PAHs, B[a]P, TBT, and the biological effects parameters imposex (VSDI), PYR10, ALA-D and EROD.

#### **2.1.4. The effect of depuration and freezing on mussels**

Based on samples collected in the Sørnfjord and Hardangerfjord, the JAMP-method of pre-treatment of mussels (i.e., depuration and then cleaning) contrasted significantly to the Index-method (freezing then cleaning) (Green *et al.* 2001a). Using the JAMP-method, cadmium concentrations were significantly higher (24%), whereas significant lower concentrations were found for lead (45%), zinc (14%), PCBs (CB101, -118, -138, -153 27-52%) and DDTs (50-64%). Lower concentrations indicated that these contaminants are lost by depuration and gut emptying.

The results from a previous study from this region indicated no significant difference between the methods for mercury, cadmium, copper, lead and zinc (Green 1989). A study on mussels from the mouth of the Glomma River in Southern Norway showed that lead and copper were significantly lower in depurated samples (Green *et al.* 1996); however, no differences were found for PCBs or DDTs (on a lipid basis). The PCB concentrations found in the Glomma study were 3-4 times higher than Sørnfjord/Hardangerfjord.

Mercury was the only contaminant common to all three studies that had consistent results; that there is no significant difference between the two methods.

The difference in methods has indicated an effect on the concentration of contaminants in mussels. However, with the exception of mercury, the results for Sørnfjord/Hardangerfjord 2003 are inconsistent with two other studies in Norway. Revision of JAMP guidelines and assessment of data should take these results into consideration.

## **2.2. Information on Quality Assurance**

NIVA has participated in all the QUASIMEME international intercalibration exercises, including Round 42. These exercises have included nearly all the contaminants analysed for JAMP. Quality assurance programme for NIVA is similar to the 2003 programme (cf. Green *et al.* 2005). In addition, NIVA was accredited in 1993 and are now accredited in accordance with the NS-EN ISO/IEC 17025 standard by the Norwegian Accreditation (reference P009). A summary of the quality assurance programme at NIVA is given in Appendix A.

## 2.3. Description of the Programme

The sampling for 2004 involved blue mussel at 58 stations, dogwhelk at 20, cod at 9 and flatfish at 11 stations (cf. Appendix E). The Norwegian JAMP has been expanded since 1989 to include monitoring in more diffusely polluted areas. Though new stations are initially intended for annual monitoring (temporal trends), there has not always been sufficient funds to do this for every station. Sample/station reduction measures have been taken to reduce costs. Furthermore, sufficient samples have not always been practical to obtain. When this applies to mussels, a new site in the vicinity is often chosen. As for fish, the quota of 25 individuals ( $\pm 10\%$ ), indicated in Appendix E, as either 25 individuals or 5 bulked samples consisting of 5 fish per bulked sample, was met for all stations in 2004 except witch from st.67F (10 individuals), flounder st.21F (5 individuals). Plaice st.98F (25 individuals) and 10F (20 individuals).

Concentrations of metals, organochlorines (including pesticides) and polycyclic aromatic hydrocarbons in mussels and fish were determined at the Norwegian Institute for Water Research (JAMP code NIVA).

Analytical methods have been described previously (Green *et al.* (2001b). An overview of the samples collected from 1981 to 2003 is given in Appendix E. An overview of analyses applied from 1981 to 2003 for biological material is given in Appendix C. Parameter abbreviations are given in Appendix B.

The data is stored at NIVA in MS ACCESS 1997. The tables are generated using MS ACCESS 97 and MS EXCEL 97.

### 3. References

- Titles translated to English in square brackets [ ] are not official.
- Ahlborg, U.G., 1989. Nordic risk assessment of PCDDs and PCDFs. *Chemosphere* 19:603-608.
- Ahlborg, U.G., Becking G.B., Birnbaum, L.S., Brouwer, A., Derkx, H.J.G.M., Feely, M., Golor, G., Hanberg, A., Larsen, J.C., Liem, A.K.G., Safe, S.H., Schlatter, C., Wärn, F., Younes, M., Yrjänheikki, E., 1994. Toxic equivalency factors for dioxin-like PCBs. Report on a WHO-ECEH and IPSC consultation , December 1993. *Chemosphere* 28:1049-1067.
- ASMO, 1994. Draft assessment of temporal trends monitoring data for 1983-91: Trace metals and organic contaminants in biota. Environmental Assessment and Monitoring Committee (ASMO). Document ASMO(2) 94/6/1.
- ASMO, 1997. Summary Record. Environmental Assessment and Monitoring Committee (ASMO). Document ASMO 97/9/1.
- Følsvik N., Berge J.A., Brevik E. M. & M. Walday. 1999. Quantification of organotin compounds and determination of imposex in populations of dogwhelk (*Nucella lapillus*) from Norway. *Chemosphere*. 38 (3): 681-691.
- Green, N.W., 1987. "Joint Monitoring Group" (JMG). Felles monitoring program i Norge: Oslofjord-området, Sørkjorden og Hardangerfjorden, og Orkdalsfjorden. Programforslag for 1988. 4.Dec.1987. NIVA-project 80106, 12 pp..
- Green, N.W., 1989. The effect of depuration on mussels analyses. Report of the 1989 meeting of the working group on statistical aspects of trend monitoring. The Hague, 24-27 April 1989. ICES-report C.M.1989/E:13 Annex 6:52-58.
- Green, N.W., 1991. Joint Monitoring Programme. National Comments to the Norwegian Data for 1989. Norwegian Institute for Water Research (NIVA) memo 27pp.. JMG 16/info 13-E.
- Green, N.W., 1997. Joint Assessment and Monitoring Programme (JAMP) National Comments to the Norwegian Data for 1995. Norwegian Pollution Control Authority, Monitoring report no. 685/97 TA no. 1405/1997. Norwegian Institute for Water Research project 80106, report number 3597-97, 124 pp.. ISBN number 82-577-3152-8. (Also as document SIME 97/5/5).
- Green, N.W., 2003 Joint Assessment and Monitoring Programme in Norway 2003 Contaminants - Oslofjord area, Lista, Sørkjorden, Hardangerfjorden, Bømlo, Lofoten area and Varangerfjord. NIVA project 80106, 15.5.2003. 55 sider.
- Green, N.W., Knutzen, J., 2001. Joint Assessment and Monitoring Programme (JAMP). Forurensnings- og referanseindeks basert på observasjoner av miljøgifter i blåskjell fra utvalgte områder 1995-1999.[Joint Assessment and Monitoring Programme (JAMP). Pollution and reference indicies based on contaminants in blue mussel 1995-1999]. Norwegian Pollution Control Authority, Monitoring report no. 821/01 TA no. 1799/2001. NIVA project O-80106, (report number 4342-2001) 35 pp.. ISBN number 82-577-3977-4.
- Green, N.W., Knutzen, J., 2003. Organohalogens and metals in marine fish and mussels and some relationships to biological variables at reference localities in Norway. *Marine Pollution Bulletin* 46(3):362-374.
- Green, N.W., Knutzen, J., Helland, A., Brevik, E.M., 1995. Overvåking av miljøgifter i sedimenter og organismer 1981-92. "Joint Monitoring Programme (JMP)". [Monitoring av contamiants in sediment and organisms 1981-92 "Joint Assessment and Monitoring Programme (JMP)".] Norwegian Pollution Control Authority, Monitoring report no. 593/95. Norwegian Institute for Water Research project 80106, report number 3184, 195 pp.. ISBN-82-577-2676-1.
- Green, N.W., Bjerkeng B., Berge J.A., 1996. Depuration (12h) of metals, PCB and PAH concentrations by blue mussel (*Mytilus edulis*). Report of the Working Group on the Statistical Aspects of Environmental Monitoring. Stockholm 18-22 March 1996. ICES C.M.1996/D:1 Annex 13:108-117.
- Green, N.W., Berge, J.A, Helland, A., Hylland, K., Knutzen, J., Walday, M., 1999. Joint Assessment and Monitoring Programme (JAMP) National Comments regarding the Norwegian Data for 1997. Norwegian Pollution Control Authority, Monitoring report no. 752/99 TA no. 1611/1999. Norwegian Institute for Water Research project 80106, report number 3980-99, 129 pp.. ISBN number 82-577-3576-0. (Also presented as SIME document (1999)).
- Green, N.W., Bjerkeng, B, Helland, A., Hylland, K., Knutzen, J., Walday, M., 2000. Joint Assessment and Monitoring Programme (JAMP) National Comments regarding the Norwegian Data for 1998 and supplementary investigations on cod (1996) and sediment (1996-1997). Norwegian Pollution Control Authority, Monitoring report no. 788/00 TA no. 1702/2000. Norwegian Institute for Water Research project 80106, report number 4171-2000, 206 pp.. ISBN number 82-577-3787-9. (Also presented as SIME 2000 document 00/3/6).
- Green, N.W., Hylland, K., Walday, M., 2001a. Joint Assessment and Monitoring Programme (JAMP). National Comments regarding the Norwegian Data for 1999. SFT Statlig overvåkingsprogram 812/01 TA no. 1780/2001. NIVA, rapport nr. 4335-2001, 181 s. ISBN nr. 82-577-3969-3.

- Green, N.W., Følsvik, N., Oredalen, T.J., Prestbakmo, G., 2001b. Joint Assessment and Monitoring Programme (JAMP). Overview of analytical methods 1981-2000. SFT Statlig overvåkingsprogram rapport nr. 822/01, TA nr. 1800/2001. NIVA, rapport nr. 4353-2001, 68 s. ISBN nr. 82-577-3989-8.
- Green, N.W., Helland, A., Hylland, K., Knutzen, J., Walday, M., 2001c. Joint Assessment and Monitoring Programme (JAMP). Overvåking av miljøgifter i marine sedimenter og organismer 1981-1999. SFT Statlig overvåkingsprogram rapport nr. 819/01, TA nr. 1797/2001. NIVA, rapport nr. 4358-2001, 191 s. ISBN nr. 82-577-3995-2.
- Green, N.W., Severinsen G., Rogne, Å.K., 2002a. Joint Assessment and Monitoring Programme (JAMP). Contaminant data for sediments 1986-1997. Norwegian Pollution Control Authority, Monitoring report no. 861/02 TA no. 1918/2002. Norwegian Institute for Water Research project 80106, report number 4599-2002, 230 pp.. ISBN number 82-577-4259-7.
- Green, N.W., Severinsen G., Rogne, Å.K., 2002b. Joint Assessment and Monitoring Programme (JAMP). Contaminant data for shellfish 1998-2001. Norwegian Pollution Control Authority, Monitoring report no. 862/02 TA no. 1919/2002. Norwegian Institute for Water Research project 80106, report number 4600-2002, 269 pp.. ISBN number 82-577-4260-0.
- Green, N.W., Severinsen G., Rogne, Å.K., 2002c. Joint Assessment and Monitoring Programme (JAMP). Contaminant data for fish 1998-2001. Norwegian Pollution Control Authority, Monitoring report no. 863/02 TA no. 1920/2002. Norwegian Institute for Water Research project 80106, report number 4601-2002, 336 pp.. ISBN number 82-577-4261-9.
- Green, N.W., Severinsen G., Rogne, Å.K., 2002d. Joint Assessment and Monitoring Programme (JAMP). Summary statistics for contaminants in shellfish and fish 1981-2001. Norwegian Pollution Control Authority, Monitoring report no. 864/02 TA no. 1921/2002. Norwegian Institute for Water Research project 80106, report number 4602-2002, 422 pp.. ISBN number 82-577-4262-7.
- Green, N.W., Hylland, K., Ruus, A., Walday, M., 2002e. Joint Assessment and Monitoring Programme (JAMP). National Comments regarding the Norwegian Data for 2000. Norwegian Pollution Control Authority, Monitoring report no. 842/02 TA no. 1854/2002. Norwegian Institute for Water Research project 80106, report number 4468-2002, 197 pp.. ISBN number 82-577-4115-9. (Also presentert som SIME 2002 document 02/2/info. 2).
- Green, N.W., Hylland, K., Ruus, A., Walday, M., 2003. Joint Assessment and Monitoring Programme (JAMP). National Comments regarding the Norwegian Data for 2001. Norwegian Pollution Control Authority, Monitoring report no. 867/02 TA no. 1926/2002. Norwegian Institute for Water Research project 80106, report number 4618-2002, 217 pp.. ISBN number 82-577-4279-1.
- Green, N.W., Hylland, K., Ruus, A., Walday, M., 2004a. Joint Assessment and Monitoring Programme (JAMP). National Comments regarding the Norwegian Data for 2002. Norwegian Pollution Control Authority, Monitoring report no. 894/2003 TA no. 2003/2003. Norwegian Institute for Water Research project 80106, report number 4778-2004, 223 pp.. ISBN number 82-577-4454-9. Also as Trends and Effects of Substances in the Marine Environment (SIME) London (Secretariat) 24-26 February 2004. SIME 04/02/info. 4 -E
- Green, N.W., Ruus, A., Walday, M., 2004b. Joint Assessment and Monitoring Programme (JAMP). NationalComments regarding the Norwegian Data for 2003. Norwegian Pollution Control Authority,Monitoring report no. 921/2004 TA no. 2072/2004. Norwegian Institute for Water Research project 80106, report number 4927-2004, 219 pp.. ISBN number 82-577-4618-5. Also as Trends and Effects of Substances in the Marine Environment (SIME) Vigp (Spain) 15-17 Mars 2005. SIME 05/02/info. 7 -E
- Harding M.J.C., Bailey S.K. & I.M. Davies. 1992. TBT imposex survey of the North Sea. Annex 7:Norway. Scottish Fisheries working paper No 10/92 (1992).
- ICES, 1996. ICES Environmental Data Reporting Formats. Version 2.2, revision 2 - July 1996.
- IARC, 1987. IARC [International Agency for Research on Cancer] monographs on the evaluation of the carcinogenic risk of chemicals to humans. Overall evaluations of carcinogenicity: an updating of IARC monographs. Vol., 1-42. Suppl. 7. Lyon.
- Knutzen, J., Skei, J., 1990. Kvalitetskriterier for miljøgifter i vann, sedimenter og organismer samt foreløpige forslag til klassifikasjon av miljøkvalitet. [Quality criteria for water, sediments and organisms and preliminary proposals for classification of environmental quality]. NIVA-prosjekt O-8000309 (NIVA report no. 2540), 139 sider. ISBN.82-577-1855-6.
- Knutzen, J., Green, N.W., 1995. Bakgrunnsnivåer av en del miljøgifter i fisk, blåskjell og reker. Data fra utvalgte norske prøvesteder innen den felles overvåking under Oslo-/Paris-kommisjonene 1990-1993. [Background levels of some micropollutants in fish, the blue mussel and shrimps. Data from selected Norwegian sampling sites within the joint monitoring of the Oslo-/Paris Commissions 1990-1993]. Norwegian Pollution Control Authority, Monitoring report no. 594/94 TA no. 1173/1994. NIVA project O-80106/E-91412, (report number 3302) 105 pp.. ISBN number 82-577-2678-8.

- Knutzen, J., Green, N.W., 2001a. Tiltaksorienterte miljøundersøkelser i Sørfjorden og Hardangerfjord 2000. Delrapport 2. Miljøgifter i organismer. [Investigation of micropollutants in the Sørfjord and Hardangerfjord 2000. Report 2. Contaminants in organisms.] Norwegian Pollution Control Authority, Monitoring report no. 836/01. TA no. 1833/2001. NIVA project O-800309, (report number 4445-2001) 51 pp.. ISBN number 82-577-4091-8.
- Knutzen, J., Green, N.W., 2001b. Joint Assessment and Monitoring Programme (JAMP). "Bakgrunnsnivåer" av miljøgifter i fisk og blåskjell basert på datamateriale fra 1990-1998.[Joint Assessment and Monitoring Programme (JAMP). Background levels of some contaminants in fish and blue mussel based on data from 1990-1998]. Norwegian Pollution Control Authority, Monitoring report no. 820/01 TA no. 1798/2001. NIVA project O-80106, (report number 4339) 145 pp.. ISBN number 82-577-3973-1.
- Knutzen, J., Bjerkeng, B., Green, N.W., Kringstad, M., Schlaback, M., Skåre J.U., 2001. Overvåking av miljøgifter i fisk og skalldyr fra Grenlandsfjordene 2000. [Monitoring of micropollutants in fish and shellfish from the Grenland fjords (S. Norway) 2000] Norwegian Pollution Control Authority, Monitoring report no. 835/01. TA no. 1832/2001. NIVA project O-800309, (report number 4452-2001) 230 pp.. ISBN number 82-577-4098-5.
- Molvær, J., Knutzen, J., Magnusson, J., Rygg, B., Skei J., Sørensen, J., 1997. Klassifisering av miljøkvalitet i fjorder og kystfarvann. Veileddning. *Classification of environmental quality in fjords and coastal waters. A guide.* Norwegian Pollution Control Authority. TA no. TA-1467/1997. 36 pp.
- MON, 1993. Draft Summary record. Eleventh meeting of the Ad Hoc Working Group on Monitoring, Copenhagen: 8-12 November 1993. MON 11/1/7-E.
- MON, 1998. Summary record. Ad Hoc Working Group on Monitoring, Copenhagen: 23-27 February 1998. MON 98/6/1-E.
- MON, 2001. The first draft of a new OSPAR Strategy for a Joint Assessment and Monitoring Programme (JAMP). Working Group on Monitoring (MON), Belfast: 4-7 December 2001. MON 01/7/1-E.
- Minchin, A. Davies, I.M., 1999. Effect of freezing on the length of the penix in *Nucella lapillus* (L.). *J. Environ. Monit.* 1999(1):203-205.
- Nelson, D.R., Koymans, L., Kamataki, T., Stegeman, J.J., Feyreisen, R., Waxman, D.J., Waterman, M.R., Gotoh, O., Coon, M.J., Estabrook, R.W., Gunsalus, I.C., Nebert, D.W., 1996. P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics.* 6:1-42.
- Nicholson, M., Fryer, R.J., Green, N.W., 1994. Focusing on key aspects of contaminant trend assessments. Nineteenth meeting of the Joint Monitoring Group 24-29 . January, 1994. Document JMG 19/3/3.
- Nicholson, M., Fryer, R.J., Maxwell, D.M., 1997. A study of the power of various methods for detecting trends. ICES CM 1997/Env.11.
- Nicholson, M., Fryer, R.J., Larsen, J.R., 1998. Temporal trend monitoring: A robust method for analysing trend monitoring data, ICES Techniques in Marine Environmental Sciences, No.20 September 1998.
- OSPAR, 1990. Oslo and Paris Conventions. Principles and methodology of the Joint Monitoring Programme. [Monitoring manual for participants of the Joint Monitoring Programme (JMP) and North Sea Monitoring Master Plan (NSMMP)]. March 1990.
- OSPAR, 1997. JAMP Guidelines. Oslo and Paris Commissions, Joint Assessment and Monitoring Programme (including chapter updates 1998-1999).
- Ruus, A., Green, N.W., 2002. Tiltaksorienterte miljøundersøkelser i Sørfjorden og Hardangerfjorden 2001. Delrapport 2. Miljøgifter i organismer. *Measure oriented environmental monitoring of Sørfjord and Hardangerfjord 2002. Report component 2. Contaminants in organisms.* Norwegian Pollution Control Authority, Monitoring report no. 865/02 TA no. 1922/2002. NIVA project 800309 (report number 4612-2002), 41 pp.. ISBN number 82-577-4273-2.
- Ruus, A., Green, N.W., 2003. Miljøforholdene i Sørfjorden i 2002. Delrapport 2. Overvåking av miljøforholdene i Sørfjorden. Miljøgifter i organismer i 2002. *Monitoring of environmental quality in the Sørfjord. Report component 2 Contaminants in organisms 2002.* Norwegian Pollution Control Authority, Monitoring report no. 885/03 TA no. 1983/2003. NIVA project 800309 (report number 4724-2003), 45 pp.. ISBN number 82-577-4394-1.
- Ruus, A., Green, N.W., 2004. Overvåking av miljøforholdene i Sørfjorden. Miljøgifter i organismer 2003. Delrapport 3. Miljøgifter i organismer. *Monitoring of environmental quality in the Sørfjord. Contaminants in organisms 2003. Report component 3. Contaminants in organisms.* Norwegian Pollution Control Authority, Monitoring report no. 908/04 TA no. 2045/2004. NIVA project 800309 (report number 4880-2004), 54 pp.. ISBN number 82-577-4566-9.
- Ruus, A., Green, N.W., 2005. Overvåking av miljøforholdene i Sørfjorden. Miljøgifter i organismer 2004. Delrapport 3. Miljøgifter i organismer. *Monitoring of environmental quality in the Sørfjord. Contaminants in organisms 2004. Report component 3. Contaminants in organisms.* Norwegian Pollution Control Authority, Monitoring report no. 938/05 TA no. 2123/2005. NIVA project 800309 (report number 5069-2005), 61 pp.. ISBN number 82-577-4774-2.

- Ruus, A., Hylland, K., Green, N., 2003. Joint Assessment and Monitoring Programme (JAMP). Biological Effects Methods, Norwegian Monitoring 1997-2001. Norwegian Pollution Control Authority, Monitoring report no. 869/03 TA no. 1948/2003. Norwegian Institute for Water Research project 80106, report number 4649-2003, 139 pp.. ISBN number 82-577-4313-5.
- Schlabach, M., Mariussen, E., Borgen, A.: Dye, C., Enge, E.-K., Steinnes E., Green, N., Mohn, H. 2003. Kartleggin av bromerte flammehemmere og klorerte parafiner [Investigations of brominated flame retardants and chlorinated paraffins] Norwegian Pollution Control Authority, Monitoring report no. 866/02 TA no. 1924/2002. Norwegian Institute for Atmospheric Research, report number 62/2002, 69 pp.. ISBN number 82-425-1411-9.
- SIME, 1997. Summary Record. Oslo and Paris Conventions for the Prevention of Marine Pollution. Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME), Bonn: 17-21 November 1997. SIME (2) 97/12/1-E, 29 pp. + Annexes.
- SIME, 2002, OSPAR Coordinated Environmental Monitoring Programme (CEMP). Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME), Oudenburg: 29-31 January 2002. Summary Record, Annex 5 SIME 02/10/1-E, 8 pp.
- SIME, 2003. OSPAR Convention for the Protection of the Marine Environment of the North East Atlantic. Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME) London (Secretariat) 18-20 March 2003. SIME 03/10/1-E [Summary record], 18 pp. + 10 annexes.
- Skei, J., Rygg, B., Moy, F., Molvær, J., Knutzen, J., Hylland, K., Næs, K., Green, N., Johansen, T., 1998. Forerensningensutviklingen i Sørfjorden og Hardangerfjorden i perioden 1980-1997. Sammenstilling av resultater fra overvåking av vann, sedimenter og organismer. [The development of pollution in the Sørfjord and Hardangerfjord during the period 1980-1997. Summary of results from monitoring of water, sediment and organisms.] Norwegian Pollution Control Authority, Monitoring report no. 742/98 TA no. 1581/1998. NIVA project O-800310, (report number 3922-98) 95 pp.. ISBN number 82-577-3507-8.
- Skei, J., 2000. Tiltaksorienterte miljøundersøkelser i Sørfjorden og Harangerfjorden 1999. Delrapport 1 Vannkjemi. [Investigation of Sørfjord and Hardangerfjord 1999. Report 1. Water Chemistry] Norwegian Pollution Control Authority, Monitoring report no. 796/00 TA no. 1724/2000. NIVA project O-800309, (report number 4236-2000) 23 pp.. ISBN number 82-577-3858-1.
- Skei, J., 2001. Tiltaksorienterte miljøundersøkelser i Sørfjorden og Harangerfjorden 2000. Delrapport 1 Vannkjemi. [Investigation of Sørfjord and Hardangerfjord 2000. Report 1. Water Chemistry] Norwegian Pollution Control Authority, Monitoring report no. 830/01 TA no. 1818/2001. NIVA project O-800309, (report number 4406-2001) 22 pp.. ISBN number 82-577-4048-9.
- Skei, J., Knutzen, J., 2000. Utslipp av kvikksølv til Sørfjorden som følge av uhell ved Norzink AS vinteren 1999-2000. Miljømessige konsekvenser. [Accidental mercury discharge by Norzink AS to the Sørfjord, winter 1999-2000. Environmental consequences.] NIVA project O-89083 (report number 4234-2000) 12pp.. ISBN number 82-577-3856-5.
- Van den Berg, Birnbaum, L, Bosveld, A. T. C. and co-workers, 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Hlth. Perspect. 106:775-792.
- Walday, M., Green, N., Hylland, K., 1995. Kostholds- og tilstandsindikatorer for miljøgifter i marine områder. Norwegian Institute for Water Research project 93254, report number 3280, 39 pp.. ISBN number 82-577-2691-5.
- Walday M., Berge J.A. & N. Følsvik. 1997. Imposex and levels of organotin in populations of *Nucella lapillus* in Norway (English summary). Norwegian Institute for Water Research, report no. 3665-97. 28pp.
- WGSAEM, 1993. The length effect on contaminant concentrations in mussels. Section 13.2. in the Report of the Working Group on Statistical Aspects of Environmental Monitoring, Copenhagen 27-30, April 1993. International Council for the Exploration of the Sea. C-M- 1993/ENV:6 Ref.: D and E, 61 pp.
- Økland, T.E., 2005. Kostholdersråd i norske havner og fjorder. En gjennomgang av kostholdersråd i norske havner og fjorder fra 1960-tallet til i dag. Bergfeld & Co. ISBN 82-92650-01-6. 269pp.

## **Appendix A**

### **Quality assurance programme**



## Accreditation

The laboratories at NIVA, both the chemical, microbiological and the ecotoxicological laboratories, were accredited in 1993 for quality assurance system by the National Measurement Service - Norwegian Accreditation and based on European Standard EN45000. NIVA has reference number P009. The chemical laboratoruim satisfy the requirements in NS-EN ISO/IEC 17025.

## Summary of quality control results

Standard reference materials were analysed regularly (Table 8). Dogfish muscle (DORM-2) or dogfish liver (DOLT-3) was used as SRM for the control of the determination of metals (see Table A1). Mackerel oil (350) and mussel tissue (2977) was used as SRM for controls of PCBs and PAHs, respectively. NIES 11 was used for tin organic compounds. Cyprinid fish (EDF2525) at NILU was used as SRM for control of determination of dioxins. MESS-3 was used as marine SRM for trace metals in sediments and 1944 for the determination of selected polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyl (PCB) congeners, chlorinated pesticides, in marine sediment.

NIVA has participated in QUASIMEME exercises up to Round 42 (July to October 2005) which includes:

- QTM067BT - 068BT for metals in biota
- QTM072MS – 073MS for metals in sediment
- QPH039BT-040BT for PAH in biota
- QPH046MS-047MS for PAH in sediment
- QO084BT-085 BT organochlorines in biota
- QOR082MS-083 MS organochlorines in sediment

Following results for round Quasimeme –round 40 (January-April 2004) were used. This round would apply to the 2004 samples:

- QOR082BT (no.1) and QOR083BT (no.2) for organochlorines in biota.  
The results were acceptable (z-scores between -2 and 2), only two results were not classified as satisfactory those were CB101 (no.2) and g-HCH (no.1).
- QPH037BT and QPH038BT for PAH in biota.  
The results were generally acceptable, but the results for Benzo(b)fluoranthene (no.1 and no.2) and Benzo(k)fluoranthene (no. 2) were not classified as satisfactory.

A summary of the results for the analyses of the SRM sediment are shown in Table 8.

NIVA has participated in QUASIMEME Round 40 for sediments, exercises which include:

- QTM070MS (no.1) and QTM071MS (no.2) for metals in sediments, with satisfactory results with exception of lead (no.1). This was caused by contamination during analysis.
- QOR080MS (no.1) and QOR081MS (no.2) for organochlorines in sediments about 75 % of all the results were satisfactory.  
QPH044MS (no.1) and QPH045MS (no.2) the results were all over satisfactory, but the results for Anthracene (no.2), Benzo(b)fluoranthene (no.1 -and 2) and Perylene (no.2) were not classified as satisfactory.

**Table 8.** Summary of the quality control of results for the 2004 biota samples analysed in 2004. The Standard Reference Materials (SRM) were DORM-2<sup>\*</sup> (dogfish muscle) for mussels and fish fillet, DOLT-3<sup>\*</sup> (dogfish liver) for fish liver, 350<sup>\*\*</sup> (mackerel oil) for mussels and fish liver and 2977<sup>\*\*\*</sup> (mussel tissue) for mussels. SRM was analysed in series with the JAMP-samples for analyses of metals (mg/kg d.w.), NIES 11 for organochlorines or PAH (µg/kg d.w.) and EDF2525\*\*\*\* for fish (cyprinid) was analysed for dioxin(ng/kg) by NILU(Norsk institutt for luftforskning). Tissue types were: mussel softbody (SB), fish liver (LI) and fish fillet (MU). SRMs were measured several times (N) over a number of weeks (W).

Code	Contaminant	Tissue type	SRM type	SRM value ± confidence interval	N	W	Mean value	Standard deviation
<b>Cd</b>	<b>cadmium</b>	SB	DORM	0.043 ± 0.008	6	4	0.047	0.002
		LI	DOLT	19.4 ± 0.6	11	7	18.7	0.7
<b>Cu</b>	<b>copper</b>	SB	DORM	2.34 ± 0.16	6	4	2.16	0.23
		LI	DOLT	31.2 ± 1.0	11	7	30.4	1.2
<b>Pb</b>	<b>lead</b>	SB	DORM	0.065 ± 0.007	6	4	0.058	0.006
		LI	DOLT	0.319 ± 0.045	9	7	0.31	0.04
<b>Hg</b>	<b>mercury</b>	SB	DORM	4.64 ± 0.26	9	8	4.96	0.18
<b>Zn</b>	<b>zinc</b>	SB	DORM	25.6 ± 2.3	6	4	26.3	1.4
		LI	DOLT	86.6 ± 2.4	11	7	92.2	3.8
<b>CB-28</b>	<b>PCB congener CB-28</b>	(all)	350	22.5 ± 4	21	9	15.8	1.3
<b>CB-52</b>	<b>PCB congener CB-52</b>	(all)	350	62. ± 9	21	9	58.9	5.2
<b>CB-101</b>	<b>PCB congener CB-101</b>	(all)	350	164 ± 9	18	8	172	25
<b>CB-118</b>	<b>PCB congener CB-118</b>	(all)	350	142 ± 20	21	9	130	12
<b>CB-153</b>	<b>PCB congener CB-153</b>	(all)	350	317 ± 20	21	9	303	23
<b>CB-180</b>	<b>PCB congener CB-180</b>	(all)	350	73 ± 13	18	8	66.9	6.7
<b>BAA</b>	<b>benzo[a]anthracene1)</b>	SB	2977	20.34 ± 0.78	3	2	23.3	2.1
<b>BAP</b>	<b>benzo[a]pyrene1)</b>	SB	2977	8.35 ± 0.72	3	2	7.5	1.2
<b>BBF</b>	<b>benzo[b]fluoranthene1)</b>	SB	2977	11.01 ± 0.28	3	2	19	1
<b>BEP</b>	<b>benzo[e]pyrene</b>	SB	2977	13.1 ± 1.1	3	2	7.5	1.2
<b>BGHIP</b>	<b>benzo[ghi]perylene</b>	SB	2977	9.53 ± 0.43	3	2	11.7	0.6
<b>BKF</b>	<b>benzo[k]fluoranthene</b>	SB	2977	4 ± 1	3	2	7.73	0.50
<b>CHRTR</b>	<b>chrysene+triphenylene1 2)</b>	SB	2977	88 ± 2.2	3	2	48.3	0.6
<b>FLU</b>	<b>fluoranthene</b>	SB	2977	38.7 ± 1.0	3	2	36	0.001
<b>ICDP</b>	<b>indeno[1,2,3-cd]pyrene</b>	SB	2977	4.84 ± 0.81	3	2	4.40	0.52
<b>PER</b>	<b>perylene</b>	SB	2977	3.50 ± 0.76	3	2	2.43	0.50
<b>PYR</b>	<b>pyrene</b>	SB	2977	78.9 ± 3.5	3	2	63.3	0.6
<b>TBTIN</b>	<b>Tributyl-tin</b>	SB	11	1159 ± 88	12	15	969	192
<b>TPTIN</b>	<b>Triphenyl-tin</b>	SB	11	5109 ± 363	11	15	2280	1020
<b>CB 126</b>	<b>3,3,4,4,5-PeCB</b>	MU	2525	647 ± 148	4	42	661	36.1
<b>CB 169</b>	<b>3,3,4,4,5-HxCB</b>	MU	2525	50 ± 12	4	42	58.3	4.13
<b>CB77</b>	<b>3,3,4,4-TeCB</b>	MU	2525	1945 ± 354	4	42	1895	96
<b>CDD1N</b>	<b>1,2,3,7,8-HxCDD</b>	MU	2525	4.0 ± 0.57	4	42	3.22	0.33
<b>CDD4X</b>	<b>1,2,3,4,7,8-HxCDD</b>	MU	2525	0.77 ± 0.27	4	42	0.47	0.2
<b>CDD6X</b>	<b>1,2,3,6,7,8- HxCDF</b>	MU	2525	2.7 ± 1.2	4	42	1.31	0.54
<b>CDD9X</b>	<b>1,2,3,7,8,9-HpCDF</b>	MU	2525	0.63 ± 0.23	4	42	0.56	0.2
<b>CDDO</b>	<b>OCDD</b>	MU	2525	7.2 ± 3.7	4	42	1.4	0.9
<b>CDF2N</b>	<b>2,3,4,7,8-PeCDF</b>	MU	2525	14 ± 1.3	4	42	15	1.85
<b>CDF2T</b>	<b>2,3,7,8 TCDF</b>	MU	2525	22 ± 1.6	4	42	24.9	3.18
<b>CDF4X</b>	<b>2,3,4,6,7,8 HxCDF</b>	MU	2525	2.3 ± 1.9	4	42	0.73	0.44
<b>CDF6P</b>	<b>1,2,3,4,6,7,8,-HxCDF</b>	MU	2525	4.4 ± 6.0	4	42	0.34	0.12
<b>CDF6X</b>	<b>1,2,3,6,7,8-HxCDF</b>	MU	2525	2.7 ± 1.2	4	42	1.11	0.51
<b>CDF9P</b>	<b>1,2,3,4,7,8,9-HpCDF</b>	MU	2525	0.63 ± 0.23	4	42	0.42	0.3
<b>CDFDN</b>	<b>1,2,3,7,8/1,2,3,4,8-PeCDF</b>	MU	2525	4.9 ± 0.56	4	42	4.89	0.91
<b>CDFDX</b>	<b>1,2,3,4,7,8/1,2,3,4,7,9-HxCDF</b>	MU	2525	8.2 ± 3.7	4	42	6.97	0.7
<b>CDFO</b>	<b>OCDF</b>	MU	2525	2.6 ± 1.3	4	42	0.65	0.4
<b>TCDD</b>	<b>2,3,7,8-Tetra-DiBpD(TCDD)</b>	MU	2525	17 ± 1.4	4	42	16.87	1.11

<sup>\*</sup>) National Research Council Canada, Division of Chemistry, Marine Analytical Chemistry Standards

<sup>\*\*</sup>) BCR, Community Bureau of Reference, Commission of the European Communities

<sup>\*\*\*</sup>) National Institute of Standards & Technology (NIST)

<sup>\*\*\*\*</sup>) CIL, US.

<sup>1)</sup> Not certified (see NIST certificate)

<sup>2)</sup> Calculated from separate values for chrysene and triphenylene; respectively,  $(49+39) \pm \sqrt{(2^2 + 1^2)}$

**Table 9.** Summary of the quality control of results for the 2004 biota samples analysed in 2004. The Standard Reference Materials (SRM) were MESS-3 for trace metals in marine sediment (mg/kg d.w) and 1944 for the determination of selected polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyl (PCB) congeners, chlorinated pesticides, and trace elements ( $\mu\text{g}/\text{kg}$  d.w) in marine sediment. These SRM were analysed in series with the JAMP-samples for analyses of metals and the organic compounds.

Code	Contaminant	Tissue type	SRM type	SRM value confidence interval	N	W	Mean value	Standard deviation
AL	Aluminium	SM	MESS-3	8.59 $\pm$ 0.23	3	2	80.8	1.0
AS	Arsenic	SM	MESS-3	21.2 $\pm$ 1.1	3	2	23	6.1
CD	Cadmium	SM	MESS-3	0.24 $\pm$ 0.01	2	2	0.227	0.006
CR	Chromium	SM	MESS-3	105 $\pm$ 4	3	2	92.6	1.3
CU	Copper	SM	MESS-3	33.9 $\pm$ 1.6	3	2	30.9	0.5
HG	Mercury	SM	MESS-3	0.091 $\pm$ 0.009	3	2	0.086	0.002
LI	Lithium	SM	MESS-3	73.6 $\pm$ 5.2	3	2	71.1	1.0
MN	Magnesium	SM	MESS-3	324 $\pm$ 12	3	2	304	2.5
NI	Nickel	SM	MESS-3	46.9 $\pm$ 2.2	3	2	44.8	0.47
PB	Lead	SM	MESS-3	21.1 $\pm$ 0.7	3	2	18	1
ZN	Zinc	SM	MESS-3	159 $\pm$ 8	3	2	145	0.6
ACNE	Acenaphthene	SM	1944	570 $\pm$ 30	3	4	290	10
ANT	Anthracene	SM	1944	1770 $\pm$ 330	3	4	1133	115
BAP	Benzo(a)pyrene	SM	1944	4300 $\pm$ 130	3	4	4667	757
BBF	Benzo(b)fluoranthene	SM	1944	3870 $\pm$ 420	3	4	6700	1513
BGHIP	Benzo(ghi)perylene	SM	1944	2780 $\pm$ 100	3	4	3367	252
BAA	Benzo(a)anthracene	SM	1944	4720 $\pm$ 110	3	4	5067	850
CB101	CB101(IUPAC)	SM	1944	73.4 $\pm$ 2.5	2	3	69	9.9
CB105	CB105(IUPAC)	SM	1944	24.5 $\pm$ 1.1	2	3	22	0
CB118	CB118(IUPAC)	SM	1944	58.0 $\pm$ 4.3	2	3	47	1.4
CB138	CB138(IUPAC)	SM	1944	62.1 $\pm$ 3.0	2	3	57	1.4
CB153	CB153(IUPAC)	SM	1944	74.0 $\pm$ 2.9	2	3	56.5	3.5
CB156	CB156(IUPAC)	SM	1944	6.52 $\pm$ 0.66	2	3	6.8	0.28
CB180	CB180(IUPAC)	SM	1944	44.3 $\pm$ 1.2	2	3	42.5	2.1
CB209	CB209(IUPAC)	SM	1944	6.81 $\pm$ 0.33	2	3	6.75	0.35
CB28	CB28(IUPAC)	SM	1944	80.8 $\pm$ 2.7	2	3	73.5	2.1
CB52	CB52(IUPAC)	SM	1944	79.4 $\pm$ 2.0	2	3	64	1.4
CHRTR <sup>1)</sup>	Chrysene + Triphenylene	SM	1944	5900 $\pm$ 499	3	4	4800	173
DBA3A	Dibenz(a,c,a,h)anthracene	SM	1944		3	4	893	78
DDEPP	p,p'-DDE	SM	1944		2	3	72	1.4
FLE	Fluorene	SM	1944	850 $\pm$ 30	3	4	373	15
FLU	Fluoranthene	SM	1944	8920 $\pm$ 320	3	4	9400	794
HCB	Hexachlorobenzene	SM	1944	6.03 $\pm$ 0.35	2	3	5.05	0.21
HCHA	alpha-HCH	SM	1944		2	3	0.40	0.56
ICDP	Indeno(1,2,3-cd)pyrene	SM	1944	2780 $\pm$ 100	3	4	3567	252
NAP	Naphthalene	SM	1944	1650 $\pm$ 310	3	4	1167	58
PA	Phenanthrene	SM	1944	5270 $\pm$ 220	3	4	5300	265
PYR	Pyrene	SM	1944	9700 $\pm$ 420	3	4	8600	265
TDEPP	p,p'-TDE = p,p'-DDD	SM	1944		2	3	108	18

<sup>1)</sup> Calculated from separate values for chrysene and triphenylene; respectively,  $(4.86 + 1.04) \pm \sqrt{(0.27^2 + 0.42^2)}$



## **Appendix B**

## **Abbreviations**



Abbreviation <sup>1</sup>	English	Norwegian
<b>ELEMENTS</b>		
Al	aluminium	<i>aluminium</i>
As	arsenic	<i>arsen</i>
Cd	cadmium	<i>kadmium</i>
Co	cobalt	<i>kobolt</i>
Cr	chromium	<i>krom</i>
Cu	copper	<i>kobber</i>
Fe	iron	<i>jern</i>
Hg	mercury	<i>kvikksølv</i>
Li	lithium	<i>litium</i>
Mn	manganese	<i>mangan</i>
Ni	nickel	<i>nikkel</i>
Pb	lead	<i>bly</i>
Pb210	lead-210	<i>bly-210</i>
Se	selenium	<i>selen</i>
Ti	titanium	<i>titan</i>
Zn	zinc	<i>sink</i>
<b>METAL COMPOUNDS</b>		
TBT	tributyltin	<i>tributyltinn</i>
<b>PAHs</b>		
PAH	polycyclic aromatic hydrocarbons	<i>polysyklike aromatiske hydrokarboner</i>
ACNE	acenaphthene	<i>acenaften</i>
ACNLE	acenaphthylene	<i>acenaftylen</i>
ANT	anthracene	<i>antracen</i>
BAA <sup>3</sup>	benzo[a]anthracene	<i>benzo[a]antracen</i>
BAP <sup>3</sup>	benzo[a]pyrene	<i>benzo[a]pyren</i>
BBF <sup>3</sup>	benzo[b]fluoranthene	<i>benzo[b]fluoranten</i>
BBJKF <sup>3</sup>	benzo[b,j,k]fluoranthene	<i>benzo[b,j,k]fluoranten</i>
BBJKF <sup>3</sup>	benzo[b+j,k]fluoranthene	<i>benzo[b+j,k]fluoranten</i>
BBKF <sup>3</sup>	benzo[b+k]fluoranthene	<i>benzo[b+k]fluoranten</i>
BEP	benzo[e]pyrene	<i>benzo[e]pyren</i>
BGHIP	benzo[ghi]perylene	<i>benzo[ghi]perylen</i>
BIPN <sup>2</sup>	biphenyl	<i>bifenyl</i>
BJKF <sup>3</sup>	benzo[j,k]fluoranthene	<i>benzo[j,k]fluorantren</i>
BKF <sup>3</sup>	benzo[k]fluoranthene	<i>benzo[k]fluorantren</i>
CHR	chrysene	<i>chrysen</i>
CHRTR	chrysene+triphenylene	<i>chrysen+trifenylen</i>
COR	coronene	<i>coronen</i>
DBAHA <sup>3</sup>	dibenz[a,h]anthracene	<i>dibenz[a,h]antracen</i>
DBA3A <sup>3</sup>	dibenz[a,c/a,h]anthracene	<i>dibenz[a,c/a,h]antracen</i>
DBP <sup>3</sup>	dibenzopyrenes	<i>dibenzopyren</i>
DBT	dibenzothiophene	<i>dibenzothiofen</i>
DBTC1	C <sub>1</sub> -dibenzothiophenes	<i>C<sub>1</sub>-dibenzotiofen</i>
DBTC2	C <sub>2</sub> -dibenzothiophenes	<i>C<sub>2</sub>-dibenzotiofen</i>

Abbreviation <sup>1</sup>	English	Norwegian
<b>DBTC3</b>	C <sub>3</sub> -dibenzothiophenes	<i>C<sub>3</sub>-dibenzotiofen</i>
<b>FLE</b>	fluorene	<i>fluoren</i>
<b>FLU</b>	fluoranthene	<i>fluoranten</i>
<b>ICDP</b> <sup>3</sup>	indeno[1,2,3-cd]pyrene	<i>indeno[1,2,3-cd]pyren</i>
<b>NAP</b> <sup>2</sup>	naphthalene	<i>naftalen</i>
<b>NAPC1</b> <sup>2</sup>	C <sub>1</sub> -naphthalenes	<i>C<sub>1</sub>-naftalen</i>
<b>NAPC2</b> <sup>2</sup>	C <sub>2</sub> -naphthalenes	<i>C<sub>2</sub>-naftalen</i>
<b>NAPC3</b> <sup>2</sup>	C <sub>3</sub> -naphthalenes	<i>C<sub>3</sub>-naftalen</i>
<b>NAP1M</b> <sup>2</sup>	1-methylnaphthalene	<i>1-metylnaftalen</i>
<b>NAP2M</b> <sup>2</sup>	2-methylnaphthalene	<i>2-metylnaftalen</i>
<b>NAPD2</b> <sup>2</sup>	1,6-dimethylnaphthalene	<i>1,6-dimetylnaftalen</i>
<b>NAPD3</b> <sup>2</sup>	1,5-dimethylnaphthalene	<i>1,5-dimetylnaftalen</i>
<b>NAPDI</b> <sup>2</sup>	2,6-dimethylnaphthalene	<i>2,6-dimetylnaftalen</i>
<b>NAPT2</b> <sup>2</sup>	2,3,6-trimethylnaphthalene	<i>2,3,6-trimetylnaftalen</i>
<b>NAPT3</b> <sup>2</sup>	1,2,4-trimethylnaphthalene	<i>1,2,4-trimetylnaftalen</i>
<b>NAPT4</b> <sup>2</sup>	1,2,3-trimethylnaphthalene	<i>1,2,3-trimetylnaftalen</i>
<b>NAPTM</b> <sup>2</sup>	2,3,5-trimethylnaphthalene	<i>2,3,5-trimetylnaftalen</i>
<b>NPD</b>	Collective term for naphthalenes, phenanthrenes and dibenzothiophenes	<i>Sammebetegnelse for naftalen, fenantren og dibenzotiofens</i>
<b>PA</b>	phenanthrene	<i>fenantren</i>
<b>PAC1</b>	C <sub>1</sub> -phenanthrenes	<i>C<sub>1</sub>-fanantren</i>
<b>PAC2</b>	C <sub>2</sub> -phenanthrenes	<i>C<sub>2</sub>-fanantren</i>
<b>PAM1</b>	1-methylphenanthrene	<i>1-metylfanantren</i>
<b>PAM2</b>	2-methylphenanthrene	<i>2-metylfanantren</i>
<b>PADM1</b>	3,6-dimethylphenanthrene	<i>3,6-dimetylfanantren</i>
<b>PADM2</b>	9,10-dimethylphenanthrene	<i>9,10-dimetylfanantren</i>
<b>PER</b>	perylene	<i>perylén</i>
<b>PYR</b>	pyrene	<i>pyren</i>
<b>DI-Σn</b>	sum of "n" dicyclic "PAH"s (footnote 2)	<i>sum "n" disykkliske "PAH" (fotnote 2)</i>
<b>P-Σn / P_S</b>	sum "n" PAH (DI-Σn not included)	<i>sum "n" PAH (DI-Σn ikke inkludert)</i>
<b>PK-Σn / PK_S</b>	sum carcinogen PAHs (footnote 3)	<i>sum kreftfremkallende PAH (fotnote 3)</i>
<b>PAHΣΣ</b>	DI-Σn + P-Σn etc.	<i>DI-Σn + P-Σn mm..</i>
<b>SPAH</b>	"total" PAH, specific compounds not quantified (outdated analytical method)	<i>"total" PAH, spesifikk forbindelser ikke kvantifisert (foreldret metode)</i>
<b>BAP_P</b>	% BAP of PAHΣΣ	<i>% BAP av PAHΣΣ</i>
<b>BAPPP</b>	% BAP of P-Σn	<i>% BAP av P-Σn</i>
<b>BPK_P</b>	% BAP of PK-Σn	<i>% BAP av PK-Σn</i>
<b>PKn_P</b>	% PK-Σn of PAHΣΣ	<i>% PK-Σn av PAHΣΣ</i>
<b>PKnPP</b>	% PK-Σn of P-Σn	<i>% PK-Σn av P-Σn</i>
<b>PCBs</b>		
<b>PCB</b>	polychlorinated biphenyls	<i>polyklorerte bifenyler</i>
<b>CB</b>	individual chlorobiphenyls (CB)	<i>enkelte klorobifenyler</i>
<b>CB28</b>	CB28 (IUPAC)	<i>CB28 (IUPAC)</i>
<b>CB31</b>	CB31 (IUPAC)	<i>CB31 (IUPAC)</i>
<b>CB44</b>	CB44 (IUPAC)	<i>CB44 (IUPAC)</i>
<b>CB52</b>	CB52 (IUPAC)	<i>CB52 (IUPAC)</i>
<b>CB77</b> <sup>4</sup>	CB77 (IUPAC)	<i>CB77 (IUPAC)</i>

Abbreviation <sup>1</sup>	English	Norwegian
<b>CB81</b> <sup>4</sup>	CB81 (IUPAC)	<i>CB81 (IUPAC)</i>
<b>CB95</b>	CB95 (IUPAC)	<i>CB95 (IUPAC)</i>
<b>CB101</b>	CB101 (IUPAC)	<i>CB101 (IUPAC)</i>
<b>CB105</b>	CB105 (IUPAC)	<i>CB105 (IUPAC)</i>
<b>CB110</b>	CB110 (IUPAC)	<i>CB110 (IUPAC)</i>
<b>CB118</b>	CB118 (IUPAC)	<i>CB118 (IUPAC)</i>
<b>CB126</b> <sup>4</sup>	CB126 (IUPAC)	<i>CB126 (IUPAC)</i>
<b>CB128</b>	CB128 (IUPAC)	<i>CB128 (IUPAC)</i>
<b>CB138</b>	CB138 (IUPAC)	<i>CB138 (IUPAC)</i>
<b>CB149</b>	CB149 (IUPAC)	<i>CB149 (IUPAC)</i>
<b>CB153</b>	CB153 (IUPAC)	<i>CB153 (IUPAC)</i>
<b>CB156</b>	CB156 (IUPAC)	<i>CB156 (IUPAC)</i>
<b>CB169</b> <sup>4</sup>	CB169 (IUPAC)	<i>CB169 (IUPAC)</i>
<b>CB170</b>	CB170 (IUPAC)	<i>CB170 (IUPAC)</i>
<b>CB180</b>	CB180 (IUPAC)	<i>CB180 (IUPAC)</i>
<b>CB194</b>	CB194 (IUPAC)	<i>CB194 (IUPAC)</i>
<b>CB209</b>	CB209 (IUPAC)	<i>CB209 (IUPAC)</i>
<b>CB-Σ7</b>	CB: 28+52+101+118+138+153+180	<i>CB: 28+52+101+118+138+153+180</i>
<b>CB-ΣΣ</b>	sum of CBs, includes CB-Σ7	<i>sum CBer, inkluderer CB-Σ7</i>
<b>TECBW</b>	Sum of CB-toxicity equivalents after WHO model, see <b>TEQ</b>	<i>Sum CB-toksitets ekvivalenter etter WHO modell, se <b>TEQ</b></i>
<b>TECBS</b>	Sum of CB-toxicity equivalents after SAFE model, see <b>TEQ</b>	<i>Sum CB-toksitets ekvivalenter etter SAFE modell, se <b>TEQ</b></i>
<b>DIOXINS</b>		
<b>TCDD</b>	2, 3, 7, 8-tetrachloro-dibenzo dioxin	<i>2, 3, 7, 8-tetrakloro-dibenzo dioksin</i>
<b>CDDST</b>	Sum of tetrachloro-dibenzo dioxins	<i>Sum tetrakloro-dibenzo dioksiner</i>
<b>CDD1N</b>	1, 2, 3, 7, 8-pentachloro-dibenzo dioxin	<i>1, 2, 3, 7, 8-pentakloro-dibenzo dioksin</i>
<b>CDDSN</b>	Sum of pentachloro-dibenzo dioxins	<i>Sum pentakloro-dibenzo dioksiner</i>
<b>CDD4X</b>	1, 2, 3, 4, 7, 8-hexachloro-dibenzo dioxin	<i>1, 2, 3, 4, 7, 8-heksakloro-dibenzo dioksin</i>
<b>CDD6X</b>	1, 2, 3, 6, 7, 8-hexachloro-dibenzo dioxin	<i>1, 2, 3, 6, 7, 8-heksakloro-dibenzo dioksin</i>
<b>CDD9X</b>	1, 2, 3, 7, 8, 9-hexachloro-dibenzo dioxin	<i>1, 2, 3, 7, 8, 9-heksakloro-dibenzo dioksin</i>
<b>CDDSX</b>	Sum of hexachloro-dibenzo dioxins	<i>Sum heksakloro-dibenzo dioksiner</i>
<b>CDD6P</b>	1, 2, 3, 4, 6, 7, 8-heptachloro-dibenzo dioxin	<i>1, 2, 3, 4, 6, 7, 8-heptakloro-dibenzo dioksin</i>
<b>CDDSP</b>	Sum of heptachloro-dibenzo dioxins	<i>Sum heptakloro-dibenzo dioksiner</i>
<b>CDDO</b>	Octachloro-dibenzo dioxin	<i>Oktakloro-dibenzo dioksin</i>
<b>PCDD</b>	Sum of polychlorinated dibenzo-p-dioxins	<i>Sum polyklorinaterte-dibenzo-p-dioksiner</i>
<b>CDF2T</b>	2, 3, 7, 8-tetrachloro-dibenzofuran	<i>2, 3, 7, 8-tetrakloro-dibenzofuran</i>
<b>CDFST</b>	Sum of tetrachloro-dibenzofurans	<i>Sum tetrakloro-dibenzofuraner</i>
<b>CDFDN</b>	1, 2, 3, 7, 8/1, 2, 3, 4, 8-pentachloro-dibenzofuran	<i>1, 2, 3, 7, 8/1, 2, 3, 4, 8-pentakloro-dibenzofuran</i>
<b>CDF2N</b>	2, 3, 4, 7, 8-pentachloro-dibenzofurans	<i>2, 3, 4, 7, 8-pentakloro-dibenzofuran</i>
<b>CDFSN</b>	Sum of pentachloro-dibenzofurans	<i>Sum pentakloro-dibenzofuraner</i>
<b>CDFDX</b>	1, 2, 3, 4, 7, 8/1, 2, 3, 4, 7, 9-hexachloro-dibenzofuran	<i>1, 2, 3, 4, 7, 8/1, 2, 3, 4, 7, 9-heksakloro-dibenzofuran</i>
<b>CDF6X</b>	1, 2, 3, 6, 7, 8-hexachloro-dibenzofuran	<i>1, 2, 3, 6, 7, 8-heksakloro-dibenzofuran</i>
<b>CDF9X</b>	1, 2, 3, 7, 8, 9-hexachloro-dibenzofuran	<i>1, 2, 3, 7, 8, 9-heksakloro-dibenzofuran</i>
<b>CDF4X</b>	2, 3, 4, 6, 7, 8-hexachloro-dibenzofuran	<i>2, 3, 4, 6, 7, 8-heksakloro-dibenzofuran</i>

Abbreviation <sup>1</sup>	English	Norwegian
<b>CDFSX</b>	Sum of hexachloro-dibenzofurans	<i>Sum heksakloro-dibenzofuraner</i>
<b>CDF6P</b>	1, 2, 3, 4, 6, 7, 8-heptachloro-dibenzofuran	<i>1, 2, 3, 4, 6, 7, 8-heptakloro-dibenzofuran</i>
<b>CDF9P</b>	1, 2, 3, 4, 7, 8, 9-heptachloro-dibenzofuran	<i>1, 2, 3, 4, 7, 8, 9-heptakloro-dibenzofuran</i>
<b>CDFSP</b>	Sum of heptachloro-dibenzofurans	<i>Sum heptakloro-dibenzofuraner</i>
<b>CDO</b>	Octachloro-dibenzofuran	<i>Octakloro-dibenzofuran</i>
<b>PCDF</b>	Sum of polychlorinated dibenzo-furans	<i>Sum polyklorinated dibenzo-furaner</i>
<b>CDDFS</b>	Sum of PCDD and PCDF	<i>Sum PCDD og PCDF</i>
<b>TCDDN</b>	Sum of TCDD-toxicity equivalents after Nordic model, see <b>TEQ</b>	<i>Sum TCDD- toksitets ekvivalenter etter Nordisk modell, se <b>TEQ</b></i>
<b>TCDDI</b>	Sum of TCDD-toxicity equivalents after international model, see <b>TEQ</b>	<i>Sum TCDD-toksitets ekvivalenter etter internasjonale modell, se <b>TEQ</b></i>
<b>PESTICIDES</b>		
<b>ALD</b>	aldrin	<i>aldrin</i>
<b>DIELD</b>	dieldrin	<i>dieldrin</i>
<b>ENDA</b>	endrin	<i>endrin</i>
<b>CCDAN</b>	cis-chlordane (=α-chlordane)	<i>cis-klordan (=α-klordan)</i>
<b>TCDAN</b>	trans-chlordane (=γ-chlordane)	<i>trans-klordan (=γ-klordan)</i>
<b>OCDAN</b>	oxy-chlordane	<i>oksy-klordan</i>
<b>TNONC</b>	trans-nonachlor	<i>trans-nonaklor</i>
<b>TCDAN</b>	trans-chlordane	<i>trans-klordan</i>
<b>OCS</b>	octachlorostyrene	<i>oktaklorstyren</i>
<b>QCB</b>	pentachlorobenzene	<i>pentaklorbenzen</i>
<b>DDD</b>	dichlorodiphenylchloroethane 1,1-dichloro-2,2-bis-(4-chlorophenyl)ethane	<i>diklordinfenyldikloretan 1,1-dikloro-2,2-bis-(4-klorofenyl)etan</i>
<b>DDE</b>	dichlorodiphenylchloroethylene (principle metabolite of DDT) 1,1-dichloro-2,2-bis-(4-chlorophenyl)ethylene*	<i>diklordinfenyldikloetylen (hovedmetabolitt av DDT) 1,1-dikloro-2,2-bis-(4-klorofenyl)etylen</i>
<b>DDT</b>	dichlorodiphenyltrichloroethane 1,1,1-trichloro-2,2-bis-(4-chlorophenyl)ethane	<i>diklordinfenyltrikloretan 1,1,1-trikloro-2,2-bis-(4-klorofenyl)etan</i>
<b>DDEOP</b>	o,p'-DDE	<i>o,p'-DDE</i>
<b>DDEPP</b>	p,p'-DDE	<i>p,p'-DDE</i>
<b>DDTOP</b>	o,p'-DDT	<i>o,p'-DDT</i>
<b>DDTPP</b>	p,p'-DDT	<i>p,p'-DDT</i>
<b>TDEPP</b>	p,p'-DDD	<i>p,p'-DDD</i>
<b>DDTEP</b>	p,p'-DDE + p,p'-DDT	<i>p,p'-DDE + p,p'-DDT</i>
<b>DD-nΣ</b>	sum of DDT and metabolites, n = number of compounds	<i>sum DDT og metabolitter, n = antall forbindelser</i>
<b>HCB</b>	hexachlorobenzene	<i>heksaklorbenzen</i>
<b>HCHG</b>	Lindane γ HCH = gamma hexachlorocyclohexane (γ BHC = gamma benzenehexachloride, outdated synonym)	<i>Lindan γ HCH = gamma heksaklorosykloheksan (γ BHC = gamma benzenheksaklorid, foreldret betegnelse)</i>
<b>HCHA</b>	α HCH = alpha HCH	<i>α HCH = alpha HCH</i>
<b>HCHB</b>	β HCH = beta HCH	<i>β HCH = beta HCH</i>
<b>HC-nΣ</b>	sum of HCHs, n = count	<i>sum av HCHs, n = antall</i>

Abbreviation <sup>1</sup>	English	Norwegian
<b>EOCI</b>	extractable organically bound chlorine	<i>ekstraherbart organisk bundet klor</i>
<b>EPOCI</b>	extractable persistent organically bound chlorine	<i>ekstraherbart persistent organisk bundet klor</i>
<b>NTOT</b>	total organic nitrogen	<i>total organisk nitrogen</i>
<b>CTOT</b>	total organic carbon	<i>total organisk karbon</i>
<b>CORG</b>	organic carbon	<i>organisk karbon</i>
<b>GSAMT</b>	grain size	<i>kornfordeling</i>
<b>MOCON</b>	moisture content	<i>vanninnhold</i>

**INSTITUTES**

<b>IFEN</b>	Institute for Energy Technology	<i>Institutt for energiteknikk</i>
<b>FIER</b>	Institute for Nutrition, Fisheries Directorate	<i>Fiskeridirektoratets Ernæringsinstitutt</i>
<b>FORC</b>	FORCE Institutes, Div. for Isotope Technique and Analysis [DK]	<i>FORCE Institutterne, Div. for Isotopteknik og Analyse [DK]</i>
<b>IMRN</b>	Institute of Marine Research (IMR)	<i>Havforskningsinstituttet</i>
<b>NACE</b>	Nordic Analytical Center	<i>Nordisk Analyse Center</i>
<b>NILU</b>	Norwegian Institute for Air Research	<i>Norsk institutt for luftforskning</i>
<b>NIVA</b>	Norwegian Institute for Water Research	<i>Norsk institutt for vannforskning</i>
<b>SERI</b>	Swedish Environmental Research Institute	<i>Institutionen för vatten- och luftvårdsforskning</i>
<b>VETN</b>	Norwegian Veterinary Institute	<i>Veterinærinstituttet</i>
<b>SIIF</b>	Fondation for Scientific and Industrial Research at the Norwegian Institute of Technology - SINTEF (a division, previously: Center for Industrial Research SI)	<i>Stiftelsen for industriell og teknisk forskning ved Norges tekniske høgskole- SINTEF (en avdeling, tidligere: Senter for industorforskning SI)</i>

<sup>1)</sup> After: ICES Environmental Data Reporting Formats. International Council for the Exploration of the Sea. July 1996 and supplementary codes related to non-ortho and mono-ortho PCBs and "dioxins" (ICES pers. comm.)

<sup>2)</sup> Indicates "PAH" compounds that are dicyclic and not truly PAHs typically identified during the analyses of PAH, include naphthalenes and "biphenyls".

<sup>3)</sup> Indicates PAH compounds potentially cancerogenic for humans according to IARC (1987), i.e., categories 2A+2B (possibly and probably carcinogenic).

<sup>4)</sup> Indicates non ortho- co-planer PCB compounds i.e., those that lack Cl in positions 1, 1', 5, and 5'

\*) The Pesticide Index, second edition. The Royal Society of Chemistry, 1991.

**Other abbreviations andre forkortelser**

	English	Norwegian
<b>TEQ</b>	"Toxicity equivalency factors" for the most toxic compounds within the following groups:	"Toxositetsekvivalentfaktorer" for de giftigste forbindelsene innen følgende grupper.
	<ul style="list-style-type: none"> <li>• polychlorinated dibenzo-p-dioxins and dibenzofurans (<b>PCDD/PCDFs</b>). Equivalents calculated after Nordic model (Ahlborg 1989)<sup>1</sup> or international model (Int./EPA, cf. Van den Berg <i>et al.</i>, 1998)<sup>2</sup></li>   <li>• non-ortho and mono-ortho substituted chlorobiphenyls after WHO model (Ahlborg <i>et al.</i>, 1994)<sup>3</sup> or Safe (1994, cf. NILU pers. comm.)</li> </ul>	<ul style="list-style-type: none"> <li>• polyklorerte dibenzo-p-dioksiner og dibenzofuraner (<b>PCDD/PCDF</b>). Ekvivalentberegning etter nordisk modell (Ahlborg 1989)<sup>1</sup> eller etter internasjonal modell (Int./EPA, cf. Van den Berg <i>et al.</i> 1998)<sup>2</sup></li>   <li>• non-ortho og mono-ortho substituerte klorobifenyler etter WHO modell (Ahlborg <i>et al.</i>, 1994)<sup>3</sup> eller Safe (1994, cf. NILU pers. medd.)</li> </ul>
<b>ppm</b>	parts per million, mg/kg	deler pr. milliondeler, mg/kg
<b>ppb</b>	parts per billion, µg/kg	deler pr. milliarddeler, µg/kg
<b>ppp</b>	parts per trillion, ng/kg	deler pr. tusen-milliarddeler, ng/kg
<b>d.w.</b>	dry weight basis	tørrvekt basis
<b>w.w.</b>	wet weight or fresh weight basis	våtvekt eller friskvekt basis

<sup>1</sup>) Ahlborg, U.G., 1989. Nordic risk assessment of PCDDs and PCDFs. Chemosphere 19:603-608.

<sup>2</sup>) Van den Berg, Birnbaum, L, Bosveld, A. T. C. and co-workers, 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Hlth. Perspect. 106:775-792.

<sup>3</sup>) Ahlborg, U.G., Becking G.B., Birnbaum, L.S., Brouwer, A, Derkks, H.J.G.M., Feely, M., Golor, G., Hanberg, A., Larsen, J.C., Liem, A.K.G., Safe, S.H., Schlatter, C., Wärn, F., Younes, M., Yrjänheikki, E., 1994. Toxic equivalency factors for dioxin-like PCBs. Report on a WHO-ECEH and IPSC consultation , December 1993. Chemosphere 28:1049-1067.

## **Appendix C**

### **Analytical overview, sediment and biota**

**Sorted by:**

**Contaminant, year, laboratory, intercalibration**

**Abbreviations are defined in Appendix B and Appendix D**

<b>Contamin.</b>	<b>Contaminant defined in Appendix B</b>
<b>Mon. Year</b>	<b>Monitoring year</b>
<b>Lab.</b>	<b>Analytical laboratory (cf. Appendix B)</b>
<b>Intercalibr. +basis</b>	<b>Intercalibration exercise (cf. Appendix D) and basis where W = wet weight and D = dry weight .</b>
<b>Detect limit</b>	<b>"Normal" detection limit</b>
<b>Count below d.lim</b>	<b>Number of analyses below normal detection limit</b>
<b>N (&lt;) above d.lim</b>	<b>Number of analyses where detection limit was higher than normal.</b>



## Analytical overview - sediment

Contamin.	Mon. Year	Lab. Inter- calibr. +basis	Analys method code	Detect limit (ppb)	Total value count	Count below d.lim	N (<) above d.lim
ACNE	1992-NIVA	D	369	~1	23	23	
	1994-NIVA		369	1	24	23	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
ACNE	2004-NIVA	QW	D	369	1	156	1 44
ACNLE	1992-NIVA	D	369	~1	23	23	
	1994-NIVA		369	1	24	23	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
	2004-NIVA		QW	369	1	156	1 56
AL	1987-NIVA	D	352	~0.001	28		
	1990-NIVA		352	~0.001	128		
AL	2004-NIVA	QT	D	355	10000	173	
ALD	1990-IMRN	D	760	~0.05	14	14	
ANT	1990-IMRN	D	769	~1	14	14	
	1992-NIVA		369	~1	24	24	
	1994-NIVA		369	1	24	22	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
ANT	2004-NIVA	D	369	1	156		27
AS	1994-NIVA	D	354	500	12		
AS	2004-NIVA	QT	D	355	15000	172	21
BAP	1990-IMRN	D	769	~1	14	14	
	1992-NIVA		369	~1	23	23	
	1994-NIVA		369	1	24	12	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
BAP	2004-NIVA	QW	D	369	1	156	2
BBF	1992-NIVA	D	369	~1	23	23	
	1994-NIVA		369	1	24	9	
BBF	2004-NIVA	QW	D	369	0.5	156	
BBJKF	1996-NIVA	D	369	1	10		
	1997-NIVA		369	1	18		
BBKF	1990-IMRN	D	769	~1	14	14	
BEP	1990-IMRN	D	769	~1	14	14	
	1992-NIVA		369	~1	23	23	
	1994-NIVA		369	1	24	8	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
BGHIP	1990-IMRN	D	769	~1	14	14	
	1992-NIVA		369	~1	24	24	
	1994-NIVA		369	1	24	9	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
	2004-NIVA		QW	369	1	156	
BIPN	1992-NIVA	D	369	~1	23	23	
	1994-NIVA		369	1	24	21	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
BJKF	1992-NIVA	D	369	~1	14	14	
	1994-NIVA		369	1	24	11	
BJKF	2004-NIVA	D	369	0.5	92		1
BAA	1990-IMRN	D	769	~1	14	14	
	1992-NIVA		369	~1	24	24	
	1994-NIVA		369	1	24	11	
	1996-NIVA		369	1	10		
	1997-NIVA		369	1	18		
BAA	2004-NIVA	QW	D	369	1	156	3

JAMP National Comments 2004 - Norway

Contamin.	Mon. Year	Lab. +basis	Inter- calibr.	Analys method code	Detect limit (ppb)	Total value	Count below d.lim	N (<) above d.lim
CB101	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		12
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
	2004-NIVA	QV	D	360	0.2	151		1
CB105	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		24
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
	2004-NIVA	QV	D	360	0.2	146		1
CB118	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		13
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	17		
	2004-NIVA	QV	D	360	0.2	155		1
CB128	1990-IMRN		D	760	~0.05	14	14	
CB138	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		12
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
	2004-NIVA	QV	D	360	0.2	153		1
CB149	1990-IMRN		D	760	~0.05	14	14	
CB153	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		12
	1996-NIVA		D	360	0.05	10		
	1997-NIVA		D	360	0.05	18		
	2004-NIVA	QV	D	360	0.05	81		19
CB156	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		22
	1996-NIVA		D	360	0.2	10	1	
	1997-NIVA		D	360	0.2	18	2	
	2004-NIVA		D	360	0.2	154		1
CB170	1990-IMRN		D	760	~0.05	14	14	
CB180	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		13
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
	2004-NIVA	QV	D	360	0.2	156		1
CB209	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8C	D	360	0.05	24		12
	1996-NIVA		D	360	0.2	10	1	
	1997-NIVA		D	360	0.2	18	1	
	2004-NIVA		D	360	0.2	152		5
CB28	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	23	23	
	1994-NIVA	8Z	D	360	0.05	24		2
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
CB28	2004-NIVA	QV	D	360	0.2	152		1
CB31	1990-IMRN	8B	D	760	~0.05	14	14	
CB52	1990-IMRN	8B	D	760	~0.05	14	14	
	1992-NIVA	8C	D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		2
	1996-NIVA		D	360	0.2	10		

JAMP National Comments 2004 - Norway

Contamin.	Mon. Year	Lab. +basis	Inter- calibr.	Analys method code	Detect limit (ppb)	Total value	Count below d.lim	N (<) above d.lim
	1997-NIVA		D	360	0.2	18		
CB52	2004-NIVA	QV	D	360	0.2	133		
CD	1986-NIVA	7C	D	352	~0.001	24	24	
	1987-NIVA	7C	D	352	~0.001	25	25	
	1990-NIVA		D	353	~0.001	14	14	
	1990-NIVA	7E	D	353	~0.001	114	114	
	1992-NIVA	7E	D	353	~0.001	107	107	
	1994-NIVA	7Z	D	353	1	114		
	1996-NIVA		D	353	200	23	22	
	1997-NIVA		D	353	200	27	15	
CD	2004-NIVA		D	353	50	173	3	
CHR	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	24	24	
CHRTR	1994-NIVA		D	369	0.5	24		
	1996-NIVA		D	369	0.5	10		
	1997-NIVA		D	369	0.5	18		
	2004-NIVA	QW	D	369	0.5	156	2	
COR	1992-NIVA		D	369	~1	24	24	
CORG	1986-NIVA		D	390	~1	18	18	
	1987-NIVA		D	390	~1	28	28	
	1990-NIVA		D	390	~0.2	128	128	
	1992-NIVA		D	390	~0.2	107	107	
	1994-NIVA		D	390	200000	114		
	1996-NIVA		D	390	200000	23		
	1997-NIVA		D	390	200000	27		
CORG	2004-NIVA		D	390	200000	173		
CR	1994-NIVA	7Z	D	353	5	12		
CR	2004-NIVA	QT	D	355	1500	173		
CTOT	1994-NIVA		D	390	1000000	12		
	1996-NIVA		D	390	1000000	23		
	1997-NIVA		D	390	1000000	27		
CU	1986-NIVA	7C	D	351	~0.01	24	24	
	1987-NIVA	7C	D	351	~0.01	28	28	
	1990-NIVA	7E	D	351	~0.01	128	128	
	1992-NIVA	7E	D	351	~0.01	107	107	
	1994-NIVA	7Z	D	351	10	114		
	1996-NIVA		D	351	10	23		
	1997-NIVA		D	351	10	27		
CU	2004-NIVA	QT	D	355	1000	173		
DBA3A	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	23	11	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
	2004-NIVA	QW	D	369	1	156	20	
DBAHA	1990-IMRN		D	769	~1	14	14	
DBP	1992-NIVA		D	369	~1	24	24	
DBT	1990-IMRN		D	769	~1	14	14	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
DBT	2004-NIVA	QW	D	369	1	156	40	
DBTC1	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	0.5	156	50	
DBTC2	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	0.5	156	57	
DBTC3	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	0.5	156	63	
DBTIN	2004-NIVA		D	370	0.26	141	23	
DDEOP	1990-IMRN		D	760	~0.05	14	14	
DDEPP	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		12

JAMP National Comments 2004 - Norway

Contamin.	Mon. Year	Lab. +basis	Inter- calibr.	Analys method code	Detect limit (ppb)	Total value	Count below d.lim	N (<) above d.lim
	1996-NIVA		D	360	0.05	10		
	1997-NIVA		D	360	0.05	18		
	2004-NIVA	QV	D	360	0.05	151		102
DDTOP	1990-IMRN		D	760	~0.05	14	14	
DDTPP	1990-IMRN		D	760	~0.05	14	14	
	1996-NIVA		D	999	0.7	10		5
	1997-NIVA		D	999	0.7	18		3
DPTIN	2004-NIVA		D	370	~0.22	128	128	
FLE	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24		23
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
FLE	2004-NIVA	QW	D	369	1	156		32
FLU	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24		10
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
FLU	2004-NIVA	QW	D	369	1	156		1
GSAMT	1996-NIVA		D	miss	miss	31		
	1996-VKID		D	miss	miss	35		
	1997-NIVA		D	miss	miss	45		
	1997-VKID		D	miss	miss	47		
	2004-NIVA		D	miss	miss	229		
HCB	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA		8Z	360	0.05	24		10
	1996-NIVA		D	360	0.1	10		
	1997-NIVA		D	360	0.1	18		
HCB	2004-NIVA		D	360	0.1	141		117
HCHA	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA		8Z	360	0.05	24		23
	1996-NIVA		D	360	0.2	10	2	
	1997-NIVA		D	360	0.2	18	1	
HCHA	2004-NIVA		D	360	0.2	148		4
HCHB	1990-IMRN		D	760	~0.05	14	14	
HCHG	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA		8Z	360	0.05	24		15
	1996-NIVA		D	360	0.2	10	1	
	1997-NIVA		D	360	0.2	18	1	
HCHG	2004-NIVA	QV	D	360	0.2	149		7
HG	1986-NIVA	7C	D	350	~0.01	24	24	
	1987-NIVA	7C	D	350	~0.01	28	28	
	1990-NIVA	7E	D	350	~0.01	128	128	
	1992-NIVA	7E	D	350	~0.01	107	107	
	1994-NIVA	7Z	D	350	10	114	2	
	1996-NIVA		D	350	10	23		
	1997-NIVA		D	350	10	27		
HG	2004-NIVA		D	350	10	173	7	
ICDP	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24		12
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
ICDP	2004-NIVA	QW	D	369	1	156		
LI	1990-NIVA	7E	D	353	~0.001	14	14	
	1992-NIVA	7E	D	353	~0.001	107	107	
	1994-NIVA	7E	D	353	1	114		

JAMP National Comments 2004 - Norway

Contamin.	Mon. Year	Lab. +basis	Inter- calibr.	Analys method code	Detect limit (ppb)	Total value	Count below d.lim	N (<) above d.lim
	1996-NIVA		D	353	1	23		
	1997-NIVA		D	353	1	27		
LI	2004-NIVA	QT	D	355	1000	173		
MBTIN	2004-NIVA		D	370	0.34	142	32	
MN	2004-NIVA	QT	D	355	300	172		
MOCON	1994-NIVA		D	340	~1	62		
	1996-NIVA		D	340	~1	31		
	1996-VKID		D	340	~1	35		
	1997-VKID		D	340	~1	47		
	2004-NIVA		D	340	~1	173		
MPTIN	2004-NIVA		D	370	0.3	118	65	
NAP	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24	18	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAP1M	2004-NIVA	QW	D	369	1	154	27	
	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24	19	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAP2M	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24	17	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPC1	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	2	156	15	
NAPC2	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	2	156	40	
NAPC3	1990-IMRN		D	769	~1	14	14	
	2004-NIVA		D	369	2	156	28	
NAPD2	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPD3	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPDI	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24	18	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPT2	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPT3	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPT4	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NAPTM	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24	24	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
NI	1994-NIVA	7Z	D	353	50	12		
NI	2004-NIVA	QT	D	355	2000	173		
NTOT	1994-NIVA		D	390	1000000	114		
	1996-NIVA		D	390	1000000	23		
	1997-NIVA		D	390	1000000	27		
NTOT	2004-NIVA		D	390	1000000	173		
OCS	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA		D	360	0.05	24		24
	1996-NIVA		D	360	0.1	10		
	1997-NIVA		D	360	0.1	18	1	1
OCS	2004-NIVA		D	360	0.1	152	142	
PA	1990-IMRN		D	769	~1	14	14	

JAMP National Comments 2004 - Norway

Contamin.	Mon. Year	Lab. +basis	Inter- calibr.	Analys method code	Detect limit (ppb)	Total value	Count below d.lim	N (<) above d.lim
	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24	11	
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PA	2004-NIVA	QW	D	369	1	156		2
PAC1	1990-IMRN		D	769	~1	14	14	
PAC1	2004-NIVA		D	369	2	156		14
PAC2	1990-IMRN		D	769	~1	14	14	
PAC2	2004-NIVA		D	369	2	156		28
PAC3	2004-NIVA		D	369	2	156		55
PADM1	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PADM2	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PAM1	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24		17
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PAM2	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PB	1986-NIVA	7C	D	352	~0.05	24	24	
	1987-NIVA	7C	D	352	~0.05	28	28	
	1990-NIVA		D	353	~0.05	14	14	
	1990-NIVA	7E	D	353	~0.001	114	108	
	1992-NIVA	7E	D	353	~0.001	107	107	
	1994-NIVA	7Z	D	353	1	114		
	1996-NIVA		D	353	1	23		
	1997-NIVA		D	353	1	27		
PB	2004-NIVA	QT	D	355	10000	173		
PB210	1990-VKID		D	650	~1	70	26	
	1992-VKID		D	650	~1	56	15	
	1994-VKID		D	650	~1	62	25	
	1996-VKID		D	650	~1	11		
	1997-VKID		D	650	~1	21	3	
PER	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	23	23	
	1994-NIVA		D	369	1	24		3
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PYR	1990-IMRN		D	769	~1	14	14	
	1992-NIVA		D	369	~1	24	24	
	1994-NIVA		D	369	1	24		12
	1996-NIVA		D	369	1	10		
	1997-NIVA		D	369	1	18		
PYR	2004-NIVA	QW	D	369	1	156		1
QCB	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA		D	360	0.05	24		22
	1996-NIVA		D	360	0.05	10		
	1997-NIVA		D	360	0.05	18		
QCB	2004-NIVA		D	360	0.05	142		100
SPAH	1990-IMRN		D	769	~1	14	14	
TBTIN	2004-NIVA		D	370	0.2	142		28
TDEOP	1990-IMRN		D	760	~0.05	14	14	
TDEPP	1990-IMRN		D	760	~0.05	14	14	
	1992-NIVA		D	360	~0.05	24	24	
	1994-NIVA	8Z	D	360	0.05	24		21
	1996-NIVA		D	360	0.2	10		
	1997-NIVA		D	360	0.2	18		
	2004-NIVA		D	360	0.2	155		38
TPTIN	2004-NIVA		D	370	0.17	141		69
ZN	1986-NIVA	7C	D	351	~0.1	24	24	

JAMP National Comments 2004 - Norway

Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)
	Year		calibr.	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim
1987-NIVA		7C	D	351	~0.1	28	28	
1990-NIVA		7E	D	351	~0.01	128	128	
1992-NIVA		7E	D	351	~0.1	107	107	
1994-NIVA		7Z	D	351	100	114		
1996-NIVA			D	351	100	23		
1997-NIVA			D	351	100	27		
ZN	2004-NIVA	QT	D	355	5000	173		
Sum of						17952	4147	1626

~

> converting to ppb ignored, due to missing unit

## Analytical overview - biota

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-calibr.	Analys method	Detect limit	Total value	Count below	N (<)	Analys method	Detect limit	Total value	Count below	N (<)
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
ACNE	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	72		20
	1996-NIVA		W						309	0.2	65		19
	1997-NIVA		W						309	0.5	34		
	1998-NIVA	CI	W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	38		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	46		
ACNLE	2004-NIVA	R5	W						309	0.5	58	32	1
	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	72		49
	1996-NIVA		W						309	0.2	65		42
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42		
AG	2003-NIVA	MQ	W						309	0.5	55		
	2004-NIVA	R5	W						309	0.5	58	29	
ANT	1996-NIVA		W	309	0.2	8			999	miss	3		
	2004-NIVA		W						315	miss	7		5
AS	1992-NIVA		W						309	0.2	45		
	1995-NIVA		W						309	0.2	72		28
	1996-NIVA		W						309	0.2	65		30
	1997-NIVA		W						309	0.5	35		
	1998-NIVA	CI	W						309	0.5	39		
	1999-NIVA	EK	W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
BAP	2004-NIVA	R5	W						309	0.5	58	22	
BBF	1996-NIVA		W	309	0.2	8			999	miss	3		
	2004-NIVA		W						315	miss	7		
BBJKF	1992-NIVA		W						309	0.2	45		
	1995-NIVA		W						309	0.2	72		21
	1996-NIVA		W						309	0.2	65		26
	1997-NIVA	AL	W						309	0.5	36		
	1998-NIVA	CI	W						309	0.5	39		
	1999-NIVA	EK	W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
BBJKF	2004-NIVA	R5	W						309	0.5	58	11	

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
2000-NIVA			W						309	0.2	39		10
2001-NIVA			W						309	0.2	42		
2002-NIVA			W						309	0.2	43		9
2003-NIVA			W						309	0.2	50		9
2004-NIVA			W						309	0.2	21		
BD100	2001-NILU		W		miss	6			miss	6			
	2002-NILU		W						843	0.02	2		
	2004-NIVA		W						miss	2			2
BD138	2001-NILU		W		miss	6	6		miss	6		6	
	2004-NIVA		W						miss	2		2	
BD153	1996-NILU		W		miss	4	4		miss	6			
	2001-NILU		W		miss	6	4		843	0.01	2		4
	2002-NILU		W						miss	2			
	2004-NIVA		W						843	0.01	2		2
BD154	2001-NILU		W		miss	6			miss	6			
	2002-NILU		W						843	0.01	2		4
	2004-NIVA		W						miss	2		2	
BD183	2001-NILU		W		miss	6	3		miss	6		6	
	2002-NILU		W						843	0.01	2		
BD209	2001-NILU		W		miss	6	5		miss	6		1	
	2002-NILU		W						843	0.03	2		
BDE28	2001-NILU		W		miss	6			miss	6		1	
	2002-NILU		W						843	0.01	2		
BDE47	1996-NILU		W		miss	4			miss	6			
	2001-NILU		W		miss	6			843	0.11	2		
	2002-NILU		W						miss	2			
	2004-NIVA		W						843	0.06	2		2
BDE99	1996-NILU		W		miss	4			miss	6			
	2001-NILU		W		miss	6			843	0.06	2		1
	2002-NILU		W						miss	2			
	2004-NIVA		W						843	0.06	2		2
BEP	1992-NIVA		W		309	0.2	8		309	0.2	45		
	1995-NIVA		W						309	0.2	72		5
	1996-NIVA		W						309	0.2	65		6
	1997-NIVA		W						309	0.2	36		
	1998-NIVA	CI	W						309	0.2	38		
	1999-NIVA	EK	W						309	0.2	34		
	2000-NIVA		W						309	0.2	39		10
	2001-NIVA		W						309	0.2	42		
	2002-NIVA		W						309	0.2	43		9
	2003-NIVA	MQ	W						309	0.2	56		10
	2004-NIVA	R5	W						309	0.2	55		
BGHIP	1992-NIVA		W		309	0.2	8		309	0.2	46		
	1995-NIVA		W						309	0.2	72		20
	1996-NIVA		W						309	0.2	65		10
	1997-NIVA		W						309	0.5	36		
	1998-NIVA	CI	W						309	0.5	35		
	1999-NIVA	EK	W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA	R5	W						309	0.5	58		6
BIPN	1992-NIVA		W		309	0.2	8		309	0.2	46		
	1995-NIVA		W						309	0.2	72		52
	1996-NIVA		W						309	0.2	62		39
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		1
	1999-NIVA		W						309	0.5	34		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
2000-NIVA			W						309	0.5	38		1
2001-NIVA			W						309	0.5	41		
2002-NIVA			W						309	0.5	42		
2003-NIVA			W						309	0.5	55		1
BJKF	1992-NIVA		W		309	0.2	8		309	0.2	45		
	1995-NIVA		W						309	0.2	24		21
	1996-NIVA		W						309	0.2	57		16
	2004-NIVA		W						miss		37		
BAA	1992-NIVA		W		309	0.2	8		309	0.2	44		
	1995-NIVA		W						309	0.2	72		9
	1996-NIVA		W						309	0.2	65		8
	1997-NIVA		W						309	0.5	36		
	1998-NIVA	CI	W						309	0.5	39		
	1999-NIVA	EK	W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA	R5	W						309	0.5	58		3
CB101	1987-SIIF		W						111	0.2	21		1
	1988-SIIF		D						111	0.1	6		
	1988-SIIF		W						111	0.1	22		
	1989-NACE		W		510	20	93						
	1989-SIIF		W						111	0.1	36		
	1990-NIVA	2G	W		340	1	169	1	341	0.05	58		
	1990-SIIF	2G	W						111	0.4	41		6
	1991-NIVA	2H	W		340	1	179		341	0.05	68		
	1991-SIIF	2H	W						111	0.2	35		1
	1992-NIVA	2J	W		340	5	192	3	341	0.1	146		
	1993-NIVA	2K	W		340	4	212	12	341	0.1	138		
	1994-NIVA	2Z	W		340	3	300	3	341	0.05	170		39
	1995-NIVA		W		340	3	318	10	341	0.05	231		10
	1996-NIVA		W		340	3	332	14	341	0.05	243		9
	1997-NIVA		W		340	3	260	24					
	1997-NIVA	AJ	W						341	0.05	221		4
	1998-NIVA		W		340	3	284	19	1				
	1998-NIVA	CH	W						341	0.05	203		1
	1999-NIVA		W		340	3	249	6					3
	1999-NIVA	EG	W						341	0.05	232		13
	2000-NIVA		W		340	3	230	24					
	2000-NIVA	GU	W						341	0.05	186		11
	2001-NIVA		W		340	3	250	19	4				7
	2001-NIVA	IO	W						341	0.05	211		16
	2002-NIVA		W		340	3	241	13					17
	2002-NIVA	LJ	W						341	0.05	212		
	2003-NIVA		W		340	3	239	18					
	2003-NIVA	MO	W						341	0.05	175		6
	2004-NIVA		W		340	3	272	19					
	2004-NIVA	R1	W						341	0.05	170		
CB105	1991-NIVA	2H	W		340	1	87	1	341	0.05	47		
	1992-NIVA		W		340	5	192	3	341	0.1	146		
	1993-NIVA	QM	W		340	4	212	21	341	0.1	138		
	1994-NIVA	2Z	W		340	3	300	8	341	0.05	170		53
	1995-NIVA		W		340	3	318	13	341	0.05	230		34
	1996-NIVA		W		340	3	332	22	341	0.05	237		23
	1997-NIVA		W		340	3	260	24	341	0.05	221		3
	1998-NIVA		W		340	3	284	31	19				1
	1998-NIVA	CH	W						341	0.05	207		11
	1999-NIVA		W		340	3	249	17					16
	1999-NIVA	EG	W						341	0.05	232		4
													62

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
2000-NIVA		W		340	3	230	32		341	0.05	186	21	40
2000-NIVA	GU	W		340	3	250	29	2	341	0.05	211		76
2001-NIVA		W		340	3	249	30		341	0.05	210		59
2002-NIVA	IO	W		340	3	239	23		341	0.05	183		45
2003-NIVA		W		340	3	272	44		341	0.05	241		6
2004-NIVA	MO	W		340	20	93			341	0.1	36		
2004-NIVA	R1	W		340	1	169			341	0.05	58		
CB118	1989-NACE	W		340	1	179			341	0.2	41	1	
1989-SIIF		W		340	5	192	2		341	0.05	68		
1990-NIVA	2G	W		340	4	212	10		341	0.1	146		
1990-SIIF	2G	W		340	3	300	2		341	0.05	138		
1991-NIVA	2H	W		340	3	318	2		341	0.05	231	2	
1991-SIIF	2H	W		340	3	332	6		341	0.05	243	4	
1992-NIVA	2J	W		340	3	260	5		341	0.05	221		
1993-NIVA	2K	W		340	3	284	6	1	341	0.05	209	3	1
1994-NIVA	2Z	W		340	3	249	2		341	0.05	232		7
1995-NIVA		W		340	3	230	5		341	0.05	186	6	7
1996-NIVA	AJ	W		340	3	250	1	1	341	0.05	211		21
1997-NIVA		W		340	3	249	7		341	0.05	212		22
1998-NIVA	CH	W		340	3	239	6		341	0.05	183		18
1999-NIVA	EG	W		340	3	272	7		341	0.05	241		1
CB126	1995-NILU	W		841	0.0001	4			841	2E-05	6		
1996-NILU		W							841	0.0001	18		
2002-NILU		W							841	0.0001	12		
2003-NILU		W							841	0.0001	12		
CB138	1988-SIIF	D							111	0.1	6		
1988-SIIF		W							111	0.1	21		
1989-NACE		W		510	20	93			111	0.1	36		
1989-SIIF		W		340	1	169			341	0.05	58		
1990-NIVA	2G	W		340	1	179			111	0.3	41		
1990-SIIF	2G	W		340	5	192			341	0.05	68		
1991-NIVA	LJ	W		340	4	212	3		111	0.3	35	1	
1991-SIIF		W		340	3	300			341	0.1	143		
1992-NIVA	QM	W		340	3	318	2		341	0.1	138		
1993-NIVA	2Z	W		340	3	331	1		341	0.05	241		
1994-NIVA		W		340	3	260	1		341	0.05	221	1	
1995-NIVA		W		340	3	230	3		341	0.05	209		
1996-NIVA		W		340	3	250			341	0.05	232		1

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other					
Contamin.	Mon.	Lab.	Inter-calibr. +basis	Analys method code	Detect limit (ppb)	Total value count	Count below d.lim	N (< d.lim)	Analys method code	Detect limit (ppb)	Total value count	Count below d.lim	N (< d.lim)	
2000-NIVA	2000	GU	W	340	3	250	1	1	341	0.05	186	3	3	
			W											
		IO	W										7	
			W											
			W										6	
		MO	W											
			W										4	
		R1	W											
CB153	1988-SIIF	D		510	20	93			111	0.1	6			
	1988-SIIF	W												
	1989-NACE	W											36	
	1989-SIIF	W												
	1990-NIVA	2G	W										58	
	1990-SIIF	2G	W											
	1991-NIVA	2H	W										68	
	1991-SIIF	2H	W											
	1992-NIVA	2J	W										146	
	1993-NIVA	2K	W											
	1994-NIVA	2Z	W										9	
	1995-NIVA	W												
	1996-NIVA	W											138	
	1997-NIVA	W												
	1997-NIVA	AJ	W										221	
	1998-NIVA	W												
	1998-NIVA	CH	W										1	
	1999-NIVA	W												
	1999-NIVA	EG	W										1	
	2000-NIVA	W												
	2000-NIVA	GU	W										1	
	2001-NIVA	W												
	2001-NIVA	IO	W										5	
	2002-NIVA	W												
	2002-NIVA	LJ	W										4	
	2003-NIVA	W												
	2003-NIVA	MO	W										1	
	2004-NIVA	W												
	2004-NIVA	R1	W										1	
CB156	1991-NIVA	2H	W	340	1	87	15	15	341	0.05	47	5		
	1992-NIVA	W												
	1993-NIVA	QM	W											138
	1994-NIVA	2Z	W											73
	1995-NIVA	W												
	1996-NIVA	W												68
	1997-NIVA	W												
	1997-NIVA	AJ	W											62
	1998-NIVA	W												
	1998-NIVA	CH	W											10
	1999-NIVA	W												
	1999-NIVA	EG	W											139
	2000-NIVA	W												
	2000-NIVA	GU	W											95
	2001-NIVA	W												
CB169	2001-NIVA	IO	W											134
	2002-NIVA	W												102
	2003-NIVA	W												
	2003-NIVA	MO	W											83
	2004-NIVA	W												
	2004-NIVA	R1	W											7
	1995-NILU	W												6
	1996-NILU	W		841	0.0001	4				841	2E-05	18	2	

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
2002-NILU			W						841	0.0001	12		
2003-NILU			W						841	0.0001	12	1	1
CB180	1987-SIIF		W						111	0.2	21		6
	1988-SIIF		D						111	0.1	6		
	1988-SIIF		W						111	0.1	22		
	1989-NACE		W	510	20	93	1						
	1989-SIIF		W						111	0.1	36		
	1990-NIVA	2G	W	340	1	169			341	0.05	58		
	1990-SIIF	2G	W						111	0.2	41		8
	1991-NIVA	2H	W	340	1	179			341	0.05	68		
	1991-SIIF	2H	W						111	0.2	35		
	1992-NIVA	2J	W	340	5	192	3		341	0.1	146		
	1993-NIVA	2K	W	340	4	212	15		341	0.1	138		
	1994-NIVA	2Z	W	340	3	300	3		341	0.05	167		49
	1995-NIVA		W	340	3	318	5		341	0.05	231		22
	1996-NIVA		W	340	3	332	14		341	0.05	243		25
	1997-NIVA		W	340	3	260	18						
	1997-NIVA	AJ	W						341	0.05	221	1	1
	1998-NIVA		W	340	3	284	20	14					
	1998-NIVA	CH	W						341	0.05	209	19	44
	1999-NIVA		W	340	3	249	7	1					
	1999-NIVA	EG	W						341	0.05	232	2	78
	2000-NIVA		W	340	3	230	15						
	2000-NIVA	GU	W						341	0.05	186	15	83
	2001-NIVA		W	340	3	250	17	1					
	2001-NIVA	IO	W						341	0.05	211		99
	2002-NIVA		W	340	3	249	24						
	2002-NIVA	LJ	W						341	0.05	212		104
	2003-NIVA		W	340	3	238	13						
	2003-NIVA	MO	W						341	0.05	183		71
	2004-NIVA		W	340	3	272	14						
	2004-NIVA	R1	W						341	0.05	241		6
CB209	1990-NIVA		W	340	2	169	24	11	341	0.05	58		
	1991-NIVA		W	340	2	179	11	88	341	0.05	68	5	13
	1992-NIVA		W	340	5	192	3		341	0.1	146		1
	1993-NIVA		W	340	4	212	46	14	341	0.1	138		
	1994-NIVA		W	340	3	300	29	24	341	0.05	170		96
	1995-NIVA		W	340	3	318	36		341	0.05	231	95	5
	1996-NIVA		W	340	3	332	255		341	0.05	243	107	9
	1997-NIVA		W	340	3	260	196		341	0.05	221	30	14
	1998-NIVA		W	340	3	283	120	121	341	0.05	209	54	69
	1999-NIVA		W	340	3	243	163	17	341	0.05	230	19	178
	2000-NIVA		W	340	3	228	151	18	341	0.05	178	33	111
	2001-NIVA		W	340	3	250	184	10	341	0.05	211	21	185
	2002-NIVA		W	340	3	248	207	1	341	0.05	209		114
	2003-NIVA		W	340	3	236	126		341	0.05	177		99
	2004-NIVA		W	340	3	272	228		341	0.05	241		8
CB28	1988-SIIF		D						111	0.1	6		
	1988-SIIF		W						111	0.1	22		
	1989-NACE		W	510	20	93							
	1989-SIIF		W						111	0.1	36		1
	1990-NIVA	2G	W	340	1	169	2	2	341	0.05	58		
	1990-SIIF	2G	W						111	0.2	41		7
	1991-NIVA	2H	W	340	1	179	2	52	341	0.05	68	5	4
	1991-SIIF	2H	W						111	0.3	35		
	1992-NIVA	2J	W	340	5	192	3		341	0.1	143		
	1993-NIVA	2K	W	340	4	212	44	5	341	0.1	138		
	1994-NIVA	2Z	W	340	3	282	18	4	341	0.05	168		76
	1995-NIVA		W	340	3	313	27		341	0.05	231		80
	1996-NIVA		W	340	3	332	107		341	0.05	242		70

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1997-NIVA		W		340	3	260	81		341	0.05	221	22	14
1997-NIVA	AJ	W		340	3	284	96	99	341	0.05	207	36	46
1998-NIVA		W		340	3	249	96	18	341	0.05	232	14	145
1998-NIVA	CH	W		340	3	230	110	7	341	0.05	186	26	66
1999-NIVA		W		340	3	250	146	10	341	0.05	211	17	150
1999-NIVA	EG	W		340	3	249	144	1	341	0.05	207		101
2000-NIVA		W		340	3	238	97		341	0.05	173		75
2000-NIVA	GU	W		340	3	270	160		341	0.05	240		9
2001-NIVA		W											
2001-NIVA	IO	W											
2002-NIVA		W											
2002-NIVA	LJ	W											
2003-NIVA		W											
2003-NIVA	MO	W											
2004-NIVA		W											
2004-NIVA	R1	W											
CB52	1987-SIIF	W							111	0.2	20	1	
	1988-SIIF	D							111	0.1	6		
	1988-SIIF	W							111	0.1	22		
	1989-NACE	W		510	20	93			111	0.1	36		
	1989-SIIF	W											
	1990-NIVA	2G	W	340	1	169	2	6	341	0.05	58		
	1990-SIIF	2G	W						111	0.4	41	7	
	1991-NIVA	2H	W	340	1	179	1	37	341	0.05	68	5	1
	1991-SIIF	2H	W						111	0.3	35		
	1992-NIVA	2J	W	340	5	192	3		341	0.1	143		
	1993-NIVA	2K	W	340	4	212	40		341	0.1	138		
	1994-NIVA	2Z	W	340	3	300	9		341	0.05	170	64	
	1995-NIVA		W	340	3	312	19		341	0.05	220	28	
	1996-NIVA		W	340	3	332	49		341	0.05	241	31	
	1997-NIVA		W	340	3	260	116						
	1997-NIVA	AJ	W						341	0.05	221	25	10
	1998-NIVA		W	340	3	281	47	44	341	0.05	169	12	17
	1999-NIVA		W	340	3	249	52	11	341	0.05	222	7	73
	1999-NIVA	EG	W										
	2000-NIVA		W	340	3	230	65	4	341	0.05	183	22	23
	2000-NIVA	GU	W										
	2001-NIVA		W	340	3	250	66	4	341	0.05	186	7	58
	2001-NIVA	IO	W										
	2002-NIVA		W	340	3	193	29		341	0.05	162		55
	2002-NIVA	LJ	W										
	2003-NIVA		W	340	3	239	54						
	2003-NIVA	MO	W						341	0.05	147		41
	2004-NIVA		W	340	3	267	75						
	2004-NIVA	R1	W						341	0.05	215		5
CB77	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	0.0001	4			841	0.0001	18		
	2002-NILU	W							841	0.0001	12		
	2003-NILU	W							841	0.0001	12		
CB81	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	0.0001	4			841	0.0001	18		
	2002-NILU	W							841	0.0001	12		
	2003-NILU	W							841	0.0001	12		
CD	1981-NIVA	D								miss	3		
	1981-SIIF	1E	W	130	10	28			130	5	27		
	1981-SIIF	1F	W						130	10	7		
	1982-NIVA	D								miss	3		
	1982-SIIF	1F	W						130	10	18		
	1982-VETN		W	230	10	54			130	10	17		
	1983-SIIF	1F	W										

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1983-VETN		1Z	W	230	10	46							
1984-FIER		1H	W	402	1	23							
1984-SIIF		1G	W										
1984-VETN		1Z	W	230	10	66							
1985-SIIF		1G	D										
1985-VETN		1Z	W	230	10	45		3					
1986-NIVA		1H	D	312	30	56	1		312	30	20		
1987-FIER		1G	W	402	1	37							
1987-NIVA		1H	D	312	30	57		4	312	30	42		
1988-NIVA		1H	D	312	30	61	11	1	312	30	55		
1989-NIVA		1H	D	312	30	135	11	8	312	30	3		
1989-NIVA		1H	W						312	30	36		
1990-NIVA		1H	D						312	10	6		
1990-NIVA		1H	W	312	10	189	9	2	312	30	77	5	
1991-NIVA		1H	D						312	10	6		
1991-NIVA		1H	W	312	10	190	29	2	312	10	67		
1992-NIVA		1H	D						312	10	6		
1992-NIVA		1H	W	312	10	191	4		312	10	111		
1993-NIVA		1H	D						312	50	5		
1993-NIVA		1H	W	312	50	221	98		312	50	79		
1994-NIVA		1Z	D						312	50	5		
1994-NIVA		1Z	W	312	50	302	134		312	50	81		
1995-NIVA			D						312	50	6		
1995-NIVA			W	312	50	318	129		312	50	139	2	
1996-NIVA		V1	D						312	50	6		
1996-NIVA		V1	W						312	50	125		
1996-NIVA		V2	W	312	50	368	128						
1997-NIVA			W	312	50	287	90						
1997-NIVA		AH	D						312	50	6		
1997-NIVA		AH	W						312	50	107		
1998-NIVA			D						312	50	6		
1998-NIVA			W	312	50	285	101		312	50	93		
1999-NIVA			W	312	50	235	79						
1999-NIVA		EF	D						312	50	6		
1999-NIVA		EF	W						312	50	132	15	
2000-NIVA			W	312	50	227	82						
2000-NIVA		GS	D						312	50	7		
2000-NIVA		GS	W						312	50	90		
2001-NIVA			W	312	50	261	103						
2001-NIVA		IM	D						312	50	6		
2001-NIVA		IM	W						312	50	93		
2002-NIVA			W	315	50	230	126						
2002-NIVA		LH	D						315	50	6		
2002-NIVA		LH	W						315	50	110		
2003-NIVA			W	315	50	233	121						
2003-NIVA		MM	W						315	50	99		
2004-NIVA			W	315	50	249	146		315	50	142		
CDD1N	1995-NILU		W						841	2E-05	6	1	1
	1996-NILU		W	841	1E-05	4			841	1E-05	18		2
	2002-NILU		W						841	1E-05	12		2
	2003-NILU		W						841	1E-05	12		6
	2004-NILU		W						841	1E-05	12		6
CDD4X	1995-NILU		W						841	2E-05	6	3	1
	1996-NILU		W	841	2E-05	4			841	2E-05	18		1
	2002-NILU		W						841	2E-05	12		2
	2003-NILU		W						841	2E-05	12		6
	2004-NILU		W						841	2E-05	12		5
CDD6P	1995-NILU		W						841	2E-05	6		
	1996-NILU		W	841	4E-05	4			841	4E-05	18		
	2002-NILU		W						841	4E-05	12	1	

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
2003-NILU		W							841	4E-05	12		2
2004-NILU		W							841	4E-05	12		
CDD6X	1995-NILU	W							841	2E-05	6		1
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12	2	1
	2003-NILU	W							841	2E-05	12		6
	2004-NILU	W							841	2E-05	12		5
CDD9X	1995-NILU	W							841	2E-05	6	2	1
	1996-NILU	W		841	2E-05	3	1		841	2E-05	18		1
	2002-NILU	W							841	2E-05	12	2	2
	2003-NILU	W							841	2E-05	12		8
	2004-NILU	W							841	2E-05	12		6
CDDFS	1996-NILU	W		miss	4								
CDDO	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	0.0001	4			841	0.0001	18		
	2002-NILU	W							841	0.0001	12		
	2003-NILU	W							841	0.0001	12		
	2004-NILU	W							841	0.0001	12		
CDDSN	1995-NILU	W							841	2E-05	5		
	1996-NILU	W		841	1E-05	3			841	1E-05	18		3
	2002-NILU	W							841	1E-05	10		
CDDSP	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	4E-05	4			841	4E-05	18		
	2002-NILU	W							841	4E-05	11	1	
CDDST	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		
	2002-NILU	W							841	1E-05	12		
CDDSX	1995-NILU	W							841	2E-05	5		
	1996-NILU	W		841	2E-05	3			841	2E-05	18		2
	2002-NILU	W							841	2E-05	11		
CDF2N	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		1
	2002-NILU	W							841	1E-05	12		
	2003-NILU	W							841	1E-05	12		3
	2004-NILU	W							841	1E-05	12		
CDF2T	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		
	2002-NILU	W							841	1E-05	12		
	2003-NILU	W							841	1E-05	12		
	2004-NILU	W							841	1E-05	12		
CDF4X	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12	4	
	2003-NILU	W							841	2E-05	12	1	3
	2004-NILU	W							841	2E-05	12	1	
CDF6P	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	4E-05	4			841	4E-05	18	2	1
	2002-NILU	W							841	4E-05	12	3	
	2003-NILU	W							841	4E-05	12	1	2
	2004-NILU	W							841	4E-05	12		
CDF6X	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12		1
	2003-NILU	W							841	2E-05	12	1	2
	2004-NILU	W							841	2E-05	12	1	1
CDF9P	1995-NILU	W							841	2E-05	6	2	1
	1996-NILU	W		841	8E-05	4			841	8E-05	17	3	1
	2002-NILU	W							841	8E-05	12	2	2
	2003-NILU	W							841	8E-05	12	3	4

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
2004-NILU		W							841	8E-05	12	7	
CDF9X	1995-NILU	W							841	2E-05	6	3	1
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12		3
	2003-NILU	W							841	2E-05	12		7
	2004-NILU	W							841	2E-05	12		7
CDFDN	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		1
	2002-NILU	W							841	1E-05	12		
	2003-NILU	W							841	1E-05	12		1
	2004-NILU	W							841	1E-05	12		1
CDFDX	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12		1
	2003-NILU	W							841	2E-05	12	1	4
	2004-NILU	W							841	2E-05	12	1	
CDO	1995-NILU	W							841	2E-05	6		1
	1996-NILU	W		841	0.0001	4			841	0.0001	18	3	1
	2002-NILU	W							841	0.0001	11	1	
	2003-NILU	W							841	0.0001	12	1	2
	2004-NILU	W							841	0.0001	12	1	1
CDFSN	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		1
	2002-NILU	W							841	1E-05	12		
CDFSP	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	8E-05	4			841	8E-05	18	6	1
	2002-NILU	W							841	8E-05	12	4	
CDFST	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	1E-05	4			841	1E-05	18		
	2002-NILU	W							841	1E-05	12		
CDFSX	1995-NILU	W							841	2E-05	6		
	1996-NILU	W		841	2E-05	4			841	2E-05	18		1
	2002-NILU	W							841	2E-05	12	1	
CHR	1992-NIVA	W		309	0.2	8			309	0.2	44		
	1995-NIVA	W							309	0.2	56		
	1996-NIVA	W							309	0.2	65	3	
CHRTR	1995-NIVA	W							309	0.2	15		2
	1997-NIVA	W							309	0.5	36		
	1998-NIVA	W							309	0.5	39		
	1999-NIVA	W							309	0.5	34		
	2000-NIVA	W							309	0.5	39		
	2001-NIVA	W							309	0.5	42		
	2002-NIVA	W							309	0.5	43		
	2003-NIVA	W							309	0.5	56		
	2004-NIVA	W							309	0.5	58		
CO	1996-NIVA	W							999	miss	3		
	2004-NIVA	W							315	miss	7		
COR	1992-NIVA	W		309	0.2	8			309	0.2	46		
CR	1992-NIVA	W							312	10	6		
	1996-NIVA	W							999	miss	3		
	2004-NIVA	W							315	miss	7		
CU	1981-NIVA	D								miss	3		
	1982-NIVA	D								miss	3		
	1983-SIIF	1G	W						130	10	12		
	1984-SIIF	1G	W						130	10	27		
	1986-NIVA	1H	D	311	150	56			311	150	20		
	1987-FIER	1G	W	404	50	37							
	1987-NIVA	1H	D	311	150	57			311	150	42		
	1988-NIVA	1H	D	311	150	61			311	150	55		

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	count	d.lim
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1989-NIVA		1H	D	311	150	135			311	150	3		
1989-NIVA		1H	W						311	150	36		
1990-NIVA		1H	D						311	150	6		
1990-NIVA		1H	W	311	150	189			311	150	77		
1991-NIVA		1H	D						311	50	6		
1991-NIVA		1H	W	311	50	193	2		311	50	67		
1992-NIVA		1H	D						311	10	6		
1992-NIVA		1H	W	311	10	191			311	10	111		
1993-NIVA		1H	D						311	10	5		
1993-NIVA		1H	W	311	10	221			311	10	79		
1994-NIVA		1Z	D						311	10	5		
1994-NIVA		1Z	W	311	10	302			311	10	81		
1995-NIVA			D						311	10	6		
1995-NIVA			W	311	10	318			311	10	124		
1996-NIVA		V1	D						311	10	6		
1996-NIVA		V1	W						311	10	113		
1996-NIVA		V2	W	311	10	368							
1997-NIVA			W	311	5000a	287	1		311	10	6		
1997-NIVA		AH	D						311	10	96		
1997-NIVA		AH	W										
1998-NIVA			W	311	10	285			311	10	6		
1998-NIVA		CF	D						311	10	51		
1998-NIVA		CF	W										
1999-NIVA			W	311	10	235			311	10	6		
1999-NIVA		EF	D						311	10	99		
1999-NIVA		EF	W										
2000-NIVA			W	311	10	227			311	10	7		
2000-NIVA		GS	D						311	10	51		
2000-NIVA		GS	W										
2001-NIVA			W	311	10	261							
2001-NIVA		IM	D						311	10	6		
2001-NIVA		IM	W						311	10	51		
2002-NIVA			W	315	10	230			315	10	6		
2002-NIVA		LH	D						315	10	65		
2002-NIVA		LH	W										
2003-NIVA			W	315	10	233			315	10	51		
2003-NIVA		MM	W										
2004-NIVA			W	315	10	249			315	10	101		
DBA3A	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	71	48	
	1996-NIVA		W						309	0.2	65	53	
	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA		W						309	0.5	58	26	
DBP	1992-NIVA		W	309	0.2	8			309	0.2	46		
DBT	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56	20	
	2004-NIVA	R5	W						309	0.5	58	31	
DBTC1	1995-NIVA		W						309	0.2	57	14	
	1996-NIVA		W						309	0.2	65	9	
	2004-NIVA		W						309	0.5	58	14	

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
DBTC2	1995-NIVA		W						309	0.2	56		9
	1996-NIVA		W						309	0.2	62		11
	2004-NIVA		W						309	0.5	58		1
DBTC3	1995-NIVA		W						309	0.2	57		4
	1996-NIVA		W						309	0.2	65		5
	2004-NIVA		W						309	0.5	58		5
DBTIN	1997-NIVA		D						320	5	13		
	1998-NIVA		D						320	5	15		
	1999-NIVA		D						320	5	13		
	1999-NIVA		W						320	5	6	2	
	2000-NIVA		W						320	0.5	23		
	2001-GALG		W						520	0.15	11		
	2001-NIVA		W						320	0.5	16		1
	2002-EFDH		W						720	2	33	5	
	2002-NIVA		W							miss	2		2
	2003-NIVA		W						720	2	36	14	
	2004-NIVA		W						720	2	72	40	
DBTIO	1997-NIVA		W						309	0.5	34		
DDEPP	1982-VETN		W	210	50	53							
	1983-VETN	2E	W	210	50	48			211a	50	48		
	1984-VETN	2E	W	210	50	66							
	1985-VETN	2E	W	210	50	45							
	1986-NACE	ZZ	W	510	20	56							
	1987-NACE	ZZ	W	510	40	53							
	1988-NACE	ZZ	W	510	40	61							
	1989-NACE	ZZ	W	510	20	93							
	1990-NIVA		W	340	1	169			341	0.05	58		
	1991-NIVA		W	340	1	179			341	0.05	68		
	1992-NIVA		W	340	5	192	2		341	0.1	146		
	1993-NIVA		W	340	4	212	3		341	0.1	138		
	1994-NIVA	2Z	W	340	4	300			341	0.1	170	27	
	1995-NIVA		W	340	4	318	2		341	0.1	231	30	
	1996-NIVA		W	340	4	332	2		341	0.1	243	47	
	1997-NIVA		W	340	4	260	3		341	0.1	221	1	
	1998-NIVA		W	340	4	284	6						
	1998-NIVA	CH	W						341	0.1	209	4	
	1999-NIVA		W	340	4	249							
	1999-NIVA	EG	W						341	0.1	232	2	
	2000-NIVA		W	340	4	230	7						
	2000-NIVA	GU	W						341	0.1	185	6	
	2001-NIVA		W	340	4	250		1					
	2001-NIVA	IO	W						341	0.1	211	1	7
	2002-NIVA		W	340	4	249	4		341	0.1	210	5	
	2003-NIVA	MO	W	340	4	239	4		341	0.1	183	3	
	2004-NIVA		W	340	4	272	6						
	2004-NIVA	R1	W						341	0.1	241	56	
DDTEP	1983-SIIF		W						111	0.5	12		
	1984-SIIF		W						111	0.5	24		1
	1985-SIIF		W						111	0.5	27	1	5
	1986-SIIF		W						111	0.5	21		
	1987-SIIF		W						111	0.5	21	1	
	1988-SIIF		D						111	0.5	6		
	1988-SIIF		W						111	0.5	22	1	
	1989-SIIF		W						111	0.5	36	1	
	1990-SIIF		W						111	0.2	41	1	
	1991-SIIF		W						111	0.3	35		
DDTPP	1986-NACE		W	510	40	56							
	1987-NACE		W	510	40	53							
	1988-NACE		W	510	40	61							
	1989-NACE		W	510	20	93							

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
1991-NIVA		W								miss	6		
1992-NIVA		W								miss	6		4
1993-NIVA		W								miss	5		1
1994-NIVA		W								miss	5		
1995-NIVA		W								340	0.05	78	
1996-NILU		W			miss	2							
1996-NIVA		W		340	0.05	54		4	340	0.05	51		
1997-NIVA		W		340	2	32							
1997-NIVA	AJ	W								340	0.05	48	
1998-NIVA		W		340	2	37	1	8	340	0.05	74		28
1999-NIVA		W		340	2	29		4	340	0.05	99		7
2000-NIVA		W		340	2	22			340	0.05	54		6
2001-NIVA		W		340	2	46		2	340	0.05	53		11
2002-NILU		W								miss	1		
2002-NIVA		W		340	2	32		10	340	0.05	67		21
2003-NIVA		W		340	2	35		10	340	0.05	45		22
2004-NIVA		W		340	2	33			340	0.05	123		70
DPTIN	1997-NIVA	D							320	5	13	5	
	1998-NIVA	D							320	2	15		6
	1999-NIVA	D							320	5	13	12	
	1999-NIVA	W							320	5	6	6	
	2000-NIVA	W							320	0.5	23	1	1
	2001-NIVA	W							320	0.5	16		16
	2002-NIVA	W								miss	2		2
	2003-NIVA	W							720	2	36	36	
	2004-NIVA	W							720	2	72	70	
EOCL	1989-SIIF	W							605	170	5		
EPOCL	1986-NACE	W		610	800	56							
	1986-SIIF	W							605	5000	21	21	
	1987-NACE	W		610	800	53							
	1987-SIIF	W							605	40	20		
	1988-NACE	W		610	800	60							
	1988-SIIF	W							605	40	27		
	1989-NACE	W		610	800	89	1						
	1989-SIIF	W							605	40	35		
	1990-NIVA	W		615	40	117		3					
	1990-SIIF	W							605	40	41		
	1991-NIVA	W		615	40	116		12					
	1991-SIIF	W							605	130	35		
	1997-IFEN	W							607	50	6		
	1998-IFEN	W							607	1	6		
	2000-SINT	W							607	1	6		
	2001-SINT	W							607	1	6		
	2004-IFEN	W							607	1	5		
FLE	1992-NIVA	W		309	0.2	8			309	0.2	45		
	1995-NIVA	W							309	0.2	72		22
	1996-NIVA	W							309	0.2	65		6
	1997-NIVA	AL	W						309	0.5	34		
	1998-NIVA	CI	W						309	0.5	39		
	1999-NIVA	W							309	0.5	34		
	2000-NIVA	W							309	0.5	39		
	2001-NIVA	W							309	0.5	42		
	2002-NIVA	W							309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA	R5	W						309	0.5	58	18	
FLU	1992-NIVA	W		309	0.2	8			309	0.2	44		
	1995-NIVA	W							309	0.2	72		
	1996-NIVA	W							309	0.2	65		
	1997-NIVA	AL	W						309	0.2	36		
	1998-NIVA	CI	W						309	0.2	39		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1999-NIVA		EK	W						309	0.2	34		
2000-NIVA			W						309	0.2	39		
2001-NIVA			W						309	0.2	42		
2002-NIVA			W						309	0.2	43		3
2003-NIVA		MQ	W						309	0.2	56		
2004-NIVA		R5	W						309	0.2	58		
HBCDA	2001-NILU		W		miss	4			miss	5		2	
HBCDB	2001-NILU		W		miss	4	4		miss	5		5	
HBCDG	2001-NILU		W		miss	5	4		miss	4		4	
HCB	1983-SIIF		W						111	0.5	12		
	1983-VETN	2Z	W	210	10	48			211a	10	48		
	1984-SIIF		W						111	0.2	24		1
	1984-VETN	2Z	W	210	10	66			111	0.2	30	6	2
	1985-SIIF		W										
	1985-VETN	2Z	W	210	10	45	4						
	1986-NACE	2Z	W	510	10	56							
	1986-SIIF	2Z	W						111	0.2	21	3	
	1987-NACE	2Z	W	510	40	53			111	0.2	21	4	
	1988-NACE	2Z	W	510	40	61							
	1988-SIIF	2Z	D						111	0.2	6		
	1988-SIIF	2Z	W						111	0.2	22	2	
	1989-NACE	2Z	W	510	20	93							
	1989-SIIF	2Z	W						111	0.05	36		
	1990-NIVA		W	340	1	169	2		341	0.05	58		
	1990-SIIF	2Z	W						111	0.05	41	3	
	1991-NIVA		W	340	1	179	4	13	341	0.05	68	5	1
	1991-SIIF	2Z	W						111	0.1	35		
	1992-NIVA		W	340	5	189	3		341	0.1	146		
	1993-NIVA		W	340	4	212	31		341	0.1	138		
	1994-NIVA	2Z	W	340	3	300	24	1	341	0.05	170	37	
	1995-NIVA		W	340	3	317	37		341	0.05	231	32	
	1996-NIVA		W	340	3	332	52		341	0.05	243	37	
	1997-NIVA		W	340	2	260	39						
	1997-NIVA	AJ	W						341	0.05	221	7	
	1998-NIVA		W	340	2	284	48	13	341	0.05	209	68	2
	1999-NIVA		W	340	2	249	18						
	1999-NIVA	EG	W						341	0.05	232	19	8
	2000-NIVA		W	340	2	230	40						
	2000-NIVA	GU	W						341	0.05	186	43	1
	2001-NIVA		W	340	2	250	36	1	341	0.05	211	36	3
	2002-NIVA		W	340	2	249	39		341	0.05	210	29	2
	2003-NIVA		W	340	2	239	31						
	2003-NIVA	MO	W						341	0.05	174	18	
	2004-NIVA		W	340	2	271	42						
	2004-NIVA	R1	W						341	0.05	241	109	
HCHA	1990-NIVA		W	340	1	168			341	0.05	58		
	1991-NIVA		W	340	1	179	2	111	341	0.05	68	5	10
	1992-NIVA		W	340	5	192	3		341	0.1	146		
	1993-NIVA		W	340	4	212	45	22	341	0.1	138		
	1994-NIVA	2Z	W	340	3	296	32	3	341	0.05	170	85	
	1995-NIVA		W	340	3	318	45		341	0.05	231	100	
	1996-NIVA		W	340	3	332	111		341	0.05	237	100	
	1997-NIVA		W	340	0.5	260	2	10	341	0.05	221	20	11
	1998-NIVA		W	340	0.5	284	8	208	341	0.05	208	26	121
	1999-NIVA		W	340	0.5	249	17	78	341	0.05	232	23	151
	2000-NIVA		W	340	0.5	230	31	62	341	0.05	186	42	84
	2001-NIVA		W	340	0.5	250	25	50	341	0.05	211	20	184
	2002-NIVA		W	340	0.5	249	23	149	341	0.05	210		121
	2003-NIVA		W	340	0.5	239	4	201	341	0.05	183		99

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
2004-NIVA		W		340	0.5	270	13	192	341	0.05	238	2	9
HCHG	1986-NACE	W		510	30	56	1		111	3	21		
	1986-SIIF	W							111	5	21		1
	1987-NACE	W		510	40	53							
	1987-SIIF	W											
	1988-NACE	W		510	40	61							
	1989-NACE	W		510	20	93							
	1989-SIIF	W							111	50	36		
	1990-NIVA	W		340	1	169	1	9	341	0.05	58		
	1990-SIIF	W							111	0.1	41		
	1991-NIVA	W		340	1	179	3	18	341	0.05	68	5	1
	1991-SIIF	W							111	0.3	35		
	1992-NIVA	W		340	5	192	3		341	0.1	146		
	1993-NIVA	W		340	4	212	42	17	341	0.1	138		
	1994-NIVA	2Z	W	340	3	300	24	1	341	0.05	170	46	
	1995-NIVA	W		340	3	313	31		341	0.05	219	29	
	1996-NIVA	W		340	3	330	68		341	0.05	226	8	
	1997-NIVA	W		340	2	260	47						
	1997-NIVA	AJ	W						341	0.05	221	3	9
	1998-NIVA	W		340	2	284	25	63	341	0.05	209	10	23
	1999-NIVA	W		340	2	249	52	3	341	0.05	232	19	62
	2000-NIVA	W		340	2	230	65	29	341	0.05	186	27	10
	2001-NIVA	W		340	2	250	96	20	341	0.05	211	21	160
	2002-NIVA	W		340	2	249	147	13	341	0.05	210		83
	2003-NIVA	W		340	2	239	96	85	341	0.05	181		102
	2004-NIVA	W		340	2	271	137	19	341	0.05	241		8
HG	1981-NIVA	D							miss		3		
	1981-SIIF	1E	W	120	10	15		1	120	10	35		
	1982-NIVA	D							miss		3		
	1982-SIIF	1E	W						120	10	18		
	1982-VETN	W		220	10	51			220	10	54		
	1983-SIIF	1E	W						120	10	17		
	1983-VETN	1Z	W						220	10	48		
	1984-FIER	1G	W						401	10	39		
	1984-SIIF	1G	W						120	10	27	6	
	1984-VETN	1Z	W						220	10	66		
	1985-SIIF	1G	D						120	10	30		
	1985-VETN	1Z	W						220	10	90		
	1986-NIVA	1H	D						310	10	74		
	1987-FIER	1G	W						401	10	38		
	1987-NIVA	1H	D						310	10	98	14	
	1988-NIVA	1H	D						310	10	116		
	1989-NIVA	1H	D						310	100	137		
	1989-NIVA	1H	W						310	10	36	5	
	1990-NIVA	1H	D						310	10	6		
	1990-NIVA	1H	W						310	10	266		
	1991-NIVA	1H	D						310	100	6		
	1991-NIVA	1H	W						310	100a	264	126	
	1992-NIVA	1H	D						310	100	6		
	1992-NIVA	1H	W						310	100a	303	122	
	1993-NIVA	1H	D						310	5	5		
	1993-NIVA	1H	W						310	5	300		
	1994-NIVA	1Z	D						310	5	5		
	1994-NIVA	1Z	W						310	5	381		
	1995-NIVA	D							310	5	6		
	1995-NIVA	W							310	5	442	1	
	1996-NIVA	V1	D						310	5	6		
	1996-NIVA	V1	W						310	5	481		
	1997-NIVA	AH	D						310	5	6		
	1997-NIVA	AH	W						310	5	383		

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
1998-NIVA		CF	D						310	5	6		
1998-NIVA		CF	W						310	5	381		
1999-NIVA			W										
1999-NIVA		EF	D						310	5	6		
1999-NIVA		EF	W						310	5	386		
2000-NIVA		GS	D						310	5	7		
2000-NIVA		GS	W						310	5	330		
2001-NIVA		IM	D						310	5	6		
2001-NIVA		IM	W						310	5	356		
2002-NIVA		LH	D						310	5	6		
2002-NIVA		LH	W						310	5	366		
2003-NIVA		MM	W						310	5	347	2	
2004-NIVA			W						310	5	420		
ICDP	1992-NIVA		W		309	0.2	8		309	0.2	46		
	1995-NIVA		W						309	0.2	72		29
	1996-NIVA		W						309	0.2	65		23
	1997-NIVA		W						309	0.5	36		
	1998-NIVA	CI	W						309	0.5	37	2	
	1999-NIVA	EK	W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA	R5	W						309	0.5	58	7	
MBTIN	1997-NIVA		D						320	5	13	4	
	1998-NIVA		D						320	5	15		
	1999-NIVA		D						320	5	13		
	1999-NIVA		W						320	5	6	6	
	2000-NIVA		W						320	0.5	23		
	2001-GALG		W						520	0.2	11		
	2001-NIVA		W						320	0.5	16		5
	2002-EFDH		W						720	0.8	33		15
	2002-NIVA		W							miss	2	2	
	2003-NIVA		W						720	0.8	36	1	31
	2004-NIVA		W						720	0.8	73	50	1
MN	1984-SIIF		W						132	40	27		
	1985-SIIF		D						132	40	35		
	2004-NIVA		W						315	miss	7		
MPTIN	1997-NIVA		D						320	5	13	5	
	1998-NIVA		D						320	2	15		6
	1999-NIVA		D						320	5	13		
	1999-NIVA		W						320	5	6	6	
	2000-NIVA		W						320	0.5	23	3	
	2001-NIVA		W						320	0.5	16		15
	2002-EFDH		W						720	4	1		
	2002-NIVA		W							miss	2	2	
	2003-NIVA		W						720	4	36	36	
	2004-NIVA		W						720	4	71	71	
NAP	1992-NIVA		W		309	0.2	8		309	0.2	46		
	1995-NIVA		W						309	0.2	70		21
	1996-NIVA		W						309	0.2	61		11
	1997-NIVA		W						309	0.2	34		1
	1998-NIVA	CI	W						309	0.2	37		
	1999-NIVA		W						309	0.2	34		1
	2000-NIVA		W						309	0.2	37		7
	2001-NIVA		W						309	0.2	41		4
	2002-NIVA		W						309	0.2	42		19
	2003-NIVA	MQ	W						309	0.2	55		40
	2004-NIVA	R5	W						309	0.2	58		18
NAP1M	1992-NIVA		W		309	0.2	8		309	0.2	46		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
	1995-NIVA		W						309	0.2	15		13
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	37		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42	9	
	2003-NIVA		W						309	0.5	55		1
NAP2M	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	15		13
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	37		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42	9	
	2003-NIVA		W						309	0.5	55		4
NAPC1	1995-NIVA		W						309	0.2	55		6
	1996-NIVA		W						309	0.2	61		
	2004-NIVA		W						309	2	58	23	
NAPC2	1995-NIVA		W						309	0.2	57		6
	1996-NIVA		W						309	0.2	60		
	2004-NIVA		W						309	2	58	14	
NAPC3	1995-NIVA		W						309	0.2	57		5
	1996-NIVA		W						309	0.2	60		
	2004-NIVA		W						309	2	58	3	5
NAPD2	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42		
	2003-NIVA		W						309	0.5	55		
NAPD3	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42		
	2003-NIVA		W						309	0.5	38		
NAPDI	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	15		6
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	41		
	2002-NIVA		W						309	0.5	42		
	2003-NIVA		W						309	0.5	55		
NAPT2	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
NAPT3	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
NAPT4	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
NAPTM	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	15	11	
	1997-NIVA		W						309	0.5	34		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56	9	
NI	1983-SIIF	1G	W						130	20	12		
	1992-NIVA		W						312	10	6		
	1996-NIVA		W						999	miss	3		
	2004-NIVA		W						315	miss	7		
OCS	1990-NIVA		W	340	2	169	31	24	341	0.05	58	1	
	1991-NIVA		W	340	2	179	14	81	341	0.05	62	5	8
	1992-NIVA		W	340	5	192	3		341	0.1	146		
	1993-NIVA		W	340	4	212	51	16	341	0.1	138		
	1994-NIVA		W	340	3	300	39	22	341	0.05	170	101	
	1995-NIVA		W	340	3	318	44		341	0.05	231	108	
	1996-NIVA		W	340	3	332	287		341	0.05	243	114	
	1997-NIVA		W	340	2	260	100		341	0.05	221	30	14
	1998-NIVA		W	340	2	277	132	101	341	0.05	209	188	1
	1999-NIVA		W	340	2	249	148	2	341	0.05	232	86	26
	2000-NIVA		W	340	2	230	140	21	341	0.05	186	103	59
	2001-NIVA		W	340	2	250	189	2	341	0.05	211	94	69
	2002-NIVA		W	340	2	218	183		341	0.05	201	96	6
	2003-NIVA		W	340	2	217	178		341	0.05	180	79	
	2004-NIVA		W	340	2	265	218		341	0.05	241	71	1
PA	1992-NIVA		W	309	0.2	8			309	0.2	45		
	1995-NIVA		W						309	0.2	72		
	1996-NIVA		W						309	0.2	65		
	1997-NIVA	AL	W						309	0.2	36		
	1998-NIVA	CI	W						309	0.2	39		
	1999-NIVA	EK	W						309	0.2	34		
	2000-NIVA		W						309	0.2	39		
	2001-NIVA		W						309	0.2	42		
	2002-NIVA		W						309	0.2	43		
	2003-NIVA	MQ	W						309	0.2	56		
	2004-NIVA	R5	W						309	0.2	58		
PAC1	1995-NIVA		W						309	0.2	57	1	
	1996-NIVA		W						309	0.2	65		
	2004-NIVA		W						309	2	58	8	
PAC2	1995-NIVA		W						309	0.2	56		
	1996-NIVA		W						309	0.2	65		2
	2004-NIVA		W						309	2	58		
PAC3	2004-NIVA		W						309	2	58	5	
PADM1	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
	Year		calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
PADM2	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		1
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
PAH	1987-NIVA		W	309	0.02	1							
PAM1	1992-NIVA		W	309	0.2	8			309	0.2	45		
	1995-NIVA		W						309	0.2	15		2
	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	55		9
PAM2	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA		W						309	0.5	34		
	2000-NIVA		W						309	0.5	38		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA		W						309	0.5	56		
PB	1981-NIVA		D							miss	3		
	1982-NIVA		D							miss	3		
	1983-SIIF	1G	W						130	20	12		
	1984-SIIF	1G	W						130	20	27		2
	1985-SIIF	1G	D						130	20	35		
	1986-NIVA	1Z	D	312	150	56	4		312	150	20		
	1987-FIER	1G	W	403	10	37	1						
	1987-NIVA	1Z	D	312	150	57		12	312	150	42		
	1988-NIVA	1Z	D	312	150	61	17	3	312	150	55		
	1989-NIVA	1Z	D	312	150	135	9	9	312	150	3		
	1989-NIVA	1Z	W						312	150	36		
	1990-NIVA	1Z	D						312	50	6		
	1990-NIVA	1Z	W	312	50	187	3	1	312	150	77	3	
	1991-NIVA	1Z	D						312	50	6		
	1991-NIVA	1Z	W	312	50	193	14		312	50	67		
	1992-NIVA	1Z	D						312	50	6		
	1992-NIVA	1Z	W	312	50	191	119		312	50	111	2	
	1993-NIVA	1H	D						312	30	5		
	1993-NIVA	1H	W	312	30	221	40		312	30	79		
	1994-NIVA	1Z	D						312	30	5		
	1994-NIVA	1Z	W	312	30	302	3		312	30	81		
	1995-NIVA		D						312	30	6		
	1995-NIVA		W	312	30	318	162	30	312	30	124		
	1996-NIVA	V1	D						312	30	6		
	1996-NIVA	V1	W						312	30	110		
	1996-NIVA	V2	W	312	30	368		109					
	1997-NIVA		D						312	40	6		
	1997-NIVA		W	312	40	287	10	28	312	40	92		
	1998-NIVA		W	312	40	285	126	2					
	1998-NIVA	CF	D						312	40	6		
	1998-NIVA	CF	W						312	40	90		
	1999-NIVA		W	312	40	235	118	11					
	1999-NIVA	EF	D						312	40	6		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1999-NIVA		EF	W	312	40	227	67	4	312	40	129	10	
2000-NIVA			W						312	40	7		
2000-NIVA		GS	D						312	40	87		
2000-NIVA		GS	W						312	40	6		
2001-NIVA			W	312	40	261	156	6	312	40	90		
2001-NIVA		IM	D						315	40	6		
2001-NIVA		IM	W						315	40	107		
2002-NIVA			D						315	40	96		
2002-NIVA			W	315	40	230	164		315	40	139		
2003-NIVA		MM	W	315	40	233	179	1	315	40	96		
2004-NIVA			W	315	40	249	182		315	40	6		
PBB15	1996-NILU		W	miss		4		3					
	2001-NILU		W	miss		6		6	miss		6		3
	2002-NILU		W						843	0.01	2		
PBB49	2001-NILU		W	miss		6		1	miss		6		3
	2002-NILU		W						843	0.01	2		
PBB52	1996-NILU		W	miss		4							
	2001-NILU		W	miss		6		1	miss		6		5
	2002-NILU		W						843	0.01	2		
PCB	1981-SIIF	2D	W	110	10	27			110	10	35		
	1982-SIIF	2D	W						111	5	17		
	1982-VETN		W	210	50	53			211	50	54		
	1983-SIIF	2E	W						111	5	14		
	1983-VETN	2E	W						211	50	48		
	1983-VETN	2Z	W	210	50	48							
	1984-SIIF	2E	W						111	5	24		
	1984-VETN	2E	W						211	50	66		
	1984-VETN	2Z	W	210	50	66							
	1985-SIIF	2E	W						111	5	32		6
	1985-VETN	2E	W						211	50	90		1
	1985-VETN	2Z	W	210	50	45							
	1986-NACE	2Z	W	511a	40a	56			511	20	56		
	1986-SIIF	2E	W						111	5	21		
	1987-NACE	2Z	W	510	40	53			511	20	54		
	1987-NIVA		W	340	0.1	2							
	1987-SIIF	2E	W						111	5	21		
	1988-NACE	2Z	W	510	40	61			511	20	13		
	1988-SIIF	2E	D						111	5	6		
	1988-SIIF	2E	W						111	5	22		4
	1989-NACE	2Z	W	510	20	93			511	20	17		
	1989-SIIF	2E	W						111	5	36		6
	1990-SIIF	2E	W						111	5	41		
	1991-SIIF	2E	W						111	5	35		
PCC26	1996-NILU		W						842	0.001	6		
PCC32	1996-NILU		W						842	0.003	6		4
PCC50	1996-NILU		W						842	0.001	6		
PCC62	1996-NILU		W						842	0.025	6		6
PCDD	1995-NILU		W						841	2E-05	6		
	1996-NILU		W	841	0.0001	4			841	0.0001	18		
	2002-NILU		W						841	0.0001	12		
PCDF	1995-NILU		W						841	2E-05	6		
	1996-NILU		W	841	0.0001	4			841	0.0001	18		
	2002-NILU		W						841	0.0001	11		
PER	1992-NIVA		W	309	0.2	8			309	0.2	46		
	1995-NIVA		W						309	0.2	72		32
	1996-NIVA		W						309	0.2	65		40
	1997-NIVA		W						309	0.5	36		
	1998-NIVA		W						309	0.5	39		
	1999-NIVA	EK	W						309	0.5	34		

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
	Year		calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
	2000-NIVA		W						309	0.5	39		
	2001-NIVA		W						309	0.5	42		
	2002-NIVA		W						309	0.5	43		
	2003-NIVA	MQ	W						309	0.5	56		
	2004-NIVA		W						309	0.5	55	24	
PYR	1992-NIVA		W	309	0.2	8			309	0.2	44		
	1995-NIVA		W						309	0.2	72		4
	1996-NIVA		W						309	0.2	65		1
	1997-NIVA	AL	W						309	0.2	36		
	1998-NIVA	CI	W						309	0.2	39		
	1999-NIVA	EK	W						309	0.2	34		
	2000-NIVA		W						309	0.2	39		
	2001-NIVA		W						309	0.2	42		
	2002-NIVA		W						309	0.2	43		3
	2003-NIVA	MQ	W						309	0.2	56		
	2004-NIVA	R5	W						309	0.2	58		
QCB	1990-NIVA		W	340	2	169	33	39	341	0.05	58		
	1991-NIVA		W	340	2	178	13	97	341	0.05	63	5	13
	1992-NIVA		W	340	5	192	3		341	0.1	131		
	1993-NIVA		W	340	4	212	52	24	341	0.1	138		
	1994-NIVA		W	340	3	299	38	23	341	0.05	170	98	
	1995-NIVA		W	340	3	318	45		341	0.05	231	108	
	1996-NIVA		W	340	3	332	306		341	0.05	243	109	
	1997-NIVA		W	340	2	260	79		341	0.05	221	27	10
	1998-NIVA		W	340	2	284	121	101	341	0.05	209	177	1
	1999-NIVA		W	340	2	242	185	2	341	0.05	232	88	14
	2000-NIVA		W	340	2	230	198	1	341	0.05	186	123	1
	2001-NIVA		W	340	2	232	216	1	341	0.05	211	95	63
	2002-NIVA		W	340	2	248	235		341	0.05	210	99	4
	2003-NIVA		W	340	2	186	182		341	0.05	183	79	
	2004-NIVA		W	340	2	229	227		341	0.05	241	215	
SCCP	2001-NILU		W	miss				4	miss				3
SE	1982-VETN		W	240	10	46			240	10	54		
TBA	2001-NILU		W	miss				6	miss				6
	2002-NILU		W						843	0.35	1		
TBBPA	2001-NILU		W	miss				6	miss				6
TBTIN	1997-NIVA	D							320	5	13		
	1998-NIVA	D							320	5	15		
	1999-NIVA	D							320	5	13		
	1999-NIVA	W							320	5	6		
	2000-NIVA	W							320	0.5	23		
	2001-GALG	W							520	0.12	11		
	2001-NIVA	W							320	0.5	16		
	2002-EFDH	W							720	0.2	32		
	2002-NIVA	W							miss				2
	2003-NIVA	W							720	0.2	36	1	2
	2004-NIVA	W							720	0.2	72		1
TCDD	1995-NILU		W						841	2E-05	6	1	
	1996-NILU		W	841	1E-05	4			841	1E-05	18		
	2002-NILU		W						841	1E-05	12		
	2003-NILU		W						841	1E-05	12		2
	2004-NILU		W						841	1E-05	12		
TDEPP	1991-NIVA	W		340	1	138		1	341	0.05	68		
	1992-NIVA	W		340	5	191	3		341	0.1	146		
	1993-NIVA	W		340	4	212	24	3	341	0.1	138		
	1994-NIVA	2Z	W	340	3	300	17	5	341	0.05	170	47	
	1995-NIVA	W		340	3	318	36		341	0.05	228	51	
	1996-NIVA	W		340	3	332	23		341	0.05	243	16	
	1997-NIVA	W		340	3	260	23		341	0.05	221	11	
	1997-NIVA	AJ	W										

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			+basis	method	limit	value	below	above	method	limit	value	below	above
				code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
1998-NIVA		W		340	3	278	19	26	341	0.05	209	1	44
1998-NIVA	CH	W		340	3	249	6	1	341	0.05	232	2	71
1999-NIVA		W		340	3	230	35	4	341	0.05	185	11	67
1999-NIVA	EG	W		340	3	250	24	3	341	0.05	210	1	101
2000-NIVA		W		340	3	248	24	3	341	0.05	210		124
2000-NIVA	GU	W		340	3	239	18	9	341	0.05	183		106
2001-NIVA		W		340	3	272	30		341	0.05	241		138
2002-NIVA		W											
2003-NIVA		W											
2004-NIVA		W											
TPTIN	1997-NIVA	D							320	5	13		
	1998-NIVA	D							320	10	15		
	1999-NIVA	D							320	5	13		
	1999-NIVA	W							320	5	6	4	
	2000-NIVA	W							320	0.5	23		
	2001-GALG	W							520	0.1	11		1
	2001-NIVA	W							320	0.5	16		9
	2002-EFDH	W							720	2	24	13	
	2002-NIVA	W								miss	2		2
	2003-NIVA	W							720	2	36	35	
	2004-NIVA	W							720	2	64	61	
V	1996-NIVA	W							999	miss	3		
ZN	1981-NIVA	D								miss	3		
	1982-NIVA	D								miss	3		
	1983-SIIF	1G	W						131	400	12		
	1984-SIIF	1G	W						132	400	27		
	1985-SIIF	1G	D						132	400	35		
	1986-NIVA	1H	D	311	3000	56			311	3000	20		
	1987-FIER	1G	W	405	20	37							
	1987-NIVA	1H	D	311	3000	57			311	3000	42		
	1988-NIVA	1H	D	311	3000	61			311	3000	55		
	1989-NIVA	1H	D	311	3000	135	1		311	3000	3		
	1989-NIVA	1H	W						311	3000	36		
	1990-NIVA	1H	D						311	3000	6		
	1990-NIVA	1H	W	311	3000	189			311	3000	77		
	1991-NIVA	1H	D						311	1000	6		
	1991-NIVA	1H	W	311	1000	193			311	1000	67		
	1992-NIVA	1H	D						311	1000	6		
	1992-NIVA	1H	W	311	1000	191			311	1000	111		
	1993-NIVA	1H	D						311	1000	5		
	1993-NIVA	1H	W	311	1000	221			311	1000	79		
	1994-NIVA	1Z	D						311	1000	5		
	1994-NIVA	1Z	W	311	1000	302			311	1000	81		
	1995-NIVA		D						311	1000	6		
	1995-NIVA		W	311	1000	318			311	1000	142		
	1996-NIVA	V1	D						311	1000	6		
	1996-NIVA	V1	W						311	1000	131		
	1996-NIVA	V2	W	311	1000	368							
	1997-NIVA		W	311	1000	287							
	1997-NIVA	AH	D						311	1000	6		
	1997-NIVA	AH	W						311	1000	110		
	1998-NIVA		W	311	1000	285							
	1998-NIVA	CF	D						311	1000	6		
	1998-NIVA	CF	W						311	1000	51		
	1999-NIVA		W	311	1000	235							
	1999-NIVA	EF	D						311	1000	6		
	1999-NIVA	EF	W						311	1000	99		
	2000-NIVA		W	311	1000	227							
	2000-NIVA	GS	D						311	1000	7		
	2000-NIVA	GS	W						311	1000	51		

JAMP National Comments 2004 - Norway

Tissue				Fish liver					Fish fillet, Shrimp tail, Mussel, Other				
Contamin.	Mon.	Lab.	Inter-	Analys	Detect	Total	Count	N (<)	Analys	Detect	Total	Count	N (<)
Year			calibr.	method	limit	value	below	above	method	limit	value	below	above
			+basis	code	(ppb)	count	d.lim	d.lim	code	(ppb)	count	d.lim	d.lim
2001-NIVA			W	311	1000	261			311	1000	6		
2001-NIVA		IM	D						311	1000	51		
2001-NIVA		IM	W						315	1000	6		
2002-NIVA			W	315	1000	230			315	1000	65		
2002-NIVA		LI	D						315	1000	51		
2002-NIVA		LI	W						315	1000	101		
2003-NIVA			W	315	1000	233							
2003-NIVA		MM	W	315	1000	249							
2004-NIVA			W										
Sum of counts				83176	13439	3100			81391	6609	7113		

a(7) > ambiguous value (Maximum value displayed)

## **Appendix D**

### **Participation in intercalibration exercises**



## Appendix D1

**Participation in intercalibration exercises other than QUASIMEME**

---

**Sea water:**

- 4H ICES/JMG Fifth Round Intercalibration on Trace Metals in Sea Water - Section 4, analysis for Hg - 1983 - (5/TM/SW:4).
- 4I JMG Sixth Intercalibration on Trace Metals in Estuarine Waters - 1986 - (6/TM/SW).
- 4Z Intercalibration exercise for SIIF/SERI (Cd) and NIVA/IAMK (IAMK=Chalmers Inst., Göteborg) - 1985.

**Seabed sediment:**

- 7E ICES, First Intercalibration Exercise on Trace metals in Marine Sediments - 1984 - (1/TM/MS).
- 8B ICES/OSPAR, First Intercomparison Exercise on Organochlorines (individual chlorobiphenyl congeners) in Marine Sediments - Phase 1, analysis of standard solutions - 1989 - (1/OC/MS:1).
- 8C ICES/OSPAR, First Intercomparison Exercise on Organochlorines (individual chlorobiphenyl congeners) in Marine Sediments - Phase 2, analysis of standard solutions - 1991 - (1/OC/MS:2).
- 8B ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 1 - (analysis of standard solutions) - 1989 - (1/OC/MS-1).
- 8C ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 2 - 1990 - (1/OC/MS-2).
- 8D ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3a (1/OC/MS-3a) 1991.
- 8E ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3b - (1/OC/MS-3b) 1992.
- 8F ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 4 - (1/OC/MS-4) 1993.

**Marine biota:**

- 1E ICES, Fifth Intercalibration Exercise on Trace Metals in Biological Tissues - 1978 - (5/TM/BT).
- 1F ICES, Sixth Intercalibration Exercise on Trace Metals (Cadmium and Lead only) in Biological Tissues - 1979 - (6/TM/BT).
- 1G ICES, Seventh Intercalibration Exercise on Trace Metals in Biological Tissues - Part A - 1983 - (7/TM/BT).
- 1H ICES, Seventh Intercalibration Exercise on Trace Metals in Biological Tissues - Part B - 1985 - (7/TM/BT) (preliminary report 1987).
- 1Z VETN Interlabculation exercise with VETN and SIIF 1983, mercury and cadmium in cod filet and liver.

- 1Z NIVA Interlabcalibration exercise with VETN, NACE and NIVA 1986 (Hg, Cd, Cu, Pb and Zn in 6 samples).
- 2D ICES Fourth Intercalibration Exercise on Organochlorines (mainly PCBs) in Biological Tissues (Sample No.5) - 1979 - (4/OC/BT).
- 2E ICES Fifth Intercalibration Exercise on Organochlorines (PCBs only) in Biological Tissues - 1982 - (5/OC/BT).
- 2G ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 1 - (analysis of standard solutions) - 1989 - (7/OC/BT-1).
- 2H ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 2 - 1990 - (7/OC/BT-2).
- 2I ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3a - (7/OC/BT-3a) 1991.
- 2J ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3b - (7/OC/BT-3b) 1992.
- 2K ICES/IOC/OSPAR Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 4 - (7/OC/BT-4) 1993.
- 2Z VETN Interlabcalibration exercise with VETN among others, 1983, PCB and HCB in cod liver.
- 2Z NACE Interlabcalibration exercise with NACE, VETNand SIIFF 1986 (PCB (all labs), DDE, OCS, HCB and DCB (NACE and VETN).

Appendix D2  
**Participation in QUASIMEME intercalibration exercises**

---

IC	Code	Year	No.	Group	Matrix
QM	QOR002BT	1993	80	BT-2	CB's in standard solution and biota - Fish oil
V1	QTM028BT	1996	280	BT-1	Trace metals in cod muscle and cod liver
V2	QTM029BT	1996	280	BT-1	Trace metals in cod muscle and cod liver
AJ	QOR054BT	1997	347	BT-2	Chlorobiphenyls and organochlorine pesticides in biota
AL	QPH008BT	1997	348	BT-4	PAHs in biota
AH	QTM036BT	1997	346	BT-1	Metals in biota
CI	QPH010BT	1998	394	BT-4	Polyaromatic hydrocarbons in biota
CH	QOR059BT	1998	393	BT-2	Chlorobiphenyls and organochlorine pesticides in Biota
CF	QTM042BT	1998	392	BT-1	Trace metals in Biota
EF	QTM046BT	1999	433	BT-1	Trace metals in biota
EG	QOR062BT	1999	434	BT-2	Chlorobiphenyls and organochlorine pesticides in biota
EK	QPH012BT	1999	435	BT-4	Polyaromatic hydrocarbons in biota
GU	QOR066BT	2000	473	BT-2	Chlorobiphenyls and organochlorine pesticides in biota
GS	QTM049BT	2000	472	BT-1	Trace metals in biota
IO	QOR070BT	2001	510	BT-2	Chlorobiphenyls and organochlorine pesticides in biota
IM	QTM053BT	2001	509	BT-1	Trace metals in biota
	QTM057BT	2003	549	BT-1	Trace metals in biota
	QOR074BT	2003	550	BT-2	Chlorobiphenyls and organochlorine pesticides in biota

---



## Appendix E

# Overview of localities and sample count for biota 1981-2004

**Station positions are shown on maps in Appendix G**

jmpco:JAMP area code (J99 = unclassified)

jmpst: station code

stnam: station code

Lon: Longitude

Lat: Latitude

icear: ICES area

speci: species code (English, Norwegian (Latin))

MYTI EDU - blue mussel, blåskjell (*Mytilus edulis*)

NUCE LAP - dogwhelk, purpursnegl (*Nucella lapillus*)

BROS BRO - tusk, brosme (*Brosme brosme*)

CHIM MON - rat fish, havmus (*Chimaera monstrosa*)

GADU MOR - Atlantic cod, torsk (*Gadus morhua*)

LEPI WHI - megrim, glassvar (*Lepidorhombus whiffiagonis*)

LIMA LIM - dab, sandflyndre (*Limanda limanda*)

MICR KIT - lemon sole, lomre (*Microstomus kitt*)

MOLV MOL - ling, lange (*Molva molva*)

PAND BOR - shrimp, reker (*Pandalus borealis*)

PLAT FLE - flounder, skrubbe (*Platichthys flesus*)

PLEU PLA - plaice, rødspette (*Pleuronectes platessa*)

tissu: tissue:

SB - soft body

LI - liver

MU - fillet

TM - tail muscle



## STATIONS AND SAMPLE COUNT FOR BIOTA

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04		
J26	01A	Sponvika	59° 5.10	11° 13.90	47G13	MYTI EDU	SB		3																								
J26	02A	Fugleskjær	59° 6.90	10° 59.00	47G09	MYTI EDU	SB		3																								
J26	03A	Tisler	58° 58.80	10° 57.50	46G07	MYTI EDU	SB		2																								
J26	301	Akershuskaia	59° 54.23	10° 45.47	48G07	MYTI EDU	SB																						2				
J26	302	Ormøya	59° 52.69	10° 45.46	48G07	MYTI EDU	SB																						2				
J26	303	Malmøya	59° 51.78	10° 45.95	48G07	MYTI EDU	SB																						2				
J26	304	Gåsøya	59° 51.11	10° 35.51	48G04	MYTI EDU	SB																						3				
J26	305	Lysaker	59° 54.36	10° 38.60	48G04	MYTI EDU	SB																					2					
J26	306	Håøya	59° 42.69	10° 33.35	48G05	MYTI EDU	SB																						3				
J26	30A	Gressholmen	59° 53.20	10° 42.658	48G07	MYTI EDU	SB																						3	3			
J26	30B	Oslo City area	59° 49.00	10° 33.00	48G05	GADU MOR	BI																						27	23			
J26	30B	Oslo City area	59° 49.00	10° 33.00	48G05	GADU MOR	BL																						20	30			
J26	30B	Oslo City area	59° 49.00	10° 33.00	48G05	GADU MOR	LI																						25	25			
J26	30B	Oslo City area	59° 49.00	10° 33.00	48G05	GADU MOR	MU																						25	25			
J26	30F	Oslo City area	59° 47.00	10° 34.00	48G05	PLEU PLA	LI																						2	5			
J26	30F	Oslo City area	59° 47.00	10° 34.00	48G05	PLEU PLA	MU																					2	5				
J26	30G	Spro	59° 45.80	10° 34.50	48G05	PAND BOR	TM																						1				
J26	30H	Storegrunn	59° 48.50	10° 33.50	48G05	PAND BOR	TM																						22				
J26	30X	West of Nesodden	59° 48.50	10° 36.00	48G05	GADU MOR	LI																					22					
J26	30X	West of Nesodden	59° 48.50	10° 36.00	48G05	GADU MOR	MU																						2				
J26	40C	Steilene	59° 49.00	10° 33.00	48G05	PAND BOR	TM																						1				
J26	31A	Solbergstrand	59° 36.90	10° 39.40	48G06	MYTI EDU	SB	2																					6	3			
J26	31B	Solbergstrand	59° 36.90	10° 39.40	48G06	GADU MOR	LI	10	27																					3	3		
J26	31B	Solbergstrand	59° 36.90	10° 39.40	48G06	GADU MOR	MU	10	27																					1			
J26	31B	Solbergstrand	59° 36.90	10° 39.40	48G06	PLAT FLE	LI		8																					1			
J26	31B	Solbergstrand	59° 36.90	10° 39.40	48G06	PLAT FLE	MU		8																						1		
J26	31C	Solbergstrand	59° 36.90	10° 39.40	48G06	PAND BOR	TM																							2			
J26	32A	Rødtangen	59° 31.50	10° 25.60	48G06	MYTI EDU	SB	1	3																					3			
J26	33B	Sande (east side)	59° 31.70	10° 21.00	48G06	PLAT FLE	LI																						25	1			
J26	33B	Sande (east side)	59° 31.70	10° 21.00	48G06	PLAT FLE	MU																						25	1			
J26	33C	Sande	59° 31.70	10° 21.00	48G06	PAND BOR	TM																						1				
J26	33X	Sande (west side)	59° 31.70	10° 20.40	48G06	PLAT FLE	LI																						3				
J26	33X	Sande (west side)	59° 31.70	10° 20.40	48G06	PLAT FLE	MU																						3				
J26	35A	Mølen	59° 29.20	10° 30.10	47G04	MYTI EDU	SB	1	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3			
J26	35C	Holmestrand-Mølen	59° 29.20	10° 30.10	47G04	PAND BOR	TM		1																				1	2			
J26	35C	Holmestrand-Mølen	59° 29.20	10° 30.10	47G04	PAND BOR	XX																							1			
J26	36B	Færder	59° 2.00	10° 32.00	47G06	GADU MOR	BI																							21	25		

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04					
J26	36B	Færder	59° 2.00	10° 32.00	47G06	GADU MOR	BL																				20	25	25	23	25					
J26	36B	Færder	59° 2.00	10° 32.00	47G06	GADU MOR	LI	10	27	23	24	14	25	25	25	25	24	25	25	25	25	26	25	25	25	23	28	30	25	25						
J26	36B	Færder	59° 2.00	10° 32.00	47G06	GADU MOR	MU	10	27	23	24	14	25	25	26	26	29	30	30	30	30	30	30	30	30	30	30	30	30	30	30					
J26	36F	Færder area	59° 4.00	10° 23.00	47G06	LIMA LIM	BI																				11	9	20							
J26	36F	Færder area	59° 4.00	10° 23.00	47G06	LIMA LIM	BL																			20	9	20								
J26	36F	Færder area	59° 4.00	10° 23.00	47G06	LIMA LIM	LI																			5	5	5	5	30						
J26	36F	Færder area	59° 4.00	10° 23.00	47G06	LIMA LIM	MU																			5	5	5	5	30						
J26	36A	Færder	59° 1.60	10° 31.70	47G06	MYTI EDU	SB	1	5	3	3	3	3	3	3	3	3	3	3	3	3	3	5	3	3	3	3	3	3	3	3					
J26	36G	Færder	59° 1.60	10° 31.70	47G06	NUCE LAP	SB																			1	1	1	1	1						
J26	73A	Lyngholmen	59° 2.60	10° 18.10	47G03	MYTI EDU	SB										3																			
J26	74A	Oddneskjær	58° 57.30	9° 52.10	46F97	MYTI EDU	SB																													
J26	71A	Bjørkøya (Risøyodd.)	59° 1.40	9° 45.40	47F99	MYTI EDU	SB	1	3	3	3	2	3	3	3	3	3	3	3	3	3	3	4	3	3	3	3	3	3	3	3					
J26	71G	Fugløyskjær	58° 58.95	9° 48.50	46F97	NUCE LAP	SB																				1	1	1	2						
J99	76A	Risøy	58° 43.60	9° 17.00	46F92	MYTI EDU	SB										3	3	3	3																
J99	76A	Risøy	58° 43.60	9° 17.00	46F92	NUCE LAP	SB																													
J99	76G	Risøy	58° 43.60	9° 17.00	46F92	NUCE LAP	SB																													
J99	77A	Flostafljord	58° 31.50	8° 56.90	46F89	MYTI EDU	SB										3	3																		
J99	77B	Borøy area	58° 33.00	9° 1.00	45F93	GADU MOR	LI										14	25																		
J99	77B	Borøy area	58° 33.00	9° 1.00	45F93	GADU MOR	MU										17	30																		
J99	77B	Borøy area	58° 33.00	9° 1.00	45F93	LIMA LIM	LI																													
J99	77C	Borøy area	58° 29.00	9° 10.00	45F91	PAND BOR	TM										2																			
J99	79A	Gjerdsvoldsøyen east	58° 24.80	8° 45.30	45F87	MYTI EDU	SB										3	3																		
J99	13A	Langøysund	57° 59.80	7° 34.60	45F74	MYTI EDU	SB										1	4																		
J99	14A	Aavigen	58° 2.20	7° 13.20	45F73	MYTI EDU	SB										3	4																		
J99	15A	Gåsøy (Ullerø)	58° 3.07	6° 53.16	45F69	MYTI EDU	SB										4	4																		
J99	15B	Ullerø area	58° 3.00	6° 43.00	45F69	GADU MOR	BI																													
J99	15B	Ullerø area	58° 3.00	6° 43.00	45F69	GADU MOR	BL																													
J99	15B	Ullerø area	58° 3.00	6° 43.00	45F69	GADU MOR	LI																													
J99	15B	Ullerø area	58° 3.00	6° 43.00	45F69	GADU MOR	MU																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	LIMA LIM	BI																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	LIMA LIM	BL																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	LIMA LIM	LI																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	LIMA LIM	MU																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	MICR KIT	LI																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	MICR KIT	MU																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	PLEU PLA	LI																													
J99	15F	Ullerø area	58° 3.00	6° 43.00	45F69	PLEU PLA	MU																													
J99	15G	Gåøy	58° 3.10	6° 43.30	45F69	NUCE LAP	SB										3	3																		
J63	51A	Byrkjenes	60° 5.10	6° 33.10	49F66	MYTI EDU	SB																													

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04
J63	52A	Eitrheimsneset	60° 5.80	6° 32.20	49F66	MYTI EDU	SB									3	3	3	3	2	3	3	3	3	3	6	3	3	3	3	3
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GADU MOR	BI																		15	28	24	25	25	25	
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GADU MOR	BL																		15	30	25	25	25	25	
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GADU MOR	LI																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GADU MOR	MU																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GLYP CYN	LI																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	GLYP CYN	MU																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	PLAT FLE	BI																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	PLAT FLE	BL																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	PLAT FLE	LI																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	PLAT FLE	MU																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	SALM TRU	LI																								
J63	53B	Inner Sørfjord	60° 10.00	6° 34.00	49F65	SALM TRU	MU																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	BROS BRO	LI																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	BROS BRO	MU																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	CHIM MON	LI																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	CHIM MON	MU																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	MOLV MOL	LI																								
J63	53D	Digraneset	60° 11.00	6° 34.05	49F65	MOLV MOL	MU																								
J63	56A	Kvalnes	60° 13.231	6° 36.120	49F65	MYTI EDU	SB																								
J63	56A1	Kvalnes, north	60° 13.514	6° 36.255	49F65	MYTI EDU	SB																								
J63	56A2	Kjeken	60° 20.329	6° 39.274	49F64	MYTI EDU	SB																								
J63	56A3	Sekse	60° 15.683	6° 37.396	49F65	MYTI EDU	SB																								
J63	56A4	Rosstadnes	60° 17.219	6° 37.428	49F65	MYTI EDU	SB																								
J63	56A5	Lofthus, south	60° 19.351	6° 39.121	49F65	MYTI EDU	SB																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	BROS BRO	LI																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	BROS BRO	MU																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	CHIM MON	LI																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	CHIM MON	MU																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	MOLV MOL	LI																								
J63	56D	Kvalnes	60° 15.00	6° 36.00	49F65	MOLV MOL	MU																								
J63	57A	Krossanes	60° 23.225	6° 41.353	49F67	MYTI EDU	SB																								
J63	57A1	Urdheim	60° 22.346	6° 40.689	49F67	MYTI EDU	SB																								
J63	57A2	Ernes	60° 21.188	6° 39.738	49F64	MYTI EDU	SB																								
J62	63A	Ranaskjær	60° 25.10	6° 24.50	49F64	MYTI EDU	SB																								
J62	65A	Vikingneset	60° 14.50	6° 9.60	49F62	MYTI EDU	SB																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	GADU MOR	BI																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	GADU MOR	BL																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	GADU MOR	LI																								

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	GADU MOR	MU								22	26	23	16	24	9	14	22	30	40	30	30	30	30	30	30	
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	LEPI WHI	LI								19	1	26	30	5	5	3	5	5	5	5	5	5	30	30	28	12
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	LEPI WHI	MU								19	1	26	30	5	5	3	5	5	5	5	5	5	30	30	28	12
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	LIMA LIM	LI																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	LIMA LIM	MU																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	PLAT FLE	BI																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	PLAT FLE	BL																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	PLAT FLE	LI																								
J62	67B	Strandebarm	60° 16.00	6° 2.00	49F62	PLAT FLE	MU																								
J62	69A	Lille Terøy	59° 58.79	5° 45.35	48F57	MYTI EDU	SB																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	BROS BRO	LI																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	BROS BRO	MU																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	CHIM MON	LI																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	CHIM MON	MU																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	MOLV MOL	LI																								
J99	21D	Åkrafjord	59° 48.00	6° 11.00	48F62	MOLV MOL	MU																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	LEPI WHI	LI																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	LEPI WHI	MU																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	LIMA LIM	LI																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	LIMA LIM	MU																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	PLAT FLE	BI																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	PLAT FLE	BL																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	PLAT FLE	LI																								
J99	21F	Åkrefjord	59° 45.00	6° 7.00	48F62	PLAT FLE	MU																								
J99	220G	Smørstakk	59° 15.21	5° 21.14	47F55	NUCE LAP	SB																								
J99	221G	Stangeland	59° 16.02	5° 19.70	47F52	NUCE LAP	SB																								
J99	222A	Kopervik harbour	59° 17.02	5° 18.94	47F52	MYTI EDU	SB																								
J99	224G	Heggjelen	59° 25.00	5° 13.90	47F51	NUCE LAP	SB																								
J99	226A	Karmsund bridge (west)	59° 22.68	5° 17.70	47F51	NUCE LAP	SB																								
J99	226G	Karmsund bridge (east)	59° 22.68	5° 17.91	47F51	NUCE LAP	SB																								
J99	227G	Melandholmen	59° 20.04	5° 18.90	47F51	NUCE LAP	SB																								
J99	227X	Høievarde	59° 19.43	5° 19.11	47F52	MYTI EDU	SB																								
J99	22G	Espevær vest	58° 34.75	5° 8.90	46F53	NUCE LAP	SB																								
J99	22A	Espevær, west	59° 35.20	5° 8.50	48F53	MYTI EDU	SB																								
J99	22C	Bømlofjord	59° 34.00	5° 11.00	48F53	PAND BOR	TM																								
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	LIMA LIM	LI																								
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	LIMA LIM	MU																								
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	MICR KIT	LI																								
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	MICR KIT	MU																								

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04		
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	PLEU PLA	LI																		5	5	5						
J99	22F	Borøyfjorden	59° 43.00	5° 21.00	48F55	PLEU PLA	MU																		5	5	5						
J99	221A	Stangeland	59° 16.62	5° 19.70	47F52	MYTI EDU	SB																		3	3							
J99	226X	Karmsund bridge (east)	59° 22.68	5° 17.91	47F51	MYTI EDU	SB																		1	3	3						
J99	227A	Melandholmen	59° 20.04	5° 18.90	47F51	MYTI EDU	SB																		3	3				1			
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	GADU MOR	BI																		22	23	24	23	25	25	25		
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	GADU MOR	BL																		25	25	25	24	25	25	25		
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	GADU MOR	LI																		25	25	25	25	25	25	25		
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	GADU MOR	MU																		30	30	30	30	30	30	30		
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	MICR KIT	LI																		1	4							
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	MICR KIT	MU																		1	4					3		
J99	23A	Austvik	59° 52.20	5° 6.60	48F51	MYTI EDU	SB																		1	1							
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	PLAT FLE	LI																		1	1							
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	PLAT FLE	MU																		3	3							
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	PLEU PLA	LI																		3	3							
J99	23B	Karihavet area	59° 54.00	5° 8.00	48F51	PLEU PLA	MU																		1	1							
J99	24A	Vardøy	60° 10.20	5° 0.80	49F52	MYTI EDU	SB																		3	3					3		
J65	80A	Østmarknes	63° 27.50	10° 27.50	55G04	MYTI EDU	SB																										
J65	81A	Biologisk Stasjon	63° 26.50	10° 21.40	55G04	MYTI EDU	SB																										
J65	82A	Flakk	63° 27.00	10° 12.60	55G01	MYTI EDU	SB																			1	2						
J65	83A	Frøsetskjær	63° 25.50	10° 7.80	55G01	MYTI EDU	SB																			1	2						
J65	84A	Trossavika	63° 20.80	9° 57.80	55F97	MYTI EDU	SB																			3	2	2	3	3		3	
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	GADU MOR	LI																			1	1						
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	GADU MOR	MU																		13	1	1	1	5				
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MELA AEG	LI																		13	10	1	1	5				
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MELA AEG	MU																		14	1	4						
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MERL MNG	LI																		1	1	5						
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MERL MNG	MU																		1	7							
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MICR KIT	LI																		3								
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	MICR KIT	MU																		3								
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	POLL POL	LI																		1	1	8						
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	POLL POL	MU																		16	1	8						
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	POLL VIR	LI																		4								
J65	84B	Trossavika	63° 20.80	9° 57.80	55F97	POLL VIR	MU																		4								
J99	82G	Flakk	63° 27.00	10° 12.60	55G01	NUCE LAP	SB																		1	1							
J99	87G	Ingdalsbukt	63° 27.80	9° 54.80	55F97	NUCE LAP	SB																		1	1	1	1	1	1			
J65	85A	Geitstrand	63° 21.90	9° 56.30	55F97	MYTI EDU	SB																		1	2	1						
J65	86A	Geitnes	63° 26.60	9° 59.20	55F97	MYTI EDU	SB																		2	2							
J65	87A	Ingdalsbukt	63° 27.80	9° 54.80	55F97	MYTI EDU	SB																								1		

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	
J65	88A	Rødberg	63° 29.20	10° 0.00	55G01	MYTI EDU	SB				1	1																				
J99	24G	Vardøy	60° 10.20	5° 0.80	49F52	NUCE LAP	SB																						1			
J99	25G	Hinnøy	61° 22.20	4° 52.80	51F47	NUCE LAP	SB																						2			
J99	25A	Hinnøy	61° 22.20	4° 52.80	51F47	MYTI EDU	SB																						3			
J99	26A	Hamnen	61° 52.70	5° 13.60	52F51	MYTI EDU	SB																						3			
J99	26G	Hamnen	61° 52.60	5° 13.30	52F51	NUCE LAP	SB																						1			
J99	27A	Grinden	62° 12.20	5° 25.40	53F55	MYTI EDU	SB													2									3			
J99	27G	Røydeskjær	62° 11.00	5° 44.30	53F58	NUCE LAP	SB																						2			
J99	28A	Eiksundet	62° 15.00	5° 51.60	53F58	MYTI EDU	SB													6	3								3			
J99	28G	Grønevikholt.,Ulsteinvik	62° 14.80	5° 53.00	53F58	NUCE LAP	SB																						2			
J99	91A	Nerdvika	63° 21.20	8° 9.60	55F81	MYTI EDU	SB																									
J99	92A	Stokken	64° 2.21	10° 1.10	57G03	MYTI EDU	SB																									
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	GADU MOR	LI																							25		
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	GADU MOR	MU																							30		
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	LIMA LIM	LI																									
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	LIMA LIM	MU																									
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	PLEU PLA	LI																									
J99	92B	Stokken area	64° 10.28	9° 53.24	57F98	PLEU PLA	MU																									
J99	93A	Sætervik	64° 23.68	10° 29.00	57G04	MYTI EDU	SB													7	3									3		
J99	94A	Landfast	65° 38.40	12° 0.50	60G23	MYTI EDU	SB													3	3									3		
J99	94G	Steinskjær, Landfast	65° 38.40	12° 0.10	60G23	NUCE LAP	SB																							2		
J99	95A	Flatskjær	66° 42.60	13° 15.80	62G32	MYTI EDU	SB													3	3									3		
J99	95G	Flatskjær	66° 42.40	13° 14.30	62G32	NUCE LAP	SB																							2		
J99	96A	Breiviken	66° 17.60	12° 50.50	61G28	MYTI EDU	SB													6	3									3		
J99	97A	Klakholmen	67° 39.90	14° 44.60	64G49	MYTI EDU	SB													4	3									3		
J99	97G	Varnesodden	67° 48.10	14° 45.10	64G48	NUCE LAP	SB																							2		
J99	98A	Svolvær området	68° 14.942	14° 39.752	65G45	MYTI EDU	SB													4	3									3		
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	GADU MOR	BI																							14	22	
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	GADU MOR	BL																							5	25	
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	GADU MOR	LI													25	29	25	24	26	25	25	25	25	25	21	25	
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	GADU MOR	MU													30	29	30	29	30	30	30	30	30	30	25	30	
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	LIMA LIM	LI													4												
J99	98B	Lille Molla	68° 12.00	14° 48.00	65G48	LIMA LIM	MU													4												
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	GLYP CYN	LI														1											
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	GLYP CYN	MU														1											
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	LIMA LIM	LI														1	1	5									
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	LIMA LIM	MU														1	1	5									
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	MICR KIT	LI													1	1											
J99	98F	Lille Molla	68° 12.00	14° 48.00	65G48	MICR KIT	MU													1	1											

JAMP National Comments 2004 - Norway

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04								
J99	10F	Skogerøy	69° 55.00	29° 51.00	68H97	PLEU PLA	BI																			15	25												
J99	10F	Skogerøy	69° 55.00	29° 51.00	68H97	PLEU PLA	BL																			11	24												
J99	10F	Skogerøy	69° 55.00	29° 51.00	68H97	PLEU PLA	LI																			5	4	18	30	30	22	24							
J99	10F	Skogerøy	69° 55.00	29° 51.00	68H97	PLEU PLA	MU																			5	4	18	30	30	22	24							
J99	11A	Sildkroneset,Bøkfj	69° 47.02	30° 11.10	68J02	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	11X	Brashavn	69° 53.92	29° 44.65	68H97	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	11G	Brashavn	69° 53.92	29° 44.65	68H97	NUCE LAP	SB																								1	1	2						
J26	I001	Sponvikskansen	59° 5.40	11° 12.50	47G13	MYTI EDU	SB																			3	3												
J26	I011	Kråkenebbet	59° 6.10	11° 17.30	47G13	MYTI EDU	SB																			3	3												
J26	I021	Kjøkø,south	59° 7.79	10° 57.11	47G09	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I022	West Damholmen	59° 6.20	10° 57.90	47G09	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I023	Singlekalven, south	59° 5.70	11° 8.20	47G13	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I024	Kirkøy, north west	59° 4.90	10° 59.20	47G09	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I301	Akershuskaia	59° 54.23	10° 45.47	48G07	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I304	Gåsøya	59° 51.11	10° 35.51	48G04	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I306	Håøya	59° 42.69	10° 33.35	48G05	MYTI EDU	SB																			3	3	3	3	3	3	3							
J26	I307	Ramtonholmen	59° 44.70	10° 31.40	48G05	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I711	Steinholmen	59° 3.15	9° 40.70	47F99	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I712	Gjemesholmen	59° 2.75	9° 42.47	47F99	MYTI EDU	SB																			3	4	3	3	3	3	3							
J99	I713	Strømtangen	59° 3.22	9° 41.500	47F99	MYTI EDU	SB																			3	4	3	3	3	3	3							
J99	I131	Lastad	58° 3.30	7° 42.40	45F79	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I131	Lastad	58° 3.30	7° 42.40	45F79	NUCE LAP	SB																								1								
J99	I131G	Lastad	58° 3.30	7° 42.40	45F79	NUCE LAP	SB																							1	1	2							
J99	I132	Fiskåtangen	58° 7.75	7° 58.60	45F79	MYTI EDU	SB																			4	4	3	3	3	3	3							
J99	I133	Odderø,west	58° 7.90	8° 0.15	45F83	MYTI EDU	SB																			4	4	3	3	3	3	3							
J99	I201	Ekkjegrunn (G1)	59° 38.65	6° 21.38	48F66	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I205	Bølsnes (G5)	59° 35.50	6° 18.30	48F63	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I241	Nordnes	60° 24.10	5° 18.20	49F51	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I242	Valheimneset	60° 23.70	5° 16.10	49F51	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I243	Hegreneset	60° 24.90	5° 18.50	49F51	MYTI EDU	SB																			3	3	3	3	3	3	3							
J99	I911	Horvika	62° 44.10	8° 31.40	54F85	MYTI EDU	SB																																
J99	I913	Fjøseid	62° 48.59	8° 16.48	54F82	MYTI EDU	SB																																
J99	I912	Honnhammer	62° 51.20	8° 9.70	54F81	MYTI EDU	SB																				3	3	3	3	3	3	3						
J99	I914	Flåøya, southeast	62° 45.354	8° 26.697	54F85	MYTI EDU	SB																																
J99	I915	Flåøya, northeast	62° 45.483	8° 26.391	54F85	MYTI EDU	SB																																
J99	I916	Sundalsfjord, Hydro kai	62° 41.108	8° 33.126	54F85	MYTI EDU	SB																																
J65	I080	Østmerknes	63° 27.50	10° 27.50	55G04	MYTI EDU	SB																																
J99	I964	Toraneskaien	66° 19.30	14° 7.97	61G42	MYTI EDU	SB																																
J99	I965	Moholmen (B5)	66° 18.72	14° 7.55	61G42	MYTI EDU	SB																																

jmpco	jmpst	stnam	lat	lon	icear	speci	tissu	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04
J99	I962	Koksverktomta (B2)	66° 19.57	14° 8.38	61G42	MYTI EDU	SB															3	3	2	3						
J99	I969	Bjørnbærviken (B9)	66° 16.79	14° 2.13	61G42	MYTI EDU	SB															3	3	3	3	3	3	3	3	3	
J99	R096	Breiviken, Tomma	66° 17.60	12° 50.50	61G28	MYTI EDU	SB															3	3								
J26	A3*	Svartskjær	58° 58.90	9° 49.90	46F97	MYTI EDU	SB	1																							



## **Appendix F**

# **Overview of localities and sample count for sediment 1987-2004**

**Station positions are shown on maps in Appendix G**

jmpco:JAMP area code (J99 = unclassified)  
jmpst: station code  
stnam: station code  
Lon: Longitude  
Lat: Latitude  
icear: ICES area



## STATIONS AND SAMPLE COUNT FOR SEDIMENT

jmpco	jmpst	stnam	lat	lon	icear	1986	1987	1990	1992	1994	1996	1997	2004
J26	30S	Steilene	59° 49.1	10° 33.8	48G05	8		34			5		
J26	35S	Holmestrand-Mølen	59° 28.96	10° 31.74	47G04	6					5		
J26	35S	Holmestrand-Mølen	59° 30	10° 35.7	47G04	2		3					
J26	36S	Færder area	59° 0.4	10° 41.6	47G09	2		40					
J26	36S	Færder area	59° 1.55	10° 32.99	47G09	6							
J26	36S	Færder area	59° 2.5	10° 46.6	47G09					56			
J99	77S	Arendal area	58° 24.2	9° 1.8	45F91		43				29		
J99	15S	Lista area	58° 1	6° 34.3	45F66		32				5		
J63	52S	Tyssedal	60° 6.9	6° 32.9	49F66		3				5		
J63	52S	Tyssedal	60° 6.92	6° 32.6	49F66						3		
J63	56S	Kvalnes	60° 13.7	6° 35.6	49F65		29				5		
J63	56S	Kvalnes	60° 13.72	6° 35.6	49F65						3		
J63	57S	Krossanes	60° 23.1	6° 40.7	49F67		3				5		
J63	57S	Krossanes	60° 23.1	6° 40.1	49F67						3		
J62	63S	Ranaskjær	60° 23.34	6° 26.7	49F64						1		
J62	63S	Ranaskjær	60° 23.34	6° 27.1	49F64						2		
J62	63S	Ranaskjær	60° 23.6	6° 27.1	49F64		3				5		
J62	67S	Strandebar	60° 13.12	6° 4.6	49F62						3		
J62	67S	Strandebar	60° 13.5	6° 5.1	49F62		28				28		
J62	69S	Kvinnheradsfjorden	60° 1.3	5° 56.1	49F59		3				5	3	
J99	22S	Børmlø area	59° 25.9	4° 50.2	47F47		29				5	3	
J99	24S	Sotra	60° 15.1	4° 33.3	49F45		3				3		
J65	82S	Flakk	63° 27.05	10° 11.08	55G01	8							
J65	82S	Flakk	63° 27.5	10° 11.8	55G01						3		
J65	89S	Thamshavn	63° 19.08	9° 52.05	55F98	4		3					
J65	89S	Thamshavn	63° 19.7	9° 52.5	55F98						3		
J65	84S	Trossavika	63° 21.7	9° 57.4	55F97	8		3			3		
J65	90S	Outer Orkdalsfjord	63° 27.3	10° 3.06	55G01						3		
J65	90S	Outer Orkdalsfjord	63° 27.4	10° 3	55G01	8		30					
J99	27S	Stadtlandet (east of)	62° 9.3	5° 21.3	53F56			30			3		
J99	93S	Raudøya (norteast of)	64° 22.7	10° 27.8	57G04			30			3		
J99	95S	Rodø (east of)	66° 41.8	13° 10	62G32						3		
J99	95S	Rodø (east of)	66° 41.8	13° 9.9	62G32			31					
J99	98S	Skrova (south of)	68° 7	14° 41	65G49			30			3		
J99	99S	Lundøy (north of)	68° 5.8	15° 10.1	65G53			30			3		

jmpco	jmpst	stnam	lat	lon	icear	1986	1987	1990	1992	1994	1996	1997	2004
J99	41S	Vågsfjorden	68° 56.25	17° 5.24	66G71					34			
J99	42S	Malangen	69° 30.38	18° 6.77	68G83					3			
J99	43S	Kvænangen	70° 3.31	21° 7.94	69H13					34			
J99	44S	Sørøysund	70° 25.91	22° 31.83	69H24					3			
J99	45S	Revbotn	70° 42.86	24° 26.65	70H45					34			
J99	46S	Porsangerfjorden	70° 52.93	26° 11.89	70H61					28			
J99	47S	Laksfjord	70° 54.96	26° 55.11	70H67					3			
J99	48S	Tanafjord	70° 52.54	28° 38.53	70H84					33			
J99	49S	Syltefjord	70° 33.94	30° 19.91	70J03					3			
J99	10S	Varangfjorden	69° 56.07	30° 6.7	68J01					29			

## **Appendix G Map of stations**

**Station positions 1981-2004  
(cf. Appendix H and Appendix L)**



## Appendix G (cont.) Map of stations


### NOTES

For a few stations the geolocation has varied somewhat in order to collect sufficient material (e.g., st. 36B and 98A) or investigate local geographical variations (e.g., in the inner Oslofjord and Sørfjord). Hence, the same station name may appear more than once on a map.


The letter A following the station identification number indicates that blue mussel was sampled. The letter B indicates sampling for cod and the letter F indicates sampling for flatfish. This system for fish is not consistent for some older stations (30, 33, 52 and 67) where only the letter B is used indicating that either cod or flatfish or both were sampled. An encircled dot indicates a mussel, shrimp or fish station. The letter G indicates sampling for dogwhelks and S indicates sampling for sediment. An encircled dot indicates the position for sampling mussels, shrimp or fish. A square and pentagon symbol indicates the position for sampling dogwhelks or sediment, respectively. A triangle indicates the position of a town or city.

The letter "I" preceding the station identification number indicates an INDEX station for determining a "pollution" index. The letter R indicates a station for evaluating a "reference" index. Only blue mussel is used for these indices. The indices are based on a selection of JAMP and INDEX stations (cf. Green *et al.* 2001).


The maps are generated using ArcView GIS version 3.3.




JAMP stations Norway. Numbers indicate map reference




MAP 1




MAP 2




MAP 3




MAP 4




MAP 5




MAP 6




MAP 7



MAP 8




MAP 10




MAP 11





MAP 12




MAP 13




MAP 14










MAP 17




MAP 18







MAP 23



MAP 24

## **Appendix H**

### **Overview of materials and analyses 2004**

**Station positions are shown on maps in Appendix G**



**Appendix H1.** Sampling and analyses for 2003, L-liver, F-fillet. (See Appendix H2 for descriptions of codes for analysis (M0, M1, M3, M4, M5, C2, C4, A1, G1), fish (P, F, D, L, M, C) and counts). Analytical overview for liver (-L) or fillet (-F) tissue is distinguished.

JAMP area	STATION	WATER	SEDIMENT	MUSSEL/ OTHER				FISH																		
								FLAT- (P,F,D,M)				COD- (C)														
				M0	M1	C1	A1	G1	M3	C2	A1	M3	C2	-L	M4	C2	A1	-L	M4	C2	A1	-F	M5	C2	A1	
<b>OSLOFJORD AREA CENTRAL, Oslofjord proper</b>																										
26	30A Gressholmen	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
26	30B Oslo city Area / Håøya	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B	
26	31A Solbergstrand	1	.	.	.	.	.	3	3	.	.	.	.	.	F-L	5B	5B	.	.	.	.	.	.	.		
26	33B Sande, east side	.	.	.	.	.	.	.	.	.	.	.	.	.	F-F	5B	5B	.	.	.	.	.	.	.		
26	35A Mølen	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
26	36A Færder	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B	
26	36B Færder area	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B	
20	36F Færder area	.	.	.	.	.	.	.	.	.	.	.	.	.	D-L	5B	5B	.	.	.	.	.	.	.		
26	<b>OSLOFJORD AREA WEST, outer Sandefjord-Langesundsfjord</b>																									
26	71A Bjørkøya	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
<b>ARENDAL AREA</b>																										
26	76A Risøy	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.			
<b>LISTA AREA</b>																										
15A	Ullerø area	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B	
15B	Ullerø area	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
15F	Ullerø area	.	.	.	.	.	.	.	.	.	.	.	.	.	D-L	5B	5B	.	.	.	.	.	.	.		
15F	Ullerø area	.	.	.	.	.	.	.	.	.	.	.	.	.	D-F	5B	5B	.	.	.	.	.	.	.		
<b>BØMLO-SOTRA AREA</b>																										
21F	Kyrping (Åkrafjord 2000)	.	.	.	.	.	.	.	.	.	.	.	.	.	D-L	5B	5B	.	.	.	.	.	.			
21F	Kyrping (Åkrafjord 2000)	.	.	.	.	.	.	.	.	.	.	.	.	.	D-F	5B	5B	.	.	.	.	.	.			
21F	Kyrping (Åkrafjord 2000)	.	.	.	.	.	.	.	.	.	.	.	.	.	F-L	5B	5B	.	.	.	.	.	.			
21F	Kyrping (Åkrafjord 2000)	.	.	.	.	.	.	.	.	.	.	.	.	.	F-F	5B	5B	.	.	.	.	.	.			
22A	Espevær, west	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B	
23B	Karihavet	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	C-L	25	25	.	C-F	25	5B
<b>HARDANGERFJORDEN</b>																										
62	69A Lille Terøy	1	.	.	.	.	.	3	3	.	.	.	.	.	ML	5B	5B	.	C-L	25	25	.	C-F	25	5B	
62	67B Strandebarm	.	.	.	.	.	.	.	.	.	.	.	.	.	MF	5B	5B	.	.	.	.	.	.	.		
62	67B Strandebarm	.	.	.	.	.	.	.	.	.	.	.	.	.	F-L	5B	5B	.	.	.	.	.	.	.		
62	67B Strandebarm	.	.	.	.	.	.	.	.	.	.	.	.	.	F-F	5B	5B	.	.	.	.	.	.	.		
62	65A Vikingneset	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
62	63A Ranaskjær	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
<b>SØRFJORDEN</b>																										
63	52A Eitrheimsneset	1	.	.	.	.	.	3	3	.	.	.	.	.	P-L	5B	5B	.	C-L	25	25	.	C-F	25	5B	
63	53B Inner Sørfjord	.	.	.	.	.	.	.	.	.	.	.	.	.	P-F	5B	5B	.	C-F	25	5B	.	.	.	.	
63	56A Kvalnes	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
63	57A Krossanes	1	.	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		

## Appendix H1 (cont.)

JAMP area	STATION	WATER	SEDIMENT	MUSSEL/	OTHER	FISH											
						FLAT- (P,F,D,M)		COD- (C)									
		M0	M1	C4	A1	G1	M3	C2	A1	-L	M4	C2	A1	-L	M4	C2	A1
<b>LOFOTEN AREA</b>																	
98A	Husvågen (1997)	1	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.
98B	Lille Molla	.	.	.	.	.	.	.	.	.	.	.	.	C-L	25	25	.
98F	Lille Molla	.	.	.	.	.	.	.	.	.	.	.	.	C-F	25	5B	.
		.	.	.	.	.	.	.	.	P-L	5B	5B	.	.	.	.	.
		.	.	.	.	.	.	.	.	P-F	5B	5B	.	.	.	.	.
<b>VARANGER PENINSULA AREA</b>																	
10A	Skalneset	1	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.
10B	Varangerfjorden	.	.	.	.	.	.	.	.	.	.	.	.	C-L	25	25	.
		.	.	.	.	.	.	.	.	.	.	.	.	C-F	25	5B	.
		.	.	.	.	.	.	.	.	P-L	5B	5B	.	.	.	.	.
		.	.	.	.	.	.	.	.	P-F	5B	5B	.	.	.	.	.
10F	Varangerfjorden	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
11X	Brashavn (1997)	1	.	.	.	.	3	3	.	.	.	.	.	.	.	.	.

**Appendix H2:** Key to analysis codes and sample counts used in Appendix H1.**ANALYSIS CODES:**

Code	Analyses
M0	suspended matter
M1	Hg, Cd, Cu, Pb, Zn, Li (normalising element) total organic carbon (TOC)
M3	Hg, Cd, Cu, Pb, Zn
M4	Cd Cu Pb Zn (for fish liver)
M5	Hg (for fish fillet)
C1	CB-28,-52,-101,-105,-118,-138,-153,-156,-180, 209, 5-CB, OCS, a+gHCH, HCB, DDE, DDD, EPOCI (optional), dry weight percent
C2	CB-28,-52,-101,-105,-118,-138,-153,-156,-180, 209, 5-CB, OCS, a+gHCH, HCB, DDE, DDD, EPOCI (optional), fat and dry weight percent
A1	PAH
G1	Sediment core geological dating

**SAMPLE COUNT CODES:**

Medium	Count	Explanation
SEAWATER	1	sample for suspended matter determination
SEDIMENT	17	17 samples for metal analyses; two cores each with samples from 0-1, 1-2, 2-4, 4-6, 6-10, 10-15, 15-20 cm and deepest 5 cm slice plus one core with sample from 0-1 cm.
	4	4 samples for PCB or PAH analyses; two each cores with samples from 0-1cm and deepest 5cm slice.
	3	3 samples for metal analyses; three cores each with samples from 0-1cm.
MUSSEL	3/6	3 size groups (2-3, 3-4, 4-5 cm) each a bulk of ca.50 individuals and/or 1 size group (3-4 or 4-5 cm), 3 replicate samples each a bulk of 20 individuals.
	1/2	1 size group (2-3 or 3-4 cm), 2 replicate samples each a bulk of 50 individuals.
SHRIMP	2	2 samples of 100 individuals (edible size)
FISH		The number of individual fish or bulk samples of fish (-B) for analyses is shown. Bulk samples of fish consist of 5 fish. The five longest fish make up one bulk sample, the next five longest fish make up the another bulk sample and so on. The letter following the number indicates the fish type: D=dab, F=flounder, L=lemon sole, M=megrim, P=plaice, W=witch, C=cod, T=tusk, R=Rat fish, and N=Ling.



# Appendix I

## Temporal trend analyses of contaminants and biomarkers in biota 1981-2004

Sorted by contaminant, species and area/station:

Cadmium (Cd)
Mercury (Hg)
Lead (Pb)
Sum PCB-7 or CB_S7 (CB: 28+52+101+118+138+153+180)
DDEPP (ppDDE)
$\gamma$ HCH (HCHG)
HCB
BAP (benzo[a]pyrene)
<b>PK-Σn or PK_S</b> (sum carcinogen PAHs, cf. Appendix B)
<b>P-Σn or P_S</b> (sum of PAHs, dicyclic "PAHs" not included, cf. Appendix B)
<b>TBT</b> (Tributyltin)
VDSI (measurement of imposex)
<b>OH-pyrene or PYR10</b> (Pyrene metabolite)
<b>ALA-D</b> ( $\delta$ -amino levulinic acid dehydrase inhibition)
<b>EROD</b> (Cytochrome P4501A-activity)
<b>MYTI EDU</b> - Blue Mussel ( <i>Mytilus edulis</i> )
JAMP-stations
"Index"-stations
<b>GADU MOR</b> - Atlantic cod ( <i>Gadus morhua</i> )
<b>LEPI WHI</b> - Megrim ( <i>Lepidorhombus whiffiagonis</i> )
<b>LIMA LIM</b> - Dab ( <i>Limanda limanda</i> )
<b>PLAT FLE</b> - Flounder ( <i>Platichthys flesus</i> )
(s) - Small fish
(l) - Large fish
Tsu -tissue:
SB - Soft body tissue
LI - Liver tissue
MU - Muscle tissue
BL - Blood
BI - Bile
<b>OC</b> Overconcentration expressed as quotient of median of last year and "high background" ("?" missing background value)
<b>TRD</b> trend
D- Significant linear trend, downward
U- Significant linear trend, upward
-- No significant trend
-? No significant linear trend, systematic non-linear trend can not be tested because of insufficient data (<6 years)
-Y No significant linear trend, but a systematic non-linear trend
DY or UY Significant linear trend (downward or upward) and a significant non-linear trend. This is considered the same as "-Y"
<b>SIZE</b> length effect (mercury in fillet)
L Significant difference in concentration levels but pattern of variation same
D As "L" but pattern of variation significantly different
- No significant difference between "small" and "large" fish
<b>SM3</b> Projected smoothed median for three years expressed as quotient of value and "high background" ("?" if missing background or if number of years is less than seven)
<b>PWR</b> POWER; estimated number of years to detect a hypothetical situation of 10% trend a year with a 90% power



Annual median concentration of CD (ppm)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30A	MYTI EDU	SB	d.w.				1.07	0.81	1.41	0.60	0.61	0.74	0.77	0.77	1.12	1.26	1.17	0.78	0.80	0.86	1.27	1.16	1.13	0.91	1.75	1.56	2.16	1.1	U-	1.4	10	
31A	MYTI EDU	SB	d.w.				1.39	1.31	0.89	1.93	0.40	0.43	0.41	0.72	0.73	0.91	0.93	0.78	1.32	0.79	0.85	1.07	1.25	1.00	1.76	1.14	1.33	1.40	no	UY	no	13
35A	MYTI EDU	SB	d.w.				1.35	0.95	1.17	1.30	0.52	0.66	0.65	0.93	1.05	1.35	1.11	0.96	0.89	0.77	0.97	1.14	1.53	1.33	1.53	1.16	1.06	1.23	no	UY	no	10
36A	MYTI EDU	SB	d.w.				0.85	1.19	0.84	1.38	0.59	0.56	0.50	0.41	1.22	1.06	0.90	1.22	1.17	1.60	1.84	0.97	1.01	1.65	2.59	1.00	0.85	0.96	no	--	no	13
71A	MYTI EDU	SB	d.w.				2.52	1.98	1.42	2.00	0.98	2.11	2.02	0.97	1.09	1.66	1.89	1.97	2.25	1.50	2.44	1.53	1.76	1.99	1.43	1.69	1.44	1.57	no	--	no	11
76A	MYTI EDU	SB	d.w.																								no	--	no	10		
15A	MYTI EDU	SB	d.w.																								no	--	no	15		
51A	MYTI EDU	SB	d.w.																								2.6	--	no	21		
52A	MYTI EDU	SB	d.w.																								2.7	D-	1.7	19		
56A	MYTI EDU	SB	d.w.																								3.1	D-	no	16		
57A	MYTI EDU	SB	d.w.																								1.6	D-	no	13		
63A	MYTI EDU	SB	d.w.																								no	D-	no	16		
65A	MYTI EDU	SB	d.w.																								no	D-	no	16		
69A	MYTI EDU	SB	d.w.																								no	--	no	13		
22A	MYTI EDU	SB	d.w.																								no	--	no	13		
23A	MYTI EDU	SB	d.w.																								0.83	no	-?	?	18	
24A	MYTI EDU	SB	d.w.																								0.96	no	-?	?	12	
82A	MYTI EDU	SB	d.w.				1.41	1.15	2.31	0.99	0.40	1.26		1.20	1.21	1.15		0.98	1.22									0.94	no	--	no	15
84A	MYTI EDU	SB	d.w.				1.39	1.86	2.38	2.10	0.96	1.19		1.82	2.11	1.60		1.64	1.29									1.54	no	--	no	12
87A	MYTI EDU	SB	d.w.				0.97	1.02	1.93	0.77	0.69	0.76		0.87	0.98	0.93		1.15	1.27									1.00	no	--	no	11
25A	MYTI EDU	SB	d.w.																								1.25	no	-?	?	9	
26A	MYTI EDU	SB	d.w.																								1.24	no	-?	?	<=5	
28A	MYTI EDU	SB	d.w.																								0.95	no	-?	?	9	
91A	MYTI EDU	SB	d.w.																								no	-?	?	11		
92A	MYTI EDU	SB	d.w.																								no	--	no	11		
93A	MYTI EDU	SB	d.w.																								0.89	no	-?	?	15	
94A	MYTI EDU	SB	d.w.																								1.13	no	-?	?	<=5	
95A	MYTI EDU	SB	d.w.																								1.00	no	-?	?	8	
96A	MYTI EDU	SB	d.w.																								0.72	no	-?	?	8	
97A	MYTI EDU	SB	d.w.																								1.44	no	-?	?	7	
98A	MYTI EDU	SB	d.w.																								no	--	no	12		
98X	MYTI EDU	SB	d.w.																								no	-?	?	6		
99A	MYTI EDU	SB	d.w.																								1.35	no	-?	?	6	
41A	MYTI EDU	SB	d.w.																								no	-?	?	12		
43A	MYTI EDU	SB	d.w.																								1.9	-?	?	8		
44A	MYTI EDU	SB	d.w.																								no	-?	?	12		
46A	MYTI EDU	SB	d.w.																								1.4	-?	?	10		
48A	MYTI EDU	SB	d.w.																								no	-?	?	<=5		
10A	MYTI EDU	SB	d.w.																								no	--	no	11		
11A	MYTI EDU	SB	d.w.																								no	-?	?	9		
11X	MYTI EDU	SB	d.w.																								no	--	no	12		

Annual median concentration of CD (ppm)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
I021	MYTI EDU	SB	d.w.														1.73	2.26	2.48	3.31	1.83	2.53	2.41	1.2	--	1.1	10					
I022	MYTI EDU	SB	d.w.														1.43	1.36	1.26	2.09	1.94	1.33	1.70	2.69	1.61	1.25	no	--	no	11		
I023	MYTI EDU	SB	d.w.														1.61	1.40	1.77	2.04	1.45	0.95	0.87	1.55	1.48	1.03	no	--	no	11		
I024	MYTI EDU	SB	d.w.														1.31	1.63	2.04	2.56	2.45	1.83	2.53	2.70	2.03	1.57	no	--	no	9		
30A	MYTI EDU	SB	d.w.														0.78	0.80	0.86	1.27	1.16	1.13	0.91	1.75	1.56	2.16	1.1	U-	1.4	10		
I301	MYTI EDU	SB	d.w.														0.82	0.80	0.82	1.03	1.29	0.72	0.90	0.89	1.15	1.32	no	--	no	9		
I304	MYTI EDU	SB	d.w.														1.33	0.72	0.78	1.05	0.99	0.92	1.16	1.30	1.37	1.10	no	--	no	10		
I306	MYTI EDU	SB	d.w.														0.81	0.78	0.65	0.71	0.84	0.59	0.73	0.87	1.28	0.99	no	--	no	9		
I307	MYTI EDU	SB	d.w.														0.94	0.82	0.69	0.72	0.83	0.72	0.90	1.46	1.44	1.14	no	U-	no	9		
I131	MYTI EDU	SB	d.w.														1.24	0.88	1.14	1.31	1.18	1.98	2.48	1.13	0.86	1.34	no	--	no	12		
I201	MYTI EDU	SB	d.w.														0.80	0.86	1.06	0.93	1.27	1.42	1.49	2.80	0.71	0.96	no	--	no	13		
I205	MYTI EDU	SB	d.w.														0.82		1.37	0.86	1.49	1.99	1.42	2.43	1.25	1.02	no	--	no	12		
51A	MYTI EDU	SB	d.w.														42.80	58.20									2.6	--	no	21		
52A	MYTI EDU	SB	d.w.														94.40	10.20	80.10	43.10	14.70	8.71	19.80	18.40	9.14	11.40	10.50	5.59	5.00	7.38	5.37	
I965	MYTI EDU	SB	d.w.																						2.02	2.15	0.81	1.90	no	-?	?	17
I962	MYTI EDU	SB	d.w.																								no	-?	?	6		
I964	MYTI EDU	SB	d.w.																								no	-?	?	16		
I969	MYTI EDU	SB	d.w.																								no	--	no	11		
10A	MYTI EDU	SB	d.w.																								no	--	no	11		

## Annual median concentration of CD (ppm)

St	Species	tis	Base																					ANALYSIS										
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
30B	GADU MOR	LI	w.w.				0.010	0.050	0.062	0.071	0.022	0.027	0.035	0.027	0.100	0.065	0.063	0.049	0.045	0.045	0.107	0.165	0.078	0.111	0.106	0.114	0.100	1	U-	1	16			
36B	GADU MOR	LI	w.w.	0.078	0.060	0.220	0.070	0.050	0.137	0.061	0.031	0.028	0.024	0.010	0.021	0.034	0.021	0.042	0.033	0.074	0.036	0.065	0.041	0.029	0.025	0.009	0.007	no	DY	no	15			
15B	GADU MOR	LI	w.w.										0.026	0.009	0.025	0.016	0.014	0.016	0.024	0.031	0.030	0.026	0.033	0.026	0.018	0.037	0.010	no	--	no	14			
53B	GADU MOR	LI	w.w.										0.058	0.093	0.045	0.149	0.215	0.038	0.007	0.180	0.143	0.228	0.726	0.829	0.565	0.431	0.253	0.368	3.7	--	no	25		
67B	GADU MOR	LI	w.w.										0.145	0.052	0.047	0.069	0.077	0.051	0.115	0.099	0.033	0.111	0.277	0.019	0.072	0.059	0.032	0.020	0.010	0.016	no	--	no	20
23B	GADU MOR	LI	w.w.										0.022	0.024	0.020	0.025	0.015	0.026	0.014	0.029	0.025	0.033	0.019	0.025	0.021	0.016	0.023	no	--	no	11			
84B	GADU MOR	LI	w.w.										0.130	0.095	0.069	0.029												no	D?	?	6			
92B	GADU MOR	LI	w.w.																									no	--?	?	16			
98B	GADU MOR	LI	w.w.																									no	--	no	24			
43B	GADU MOR	LI	w.w.																									no	-?	?	12			
10B	GADU MOR	LI	w.w.																									1	--	no	12			

## Annual median concentration of CD (ppm)

St	Species	tis	Base																					ANALYSIS										
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
33B	PLAT FLE	LI	w.w.				0.190		0.195	0.176	0.251	0.061	0.106	0.234	0.196	0.160	0.184	0.087	0.091	0.110	0.107	0.108	0.126	0.071	0.091	0.057	0.063	0.031	no	D-	no	13		
53B	PLAT FLE	LI	w.w.							2.240	1.530	1.540	1.720	1.790	0.789		0.135	2.530	0.892	1.470	2.550	1.770	2.740	2.740	4.560	1.460	4.9	--	8	20				
67B	PLAT FLE	LI	w.w.																2.480		0.187	0.185	0.148	0.059	0.065	0.080	0.061	no	D-	no	19			
21F	PLAT FLE	LI	w.w.																								no	-?	?	21				
36F	LIMA LIM	LI	w.w.													0.106	0.112	0.230	0.295	0.135	0.147	0.139	0.123	0.202	0.227	0.139	0.232	0.127	0.142	0.188	no	--	no	12
15F	LIMA LIM	LI	w.w.														0.099	0.136	0.125	0.153	0.076	0.181	0.167	0.313	0.129	0.110	0.189	no	--	no	14			
22F	LIMA LIM	LI	w.w.													0.095	0.091	0.128	0.169	0.125									no	-?	?	9		
98F	LIMA LIM	LI	w.w.														0.980	0.182	0.225										no	-?	?	21		
30F	PLEU PLA	LI	w.w.														0.110	0.101	0.222	0.230	0.231	0.244						1.1	--?	?	<5			
22F	PLEU PLA	LI	w.w.															0.100	0.747	0.324	0.203	0.214	0.821	0.521	0.217	0.218	0.073	no	--	no	21			
98F	PLEU PLA	LI	w.w.															0.571		0.141	0.248	0.302	0.204	0.316	0.307	1.5	--	1.8	16					
10F	PLEU PLA	LI	w.w.															0.097	0.033	0.051	0.037	0.049	0.034	0.054	0.049	m	DY	m	14					
67B	LEPI WHI	LI	w.w.				0.181		0.180	0.109	0.066	0.197	0.085	0.100	0.120	0.304	0.259	0.200	0.097	0.033	0.051	0.037	0.049	0.034	0.054	0.049	m	DY	m	14				

Annual median concentration of HG (ppm)

St	Species	tis	Base	ANALYSIS																															
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR				
30A	MYTI EDU	SB	d.w.		0.118	0.073	0.147	0.050	0.130	0.044	0.064	0.053	0.051	0.070	0.087	0.057	0.070	0.060	0.078	0.114	0.060	0.059	0.095	0.071	0.153	no	--	no	13						
31A	MYTI EDU	SB	d.w.		0.076	0.164	0.086	0.120	0.050	0.090	0.023	0.060	0.049	0.051	0.045	0.050	0.062	0.044	0.052	0.070	0.088	0.046	0.051	0.058	0.058	0.094	no	--	no	13					
35A	MYTI EDU	SB	d.w.		0.093	0.074	0.084	0.170	0.050	0.180	0.050	0.062	0.059	0.058	0.054	0.061	0.037	0.038	0.035	0.067	0.101	0.028	0.047	0.058	0.057	0.094	no	--	no	15					
36A	MYTI EDU	SB	d.w.		0.052	0.043	0.084	0.140	0.050	0.140	0.034	0.045	0.048	0.039	0.032	0.048	0.033	0.044	0.074	0.030	0.046	0.038	0.025	0.034	0.053	0.108	no	--	no	14					
71A	MYTI EDU	SB	d.w.		0.393	0.242	0.218	0.247	0.120	0.340	0.249	0.182	0.145	0.178	0.140	0.212	0.201	0.222	0.312	0.110	0.155	0.132	0.123	0.150	0.154	0.189	no	--	1.1	12					
76A	MYTI EDU	SB	d.w.								0.071	0.068	0.050	0.021			0.057	0.082	0.063	0.101	0.033	0.063	0.059	0.084	0.101	no	--	no	15						
15A	MYTI EDU	SB	d.w.								0.056	0.052		0.024	0.050	0.022	0.049	0.056	0.053	0.044	0.163	0.035	0.045	0.060	0.095	no	--	no	16						
51A	MYTI EDU	SB	d.w.								0.240	0.250					1.510	0.901	0.175	0.577	2.890	3.860	0.774	1.450	1.470	0.304	1.5	--	no	24					
52A	MYTI EDU	SB	d.w.									2.350	0.321	3.010	0.976	0.372	0.282	0.437	0.178	0.260	0.258	0.580	0.340	0.298	0.264	0.195	0.228	1.1	--	no	19				
56A	MYTI EDU	SB	d.w.									0.530	0.370	1.090	0.710	1.540	0.935	1.220	0.352	0.679	0.365	0.526	0.282	0.917	0.982	0.611	0.602	0.346	0.235	1.2	--	no	15		
57A	MYTI EDU	SB	d.w.									0.170	0.210	0.269	0.411	0.758	0.576	0.349	0.350	0.260	0.155	0.319	0.166	0.467	0.451	0.349	0.277	0.193	0.115	no	DY	no	12		
63A	MYTI EDU	SB	d.w.									0.310	0.140	0.177	0.394	0.468	0.294	0.143	0.190	0.252	0.172	0.203	0.226	0.268	0.299	0.365	0.289	0.213	0.070	no	--	no	14		
65A	MYTI EDU	SB	d.w.									0.100	0.150	0.104	0.312	0.328	0.124	0.119	0.134	0.148	0.118	0.136	0.079	0.142	0.155	0.189	0.132	0.135	0.064	no	--	no	13		
69A	MYTI EDU	SB	d.w.										0.100	0.150	0.104	0.312	0.328	0.124	0.119	0.134	0.148	0.118	0.136	0.079	0.142	0.155	0.189	0.132	0.135	0.064	no	--	no	15	
22A	MYTI EDU	SB	d.w.										0.053	0.073		0.106	0.026	0.083	0.070	0.104	0.111	0.077	0.161	0.107	0.146	0.106	0.099	0.059	no	--	no	15			
23A	MYTI EDU	SB	d.w.										0.054	0.076											0.086	no	-?	?	10						
24A	MYTI EDU	SB	d.w.										0.058	0.075											0.090	no	-?	?	9						
82A	MYTI EDU	SB	d.w.		0.051	0.110	0.170	0.080	0.120	0.067	0.074	0.052	0.079		0.049	0.069									0.051	no	--	no	13						
84A	MYTI EDU	SB	d.w.		0.077	0.112	0.150	0.080	0.240	0.057	0.066	0.090	0.057	0.054	0.043										0.051	no	--	no	14						
87A	MYTI EDU	SB	d.w.		0.178	0.150	0.050	0.260	0.046		0.056	0.054	0.049	0.044	0.062									0.047	no	--	no	17							
25A	MYTI EDU	SB	d.w.											0.141	0.037										0.095	no	-?	?	24						
26A	MYTI EDU	SB	d.w.											0.096	0.046										0.084	no	-?	?	17						
28A	MYTI EDU	SB	d.w.											0.077	0.062										0.085	no	-?	?	8						
91A	MYTI EDU	SB	d.w.											0.054	0.076	0.094									no	-?	?	<=5							
92A	MYTI EDU	SB	d.w.											0.055	0.034	0.052	0.041	0.023	0.067							no	--	no	14						
93A	MYTI EDU	SB	d.w.											0.068	0.118										0.202	1	-?	?	13						
94A	MYTI EDU	SB	d.w.											0.082	0.063										0.051	no	-?	?	8						
95A	MYTI EDU	SB	d.w.											0.077	0.060										0.053	no	-?	?	8						
96A	MYTI EDU	SB	d.w.											0.048	0.030										0.039	no	-?	?	12						
97A	MYTI EDU	SB	d.w.											0.077	0.070										0.059	no	-?	?	<=5						
98A	MYTI EDU	SB	d.w.											0.087	0.086										0.109	0.115	0.140	0.082	no	--	no	12			
98X	MYTI EDU	SB	d.w.												0.335	0.340	0.328									1.6	-?	?	<=5						
99A	MYTI EDU	SB	d.w.												0.088	0.049										0.055	no	-?	?	14					
41A	MYTI EDU	SB	d.w.												0.069	0.064	0.064	0.085								no	-?	?	8						
43A	MYTI EDU	SB	d.w.												0.084	0.095		0.104								no	-?	?	<=5						
44A	MYTI EDU	SB	d.w.												0.055	0.050	0.052	0.059								no	-?	?	6						
46A	MYTI EDU	SB	d.w.												0.039	0.062	0.056									no	-?	?	10						
48A	MYTI EDU	SB	d.w.												0.073	0.060	0.052									no	-?	?	<=5						
10A	MYTI EDU	SB	d.w.												0.053	0.049	0.059	0.062	0.058	0.063	0.050	0.052	0.049	0.055	0.050	no	--	no	6						
11A	MYTI EDU	SB	d.w.												0.182	0.145	0.086	0.146								no	-?	?	13						
11X	MYTI EDU	SB	d.w.																					0.081	0.037	0.056	0.067	0.065	0.037	0.037	0.037	no	--	no	12

Annual median concentration of HG (ppm)

St	Species	tis	Base	ANALYSIS																																
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR					
I021	MYTI EDU	SB	d.w.														0.212	0.397	0.496	0.859	0.356	0.436	0.319	1.6	--	1.2	13									
I022	MYTI EDU	SB	d.w.														0.130	0.134	0.321	0.404	0.415	0.182	0.238	0.289	0.197	0.155	no	--	no	13						
I023	MYTI EDU	SB	d.w.														0.140	0.143	0.295	0.310	0.263	0.094	0.096	0.150	0.142	0.129	no	--	no	14						
I024	MYTI EDU	SB	d.w.														0.107	0.180	0.450	0.543	0.425	0.120	0.295	0.238	0.233	0.210	1	--	no	16						
30A	MYTI EDU	SB	d.w.														0.057	0.070	0.060	0.078	0.114	0.060	0.059	0.095	0.071	0.153	no	--	no	13						
I301	MYTI EDU	SB	d.w.		0.118	0.073	0.147	0.050	0.130	0.044	0.064	0.053	0.051	0.070	0.087		0.066	0.068	0.058	0.068	0.063	0.041	0.068	0.050	0.073	0.114	no	--	no	10						
I304	MYTI EDU	SB	d.w.														0.047	0.069	0.040	0.054	0.050	0.029	0.051	0.046	0.049	0.063	no	--	no	11						
I306	MYTI EDU	SB	d.w.														0.045	0.062	0.039	0.061	0.051	0.036	0.035	0.040	0.051	0.074	no	--	no	10						
I307	MYTI EDU	SB	d.w.														0.038	0.071	0.034	0.047	0.054	0.033	0.049	0.054	0.062	0.089	no	--	no	11						
I711	MYTI EDU	SB	d.w.															0.382	0.287	0.198	0.200						no	-?	?	9						
I712	MYTI EDU	SB	d.w.															0.181	0.257	0.214	0.218	0.211	0.145	0.178				no	--	no	9					
I713	MYTI EDU	SB	d.w.																			0.111	0.177	0.157			no	-?	?	10						
I131	MYTI EDU	SB	d.w.														0.127	0.069	0.060	0.144	0.064	0.034	0.078	0.050	0.065	0.100	no	--	no	15						
I201	MYTI EDU	SB	d.w.															0.101	0.132	0.157	0.169	0.313	0.096	0.130				no	--	no	14					
I205	MYTI EDU	SB	d.w.															0.097	0.171	0.205	0.167	0.218	0.151	0.142				no	--	no	10					
51A	MYTI EDU	SB	d.w.		0.240	0.250											1.510	0.901	0.175	0.577	2.890	3.860	0.774	1.450	1.470	0.304	1.5	--	no	24						
52A	MYTI EDU	SB	d.w.														2.350	0.321	3.010	0.976	0.372	0.282	0.437	0.178	0.260	0.258	0.580	0.340	0.298	0.264	0.195	0.228	1.1	--	no	19
I965	MYTI EDU	SB	d.w.																					0.200	0.090	0.168	no	-?	?	17						
I964	MYTI EDU	SB	d.w.																					0.136	0.059	0.049	no	-?	?	11						
I969	MYTI EDU	SB	d.w.																					0.063	0.042	0.036	no	-?	?	7						
10A	MYTI EDU	SB	d.w.															0.053	0.049	0.059	0.062	0.058	0.063	0.050	0.052	0.049	0.055	0.050	no	--	no	6				

Annual median concentration of HG (ppm)  
 Cursive values in shaded area indicate temporal  
 trend analysis based data since 1998

St	Species	tis	Base																					ANALYSIS										
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
30B	GADU MOR (s)	MU	w.w.				0.125	0.089	0.079	0.040	0.059	0.121	0.120	0.090	0.110	0.122	0.102	0.080	0.108	0.131	0.117	0.153	0.173	0.224	0.131	0.137	0.105	1.1	U-L	no	11			
30B	GADU MOR (s)	MU	w.w.																									1.1	--L	no	10			
30B	GADU MOR (I)	MU	w.w.				0.155	0.090	0.074	0.038	0.147	0.166	0.130	0.108	0.150	0.155	0.129	0.119	0.142	0.190	0.232	0.351	0.252	0.340	0.207	0.194	0.185	1.8	U-L	1.1	13			
30B	GADU MOR (I)	MU	w.w.																									1.8	--L	1.1	10			
36B	GADU MOR (s)	MU	w.w.	0.069	0.080	0.110	0.075	0.080	0.061	0.032	0.053	0.069	0.060	0.060	0.059	0.067	0.054	0.076	0.067	0.089	0.078	0.060	0.072	0.052	0.057	0.037	0.046	no	D-L	no	10			
36B	GADU MOR (I)	MU	w.w.	0.079	0.160	0.180	0.195	0.120	0.112	0.039	0.083	0.074	0.115	0.100	0.080	0.083	0.060	0.095	0.070	0.157	0.088	0.186	0.075	0.123	0.108	0.073	0.074	no	--L	no	13			
15B	GADU MOR (s)	MU	w.w.										0.065	0.040	0.026	0.018	0.045	0.044	0.059	0.076	0.044	0.023	0.038	0.064	0.038	0.033	0.024	no	--L	no	14			
15B	GADU MOR (I)	MU	w.w.										0.120	0.070	0.063	0.039	0.081	0.046	0.087	0.108	0.090	0.027	0.047	0.087	0.048	0.037	0.027	no	--L	no	15			
53B	GADU MOR (s)	MU	w.w.										0.223	0.105	0.160	0.184	0.204	0.360	0.090	0.054	0.229	0.128	0.151	0.175	0.209	0.257	0.196	0.094	0.161	1.6	--L	no	16	
53B	GADU MOR (I)	MU	w.w.										0.196	0.105	0.203	0.170	0.269	0.396	0.141	0.090	0.277	0.243	0.298	0.285	0.395	0.715	0.375	0.168	0.186	1.9	--L	no	15	
67B	GADU MOR (s)	MU	w.w.										0.100	0.085	0.090	0.079	0.100	0.085	0.093	0.120	0.071	0.073	0.117	0.051	0.058	0.074	0.045	0.048	0.033	0.033	no	D-L	no	10
67B	GADU MOR (I)	MU	w.w.										0.170	0.085	0.102	0.255	0.130	0.141	0.083	0.106	0.072	0.089	0.160	0.068	0.060	0.107	0.092	0.094	0.056	0.040	no	--L	no	14
23B	GADU MOR (s)	MU	w.w.										0.065	0.070	0.060	0.042	0.052	0.069	0.060	0.073	0.075	0.061	0.083	0.073	0.063	0.041	0.078	no	--L	no	10			
23B	GADU MOR (I)	MU	w.w.										0.170	0.110	0.084	0.098	0.074	0.109	0.057	0.105	0.116	0.113	0.107	0.097	0.102	0.088	0.109	1.1	--L	no	10			
84B	GADU MOR (s)	MU	w.w.				0.035	0.040	0.025				0.044														no	-?	?	12				
84B	GADU MOR (I)	MU	w.w.				0.060	0.040	0.025				0.044														no	-?	?	14				
92B	GADU MOR (s)	MU	w.w.													0.046	0.079	0.080	0.077									0.054	no	-?	?	11		
92B	GADU MOR (I)	MU	w.w.													0.058	0.091	0.074	0.117									0.069	no	-?	?	12		
98B	GADU MOR (s)	MU	w.w.													0.067	0.054	0.069	0.069	0.040	0.095	0.066	0.047	0.095	0.047	0.047	0.067	0.099	no	---	no	13		
98B	GADU MOR (I)	MU	w.w.													0.065	0.064	0.069	0.086	0.037	0.128	0.090	0.043	0.090	0.072	0.059	0.059	0.162	1.6	---	1.4	15		
43B	GADU MOR (s)	MU	w.w.													0.065	0.054	0.047										no	-?	?	<=5			
43B	GADU MOR (I)	MU	w.w.													0.050	0.059	0.057										no	-?	?	6			
10B	GADU MOR (s)	MU	w.w.													0.044	0.034	0.029	0.011	0.014	0.017	0.011	0.009	0.009	0.016	0.014	0.014	no	DYL	no	11			
10B	GADU MOR (I)	MU	w.w.													0.056	0.053	0.040	0.020	0.019	0.032	0.015	0.011	0.012	0.017	0.018	0.018	no	D-L	no	12			

Annual median concentration of HG (ppm)  
 Cursive values in shaded area indicate temporal  
 trend analysis based data since 1998

St	Species	tis	Base	ANALYSIS																																
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR					
33B	PLAT FLE (s)	MU	w.w.	0.110	0.090	0.077	0.019	0.069		0.175	0.088	0.116	0.092	0.069	0.053	0.048	0.076	0.038	0.046	0.050	0.029	0.067	0.023	0.035	no	---	no	16								
33B	PLAT FLE (l)	MU	w.w.		0.139	0.100	0.077	0.024	0.069	0.195	0.135	0.196	0.103	0.088	0.049	0.060	0.087	0.070	0.119	0.078	0.059	0.028	0.062	0.091	no	---	no	16								
53B	PLAT FLE (s)	MU	w.w.			0.111	0.074	0.139	0.154	0.141	0.071			0.035	0.165	0.130	0.165	0.249	0.289	0.333	0.553	0.521	0.103	1	---	1.3	17									
53B	PLAT FLE (l)	MU	w.w.			0.111	0.128	0.090	0.124	0.100	0.116			0.036	0.208	0.221	0.257	0.157	0.233	0.438	0.450	0.490	0.443	4.4	U-	7.5	15									
53B	PLAT FLE (l)	MU	w.w.															0.257	0.157	0.233	0.438	0.450	0.490	0.443	4.4	---	7.5	12								
67B	PLAT FLE (s)	MU	w.w.															0.100	0.043	0.036	0.064	0.044	0.051	0.057	0.049	no	---	no	12							
67B	PLAT FLE (l)	MU	w.w.															0.246	0.061	0.034	0.082	0.068	0.071	0.066	0.065	no	---	no	17							
21F	PLAT FLE (s)	MU	w.w.																0.075	0.081	0.039	0.038		0.021	no	D?-	?	9								
21F	PLAT FLE (l)	MU	w.w.																0.119	0.080	0.034	0.044		0.021	no	D?-	?	12								
36F	LIMA LIM (s)	MU	w.w.															0.045	0.071	0.066	0.070	0.050	0.054	0.049	0.031	0.062	0.038	0.056	0.041	0.062	0.058	0.050	no	--L	no	10
36F	LIMA LIM (l)	MU	w.w.															0.098	0.074	0.133	0.101	0.076	0.100	0.066	0.091	0.092	0.068	0.102	0.099	0.114	0.083	0.117	1.2	--L	1.2	10
15F	LIMA LIM (s)	MU	w.w.															0.090	0.038	0.037	0.025	0.037	0.048	0.042	0.036	0.055	0.056	0.034	0.055	no	--L	no	12			
15F	LIMA LIM (l)	MU	w.w.															0.150	0.034	0.036	0.056	0.108	0.073	0.088	0.059	0.165	0.105	0.081	0.124	1.2	--L	1.3	17			
22F	LIMA LIM (s)	MU	w.w.															0.084	0.040	0.207		0.045	0.063					no	-?-	?	20					
22F	LIMA LIM (l)	MU	w.w.															0.174	0.152	0.282	0.223	0.372						3.7	-?-	?	11					
30F	PLEU PLA (s)	MU	w.w.																0.058	0.028	0.037					no	-?-	?	13							
30F	PLEU PLA (l)	MU	w.w.																0.035	0.056	0.048					no	-?-	?	10							
22F	PLEU PLA (s)	MU	w.w.																0.029	0.043	0.050					no	-?-	?	7							
22F	PLEU PLA (l)	MU	w.w.																0.051	0.051	0.083					no	-?-	?	9							
98F	PLEU PLA (s)	MU	w.w.																0.017	0.048	0.038	0.029	0.049	0.058	0.059	0.033	0.018	0.016	no	---	no	13				
98F	PLEU PLA (l)	MU	w.w.																0.025	0.075	0.026	0.059	0.049	0.164	0.090	0.031	0.041	0.016	no	---	no	18				
10F	PLEU PLA (s)	MU	w.w.																	0.032	0.014	0.029	0.019	0.019	0.015	0.016	no	--L	no	12						
10F	PLEU PLA (l)	MU	w.w.																	0.042	0.034	0.031	0.024	0.035	0.019	0.035	no	--L	no	11						
67B	LEPI WHI (s)	MU	w.w.	0.235		0.350	0.329	0.210	0.343	0.075	0.174	0.187	0.305	0.364	0.398	0.172	0.066	0.110	0.104	0.094	0.208	0.079	0.119	m	--L	m	16									
67B	LEPI WHI (l)	MU	w.w.	0.499		0.350	0.329	0.320	0.589	0.147	0.327	0.336	0.422	0.341	0.372	0.331	0.275	0.392	0.330	0.237	0.091	0.186	0.334	m	--L	m	14									

Annual median concentration of PB (ppm)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30A	MYTI EDU	SB	d.w.										1.86	1.36	3.95	2.27	2.54	1.58	2.12	2.69	36.70	2.13	1.74	1.76	2.24	2.58	3.74	1.2	--	1.3	22	
31A	MYTI EDU	SB	d.w.										1.38	1.21	1.26	1.03	1.37	1.68	1.79	0.73	1.54	0.63	0.63	0.51	1.43	0.81	0.78	no	--	no	13	
35A	MYTI EDU	SB	d.w.										1.44	1.07	1.68	1.20	1.28	0.51	0.63	0.66	0.76	0.71	0.52	0.87	0.57	0.57	0.81	no	D-	no	11	
36A	MYTI EDU	SB	d.w.										1.01	0.85	0.79	1.12	1.39	1.24	2.04	2.17	1.57	1.00	0.94	0.62	0.45	0.59	0.96	no	DY	no	11	
71A	MYTI EDU	SB	d.w.										1.16	0.75	1.72	1.42	1.92	1.49	2.21	2.83	0.87	0.90	0.77	1.45	0.92	0.92	0.87	no	--	no	13	
76A	MYTI EDU	SB	d.w.										1.77	0.97	1.50	0.91			0.80	1.84	1.23	1.99	0.60	0.83	0.77	0.94	1.48	no	--	no	14	
15A	MYTI EDU	SB	d.w.										1.46	0.78		0.98	1.05	0.52	0.67	1.12	1.28	1.66	2.20	0.96	0.71	0.76	0.86	no	--	no	12	
51A	MYTI EDU	SB	d.w.														149.00	60.30	17.20	29.60	37.10	91.70	32.40	98.40	108.00	42.20	14.1	--	25.3	19		
52A	MYTI EDU	SB	d.w.										12.10	313.00	189.00	65.50	16.40	17.50	9.84	20.60	14.70	11.60	11.00	21.80	21.80	16.90	16.30	5.4	--	7.1	23	
56A	MYTI EDU	SB	d.w.										20.70	23.40	121.00	109.00	24.70	46.40	27.80	37.50	15.70	30.30	28.50	30.50	42.90	27.80	24.70	8.2	--	8.8	17	
57A	MYTI EDU	SB	d.w.										10.50	12.10	33.30	19.20	15.10	13.20	5.60	13.70	6.15	10.40	10.30	11.90	9.59	7.02	5.79	1.9	--	1.1	14	
63A	MYTI EDU	SB	d.w.										12.10	10.10	15.40	10.90	7.22	12.10	7.60	6.10	6.39	4.84	4.52	7.05	6.57	6.30	1.92	no	D-	no	13	
65A	MYTI EDU	SB	d.w.										5.61	3.78	5.19	6.53	3.28	4.73	2.41	3.00	1.77	1.63	2.45	2.84	3.05	2.82	1.25	no	D-	no	12	
69A	MYTI EDU	SB	d.w.													4.62	3.42	2.80	3.17	4.02	3.66	1.98	3.40	2.27	3.91	2.76	2.53	0.96	no	--	no	12
22A	MYTI EDU	SB	d.w.										1.37	1.46		2.78	1.87	1.39	1.18	1.51	1.37	1.21	1.70	1.30	1.21	0.88	1.46	1.79	no	--	no	11
23A	MYTI EDU	SB	d.w.										1.42	1.47												0.88	no	-?	?	<=5		
24A	MYTI EDU	SB	d.w.										1.42	1.21													2.01	no	-?	?	8	
82A	MYTI EDU	SB	d.w.										1.28	0.93	0.92			0.62	0.67								0.56	no	D-	no	9	
84A	MYTI EDU	SB	d.w.										1.01	1.15	1.38			1.38	0.83								1.11	no	--	no	10	
87A	MYTI EDU	SB	d.w.										0.97	0.87	0.63			1.40	2.47								0.42	no	--	no	19	
25A	MYTI EDU	SB	d.w.												2.68	1.77												2.01	no	-?	?	12
26A	MYTI EDU	SB	d.w.												1.42	1.38												1.87	no	-?	?	<=5
28A	MYTI EDU	SB	d.w.												1.39	1.87												0.96	no	-?	?	10
91A	MYTI EDU	SB	d.w.										0.90	1.46	2.01												no	-?	?	6		
92A	MYTI EDU	SB	d.w.										0.93	0.63	1.09	0.66	0.65	2.18									no	--	1.4	16		
93A	MYTI EDU	SB	d.w.											1.14	1.62												2.13	no	-?	?	10	
94A	MYTI EDU	SB	d.w.											0.77	1.18												0.31	no	-?	?	13	
95A	MYTI EDU	SB	d.w.											1.04	2.16												0.59	no	-?	?	17	
96A	MYTI EDU	SB	d.w.											1.13	0.99												0.49	no	D?	?	<=5	
97A	MYTI EDU	SB	d.w.												1.37	1.26											0.50	no	D?	?	<=5	
98A	MYTI EDU	SB	d.w.										1.87	1.85													1.34	0.89	1.16	1.09	9	
98X	MYTI EDU	SB	d.w.													4.34	3.12	4.11										1.4	-?	?	11	
99A	MYTI EDU	SB	d.w.												1.20	0.75												0.70	no	-?	?	12
41A	MYTI EDU	SB	d.w.														1.29	0.90	0.79	0.65								no	D?	?	6	
43A	MYTI EDU	SB	d.w.														1.56	1.51		0.86								no	-?	?	8	
44A	MYTI EDU	SB	d.w.														2.81	2.57	1.66	1.15								no	D?	?	7	
46A	MYTI EDU	SB	d.w.														1.26	1.57	1.38								no	-?	?	8		
48A	MYTI EDU	SB	d.w.														0.68	1.08	0.33								no	-?	?	19		
10A	MYTI EDU	SB	d.w.														1.94	2.78	0.74	0.81	2.34	1.57	1.44	1.39	1.80	1.65	1.02	no	--	no	15	
11A	MYTI EDU	SB	d.w.														1.54	1.48	0.34	0.37								no	-?	?	16	
11X	MYTI EDU	SB	d.w.																	0.74	0.52	0.31	1.09	2.32	0.74	0.28	0.37	no	--	no	20	

Annual median concentration of PB (ppm)

St	Species	tis	Base	ANALYSIS																																															
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR																				
I021	MYTI EDU	SB	d.w.															1.06	2.29	1.65	2.12	0.99	1.65	1.19			no	--	no	13																					
I022	MYTI EDU	SB	d.w.															1.00	0.60	1.18	1.31	1.94	1.05	0.95	1.27	1.36	3.51	1.2	--	1.1	14																				
I023	MYTI EDU	SB	d.w.															0.77	1.27	1.38	1.70	1.38	0.64	0.62	0.75	1.28	2.34	no	--	no	13																				
I024	MYTI EDU	SB	d.w.															0.97	1.10	1.16	1.70	1.79	0.62	1.33	1.10	1.73	2.68	no	--	1.1	13																				
30A	MYTI EDU	SB	d.w.															1.86	1.36	3.95	2.27	2.54	1.58	2.12	2.69	36.70	2.13	1.74	1.76	2.24	2.58	3.74	1.2	--	1.3	22															
I301	MYTI EDU	SB	d.w.																									2.47	2.11	1.32	3.16	1.98	1.77	3.07	1	--	no	13													
I304	MYTI EDU	SB	d.w.																										2.23	1.19	0.77	1.88	1.30	1.16	1.73	no	--	no	14												
I306	MYTI EDU	SB	d.w.																										1.34	0.68	0.54	1.03	0.66	0.70	1.08	no	--	no	13												
I307	MYTI EDU	SB	d.w.																										1.05	0.80	0.51	1.01	1.26	1.01	1.07	no	--	no	12												
I201	MYTI EDU	SB	d.w.																										3.54	4.39	4.77	4.67	4.43	6.41	3.78	8.21	1.87	3.33	1.1	--	no	14									
I205	MYTI EDU	SB	d.w.																										4.77	6.96	4.00	5.97	7.09	6.15	9.27	3.40	2.76	no	--	no	13										
51A	MYTI EDU	SB	d.w.																										12.10	313.00	189.00	65.50	16.40	17.50	149.00	60.30	17.20	29.60	37.10	91.70	32.40	98.40	108.00	42.20	14.1	--	25.3	19			
52A	MYTI EDU	SB	d.w.																																			5.4	--	7.1	23										
I965	MYTI EDU	SB	d.w.																																			4.4	-?	?	16										
I962	MYTI EDU	SB	d.w.																																			no	-?	?	9										
I964	MYTI EDU	SB	d.w.																																			20.00	12.70	6.45	13.10										
I969	MYTI EDU	SB	d.w.																																			8.75	3.49	6.57	2.2	-?	?	18							
10A	MYTI EDU	SB	d.w.																																			no	--	no	12										
																																					1.94	2.78	0.74	0.81	2.34	1.57	1.44	1.39	1.80	1.65	1.02	no	--	no	15

Annual median concentration of PB (ppm)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30B	GADU MOR	LI	w.w.										0.200	0.115	0.249	0.105	0.120	0.110	0.060	0.100	0.163	0.850	0.240	0.220	0.513	0.240	0.170	1.7	--	1.1	17	
36B	GADU MOR	LI	w.w.										0.115	0.050	0.030	0.020	0.030	0.020	0.030	0.040	0.030	0.040	0.040	0.030	0.006	0.020	0.020	no	--	no	16	
15B	GADU MOR	LI	w.w.										0.170	0.060	0.030	0.030	0.030	0.020	0.030	0.040	0.030	0.030	0.040	0.030	0.010	0.020	0.020	no	D-	no	14	
53B	GADU MOR	LI	w.w.										0.190	0.260	0.140	0.030		0.020	0.075	0.070	0.105	0.115	0.130	0.130	0.142	0.040	0.090	no	--	no	17	
67B	GADU MOR	LI	w.w.										0.130	0.180	0.030	0.075	0.090	0.040	0.040	0.090	0.030	0.040	0.040	0.030	0.015	0.020	0.020	no	D-	no	16	
23B	GADU MOR	LI	w.w.										0.060	0.080	0.030	0.030	0.030	0.020	0.030	0.040	0.030	0.040	0.030	0.030	0.006	0.020	0.020	no	--	no	15	
92B	GADU MOR	LI	w.w.											0.020		0.030	0.030	0.040										0.020	no	-?	?	12
98B	GADU MOR	LI	w.w.										0.030	0.030	0.030	0.040	0.040	0.050	0.030	0.030	0.040	0.030	0.040	0.030	0.010	0.020	0.020	no	--	no	13	
43B	GADU MOR	LI	w.w.											0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.040	0.040	0.030	0.020	0.020	0.020	no	-?	?	<=5	
10B	GADU MOR	LI	w.w.											0.030	0.020	0.040	0.040	0.040	0.030	0.040	0.030	0.040	0.030	0.020	0.020	0.020	0.020	no	--	no	10	

Annual median concentration of PB (ppm)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
33B	PLAT FLE	LI	w.w.										0.240	0.350	0.060	0.030	0.030	0.020	0.030	0.040	0.040	0.040	0.030	0.030	0.030	0.020	no	DY	no	13	
53B	PLAT FLE	LI	w.w.										0.710	0.810	0.410	0.230		0.025	0.460	0.350	0.520	0.460	0.357	0.570	1.290	0.730	0.440	1.5	--	2.5	22
67B	PLAT FLE	LI	w.w.															0.350		0.030	0.030	0.030	0.008	0.020	0.020	0.020	no	--	no	21	
21F	PLAT FLE	LI	w.w.																	0.040	0.045	0.060	0.046		0.040	0.040	0.040	no	-?	?	9
36F	LIMA LIM	LI	w.w.										0.600	0.070	0.040	0.070	0.030	0.020	0.030	0.050	0.050	0.050	0.060	0.040	0.048	0.030	0.050	no	DY	no	16
15F	LIMA LIM	LI	w.w.											0.070		0.041	0.030	0.020	0.030	0.050	0.040	0.035		0.050	0.021	0.020	0.030	no	--	no	13
22F	LIMA LIM	LI	w.w.										0.250	0.160	0.042			0.060	0.070								no	-?	?	18	
98F	LIMA LIM	LI	w.w.														0.020	0.040	0.030									no	-?	?	14
30F	PLEU PLA	LI	w.w.													0.739	0.540	0.570										2.9	-?	?	7
22F	PLEU PLA	LI	w.w.															0.280	0.280	0.460								2.3	-?	?	9
98F	PLEU PLA	LI	w.w.														0.030	0.310		0.050	0.040	0.220	0.104	0.040	0.068	0.040	0.020	no	--	no	23
10F	PLEU PLA	LI	w.w.																0.150		0.065	0.080	0.050	0.058	0.045	0.045	0.045	no	D-	no	10
67B	LEPI WHI	LI	w.w.										0.190	0.070	0.060	0.070	0.040	0.070	0.030	0.040	0.040	0.030	0.040	0.031	0.020	0.025	m	D-	m	12	

Annual median concentration of CU (ppm)

St	Species	Tise	Base	ANALYSIS																														
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
30A	MYTI EDU	SB	d.wt				4.6		7.5	5.0	5.5	6.0	10.3	10.5	5.8	6.7	8.6	6.9	7.7	9.5	7.7	8.0	6.5	5.8	6.3	7.7	8.6	no	--	no	10			
31A	MYTI EDU	SB	d.wt				7.0	6.6		4.4	4.5	4.5	9.0	11.0	5.5	5.7	6.2	7.3	6.6	6.1	8.1	7.6	5.5	5.8	5.5	6.3	6.7	no	--	no	10			
35A	MYTI EDU	SB	d.wt				6.3	3.6	8.1	4.9	4.6	5.3	8.0	10.1	6.6	6.3	6.6	6.4	6.9	6.8	7.2	7.1	5.5	6.2	6.5	7.4	8.3	no	--	no	10			
36A	MYTI EDU	SB	d.wt				6.3	3.6	6.1	4.5	4.9	4.3	5.5	9.2	5.2	5.5	5.6	7.7	9.1	6.9	6.8	6.2	5.5	5.8	5.2	6.3	7.0	no	--	no	10			
71A	MYTI EDU	SB	d.wt				8.5	5.2		6.1	8.4	7.0	8.3	10.3	7.4	7.9	7.2	8.1	7.7	9.4	7.5	7.6	8.8	7.5	7.1	11.2	8.8	no	--	1.1	9			
76A	MYTI EDU	SB	d.wt									8.5	10.8	5.7	5.6				6.7	7.8	9.1	10.2	6.9	7.7	6.4	9.8	8.0	no	--	no	10			
15A	MYTI EDU	SB	d.wt									5.7	7.2			5.5	5.3	5.3	7.3	6.1	7.6	7.3	7.6	9.5	5.7	8.1	6.5	no	--	no	9			
51A	MYTI EDU	SB	d.wt									7.1	6.1					10.2	10.2	7.4		8.2					no	--	no	9				
52A	MYTI EDU	SB	d.wt										8.4	7.5	72.1	9.4	8.5	7.0	7.0	6.3	7.7	6.5	5.5	7.0	5.9	7.5	9.2	6.2	no	--	no	18		
56A	MYTI EDU	SB	d.wt										8.1	7.9	8.8	5.4	7.5	7.4	9.2	6.4	7.7	6.6	7.8	7.2	8.2	8.7	5.8	6.0	9.1	5.4	no	--	no	9
57A	MYTI EDU	SB	d.wt										8.2	6.3	6.0	6.2	7.3	6.6	6.8	6.4	5.6	5.5	7.0	6.9	6.4	5.1	5.8	7.2	6.0	3.3	no	--	no	9
63A	MYTI EDU	SB	d.wt										9.8	6.2	5.1	6.8	11.1	6.3	6.1	6.7	6.8	6.6	6.1	5.6	5.1	5.7	5.5	6.7	8.0	2.4	no	--	no	12
65A	MYTI EDU	SB	d.wt										8.0	4.9	5.2	12.2	8.1	5.5	6.0	5.1	5.7	5.8	5.6	5.1	5.3	5.9	6.5	7.7	2.8	no	--	no	12	
69A	MYTI EDU	SB	d.wt															6.2	5.3	5.4	6.3	7.3	5.6	5.2	7.2	8.5	6.8	5.8	12.6	3.1	no	--	no	13
22A	MYTI EDU	SB	d.wt															6.4	6.7									14.9	1.5	--	1.4	12		
23A	MYTI EDU	SB	d.wt															5.6	6.5									6.6	no	-?	?	7		
24A	MYTI EDU	SB	d.wt															5.8	7.0									6.8	no	-?	?	8		
82A	MYTI EDU	SB	d.wt				6.4		4.7	5.8	7.5		11.8	6.9	9.6			7.4	7.9									5.5	no	--	no	10		
84A	MYTI EDU	SB	d.wt				10.3	96.8	56.8	39.3	26.8		17.1	22.2	24.0			9.5	7.2									8.1	no	--	no	19		
87A	MYTI EDU	SB	d.wt				4.6		20.1	8.4	5.9		7.3	6.3	6.9			7.6	7.5									6.9	no	--	no	15		
25A	MYTI EDU	SB	d.wt															6.5	5.8									6.5	no	-?	?	6		
26A	MYTI EDU	SB	d.wt															7.0	6.3									6.4	no	-?	?	6		
28A	MYTI EDU	SB	d.wt															5.4	5.2									5.7	no	-?	?	<=5		
91A	MYTI EDU	SB	d.wt															6.6	6.4	8.4								no	-?	?	7			
92A	MYTI EDU	SB	d.wt															6.7	6.0	7.6	6.3	7.5	6.4	6.1	6.6	7.2	8.3	no	--	no	7			
93A	MYTI EDU	SB	d.wt															6.5	6.8									7.9	no	-?	?	<=5		
94A	MYTI EDU	SB	d.wt															6.8	6.7									6.7	no	-?	?	<=5		
95A	MYTI EDU	SB	d.wt															7.8	6.4									8.0	no	-?	?	8		
96A	MYTI EDU	SB	d.wt															7.2	6.2									6.5	no	-?	?	7		
97A	MYTI EDU	SB	d.wt															8.1	7.0									6.8	no	-?	?	7		
98A	MYTI EDU	SB	d.wt															8.9	6.6									5.5	no	--	no	7		
98X	MYTI EDU	SB	d.wt															9.0	8.9	7.7								no	-?	?	<=5			
99A	MYTI EDU	SB	d.wt															8.7	7.3									6.7	no	-?	?	7		
41A	MYTI EDU	SB	d.wt															6.8	8.2	7.6	4.8						no	-?	?	10				
43A	MYTI EDU	SB	d.wt															7.8	9.3		6.3						no	-?	?	9				
44A	MYTI EDU	SB	d.wt															6.6	9.6	7.8	6.9						no	-?	?	10				
46A	MYTI EDU	SB	d.wt															6.6	7.4	8.0							no	-?	?	<=5				
48A	MYTI EDU	SB	d.wt															6.6	7.7	7.1							no	-?	?	7				
10A	MYTI EDU	SB	d.wt															7.3	10.9	7.8	5.7	7.9	6.9	6.2	5.4	6.4	7.4	7.0	no	--	no	9		
11A	MYTI EDU	SB	d.wt															9.9	9.7	11.6	8.4						5.5	no	-?	?	8			
11X	MYTI EDU	SB	d.wt																								5.5	no	--	no	7			

Annual median concentration of ZN (ppm)

St	Species	Tis	Base	ANALYSIS																																					
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR										
30A	MYTI EDU	SB	d.wt				138	91	140	120	93	76	161	116	147	104	117	109	114	126	173	106	93	92	116	123	141	no	--	no	10										
31A	MYTI EDU	SB	d.wt				88	132	77	106	66	68	58	181	128	125	96	97	151	103	128	120	112	84	83	93	127	128	no	--	no	11									
35A	MYTI EDU	SB	d.wt				92	80	76	90	68	82	83	166	139	131	119	98	83	94	103	112	111	66	72	85	106	131	no	UY	no	9									
36A	MYTI EDU	SB	d.wt				67	86	66	58	62	74	65	126	127	104	84	121	115	137	145	105	96	125	100	102	125	99	no	U-	no	9									
71A	MYTI EDU	SB	d.wt				124	125	77	115	101	169	128	162	143	166	120	157	150	122	192	114	114	99	98	134	121	117	no	--	no	9									
76A	MYTI EDU	SB	d.wt										158	127	124	58				112	135	137	132	83	116	123	127	128	no	--	no	11									
15A	MYTI EDU	SB	d.wt										157	144		71	89	63	232	107	127	105	127	83	110	101	114	no	--	no	13										
51A	MYTI EDU	SB	d.wt										378	253					386	223	120			170					no	--	no	14									
52A	MYTI EDU	SB	d.wt											824	272	453	408	218	141	196	183	247	160	143	134	180	91	137	106	no	D-	no	12								
56A	MYTI EDU	SB	d.wt										869	410	1170	572	479	418	388	211	290	246	377	143	271	225	158	146	151	178	no	D-	no	13							
57A	MYTI EDU	SB	d.wt										378	263	441	520	292	256	147	173	182	115	223	121	207	167	124	108	99	84	no	D-	no	11							
63A	MYTI EDU	SB	d.wt										579	216	241	509	392	207	122	122	189	147	170	129	115	115	127	106	119	55	no	D-	no	13							
65A	MYTI EDU	SB	d.wt										191	156	199	424	308	131	139	118	166	147	184	121	152	154	155	145	151	70	no	--	no	12							
69A	MYTI EDU	SB	d.wt																135	102	135	163	161	218	144	201	133	190	136	177	84	no	--	no	10						
22A	MYTI EDU	SB	d.wt																172	162	135	116	98	144	221	110	128	122	72	117	90	134	142	no	--	no	11				
23A	MYTI EDU	SB	d.wt																136	133										105	no	D?	?	<=5							
24A	MYTI EDU	SB	d.wt																175	133											130	no	-?	?	9						
82A	MYTI EDU	SB	d.wt										127	106	132	109	76	129	145	123	112	109	88						81	no	--	no	9								
84A	MYTI EDU	SB	d.wt										118	160	163	133	132	142	185	180	113	121	86						93	no	D-	no	9								
87A	MYTI EDU	SB	d.wt										100	93	98	102	105	97	117	114	90	109	97						72	no	--	no	7								
25A	MYTI EDU	SB	d.wt																184	117											119	no	-?	?	12						
26A	MYTI EDU	SB	d.wt																125	118											146	no	-?	?	<=5						
28A	MYTI EDU	SB	d.wt																141	96											103	no	-?	?	11						
91A	MYTI EDU	SB	d.wt															96	102	116											no	-?	?	<=5							
92A	MYTI EDU	SB	d.wt															89	61	90	78	59	98								no	--	no	10							
93A	MYTI EDU	SB	d.wt															94	96												103	no	-?	?	<=5						
94A	MYTI EDU	SB	d.wt															74	81												66	no	-?	?	6						
95A	MYTI EDU	SB	d.wt															95	81												77	no	-?	?	7						
96A	MYTI EDU	SB	d.wt															102	72												68	no	-?	?	10						
97A	MYTI EDU	SB	d.wt															92	92												90	no	D?	?	<=5						
98A	MYTI EDU	SB	d.wt															117	105												88	no	--	no	6						
98X	MYTI EDU	SB	d.wt																187	182	146											no	-?	?	6						
99A	MYTI EDU	SB	d.wt															107	70												84	no	-?	?	12						
41A	MYTI EDU	SB	d.wt																85	89	95	91										no	-?	?	<=5						
43A	MYTI EDU	SB	d.wt																99	97		89										no	-?	?	<=5						
44A	MYTI EDU	SB	d.wt																67	91	81	79									no	-?	?	8							
46A	MYTI EDU	SB	d.wt																88	112	93										no	-?	?	9							
48A	MYTI EDU	SB	d.wt																75	87	69										no	-?	?	8							
10A	MYTI EDU	SB	d.wt																127	120	95	114	102	103	125	114	134	100	149	no	--	no	8								
11A	MYTI EDU	SB	d.wt																87	98	77	88									no	-?	?	7							
11X	MYTI EDU	SB	d.wt																											104	70	81	101	121	90	72	78	no	--	no	9

Annual median concentration of CB\_S7 (ppb)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
30A	MYTI EDU	SB	d.w.		77.50	96.50	116.00	89.60	97.00		89.30	90.40	110.00	128.00	58.50	71.10	49.90	29.60	33.90	43.40	20.70	1.4	DY	no	11						
31A	MYTI EDU	SB	d.w.		21.70	24.90	37.10	24.70	34.60		52.20	49.00	63.80	24.60	12.90	18.00	6.49	8.87	8.97	7.58	3.79	no	DY	no	13						
35A	MYTI EDU	SB	d.w.		21.50	33.60	27.50	14.20	22.10		13.40	13.60	10.70	16.50	12.50	14.60	5.52	7.32	6.97	8.20		no	D-	no	12						
36A	MYTI EDU	SB	d.w.		11.00	17.90	19.30	7.94	11.20		5.69	10.50	12.30	12.70	8.62	12.10	5.28	5.54	6.03	5.75	5.58	no	--	no	12						
71A	MYTI EDU	SB	d.w.		17.00	34.40	25.00	14.20	15.30		16.50	10.50		9.27	11.80	13.60	8.52	12.70	7.55	9.74		no	D-	no	12						
76A	MYTI EDU	SB	d.w.				16.60	6.49	7.21				16.30	19.10	14.40	16.40	6.34	6.78	5.12	5.06		no	--	no	14						
15A	MYTI EDU	SB	d.w.				11.80				6.29	3.06	2.41	3.88	4.72	5.28	2.56	4.19	3.15	2.73	2.74	no	--	no	13						
51A	MYTI EDU	SB	d.w.								26.20	9.69	14.70	10.50	11.50	12.00	28.00	16.90	16.00	10.60		no	--	no	13						
52A	MYTI EDU	SB	d.w.					40.20	14.90		11.30	11.30	17.10	16.90	10.00	19.00	10.60	11.20	7.19	74.20	12.50	12.00	10.30	no	--	no	19				
56A	MYTI EDU	SB	d.w.					12.50	45.80	37.70	12.10	12.00	9.41	13.80	11.90	16.80	9.55	11.20	5.98	216.00	13.00	13.10	6.73	no	--	no	24				
57A	MYTI EDU	SB	d.w.							28.00	7.63	7.55	4.74	8.38	6.54	4.18	8.41	10.30	8.16	3.89	55.90	5.89	6.16	2.89	no	--	no	21			
63A	MYTI EDU	SB	d.w.							21.80	9.71	6.45	3.68	5.70	5.72		4.15	7.95	7.26	4.09	13.80	3.54	6.05	1.25	no	--	no	16			
65A	MYTI EDU	SB	d.w.							6.05	11.10	33.40	9.29	5.59	3.69	5.55	3.37	5.19	3.76	7.62	6.44	3.00	8.31	2.73	3.73	1.24	no	--	no	17	
69A	MYTI EDU	SB	d.w.									4.82		4.97	4.51	2.77	5.41	12.60	5.83	2.53	5.70	3.18	4.01	1.27	no	--	no	16			
22A	MYTI EDU	SB	d.w.								18.90	8.23	8.61		7.97	6.84	5.19	4.69	11.50	6.01	5.14	4.69	3.24	8.50	5.51	no	--	no	13		
82A	MYTI EDU	SB	d.w.								8.40	15.60												2.39	no	-?	? 16				
84A	MYTI EDU	SB	d.w.								5.25	20.50	5.05	8.44		3.60	6.37							0.85	no	--	no 21				
92A	MYTI EDU	SB	d.w.										4.46	2.49	5.83	4.05	2.89	7.74							no	--	no	15			
96A	MYTI EDU	SB	d.w.										3.35	1.52										0.79	no	-?	? 16				
98A	MYTI EDU	SB	d.w.										20.50	5.68					10.70	8.40	4.14	3.54	4.56	3.23	3.62	1.85	no	D-	no	14	
98X	MYTI EDU	SB	d.w.											87.30	78.40	46.40									3.1	-?	? 9				
99A	MYTI EDU	SB	d.w.											3.42	1.97										0.87	no	-?	? 12			
41A	MYTI EDU	SB	d.w.												3.49	4.26	2.39	2.58							no	-?	? 10				
43A	MYTI EDU	SB	d.w.												2.92	3.10		3.02							no	-?	? <=5				
44A	MYTI EDU	SB	d.w.													7.31	8.46	29.40							2	-?	? 15				
46A	MYTI EDU	SB	d.w.													5.74	4.16	3.11							no	D?	? <=5				
48A	MYTI EDU	SB	d.w.													6.22	4.04	3.10							no	-?	? 6				
10a	MYTI EDU	SB	d.w.													6.03	4.29	4.66	6.29		5.11	4.33	3.03	2.13	2.58	1.28	no	D-	no	10	
11A	MYTI EDU	SB	d.w.													7.48	6.92	4.32	5.75		3.34	3.56	4.48	2.79	3.10	1.93	1.93	1.28	no	-?	? 10
11X	MYTI EDU	SB	d.w.																							no	D-	no	9		

Annual median concentration of CB\_S7 (ppb)

St	Species	tis	Base	ANALYSIS																																
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR					
I021	MYTI EDU	SB	d.w.														43.1	31.8	32.2	24.1	22.2	20.0	25.1	1.7	D-	no	8									
I022	MYTI EDU	SB	d.w.														32.1	25.9	41.2	22.4	28.9	19.2	22.4	20.8	15.2	17.1	no	--	no	10						
I023	MYTI EDU	SB	d.w.														19.6	20.9	26.0	15.0	22.2	10.8	17.4	15.9	12.3	12.6	no	--	no	11						
I024	MYTI EDU	SB	d.w.														31.8	36.1	45.6	36.6	28.7	16.8	17.7	26.0	15.0	15.8	no	D-	no	11						
30A	MYTI EDU	SB	d.w.														77.5	96.5	116.0	89.6	97.0	89.3	90.4	110.0	128.0	58.5	71.1	49.9	29.6	33.9	43.4	20.7	1.4	DY	no	11
I301	MYTI EDU	SB	d.w.														118.0	113.0	182.0	86.5	125.0	58.7	64.6	62.6	70.4	57.9	3.9	--	2.4	11						
I304	MYTI EDU	SB	d.w.														35.2	23.8	44.4	35.9		19.9	25.0	24.4	27.5	30.0	2	--	1.5	11						
I306	MYTI EDU	SB	d.w.														16.4	15.7	54.2	26.1		21.8	17.2	15.7	15.4	17.9	no	--	no	14						
I307	MYTI EDU	SB	d.w.														20.6	28.5	40.2	17.3		20.3	16.9	17.5	15.4	13.0	no	--	no	11						
I711	MYTI EDU	SB	d.w.														24.8	13.3	13.3	20.6	21.6	18.4		13.4			no	--	no	12						
I712	MYTI EDU	SB	d.w.														33.3	31.2	25.3	22.4	24.9	13.9	12.5	10.9	16.9		no	D-	no	10						
I713	MYTI EDU	SB	d.w.																				12.5	15.0	21.4	1.4	-?	?	6							
I131	MYTI EDU	SB	d.w.														7.9	11.7	13.1	22.4	12.7	10.1	14.0	29.4	8.1	4.0	no	--	no	16						
I132	MYTI EDU	SB	d.w.														27.5	33.7	32.0	31.1	22.5	10.2	15.8	11.8	13.3		no	D-	no	11						
I133	MYTI EDU	SB	d.w.														22.8	22.3	21.5	24.7	23.0	10.4	11.7	9.2	9.2	12.5	no	D-	no	10						
I241	MYTI EDU	SB	d.w.														54.3	78.9	47.2	55.2	80.8	55.5	36.3	96.4	125.0	118.0	7.9	--	9.3	13						
I242	MYTI EDU	SB	d.w.														63.0	81.6	29.6	45.6	59.5	36.6	26.2	44.6	81.9	55.9	3.7	--	4.2	14						
I243	MYTI EDU	SB	d.w.														115.0	169.0	122.0	78.2	92.4	47.9	29.3	52.5	326.0	288.0	19.2	--	24.7	17						
51A	MYTI EDU	SB	d.w.															26.2	9.7	14.7	10.5	11.5	12.0	28.0	16.9	16.0	10.6	no	--	no	13					
52A	MYTI EDU	SB	d.w.														40.2	14.9		11.3	17.1	16.9	10.0	19.0	10.6	11.2	7.2	74.2	12.5	12.0	10.3	no	--	no	19	
10A	MYTI EDU	SB	d.w.															6.0	4.3	4.7	6.3		5.1	4.3	3.0	2.1	2.6	1.3	no	D-	no	10				

Annual median concentration of CB\_S7 (ppb)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30B	GADU MOR	LI	w.w.										1240.0	3430.0	2800.0	2500.0	2910.0	2350.0	2790.0	3240.0	3660.0	3520.0	2080.0	2440.0	2230.0	2140.0	2620.0	5.2	--	4.4	11	
36B	GADU MOR	LI	w.w.										441.0	344.0	396.0	636.0	376.0	1650.0	974.0	720.0	736.0	766.0	482.0	288.0	269.0	425.0	535.0	1.1	--	no	14	
15B	GADU MOR	LI	w.w.										182.0	349.0	266.0	182.0	295.0	307.0	274.0	399.0	279.0	257.0	153.0	377.0	244.0	213.0	154.0	no	--	no	12	
53B	GADU MOR	LI	w.w.										435.0	524.0	1760.0	166.0	162.0	701.0	576.0	2370.0	487.0	1520.0	842.0	956.0	317.0	463.0	no	--	no	22		
67B	GADU MOR	LI	w.w.										316.0	293.0	268.0	226.0	329.0	210.0	269.0	627.0	206.0	273.0	148.0	225.0	145.0	92.6	94.4	no	D-	no	13	
23B	GADU MOR	LI	w.w.										222.0	244.0	228.0	208.0	128.0	193.0	196.0	125.0	179.0	229.0	207.0	167.0	111.0	114.0	202.0	no	--	no	11	
92B	GADU MOR	LI	w.w.																									119.0	no	-?	?	17
98B	GADU MOR	LI	w.w.																									no	--	no	14	
43B	GADU MOR	LI	w.w.																									no	-?	?	13	
10B	GADU MOR	LI	w.w.																									no	D-	no	12	
30B	GADU MOR	MU	w.w.																									1.6	--	1.3	19	
36B	GADU MOR	MU	w.w.																									no	--	no	21	
15B	GADU MOR	MU	w.w.																									no	--	no	16	
53B	GADU MOR	MU	w.w.																									no	--	no	>25	
67B	GADU MOR	MU	w.w.																									no	--	no	24	
23B	GADU MOR	MU	w.w.																									no	--	no	17	
92B	GADU MOR	MU	w.w.																									0.2	no	-?	?	18
98B	GADU MOR	MU	w.w.																									no	--	no	22	
43B	GADU MOR	MU	w.w.																									no	-?	?	16	
10B	GADU MOR	MU	w.w.																									no	--	no	17	

Annual median concentration of CB\_S7 (ppb)

St	Species	tis	Base	ANALYSIS																													
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR		
33B	PLAT FLE	LI	w.w.										36.0	31.1	97.5	69.0	57.0	86.0	38.3	40.5	30.5	47.2	90.7	158.0	53.0	60.1	62.8	no	--	no	15		
53B	PLAT FLE	LI	w.w.										509.0	517.0	309.0	36.0		22.8	115.0	113.0	111.0	156.0	95.8	95.1	158.0	165.0	108.0	1.1	--	1.5	19		
67B	PLAT FLE	LI	w.w.															70.0			96.9	45.8	44.0	36.3	32.0	27.6	33.1	no	D-	no	10		
21F	PLAT FLE	LI	w.w.																		22.9	7.0	33.6	48.9		121.0	1.2	-?	?	19			
33B	PLAT FLE	MU	w.w.										2.0	4.0	1.8	1.0	0.5	1.4	1.4	1.0	0.9	5.3	1.1	1.8	1.5	1.3	0.7	no	--	no	18		
53B	PLAT FLE	MU	w.w.										27.4	33.2	14.2	1.5		0.8	3.2	2.7	2.2	2.7	3.0	2.7	2.0	1.1	2.0	no	DY	no	19		
67B	PLAT FLE	MU	w.w.																0.8			1.8	1.7	1.5	1.0	0.8	0.8	0.3	no	--	no	14	
21F	PLAT FLE	MU	w.w.																		0.8	0.2	0.6	1.3		1.6	no	-?	?	20			
36F	LIMA LIM	LI	w.w.										301.0	217.0	339.0	418.0	404.0	433.0	386.0	387.0	236.0	412.0	838.0	527.0	297.0	272.0	538.0	1.1	--	no	13		
15F	LIMA LIM	LI	w.w.										124.0		58.2	77.0	74.0	62.5	64.5	51.1	69.6		106.0	50.2	49.8	66.8	no	--	no	12			
22F	LIMA LIM	LI	w.w.										170.0	127.0	140.0		60.0	88.7									no	-?	?	11			
36F	LIMA LIM	MU	w.w.										2.8	7.1	5.6	7.8	5.9	8.2	9.6	5.2	9.4	3.9	8.4	7.7	6.6	3.7	4.3	no	--	no	13		
15F	LIMA LIM	MU	w.w.										3.7		0.8	0.4	1.1	1.8	1.3	1.4	1.0			1.2	0.7	0.9	0.3	no	--	no	18		
22F	LIMA LIM	MU	w.w.										2.0	5.0	3.8		1.2	5.1									1	-?	?	20			
98F	LIMA LIM	MU	w.w.															0.8	1.3	3.3						no	-?	?	9				
30F	PLEU PLA	LI	w.w.												313.0		207.0	216.0									4.3	-?	?	8			
22F	PLEU PLA	LI	w.w.																21.0	20.1	14.5						no	-?	?	7			
98F	PLEU PLA	LI	w.w.													9.4		37.5		27.8	24.1	24.7	40.8	25.5	10.3	10.5	8.4	no	--	no	15		
10F	PLEU PLA	LI	w.w.																42.1		62.8	45.0	24.9	86.2	16.6	22.3	no	--	no	18			
30F	PLEU PLA	MU	w.w.													6.8		3.0	1.5								no	-?	?	9			
22F	PLEU PLA	MU	w.w.																1.4	1.0	3.3						1.7	-?	?	19			
98F	PLEU PLA	MU	w.w.															0.5		2.5		0.6	0.8	0.4	1.5	0.6	0.3	0.3	0.2	no	--	no	19
10F	PLEU PLA	MU	w.w.																	1.0		2.6	1.8	1.1	1.0	0.5	0.2	no	DY	no	16		
67B	LEPI WHI	LI	w.w.										111.0	100.0	143.0	101.0	172.0	166.0	97.2	91.5	118.0	82.3	83.9	63.8	105.0	85.2	73.3	m	--	m	10		
67B	LEPI WHI	MU	w.w.										0.8	0.9	1.4	0.6	0.5	1.5	0.4	0.7	0.4	1.0	0.8	0.7	0.4	0.4	0.5	m	--	m	15		

Annual median concentration of DDEPP (ppb)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30A	MYTI EDU	SB	d.w.												5.24	3.86	7.08	5.70	2.56	5.88	3.87	5.91	3.47	1.99	1.97	2.08	1.22	no	D-	no	13	
31A	MYTI EDU	SB	d.w.												3.30	1.89	3.45	1.84	0.51	3.37	3.49	5.47	1.19	2.10	1.79	1.01	1.25	no	--	no	19	
35A	MYTI EDU	SB	d.w.												4.91	2.08	3.13	2.84	0.57	3.91	3.73	5.93	1.61	3.29	2.17	1.80	2.94	no	--	no	19	
36A	MYTI EDU	SB	d.w.												2.76	1.06	1.03	1.76	0.44	2.11	1.79	2.98	1.48	1.51	1.34	0.76	1.47	no	--	no	17	
71A	MYTI EDU	SB	d.w.												2.61	1.58	3.21	1.29	0.74	1.02	2.20	2.41	2.26	3.58	1.10	1.67	0.76	no	--	no	15	
76A	MYTI EDU	SB	d.w.												1.40	0.79			0.36	1.21	2.29	2.49	0.78	0.83	0.75	0.62	1.10	no	--	no	18	
15A	MYTI EDU	SB	d.w.												0.98	1.72	0.74	0.29	1.02	1.41	2.05	0.54	0.62	0.85	0.67	0.86	no	--	no	18		
51A	MYTI EDU	SB	d.w.														33.90	6.67	14.70	17.10	13.20	16.90	5.48	9.52	10.00	10.50	1	--	no	17		
52A	MYTI EDU	SB	d.w.												12.30	25.50	19.40	18.50	9.53	13.10	16.70	13.70	11.90	6.47	6.82	8.86	10.50	1.1	--	no	12	
56A	MYTI EDU	SB	d.w.												50.00	47.50	115.00	40.80	33.90	72.30	52.60	39.80	26.20	60.60	40.00	55.10	49.30	4.9	--	6.4	14	
57A	MYTI EDU	SB	d.w.												25.90	18.30	35.00	25.30	15.80	50.00	82.90	35.20	27.50	24.70	14.70	27.80	16.60	1.7	--	1	15	
63A	MYTI EDU	SB	d.w.												12.90	9.29	9.68	8.36	5.53	13.00	15.50	11.40	10.20	7.09	4.76	11.30	3.82	no	--	no	14	
65A	MYTI EDU	SB	d.w.												7.60	5.19	7.79	4.12	5.00	6.90	11.90	7.38	6.76	5.43	3.61	6.47	2.55	no	--	no	13	
69A	MYTI EDU	SB	d.w.												3.55	3.16	3.54	2.91	0.40	3.69	6.52	2.61	2.70	2.25	1.61	2.62	0.91	no	--	no	21	
22A	MYTI EDU	SB	d.w.												2.22	1.31	1.88	1.45	0.39	1.37	5.11	1.96	1.49	0.91	0.73	1.46	0.86	no	--	no	19	
84A	MYTI EDU	SB	d.w.												3.13	2.23			0.99	0.74						0.51	no	-?	?	15		
25A	MYTI EDU	SB	d.w.												1.29	1.03										0.88	no	-?	?	8		
28A	MYTI EDU	SB	d.w.												1.11	0.86										0.38	no	-?	?	7		
92A	MYTI EDU	SB	d.w.												0.68	2.09	1.41	0.77	0.28	1.93							no	--	no	22		
96A	MYTI EDU	SB	d.w.												1.03	0.44										0.26	no	-?	?	17		
98A	MYTI EDU	SB	d.w.												5.81	2.29					1.59	1.87	0.87	0.58	1.31	0.63	0.73	0.42	no	D-	no	14
98X	MYTI EDU	SB	d.w.												0.62	0.87			31.60	22.90	5.16						0.25	no	-?	?	15	
99A	MYTI EDU	SB	d.w.												0.62	0.87			0.62	0.42	0.29	0.61					no	-?	?	12		
41A	MYTI EDU	SB	d.w.												0.62	0.42	0.29	0.61								no	-?	?	15			
44A	MYTI EDU	SB	d.w.												0.49	0.34	1.41									no	-?	?	20			
46A	MYTI EDU	SB	d.w.												1.05	0.76	0.27									no	-?	?	11			
48A	MYTI EDU	SB	d.w.												1.71	1.13	0.29									no	-?	?	14			
10A	MYTI EDU	SB	d.w.												0.85	0.78	0.44	1.49		1.45	0.61	0.87	0.61	0.58	0.27	no	--	no	15			
11A	MYTI EDU	SB	d.w.												1.30	1.88	0.41	1.23		0.81	1.04	1.04	0.77	0.76	0.47	0.47	0.32	no	-?	?	21	
11X	MYTI EDU	SB	d.w.																							no	D-	no	9			

Annual median concentration of DDEPP (ppb)

St	Species	tis	Base	ANALYSIS																														
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
I021	MYTI EDU	SB	d.w.															4.60	1.45	4.80	3.25	5.19	2.73	4.73			no	--	no	17				
I022	MYTI EDU	SB	d.w.															3.95	1.38	7.13	4.58	7.96	4.92	3.73	4.51	2.54	2.70	no	--	no	16			
I023	MYTI EDU	SB	d.w.															1.81	1.32	3.79	2.32	6.10	2.39	2.91	3.31	1.42	1.29	no	--	no	15			
I024	MYTI EDU	SB	d.w.															3.50	3.52	8.91	7.17	8.96	4.94	2.52	5.15	2.40	2.32	no	--	no	14			
30A	MYTI EDU	SB	d.w.															5.24	3.86	7.08	5.70	2.56	5.88	3.87	5.91	3.47	1.99	1.97	2.08	1.22	no	D-	no	13
I301	MYTI EDU	SB	d.w.															2.59	3.75	17.80	5.96	7.45	5.58	4.51	5.06	3.54	4.47	no	--	no	16			
I304	MYTI EDU	SB	d.w.															2.14	0.75	3.42	3.89		1.95	2.71	2.62	1.73	2.71	no	--	no	17			
I306	MYTI EDU	SB	d.w.															1.84	0.46	4.25	3.31		2.37	1.88	1.92	1.12	1.74	no	--	no	19			
I307	MYTI EDU	SB	d.w.															2.18	1.03	3.42	2.74		2.12	4.13	2.28	1.09	1.34	no	--	no	15			
I711	MYTI EDU	SB	d.w.															3.46	0.72	1.49	2.19	2.18	3.85		1.26			no	--	no	19			
I712	MYTI EDU	SB	d.w.															2.43	1.34	3.14	3.09	3.49	3.46	2.48	1.60	2.65	2.94	no	--	no	12			
I713	MYTI EDU	SB	d.w.																					1.61	2.26	2.96	no	U?	?	<=5				
I131	MYTI EDU	SB	d.w.															1.46	0.69	1.89	2.06	1.67	1.11	0.92	1.11	0.94	1.37	no	--	no	13			
I132	MYTI EDU	SB	d.w.															1.88	1.36	2.15	2.02	2.06	1.19	1.15	1.26	1.17	0.92	no	D-	no	10			
I133	MYTI EDU	SB	d.w.															2.16	0.88	1.62	1.93	2.73	1.16	1.11	1.03	0.93	0.94	no	--	no	13			
I241	MYTI EDU	SB	d.w.															6.40	6.21	2.30	6.49	5.59	4.45	2.93	4.40	4.37	5.49	no	--	no	14			
I242	MYTI EDU	SB	d.w.															6.52	9.74	1.58	3.53	9.47	3.52	2.22	2.88	3.38	4.41	no	--	no	19			
I243	MYTI EDU	SB	d.w.															7.47	6.12	1.72	5.43	5.11	4.01	1.99	3.32	3.88	6.41	no	--	no	16			
51A	MYTI EDU	SB	d.w.																33.90	6.67	14.70	17.10	13.20	16.90	5.48	9.52	10.00	10.50	1	--	no	17		
52A	MYTI EDU	SB	d.w.															12.30	25.50	19.40	18.50	9.53	13.10	16.70	13.70	11.90	6.47	6.82	8.86	10.50	1.1	--	no	12
10A	MYTI EDU	SB	d.w.																0.85	0.78	0.44	1.49		1.45	0.61	0.87	0.61	0.58	0.27	no	--	no	15	

Annual median concentration of DDEPP (ppb)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30B	GADU MOR	LI	w.w.										163.0	440.0	182.0	159.0	191.0	194.0	312.0	383.0	260.0	230.0	160.0	180.0	180.0	240.0	210.0	1.1	--	1.2	12	
36B	GADU MOR	LI	w.w.										91.9	51.0	50.0	75.0	55.0	105.0	141.0	129.0	45.0	86.0	47.0	46.0	39.0	32.0	34.0	no	D-	no	13	
15B	GADU MOR	LI	w.w.										50.0	136.0	48.0	57.0	86.0	33.5	75.0	140.0	72.5	76.0	46.0	60.0	78.0	74.0	50.0	no	--	no	15	
53B	GADU MOR	LI	w.w.										637.0	806.0	939.0	85.0		42.0	491.0	936.0	490.0	160.0	380.0	260.0	200.0	145.0	300.0	1.5	--	no	24	
67B	GADU MOR	LI	w.w.										776.0	554.0	347.0	392.0	471.0	109.0	460.0	2060.0	270.0	200.0	177.0	140.0	110.0	89.0	74.0	no	--	no	20	
23B	GADU MOR	LI	w.w.										68.0	85.4	42.0	41.0	35.0	31.0	49.0	33.0	49.0	48.0	59.0	52.9	24.0	37.0	52.0	no	--	no	12	
92B	GADU MOR	LI	w.w.																									50.0	no	-?	?	20
98B	GADU MOR	LI	w.w.																									no	--	no	16	
43B	GADU MOR	LI	w.w.																									no	-?	?	9	
10B	GADU MOR	LI	w.w.																									no	--	no	14	
30B	GADU MOR	MU	w.w.										0.5	1.2	2.0	1.0	0.3	0.3	1.0	1.0	1.5	1.5	0.4	0.7	0.7	0.7	0.7	no	--	no	18	
36B	GADU MOR	MU	w.w.										0.3	0.3	0.2	0.5	0.1	0.9	0.6	0.9	0.3	0.3	0.2	0.2	0.2	0.2	0.3	no	--	no	18	
15B	GADU MOR	MU	w.w.										0.5	0.4	0.3	0.2	0.1	0.3	0.4	0.5	0.5	0.2	0.3	0.3	0.2	0.2	0.4	0.1	no	--	no	16
53B	GADU MOR	MU	w.w.										2.4	2.2	6.8	1.8		0.1	4.1	4.6	4.6	3.2	2.5	0.6	1.8	2.4	1.9	1.9	--	1.9	>25	
67B	GADU MOR	MU	w.w.										2.3	3.0	1.4	1.0	2.5	1.1	7.0	19.0	1.0	1.1	1.1	1.8	0.4	0.2	0.3	no	--	no	23	
23B	GADU MOR	MU	w.w.										0.2	0.6	0.1	0.2	0.0	0.2	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.1	no	--	no	17	
92B	GADU MOR	MU	w.w.																									0.1	no	-?	?	21
98B	GADU MOR	MU	w.w.																									no	--	no	23	
43B	GADU MOR	MU	w.w.																									no	-?	?	9	
10B	GADU MOR	MU	w.w.																									no	--	no	17	

Annual median concentration of DDEPP (ppb)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
33B	PLAT FLE	LI	w.w.										13.0	9.1	24.0	14.0	13.0	7.0	10.2	9.7	8.6	6.8	27.0	27.0	17.0	22.0	19.0	no	--	no	14
53B	PLAT FLE	LI	w.w.										94.0	70.1	32.0	41.0		8.0	25.0	45.0	38.0	44.0	17.5	39.0	42.0	40.0	29.0	no	--	1.3	16
67B	PLAT FLE	LI	w.w.															27.0		84.5	40.0	35.0	25.0	24.0	20.0	21.0	no	--	no	14	
21F	PLAT FLE	LI	w.w.																	11.0	2.6	16.0	7.4		90.0	3	-?	?	25		
33B	PLAT FLE	MU	w.w.										0.9	1.9	0.6	0.2	0.2	0.3	0.4	0.3	0.2	1.5	0.3	0.6	0.4	0.5	0.3	no	--	no	19
53B	PLAT FLE	MU	w.w.										4.7	5.3	3.8	1.3		0.4	1.8	1.4	1.0	0.9	0.6	0.9	0.7	0.8	0.6	no	D-	no	16
67B	PLAT FLE	MU	w.w.																0.9		1.3	1.4	1.2	0.5	0.6	0.7	0.1	no	--	no	17
21F	PLAT FLE	MU	w.w.																	0.3	0.1	0.4	0.2		1.2	1.2	-?	?	23		
36F	LIMA LIM	LI	w.w.										28.0	34.4	28.0	21.0	50.0	40.0	40.0	22.0	18.0	52.0	45.0	27.0	31.0	36.0	17.0	no	--	no	14
15F	LIMA LIM	LI	w.w.										39.0		13.4	23.5	9.0	20.7	20.0	13.0	32.0		41.0	15.0	17.0	23.0	no	--	no	16	
22F	LIMA LIM	LI	w.w.										68.9	48.0	39.9		21.0	9.2									no	D?	?	10	
36F	LIMA LIM	MU	w.w.										0.4	1.2	0.7	0.5	1.0	0.9	0.9	0.5	0.7	0.5	0.5	0.5	0.6	0.5	0.1	no	--	no	15
15F	LIMA LIM	MU	w.w.											1.2		0.2	0.1	0.3	0.6	0.4	0.4	0.3		0.6	0.2	0.3	0.1	no	--	no	19
22F	LIMA LIM	MU	w.w.										1.1	2.0	1.2			0.6	0.8								no	-?	?	14	
98F	LIMA LIM	MU	w.w.															0.6	0.3	1.6						no	-?	?	23		
30F	PLEU PLA	LI	w.w.												21.2		13.0	12.0										1.2	-?	?	6
22F	PLEU PLA	LI	w.w.																7.8	12.0	2.8						no	-?	?	21	
98F	PLEU PLA	LI	w.w.													3.0		8.0		15.5	6.2	7.8	10.8	8.0	5.1	3.6	2.8	no	--	no	15
10F	PLEU PLA	LI	w.w.																15.0		34.7	28.0	8.9	19.0	4.7	5.8	no	--	no	18	
30F	PLEU PLA	MU	w.w.													0.7		0.3	0.2		0.5	0.3	0.8					no	-?	?	8
22F	PLEU PLA	MU	w.w.																0.5	0.3	0.8						no	-?	?	15	
98F	PLEU PLA	MU	w.w.															0.1		0.5	0.3	0.2	0.1	0.1	0.1	no	--	no	18		
10F	PLEU PLA	MU	w.w.																0.4		0.9	1.1	0.3	0.4	0.1	0.1	no	D-	no	18	
67B	LEPI WHI	LI	w.w.										294.0	240.0	183.0	163.0	250.0	145.0	143.0	167.0	160.0	160.0	130.0	58.0	64.0	73.0	71.1	m	D-	m	10
67B	LEPI WHI	MU	w.w.										2.6	1.5	2.5	0.8	0.8	3.0	0.8	1.3	0.6	1.4	1.1	0.5	0.4	0.6	0.5	m	--	m	17

Annual median concentration of HCB (ppb)

Cursive values in shaded area indicate temporal trend analysis based data since 1990

St	Species	tis	Base																					ANALYSIS								
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30A	MYTI EDU	SB	d.w.		1.18	0.88	2.06	0.92	1.15	0.87	0.35	0.59	0.95	0.54	0.27	0.24	0.25	0.28	0.30	0.36	0.23	0.34	0.32	0.24	no	D-	no	13				
31A	MYTI EDU	SB	d.w.		13.40	1.38	3.83	1.89	0.93	0.89	0.36	0.32	0.61	0.55	0.45	0.24	0.31	0.22	0.26	0.21	0.23	0.27	0.32	0.33	0.22	no	DY	no	15			
35A	MYTI EDU	SB	d.w.		12.80	0.95	3.33	0.79	0.98	1.12	0.47	0.42	0.59	0.58	0.51	0.23	0.28	0.22	0.52	0.20	0.34	0.36	0.30	0.29	0.27	0.31	no	D-	no	17		
36A	MYTI EDU	SB	d.w.		15.00	0.95	3.83	2.90	2.37	0.96	0.43	0.33	0.55	0.39	0.53	0.24	0.33	0.28	0.31	0.15	0.25	0.20	0.21	0.29	0.22	no	D-	no	17			
71A	MYTI EDU	SB	d.w.		15.30	10.40	91.40	11.10	207.00	1.83	149.00	8.48	6.91	4.14	3.91	1.47	2.13	4.48	2.04	1.78	3.10	1.85	2.42	0.81	0.33	0.61	1.2	D-	no	>25		
71A	MYTI EDU	SB	d.w.									8.48	6.91	4.14	3.91	1.47	2.13	4.48	2.04	1.78	3.10	1.85	2.42	0.81	0.33	0.61	1.2	D-	no	15		
76A	MYTI EDU	SB	d.w.									0.38	0.57	0.50	0.79			0.25	0.29	0.40	0.26	0.22	0.24	0.30	0.31	0.97	1.9	--	2	14		
15A	MYTI EDU	SB	d.w.									0.20				0.49	0.25	0.22	0.29	0.25	0.16	0.22	0.18	0.25	0.30	0.33	0.29	no	--	no	12	
51A	MYTI EDU	SB	d.w.									0.61	0.33	0.31	0.40	0.40	0.39	0.20	0.48	0.40	0.49	no	--	1.1	12							
52A	MYTI EDU	SB	d.w.									0.86	0.38	0.81	0.81	0.28	0.32	0.26	0.33	0.21	0.33	0.20	0.38	0.32	0.46	no	--	1.1	13			
56A	MYTI EDU	SB	d.w.									0.20	0.79	0.41	0.79	0.94	1.04	0.31	0.38	0.31	0.44	0.35	0.70	0.18	0.34	0.30	0.45	0.53	1.1	--	1.2	15
57A	MYTI EDU	SB	d.w.									0.77	0.76	0.72	0.79	0.36	0.30	0.26	0.43	0.58	0.63	0.26	0.36	0.25	0.40	0.24	no	D-	no	12		
63A	MYTI EDU	SB	d.w.									1.05	0.97	0.74	0.63	0.32	0.33	0.33	0.41	0.45	0.51	0.23	0.32	0.33	0.35	0.15	no	D-	no	11		
65A	MYTI EDU	SB	d.w.									0.20	0.43	0.52	0.86	0.62	0.67	0.28	0.30	0.29	0.35	0.38	0.52	0.15	0.27	0.26	0.29	0.18	no	D-	no	13
69A	MYTI EDU	SB	d.w.									0.53	0.53	0.29	0.25	0.29	0.29	0.50	0.48	0.46	0.36	0.21	0.33	0.31	0.27	0.25	no	--	no	12		
22A	MYTI EDU	SB	d.w.									0.27	0.61	0.56	0.44	0.25	0.25	0.30	0.31	0.17	0.32	0.20	0.30	0.24	0.33	0.22	no	--	no	12		
82A	MYTI EDU	SB	d.w.		2.26	10.70	0.66	0.62	0.80	0.54															0.18	no	--	no	24			
84A	MYTI EDU	SB	d.w.		3.41	8.79	3.33	2.04	1.23	0.48		0.51	0.63	0.53		0.25	0.22								0.29	no	DY	no	15			
87A	MYTI EDU	SB	d.w.									0.92	0.42												0.14	no	-?	?	16			
25A	MYTI EDU	SB	d.w.												0.70	0.51									0.35	no	-?	?	9			
28A	MYTI EDU	SB	d.w.												0.56	0.43									0.16	no	-?	?	7			
92A	MYTI EDU	SB	d.w.												0.68	0.42	0.24	0.23	0.23	0.25						no	D-	no	12			
96A	MYTI EDU	SB	d.w.												0.53	0.42									0.15	no	-?	?	6			
98A	MYTI EDU	SB	d.w.												0.62	0.57									0.15	no	--	no	9			
98X	MYTI EDU	SB	d.w.													0.56	0.32	0.26								no	-?	?	8			
99A	MYTI EDU	SB	d.w.												0.62	0.44									0.15	no	-?	?	8			
41A	MYTI EDU	SB	d.w.													0.29	0.26	0.29	0.30						no	-?	?	6				
43A	MYTI EDU	SB	d.w.													0.33	0.34		0.45						no	-?	?	<=5				
44A	MYTI EDU	SB	d.w.													0.27	0.29	0.33							no	-?	?	<=5				
46A	MYTI EDU	SB	d.w.													0.26	0.29	0.27							no	-?	?	6				
48A	MYTI EDU	SB	d.w.													0.28	0.29	0.24							no	-?	?	7				
10A	MYTI EDU	SB	d.w.													0.29	0.27	0.25	0.31		0.28	0.28	0.29	0.31	0.28	0.28	0.28	no	--	no	6	
11A	MYTI EDU	SB	d.w.													0.35	0.43	0.34	0.46		0.34	0.21	0.25	0.24	0.25	0.24	0.28	no	-?	?	8	
11X	MYTI EDU	SB	d.w.																	0.34	0.21	0.25	0.24	0.25	0.24	0.28	no	--	no	8		

Annual median concentration of HCB (ppb)

Cursive values in shaded area indicate temporal trend analysis based data since 1990

St	Species	tis	Base	ANALYSIS																																						
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR											
I021	MYTI EDU	SB	d.w.														0.83	0.92	0.48	0.38	0.48	0.64	0.55			1.1	--	1.1	11													
I022	MYTI EDU	SB	d.w.														0.42	0.48	0.97	0.31	0.78	0.46	0.54	0.55	0.38	0.51	1	--	no	14												
I023	MYTI EDU	SB	d.w.														0.48	0.42	0.43	0.26	0.62	0.35	0.34	0.42	0.39	0.45	no	--	no	11												
I024	MYTI EDU	SB	d.w.														0.49	0.60	1.16	0.43	0.66	0.56	0.54	0.50	0.39	0.40	no	--	no	12												
30A	MYTI EDU	SB	d.w.														0.24	0.25	0.28	0.30	0.36	0.23	0.34	0.32	0.24		no	--	no	13												
I301	MYTI EDU	SB	d.w.														0.29	0.28	0.70	0.82	1.55	0.51	0.68	0.42	0.32	0.97	1.9	--	no	16												
I304	MYTI EDU	SB	d.w.														0.34	0.28	0.72	0.49	0.29	0.53	0.39	0.31	0.50	no	--	no	13													
I306	MYTI EDU	SB	d.w.														0.30	0.31	0.77	0.25	0.30	0.29	0.34	0.35	0.46	no	--	1	14													
I307	MYTI EDU	SB	d.w.														0.27	0.32	0.67	0.17	0.33	0.34	0.45	0.37	0.36	no	--	no	15													
I711	MYTI EDU	SB	d.w.														4.45	5.54	0.58	4.46	6.96	2.56	4.00			8	--	11.3	24													
I712	MYTI EDU	SB	d.w.														3.43	16.40	7.90	4.83	5.00	3.31	1.78	2.75	3.27	2.90	5.8	--	5.3	16												
I713	MYTI EDU	SB	d.w.																				3.49	2.64	4.17	8.3	-?	?	12													
71A	MYTI EDU	SB	d.w.														15.30	10.40	91.40	11.10	207.00	1.83	149.00	8.48	6.91	4.14	3.91	1.47	2.13	4.48	2.04	1.78	3.10	1.85	2.42	0.81	0.33	0.61	1.2	D-	no	>25
71A	MYTI EDU	SB	d.w.															8.48	6.91	4.14	3.91	1.47	2.13	4.48	2.04	1.78	3.10	1.85	2.42	0.81	0.33	0.61	1.2	D-	no	15						
I131	MYTI EDU	SB	d.w.															0.32	0.30	0.27	0.20	0.58	0.29	0.33	0.29	0.37	1.40	2.8	--	2.4	15											
I132	MYTI EDU	SB	d.w.															52.80	89.70	40.20	44.20	1.89	4.73	3.11	2.36	1.56	1.94	3.9	D-	no	21											
I133	MYTI EDU	SB	d.w.															18.10	43.50	8.12	28.00	1.70	6.18	2.30	1.62	2.45	3.76	7.5	--	4.7	22											
I241	MYTI EDU	SB	d.w.															1.28	0.71	0.69	0.75	0.70	0.62	0.29	0.55	0.30	1.14	2.3	--	1.6	15											
I242	MYTI EDU	SB	d.w.															1.20	0.92	0.56	0.60	0.65	0.55	0.24	0.50	0.46	0.59	1.2	--	1.1	12											
I243	MYTI EDU	SB	d.w.															1.03	0.66	0.52	1.24	0.66	0.67	0.26	0.44	0.34	0.61	1.2	--	no	14											
51A	MYTI EDU	SB	d.w.															0.61	0.33	0.31	0.40	0.40	0.39	0.20	0.48	0.40	0.49	no	--	1.1	12											
52A	MYTI EDU	SB	d.w.															0.86	0.38	0.81	0.81	0.28	0.32	0.21	0.33	0.20	0.38	0.32	0.46	no	--	1.1	13									
10A	MYTI EDU	SB	d.w.																0.29	0.27	0.25	0.31		0.28	0.28	0.29	0.31	0.28	0.28	no	--	no	6									

Annual median concentration of HCB (ppb)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
30B	GADU MOR	LI	w.w.										10.00	17.00	7.48	16.00	11.00	11.00	12.00	7.00	5.30	5.10	9.10	8.90	6.70	6.00	8.90	no	--	no	12	
36B	GADU MOR	LI	w.w.										7.00	9.00	9.00	10.00	9.00	5.00	9.00	6.00	4.40	6.50	5.40	4.60	3.10	3.30	4.00	no	D-	no	10	
15B	GADU MOR	LI	w.w.										5.00	20.50	10.00	14.00	14.00	9.00	11.00	13.00	11.50	11.00	6.20	6.60	8.20	6.40	9.70	no	--	no	13	
53B	GADU MOR	LI	w.w.										10.00	10.00	16.50	7.00		5.00	7.00	7.00	5.00	4.70	12.00	2.10	3.00	2.25	2.60	no	D-	no	15	
67B	GADU MOR	LI	w.w.										14.00	8.00	7.94	8.00	8.49	10.00	8.00	15.50	9.90	4.60	5.63	4.90	4.60	5.10	5.30	no	D-	no	11	
23B	GADU MOR	LI	w.w.										6.00	9.49	12.00	9.00	8.00	6.00	10.00	6.00	8.40	7.80	7.60	9.25	4.70	7.90	5.80	no	--	no	11	
92B	GADU MOR	LI	w.w.													17.00	11.00	14.00	13.00									9.80	no	-?	?	9
98B	GADU MOR	LI	w.w.													20.00	9.95	12.00	18.00	35.00	20.50	16.00	13.00	3.10	2.60	10.00	13.00	11.00	no	--	no	17
43B	GADU MOR	LI	w.w.															15.00	16.50	13.00								no	-?	?	8	
10B	GADU MOR	LI	w.w.															13.00	11.00	16.00	17.00	17.00	25.00	9.00	9.90	11.00	9.43	5.10	no	D-	no	12
30B	GADU MOR	MU	w.w.										0.09	0.09	0.10	0.10	0.04	0.03	0.05	0.05	0.06	0.06	0.06	0.05	0.06	0.05	0.06	no	--	no	11	
36B	GADU MOR	MU	w.w.										0.11	0.07	0.10	0.10	0.04	0.05	0.06	0.06	0.05	0.06	0.04	0.05	0.03	0.03	0.03	no	D-	no	11	
15B	GADU MOR	MU	w.w.										0.11	0.11	0.10	0.10	0.06	0.07	0.08	0.07	0.10	0.06	0.10	0.04	0.06	0.07	0.05	no	--	no	11	
53B	GADU MOR	MU	w.w.										0.10	0.03	0.10	0.10		0.03	0.06	0.06	0.05	0.05	0.09	0.04	0.05	0.08	0.03	no	--	no	17	
67B	GADU MOR	MU	w.w.										0.10	0.08	0.10	0.10	0.07	0.06	0.05	0.07	0.06	0.05	0.05	0.04	0.05	0.05	0.04	no	D-	no	8	
23B	GADU MOR	MU	w.w.										0.08	0.08	0.10	0.10	0.04	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.04	0.05	0.03	no	D-	no	10	
92B	GADU MOR	MU	w.w.														0.10	0.07	0.05	0.09								0.07	no	-?	?	12
98B	GADU MOR	MU	w.w.													0.20	0.20	0.07	0.10	0.11	0.10	0.10	0.08	0.10	0.06	0.07	0.09	0.08	no	--	no	11
43B	GADU MOR	MU	w.w.															0.09	0.13	0.06								no	-?	?	15	
10B	GADU MOR	MU	w.w.															0.16	0.11	0.09	0.20	0.17	0.26	0.09	0.11	0.13	0.09	0.09	no	--	no	13

Annual median concentration of HCB (ppb)

St	Species	tis	Base	ANALYSIS																												
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR	
33B	PLAT FLE	LI	w.w.										1.00	0.50	5.00	2.00	1.00	1.00	0.60	0.80	0.59	0.54	1.60	1.60	1.90	1.20	no	--	no	18		
53B	PLAT FLE	LI	w.w.										6.00	4.47	5.00	2.00		1.00	2.00	3.00	1.80	2.50	2.39	2.00	2.90	1.40	1.00	no	D-	no	14	
67B	PLAT FLE	LI	w.w.															3.00		6.39	3.60	4.20	4.30	3.50	3.70	3.10	no	--	no	11		
21F	PLAT FLE	LI	w.w.																	3.10	1.10	2.40	0.90		3.60	no	-?	?	20			
33B	PLAT FLE	MU	w.w.										0.06	0.07	0.10	0.10	0.03	0.03	0.03	0.05	0.03	0.06	0.04	0.04	0.05	0.06	0.03	no	--	no	14	
53B	PLAT FLE	MU	w.w.										0.45	0.30	0.20	0.10		0.08	0.05	0.10	0.06	0.06	0.09	0.06	0.05	0.13	0.03	no	D-	no	14	
67B	PLAT FLE	MU	w.w.															0.05		0.10	0.19	0.16	0.12	0.14	0.12	0.03	no	--	no	16		
21F	PLAT FLE	MU	w.w.																0.10	0.03	0.06	0.04		0.09	no	-?	?	18				
36F	LIMA LIM	LI	w.w.										5.48	3.00	5.00	2.00	3.00	2.00	2.30	3.00	1.10	2.50	3.00	2.60	2.00	2.50	1.80	no	--	no	13	
15F	LIMA LIM	LI	w.w.											4.00		4.00	4.00	2.00	3.00	3.20	3.00	3.64		5.90	2.50	4.30	3.10	no	--	no	12	
22F	LIMA LIM	LI	w.w.										6.00	3.00	5.00		1.00	1.41									no	-?	?	15		
36F	LIMA LIM	MU	w.w.										0.10	0.09	0.10	0.10	0.06	0.06	0.07	0.05	0.05	0.05	0.05	0.06	0.06	0.05	0.03	no	D-	no	9	
15F	LIMA LIM	MU	w.w.											0.20		0.10	0.04	0.07	0.09	0.07	0.09	0.09	0.08		0.15	0.04	0.09	0.03	no	--	no	16
22F	LIMA LIM	MU	w.w.										0.12	0.20	0.10		0.05	0.07									no	-?	?	14		
98F	LIMA LIM	MU	w.w.														0.07	0.13	0.16									no	-?	?	9	
30F	PLEU PLA	LI	w.w.											5.00		2.00	2.00											no	-?	?	11	
22F	PLEU PLA	LI	w.w.															0.50	0.90	0.30									no	-?	?	19
98F	PLEU PLA	LI	w.w.												1.00		1.00		1.74	1.00	2.50	1.30	1.80	0.96	0.68	1.20	no	--	no	13		
10F	PLEU PLA	LI	w.w.															6.10		8.77	6.40	2.40	1.60	2.15	1.99	no	D-	no	14			
30F	PLEU PLA	MU	w.w.											0.14		0.05	0.03		0.03	0.03	0.04							no	D?	?	<=5	
22F	PLEU PLA	MU	w.w.															0.03	0.05	0.04		0.07	0.07	0.04	0.04	0.04	0.03	no	-?	?	7	
98F	PLEU PLA	MU	w.w.														0.10		0.13	0.05	0.04	0.07	0.07	0.04	0.04	0.04	0.03	no	--	no	12	
10F	PLEU PLA	MU	w.w.															0.22		0.30	0.49	0.15	0.43	0.06	0.03	no	--	no	20			
67B	LEPI WHI	LI	w.w.										9.00	4.00	5.00	4.00	5.00	2.00	4.60	4.00	5.00	2.80	4.80	3.40	3.80	3.90	3.45	m	--	m	12	
67B	LEPI WHI	MU	w.w.										0.09	0.07	0.10	0.10	0.03	0.04	0.03	0.07	0.03	0.04	0.05	0.03	0.04	0.03	m	--	m	13		

JAMP National Comments 2004 - Norway

---

Annual median concentration of BAP (ppb)

St	Species	tis	Base	ANALYSIS																													
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR		
30A	MYTI EDU	SB	d.w.															2.5										no	--	no	8		
I301	MYTI EDU	SB	d.w.																4.4	19.3	18.8	6.0	13.1	2.6	9.8	3.1	3.2	34.2	6.8	--	3.6	24	
I304	MYTI EDU	SB	d.w.																3.4	2.8	3.3	3.4	2.8	2.9	3.8	3.9	3.1	5.3	1.1	--	1.1	9	
I306	MYTI EDU	SB	d.w.																3.0	3.1	2.9	3.1	3.1	3.0	2.9	3.4	3.5	4.8	no	UY	1.1	6	
I307	MYTI EDU	SB	d.w.																2.7	3.2	2.8	2.9	3.0	3.3	3.4	4.5	3.7	5.3	1.1	U-	1.2	7	
I131	MYTI EDU	SB	d.w.																3.3	2.7	2.7	3.6	5.0	2.4	3.3	2.8	3.7	8.0	1.6	--	1.6	12	
I132	MYTI EDU	SB	d.w.																71.8	89.0	18.6	22.6	300.0	10.8	32.7	49.6	89.7	52.4	10.5	--	16.8	>25	
I133	MYTI EDU	SB	d.w.																80.6	13.7	51.7	18.6		8.5	19.0	23.7	39.3	135.0	27	--	31.7	19	
I201	MYTI EDU	SB	d.w.																93.2	207.0	679.0	10.5	83.8	47.4	31.7	188.0	7.2	3.8	no	--	no	>25	
I205	MYTI EDU	SB	d.w.																7.4		23.1	64.5	7.5	5.6	7.6	33.0	3.2	48.2	9.6	--	6.4	>25	
I913	MYTI EDU	SB	d.w.																			5.9	15.6	3.0	13.3	3.7	1.4	no	--	no	22		
I912	MYTI EDU	SB	d.w.																7.0		9.5	5.4	16.4	135.0	4.2	20.0	4.0	1.5	no	--	no	>25	
I965	MYTI EDU	SB	d.w.																						233.0	30.8	43.6	87.7	17.5	-?	?	25	
I962	MYTI EDU	SB	d.w.																246.0	33.5		87.0								17.4	-?	?	>25
I964	MYTI EDU	SB	d.w.																							37.3	289.0	251.0	50.3	-?	?	23	
I969	MYTI EDU	SB	d.w.																14.2	10.7	17.6	8.4	10.3	17.1	23.5	3.7	46.7	34.7	6.9	--	9.3	21	

Annual median concentration of PK\_S (ppb)

St	Species	tis	Base	ANALYSIS																														
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
30A	MYTI EDU	SB	d.w.																38										no	--	no	16		
I301	MYTI EDU	SB	d.w.																73	257	197	100	175	32	112	44	38	313	6.3	--	3.5	22		
I304	MYTI EDU	SB	d.w.																14	17	14	37	33	8	21	10	8	70	1.4	--	no	21		
I306	MYTI EDU	SB	d.w.																13	15	35	31	28	7	28	12	9	53	1.1	--	no	20		
I307	MYTI EDU	SB	d.w.																12	19	20	28	42	12	27	29	10	74	1.5	--	no	19		
I131	MYTI EDU	SB	d.w.																48	18	35	61	57	27	28	16	27	127	2.5	--	1.6	19		
I132	MYTI EDU	SB	d.w.																594	586	281	581	2730	243	389	783	1520	570	11.4	--	20.3	23		
I133	MYTI EDU	SB	d.w.																602	121	647	287		150	339	476	580	1200	24	--	34	19		
I201	MYTI EDU	SB	d.w.																705	1590	7970	281	999	903	638	189	111	2.2	--	no	>25			
I205	MYTI EDU	SB	d.w.																96	470	1380	197	187	189	1680	98	808	16.2	--	13.9	>25			
I913	MYTI EDU	SB	d.w.																			107	604	77	405	29	21	no	--	no	>25			
I912	MYTI EDU	SB	d.w.																109		342	195	187	1560	59	412	28	11	no	--	no	>25		
I965	MYTI EDU	SB	d.w.																							1330	431	468	854	17.1	-?	?	18	
I962	MYTI EDU	SB	d.w.																1450	265		665									13.3	-?	?	>25
I964	MYTI EDU	SB	d.w.																								368	2370	2090	41.9	-?	?	21	
I969	MYTI EDU	SB	d.w.																139	108	230	131	285	192	169	40	625	361	7.2	--	9.2	21		

Annual median concentration of P\_S (ppb)

St	Species	tis	Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	ANALYSIS												
																										OC	TRD	SM3	PWR											
30A	MYTI EDU	SB	d.w.																								2.2	--	1.5	17										
I301	MYTI EDU	SB	d.w.																								22	--	8.4	24										
I304	MYTI EDU	SB	d.w.																								2.3	--	1.2	21										
I306	MYTI EDU	SB	d.w.																								1.7	--	no	21										
I307	MYTI EDU	SB	d.w.																								84	275	177	421	82	168	279	80	389	1.6	--	1.1	20	
I131	MYTI EDU	SB	d.w.																								207	255	191	265	360	282	118	133	214	1180	4.7	--	3.1	18
I132	MYTI EDU	SB	d.w.																								2760	2810	1390	1730	7270	1380	1170	1920	2790	2240	8.9	--	9.7	18
I133	MYTI EDU	SB	d.w.																								5690	1770	1960	1150	1080	964	1440	1260	3310	13.2	DY	13.5	13	
I201	MYTI EDU	SB	d.w.																								2660	5210	17100	861	3720	2560	1300	615	716	2.9	--	no	24	
I205	MYTI EDU	SB	d.w.																								614	1770	3540	891	658	509	444	2780	11.1	--	4.7	23		
I913	MYTI EDU	SB	d.w.																								621	2490	657	2250	190	254	1	--	no	24				
I912	MYTI EDU	SB	d.w.																								1100	1530	963	1970	7300	832	2220	280	93	no	--	no	22	
I965	MYTI EDU	SB	d.w.																								6340	1690	1850	3200	854	1480	4090	16.4	-?	?	22			
I962	MYTI EDU	SB	d.w.																								7.4	-?	?	20										
I964	MYTI EDU	SB	d.w.																								771	6970	10400	41.6	-?	?	20							
I969	MYTI EDU	SB	d.w.																								1060	986	747	629	917	1160	824	170	2550	2060	8.2	--	9.4	21

Annual median concentration of TBT (ppm)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
30A	MYTI EDU	SB	d.w.																	2.710	1.510	1.470	0.935	0.802	1.810	1.080	0.723	7.2	--	7.3	13
I301	MYTI EDU	SB	d.w.																		2.590	2.110	2.830		28.3	-?	?	10			
36A	MYTI EDU	SB	d.w.																0.034	0.179	0.217	0.083	0.059	0.079	0.103	0.060	no	--	no	19	
I712	MYTI EDU	SB	d.w.																		1.200	0.912	0.300		3	-?	?	13			
I713	MYTI EDU	SB	d.w.																	1.370	0.668	0.220		2.2	-?	?	8				
71A	MYTI EDU	SB	d.w.																1.020		0.431	0.702	0.375	0.119		1.2	-?	?	17		
76A	MYTI EDU	SB	d.w.															0.188		0.053	0.092	0.106	0.034		no	-?	?	16			
I131	MYTI EDU	SB	d.w.																	0.491	0.120		0.062		no	-?	?	17			
15A	MYTI EDU	SB	d.w.																0.098	0.081	0.062	0.018		no	-?	?	14				
22A	MYTI EDU	SB	d.w.																0.170	0.138	0.587	0.291		2.9	-?	?	18				
227X	MYTI EDU	SB	d.w.																0.375	0.417	0.672	0.709	0.314		3.1	-?	?	14			
98A	MYTI EDU	SB	d.w.																0.108	0.105	0.114	0.047		no	-?	?	13				
10A	MYTI EDU	SB	d.w.																0.022	0.012	0.007			no	D?	?	<=5				
11X	MYTI EDU	SB	d.w.																	0.035	0.020	0.004			no	-?	?	15			

Annual median concentration of TBT (ppm)

St	Species	tis	Base	ANALYSIS																										
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3
36G	NUCE LAP	SB	d.w.																0.105	0.203	0.142	0.095	0.050	0.155	0.085	0.041	m	--	m	16
71G	NUCE LAP	SB	d.w.																0.133	0.344	0.235	0.115		m	-?	m	18			
76G	NUCE LAP	SB	d.w.																0.041	0.196		0.068		m	-?	m	>25			
131G	NUCE LAP	SB	d.w.																0.033	0.064		0.051		m	-?	m	14			
15G	NUCE LAP	SB	d.w.																0.071	0.091	0.065	0.026		m	-?	m	14			
220G	NUCE LAP	SB	d.w.																0.082	0.100	0.085			0.113		m	-?	m	7	
224G	NUCE LAP	SB	d.w.															0.077	0.295		0.120				m	-?	m	24		
226G	NUCE LAP	SB	d.w.																0.844	0.225	0.210				m	-?	m	16		
227G	NUCE LAP	SB	d.w.															0.089	0.625	0.267	0.387	0.135	0.446	0.878	0.258	m	--	m	22	
22G	NUCE LAP	SB	d.w.																0.070	0.101	0.322	0.200		m	-?	m	16			
98G	NUCE LAP	SB	d.w.																0.026	0.063	0.061	0.049		m	-?	m	14			
11G	NUCE LAP	SB	d.w.																	0.013	0.026	0.010			m	-?	m	19		

Annual median concentration of ALAD (NG PBG/MIN/MG PROT)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
30B	GADU MOR	BL	w.w.																	8.98	15.60	13.00	14.60	12.70	10.40	6.91	14.20	m	--	m	12
36B	GADU MOR	BL	w.w.																	13.00	26.20	9.93	22.00	19.40				m	-?	m	15
15B	GADU MOR	BL	w.w.																	17.20	23.40	8.45		18.90				m	-?	m	17
53B	GADU MOR	BL	w.w.																	7.64	10.10	11.10	12.70	10.00	6.44	9.32	9.95	m	--	m	10
67B	GADU MOR	BL	w.w.																	7.17	28.20	16.90	22.40	19.00				m	-?	m	16
23B	GADU MOR	BL	w.w.																	15.80	24.80	18.10	19.80	24.00	19.40	16.80	19.70	m	--	m	9

Annual median concentration of EROD (PMIN/MIN/MG PROT)

St	Species	tis	Base	ANALYSIS																											
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR
30B	GADU MOR	LI	w.w.																	68.8	109.0	70.0	260.0	81.2	158.0	88.3	69.0	m	--	m	16
36B	GADU MOR	LI	w.w.																	95.1	11.4	60.2	64.9	76.2				m	-?	m	24
15B	GADU MOR	LI	w.w.																	49.9	52.3	184.0		61.0				m	-?	m	20
53B	GADU MOR	LI	w.w.																	86.5	119.0	90.1	128.0	34.7	93.9	11.7	20.0	m	--	m	19
67B	GADU MOR	LI	w.w.																	103.0	76.2	84.6	103.0	72.9				m	-?	m	9
23B	GADU MOR	LI	w.w.																	94.1	28.6	70.1	73.5	76.5	103.0	41.9	45.9	m	--	m	16

Annual median concentration of PYR1O ( $\mu$ G/KG/ABS 380 NM)

Cursive values in shaded area indicate earlier and not comparable method. These values are disregarded in temporal trend analyses

St	Species	tis	Base	ANALYSIS																										
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3
30B	GADU MOR	BI	w.w.																	115	130	12	17	29	20	44	m	-?	m	11
15B	GADU MOR	BI	w.w.																	3770	253		30	6	6	17	m	-?	m	23
53B	GADU MOR	BI	w.w.																	83	59	9	4	19	4	3	m	-?	m	21
23B	GADU MOR	BI	w.w.																	13	11	4	3	3	5	2	m	-?	m	16

#### Annual median concentration of VSDI ( )

St	Species	tis	Base	ANALYSIS																														
				1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	OC	TRD	SM3	PWR			
36G	NUCE LAP	WO	w.w.											4.10		3.90				4.00	4.00	4.00	4.00	3.95	4.00	3.96	3.65	m	--	m	7			
71G	NUCE LAP	WO	w.w.																					4.00	4.10	4.00	4.00	m	-?	m	<=5			
76G	NUCE LAP	WO	w.w.																					3.41	3.03	3.28	m	-?	m	11				
131G	NUCE LAP	WO	w.w.																					3.89	3.77	3.63	m	-?	m	<=5				
15G	NUCE LAP	WO	w.w.																					3.69	3.86	3.42	3.43	m	-?	m	9			
22G	NUCE LAP	WO	w.w.																					4.00	4.00	3.95	4.00	m	-?	m	<=5			
220G	NUCE LAP	WO	w.w.																					4.05	4.00	4.00		m	--	m	<=5			
227G	NUCE LAP	WO	w.w.																					4.00	4.15	4.09	4.50	4.30	4.50	4.13	3.92	--	m	9
98G	NUCE LAP	WO	w.w.																								3.50	3.76	3.80	4.00	m	U?	m	6
11G	NUCE LAP	WO	w.w.																									0.03	0.00	0.26	m	-?	m	7



## Appendix J Geographical distribution of contaminants and biomarkers in biota 1990-2004

Sorted by contaminant and species:

**Cadmium (Cd)**

**Mercury (Hg)**

**Lead (Pb)**

**Sum of 7 CBs (CB-28, -52, 101, -118, -138, -153 and -180)**

**DDEPP (ppDDE)**

**HCB**

**TCDDN**

**TBT**

**OH-pyrene**

**ALA-D** ( $\delta$ -amino levulinic acid dehydrase inhibition)

**EROD** (Cytochrome P4501A-activity)

**MYTI EDU** - Blue Mussel (*Mytilus edulis*)

**GADU MOR** - Atlantic cod (*Gadus morhua*)

**PLAT FLE** - Flounder (*Platichthys flesus*)

**LIMA LIM** - Dab (*Limanda limanda*)

**PLEU PLA** - Plaice (*Pleuronectes platessa*)

**MICR KIT** - Lemon sole (*Microstomus kitt*)




**LEPI WHI** - Megrime (*Lepidorhombus whiffiagonis*)

**Station positions are shown on maps in Appendix G**




Results are presented for three periods: 1990-1993, 2003 and 2004.  
The average median concentrations was used for each period. Cf. Appendix E.






**Appendix J**  
**Geographical distribution of contaminants and biomarkers in**  
**biota 1990-2004**  
**(cont.)**

**A****B****C**




**Figure 22.** Median, standard deviation and provisional "high background" concentration for cadmium in mussels (*Mytilus edulis*) 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppm (mg/kg) wet weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figures A.

**A****B****C**




**Figure 23.** Median, standard deviation and provisional "high background" concentration for cadmium in fish liver 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppm (mg/kg) wet weight (see maps in Appendix G).

**A****B****C**

**Figure 24.** Median, standard deviation and provisional "high background" concentration for mercury in mussels (*Mytilus edulis*) 1990-1993 (A), 2003 (B) and 2004 (C), ppm (mg/kg) wet weight (see maps in Appendix G).


**A****B****C**

**Figure 25.** Median, standard deviation and provisional "high background" concentration for mercury in fish fillet 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppm (mg/kg) wet weight (see maps in Appendix G).


**A****B****C**

**Figure 26.** Median, standard deviation and provisional "high background" concentration for lead in mussels (*Mytilus edulis*) 1990-1993 (A), 2003 (B) and 2004 (C), ppm (mg/kg) wet weight (see maps in Appendix G). Note: for some stations the standard deviation is off-scale in figure A.


**A**






**B**






**C**






**Figure 27.** Median, standard deviation and provisional "high background" concentration for lead in fish liver 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppm (mg/kg) wet weight (see maps in Appendix G).

**A****B****C**




**Figure 28.** Median, standard deviation and provisional "high background" concentration for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in mussels (*Mytilus edulis*) 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) wet weight (see maps in Appendix G).

**A****B****C**




**Figure 29.** Median, standard deviation and provisional "high background" concentration for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in fish liver 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb (µg/kg) wet weight (see maps in Appendix G). **Note: for some stations the standard deviation is off-scale in figures A-C.**

**A****B****C**

**Figure 30.** Median, standard deviation and provisional "high background" concentration for ppDDE (DDEPP) in mussels (*Mytilus edulis*) 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) wet weight (see maps in Appendix G). (See also footnote in Table 7). **Note: for some stations the standard deviation is off-scale in figure B.**


**A****B****C**

**Figure 31.** Median, standard deviation and provisional "high background" concentration for ppDDE (DDEPP) in fish liver 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) wet weight (see maps in Appendix G). (See also footnote in Table 7).


**A****B****C**

**Figure 32.** Median, standard deviation and provisional "high background" concentration for HCB in mussels (*Mytilus edulis*) 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) wet weight (see maps in Appendix G).


**A**





**B**



**C**




**Figure 33.** Median, standard deviation and provisional "high background" concentration for HCB in fish liver 1990-1993 (**A**), 2003 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) wet weight (see maps in Appendix G).


**A****B**

**Figure 34.** Median, standard deviation and provisional "high background" concentration for dioxin TCDD-toxicity equivalents after nordic model (TCDDN) in mussels 2003 (**A**) and 2004 (**B**), ppm (ng/kg) wet weight (see maps in Appendix G). NB: TCDDN is a sum of specific dioxin compounds of which may be of uncertain quantification.

**A**




**B**




**Figure 35.** Median and standard deviation concentration for OH-pyrene (Pyrene metabolite) in fish bile 2003 (**A**) and 2004 (**B**),  $\mu\text{g}/\text{kg}$ /ABS (absorbance) 380 nm (see maps in Appendix G).

**A**

## ALAD in fish blood

**B**


## ALAD in fish blood



**Figure 36.** Median and standard deviation activity for ALA-D ( $\delta$ -amino levulinic acid dehydrase inhibition) in fish liver 2003 (A) and 2004 (B), ng PBG (porphobilinogen)/min/mg protein (see maps in Appendix G).


**A**

EROD in fish liver





**B**



EROD in fish liver



**Figure 37.** Median and standard deviation activity for EROD (Cytochrome P4501A-activity) in fish liver 2003 (**A**) and 2004 (**B**), pmol/min/mg protein (see maps in Appendix G).

**A****B**


**Figure 38.** Median, standard deviation and provisional "high background" concentration for tributyl tin (TBT-concentration on a formulation basis) in blue mussel 2003 (**A**) and 2004 (**B**), ppm (2.44\* mg Sn/kg) dry weight (see maps in Appendix G).

**A****TBT in dogwhelk****B****TBT in dogwhelk**

**Figure 39.** Median, standard deviation and provisional "high background" concentration for tributyl tin (TBT-concentration on a formulation basis) in dogwhelk 2003 (**A**) and 2004 (**B**), ppm (2.44\* mg Sn/kg) dry weight (see maps in Appendix G).


**A**

VDSI in dogwhelk



**B**

VDSI in dogwhelk



**Figure 40.** Average VDSI in dogwhelk 2003 (**A**) and 2004 (**B**) (see maps in Appendix G).

## Appendix K Geographical distribution of contaminants in surficial sediment 1987-2004

Cadmium (Cd)  
Mercury (Hg)  
Lead (Pb)  
Copper (Cu)  
Zinc (Zn)

Sum of 7 CBs (CB-28, -52, 101, -118, -138, -153 and -180)

DDEPP (ppDDE)

$\gamma$ -HCH  
HCB

BAP (benzo[a]pyrene)

**PK- $\Sigma$ n or PK\_S** (sum carcinogen PAHs, cf. Appendix B)

**P- $\Sigma$ n or P\_S** (sum of PAHs, dicyclic "PAHs" not included, cf. Appendix B)




TBT

Station positions are shown on maps in Appendix G




Results are presented for three periods: 1987-1990, 1992-1997 and 2004.  
A station was not monitored more than once during each period, cf. Appendix F.






**Appendix K**  
**Geographical distribution of contaminants in surficial sediment**  
**1987-2004**  
**(cont.)**

**A****B****C**

**Figure 41.** Median, standard deviation and provisional "high background" concentration for cadmium in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppm (mg/kg) dry weight (see maps in Appendix G). Note: for some stations the standard deviation is off-scale in figures **A** and **C**.


**A****B****C**

**Figure 42.** Median, standard deviation and provisional "high background" concentration for mercury in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppm (mg/kg) dry weight (see maps in Appendix G). **Note: for some stations the standard deviation is off-scale in figures A and C.**


**A****B****C**

**Figure 43.** Median, standard deviation and provisional "high background" concentration for lead in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppm (mg/kg) dry weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figure A.


**A**






**B**






**C**






**Figure 44.** Median, standard deviation and provisional "high background" concentration for copper in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppm (mg/kg) dry weight (see maps in Appendix G).

**A****B****C**

**Figure 45.** Median, standard deviation and provisional "high background" concentration for zinc in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppm (mg/kg) dry weight (see maps in Appendix G).


**A****B****C**

**Figure 46.** Median, standard deviation and provisional "high background" concentration for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G).


**A****B****C**

**Figure 47.** Median, standard deviation and provisional "high background" concentration for ppDDE (DDEPP) in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G).


**A**






**B**






**C**






**Figure 48.** Median, standard deviation and provisional "high background" concentration for  $\gamma$ -HCH (Lindane) in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu$ g/kg) dry weight (see maps in Appendix G).

**A****B****C**




**Figure 49.** Median, standard deviation and provisional "high background" concentration for HCB in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figure **B**.

**A****B****C**


**Figure 50.** Median, standard deviation and provisional "high background" concentration for Benzo[a]pyrene (BAP) in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figure **B**.

**A****B****C**

**Figure 51.** Median, standard deviation and provisional "high background" concentration for sum of PAH in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figure **B**.

**A****B****C**

**Figure 52.** Median, standard deviation and provisional "high background" concentration for sum of carcinogen PAHs in surficial sediment (0-2cm) 1987-1990 (**A**), 1992-1997 (**B**) and 2004 (**C**), ppb ( $\mu\text{g}/\text{kg}$ ) dry weight (see maps in Appendix G). **Note:** for some stations the standard deviation is off-scale in figure **B**.



**Figure 53.** Median, standard deviation and provisional "high background" concentration for sum of TBT in surficial sediment (0-2cm) 2004 ppm (mg/kg) dry weight (see maps in Appendix G).

## **Appendix L**

### **Results from INDEX determinations 1995-2004**



## Introduction

The Norwegian Pollution Control Authority (SFT) has requested that a small group of indices be established to assess the quality of the environment with respect to contaminants. The target indicator medium for both indices may vary depending on what purpose is defined, however sediment, cod and mussels are considered to be the most relevant choices. Blue mussel was selected for this investigation (Appendix L1 and Appendix L2).

Two indices are calculated. One index is based on the contaminant concentrations in the blue mussel collected annually from 9 of the more contaminated fjords in Norway (Walday *et al.* 1995), herein designated "Pollution Index". This index was initiated in 1995. Initially there were 11 fjords but sampling from Orkdalsfjord and Iddefjord was discontinued in 1997. It was practical to organise sampling within JAMP. Some JAMP results could be used to calculate the index value.

In addition, a "Reference Index" was initiated in 1995 based on annual contaminant concentrations in the blue mussel. The mussels were collected at JAMP stations along the entire coast where there is presumably low levels of contamination. The importance of "reference" stations for monitoring of contaminants has been discussed earlier (cf. Green 1987). One of the main reasons for this work is to establish points of reference for contaminated fjords. Initially 8 areas were involved but since 1998 only 5 have been sampled.

## Calculation of the index

Sampling strategy and a detailed discussion of calculation of the Pollution Index has been given earlier (cf. Walday *et al.* 1995) and only a brief summary will be given here. The relevant contaminants for each of the Pollution Index fjords are summarised in Appendix L2 and J3. Their selection is based on earlier investigations. Two to five stations were sampled from each area. Three replicate samples with 20 individuals with a shell length of 3-5 cm were collected from each station. Each sample was analysed for the contaminants according to the scheme in Appendix L3. "Dioxins" were only investigated in 1995-96, but reinstated for some stations in 2002-2003. Assessment of TBT concentrations was introduced in 2002 even though it is not identified as a selection criteria by Walday *et al.* (1995).

One to three stations were sampled from selected areas for the determination of the Reference Index. Each station included three replicates which were analysed for the usual JAMP contaminants (cf. analysis code A, Appendix L3). Some samples were also analysed for PAHs and dioxins.

The strategy for sampling mussels differed depending on whether the mussels were to be used for the Index or for JAMP and Index in that stations that were exclusively to be used for Index calculations allowed a slightly greater size range (3-5 cm) compared to JAMP and that the mussels were frozen directly and not depurated.

The maximum median for each contaminant for all the stations in an area was determined. These concentrations were classified according to SFT's classification system for contaminants in the marine environment (Appendix L4). The highest class found for any contaminant measured in an area determined the index value for that area.

The SFT Classes are based on the provisional "high background" levels. This system has been revised (Molvær *et al.* 1997); where among other changes the sum of CB-28, -52, -101, -118, -138, -153, and -180 (CBΣΣe) is now a distinct parameter for classification. The sum of all PAHs excluding the dicyclic PAHs (PAH<sub>Σ</sub>) was compared to the system's "sum-PAH". Previously this was the calculation of sum-PAH that included the dicyclic PAHs. For this report PAH<sub>Σ</sub> was calculated, also for previous years. As a result, the classification may be different by a Class from what has been previously reported. "Dioxins" were assessed based on toxicity equivalency factors (TEQ) according to a Nordic model (Ahlborg 1989) which differs insignificantly from the recently revised WHO-model (van den Berg *et al.* 1998). Note that EPOCl is considered a relevant contaminant for one area but is not included in the part of the classification system based on levels in mussels. Likewise, there are contaminants which are included in the classification system but have not been measured in any area (e.g., tributyltin (TBT), arsenic, fluoride, nickel, silver).

The maximum class found for any contaminant determined the Class (I-V) of the area. The average Class for all the contaminated sub areas and all the reference localities determined the Pollution or Reference Index, respectively. The lowest Index value is 1 and means that all median values were in Class I (slightly polluted). The highest Index value is 5 and means that at least one median value from each of the areas was in Class V (extremely polluted).

## Conclusion from application of the indices

The indices have been used since 1995 based on contaminant concentrations in mussels from 14-19 areas (cf. Green *et al.* 2004b). An assessment of their application suggested that the pollution index needed mainly two improvements (Green & Knutzen 2001): 1) more stations to avoid the consequences of insufficient sample size and 2) inclusion of more relevant contaminant analyses with respect to the pollution load expected and in relation to the SFT classification system for environmental quality (Molvær *et al.* 1997). SFT provided funds to improve the index in 2002. Three additional stations have since been established: one in the Frierfjord area (I713 Strømtangen, about 800 m east of I711 Steinsholmen), one in the inner Ranfjord (I964 Toraneskaien, about 500 m north of I965 Moholmen) and one in the Sunndalsfjord area (I915 Flåøya, northwest, about halfway between I913 and the inner most part of the fjord). Dioxin and TBT analyses were added to the programme for samples collected in the Frierfjord area, inner Oslofjord and the inner Kristiansandsfjord. TBT-analyses were also included for some of the reference stations (see Annex). These changes affect the outcome of the index and comparison to previous years should be cautioned. For results up to and including 2001 SFT has presented only the results using the old method of calculation, for 2002 the results for both the old and new methods are presented, and for 2003 only the results for the new method are presented (cf. SFT's website at [www.miljostatus.no/templates/themepage\\_2699.aspx](http://www.miljostatus.no/templates/themepage_2699.aspx)). Comparison of the two methods for 2002 and 2003 has been done earlier (NIVA memorandum of 1 August 2003 and 17 July 2004).

It should also be noted that the SFT classification system is under revision and may affect calculations of the indices in the future. One likely change will be the lowering of limits to the classes for PCBs taking into consideration a lower background from 4 to 3 ppb wet weight suggested by Green & Knutzen (2003).

No special considerations were made when one but not all the stations within an area were sampled. The lack of sufficient samples has occurred several times for the Pollution Index: (st. I205 Bølsnes from Saudafjord 1996, st. I911 Horvika in the Sunndalsfjord since 1999, st. I021 in the Hvaler area 1999, st.I962 in the Inner Ranfjord since 1999, and st. I711 Steinholmen in the Frierfjord 2001).

Because insufficient amount of mussels were found at station Horvika in the Sunndalsfjord, two new stations were introduced; Fjøseid (I913) in 1999 and Flåøya, northwest (I915), in 2003, about 15 and 5 km farther out the fjord from Horvika, respectively. Because sufficient amount of mussels were not found at st. I962 Koksverktomta in the Ranfjord since 1999, a new station (I965 - Moholmen) was introduced in 2001 about 2 km south of Koksverktomta.

Based on the new calculation with the mentioned supplementary stations and supplementary analyses of dioxin and TBT, the **Pollution Index for 2004 was 3.4** compared to 3.6 for 2003 (cf., **Table 10**). A value between 3 and 4 would correspond to a classification between "marked" and "severe" in the SFT system. The index decreased one class for both Sørfjorden and Byfjord, because of lower concentrations of lead and PCBs, respectively, but increased one class for Saudafjorden because of PAHs. Statistical analyses did not reveal any significant temporal trends for these contaminants in mussels from the relevant stations in these fjords/areas.

Six fjords/areas were monitored for the Reference Index for 2004 compared to 5 for 1998-2004, 7 for 1997 and 8 for 1995-1996 (**Table 11**, Appendix L5). However, only four of these provided a common basis (cf., Table 11). Similar to the application Pollution Index, the Reference Index made no special considerations when one but not all the stations within an area were sampled. For the four common areas, this has occurred several times, all in the Varangerfjord area (st.48A since 1997 and st.11A since 1998). With Lofoten and Outer Ranfjord, and the supplementary analyses of

TBT included, the **Reference Index for 2004 was 1.3** compared to 1.8 for 2003 (**Table 11**, Appendix L5). All five fjords/areas included the TBT analyses. A value between 1 and 2, which would correspond to a classification between "moderate" and "marked" in the SFT system. The index decreased one class for the mid/outer Oslofjorden, Børnlo-Sotra area, and the Lofoten area primarily because of a decrease in TBT. No statistically significant temporal trends were found for TBT from stations in these fjords/areas. The index increased at Lista because of PAH and HCB. However, a statistical analysis on the PAH-concentrations since 1995 revealed a significant downward trend for one of the two stations from this area, due to a particularly high concentration found in 1995 (cf. Green, *et al.* 2004).

**Table 10.** Maximum environmental classification for fjords selected for Pollution INDEX. (See text and Appendix L5).

Index Area <sup>1)</sup>	'95	'96	'97 <sup>2)</sup>	'98	'99	'00	'01	'02	'02 new <sup>7)</sup>	'03	'03 new <sup>7)</sup>	'04 new <sup>7)</sup>
Hvaler/Singlefjord	2	2	2	3	2	2	2	2	2	2	2	2
Iddefjord	-	-	-	-	-	-	-	-	-	-	-	-
Inner Oslofjord	3	3	4	2	3	2	2	2	4	2	4	4
Frierfjord, Grenlandsfjords	3	4	3	3	3	3	3	5 <sup>6)</sup>	5	3 <sup>6)</sup>	5	5
Inner Kristiansandsfjord	5	5	5	5	5	4	3	3	3	4	4	4
Saudafjord	4	5	5	3	4	3	3	4	4	2	2	3
Sørfjord	5	4	3	3	4	4	3	4	4	5	5	4
Byfjorden, Bergen <sup>3)</sup>	3	3	3	2	2	2	2	3	3	4	4	3
Sunndalsfjord	3	3	3 <sup>4)</sup>	2	3	4	2	3	3	1 <sup>6)</sup>	1	1
Orkdalsfjord	-	-	-	-	-	-	-	-	-	-	-	-
Inner Ranfjord	5	3	3 <sup>5)</sup>	4	2	2	3	3 <sup>6)</sup>	3	3 <sup>8)</sup>	5	5
<b>AVERAGE (Pollution INDEX)</b>	<b>3.7</b>	<b>3.6</b>	<b>3.4</b>	<b>3.0</b>	<b>3.1</b>	<b>2.9</b>	<b>2.7</b>	<b>3.2</b>	<b>3.4</b>	<b>2.9</b>	<b>3.6</b>	<b>3.4</b>

<sup>1)</sup> Iddefjord and Orkdalsfjord not sampled since 1997, hence the indices 1995-96 do not include the local indices from these fjords

<sup>2)</sup> Copper, zinc and TCDDN excluded since 1997, hence indices for 1995-96 excludes these contaminants

<sup>3)</sup> PCB (DDTΣ, HCB, HCHΣΣ and CBΣΣ) analysed in stored samples for 1995-1996

<sup>4)</sup> Change in classification (cf. Green *et al.* 1999) due to recalculation of PAHs that excluded the dicyclic compounds

<sup>5)</sup> Change in classification (cf. Green *et al.* 1999) due to calculation error

<sup>6)</sup> Results from supplementary station would not influence the outcome of classification

<sup>7)</sup> Inclusion of supplementary a station in Frierfjord, Inner Ranfjord, and Sunndalsfjord (2003), and supplementary dioxin and TBT analyses for Inner Oslofjord, Frierfjord, and Inner Kristiansandfjord.

<sup>8)</sup> Results from supplementary station would influence the outcome of classification.

**Table 11.** Maximum environmental classification for fjords selected for Reference INDEX. (See text and Appendix L6).

Index Area	'95	'96	'97	'98	'99	'00	'01	'02	'02 new <sup>5)</sup>	'03	'03 new <sup>5)</sup>	'04 new <sup>5)</sup>
Mid and outer Oslofjord <sup>1)</sup>	2	2	2	1	1	1	2	1	1	1	2	1
Lista	1	1	1	1	2	2	2	2	2	1	1	2
Bømlo-Sotra	1	1	1	1	1	2	2	1	2	1	3	2
Outer Ranfjord, Helgeland <sup>2)</sup>	(1)	(1)	-	-	-	-	-	-	-	-	-	1
Lofoten <sup>3)</sup>	(2)	(2)	(1)	(2)	(2)	(1)	(2)	(2)	2	(2)	2	1
Finnsnes-Skjervøy <sup>2)</sup>	(2)	(1)	(1)	-	-	-	-	-	-	-	-	-
Hammerfest-Honningsvåg <sup>2)</sup>	(2)	(3) <sup>4)</sup>	(2)	-	-	-	-	-	-	-	-	-
Varanger Peninsula	1	2	1	2	1	1	1	1	1	1	1	1
<b>AVERAGE (Reference INDEX)</b>	<b>1.3</b>	<b>1.5</b>	<b>1.3</b>	<b>1.3</b>	<b>1.3</b>	<b>1.5</b>	<b>1.8</b>	<b>1.3</b>	<b>1.6</b>	<b>1.2</b>	<b>1.8</b>	<b>1.3</b>

<sup>1)</sup> Inclusion of results for arsenic, nickel and silver in 1996 did not affect the classification

<sup>2)</sup> Outer Ranfjord, Finnsnes-Skjervøy and Hammerfest-Honningsvåg stations were not sampled in 1998, hence, the index for 1995-97 did not take these results into account. See cf., Green *et al.* 2000 for more details for outer Ranfjord.

<sup>3)</sup> Inconsistency in sampling site, st.98X in 1995-96 and st.98A in 1997, hence, results from Lofoten excluded. See cf., Green *et al.* 2000 for more details for st 98X.

<sup>4)</sup> Change in classification (cf. Green *et al.* 1999) due to recalculation of PAHs that excluded the dicyclic compounds.

<sup>5)</sup> Inclusion of supplementary TBT analyses for Mid and outer Oslofjord, Lista, Bømlo-Sotra, Lofoten and Varangerfjord Peninsula.



## Appendix L1

### INDEX - Stations and programme 1995-2003

**Appendix L1.** INDEX station positions and sampling overview for blue mussel 1995-2003, where P = “Pollution Index” and R = “Reference Index” (contaminated and assumed “background” stations, respectively). Mussels were sampled from rock surfaces unless otherwise noted. See Walday *et al.* (1995) for discussion of station selection and analyses.

Station	Locality name	North latitude	East longitude	ICES position	INDEX type P/R	notes
<b>HVALER/SINGLEFJORDEN, east of outer OSLOFJORD</b>						
I021	Kjøkø, south	59°07.8'	10°57.1'	47G13	P	
I024	Kirkø, north west	59°04.9'	10°59.2'	47G09	P	
I022	West Damholmen	59°06.2'	10°57.9'	47G09	P	
I023	Kirkø, north west	59°05.7'	11°08.2'	47G09	P	
<b>IDDEFJORD, east of outer OSLOFJORD</b>						
I001	Sponvikskansen	59°05.4'	11°12.5'	47G09	P	
I011	Kråkenebbet	59°06.1'	11°17.3'	47G09	P	
<b>INNER OSLOFJORD</b>						
JAMP 30A	Gressholmen	59°52.5'	10°43.0'	48G07	P	
I301	Akershuskaia	59°54.2'	10°45.5'	48G07	P	
I304	Gåsøya	59°51.0'	10°35.5'	48G04	P	
I307	Ramtonholmen	59°44.7'	10°31.4'	48G05	P	
I306	Håøya	59°24.7'	10°33.4'	48G04	P	
<b>MID and OUTER OSLOFJORD</b>						
JAMP 31A	Solbergstrand	59°36.9'	10°39.4'	48G06	R	
JAMP 35A	Mølen	59°29.2'	10°30.1'	47G04	R	
JAMP 36A	Færder	59°01.6'	10°31.7'	47G06	R	
<b>FRIERFJORD AREA, west of outer Oslofjord</b>						
I711	Steinholmen	59°03.2'	09°40.7'	48F99	P	
I712	Gjermundsholmen	59°02.8'	09°42.5'	47F99	P	
I713	Strømtangen	59°03.2'	09°41.5'	47F99	P	
JAMP 71A	Bjørkøya (Risøyodden)	59°01.4'	09°45.4'	47F99	P	
JAMP 76A	Risøy	58°43.6'	09°17.0'	46F92	R	
<b>INNER KRISTIANSANDSFJORD</b>						
I132	Fiskatangen	57°07.7'	07°59.2'	43F79	P	
I133	Odderø, west	57°07.9'	08°00.3'	43F83	P	
<b>LISTA AREA</b>						
JAMP 15A	Gåsøy (Ullerø area)	58°03.1'	06°53.3'	45F69	R	
I131	Lastad	58°03.3'	07°42.4'	45F79	R	7
<b>SAUDAFJORD</b>						
I201	Ekkjegrunn (G1)	59°38.7'	06°21.4'	48F66	P	
** I205	Bølsnes (G5)	59°35.5'	06°18.3'	48F63	P	
<b>BØMLO AREA</b>						
JAMP 22A	Espevær, west	59°35.2'	05°58.5'	48F59	R	C, 1
<b>SØRFJORD</b>						
* 51A	Byrkjenes	60°05.1'	06°33.1'	49F66	P	
JAMP 52A	Eitrheimsneset	60°05.8'	06°32.2'	49F66	P	3

**Appendix L1 (cont'd)**

Station	Locality name	North latitude	East longitude	ICES position	INDEX type P/R	notes
<b>BYFJORDEN, Bergen</b>						
I242	Valheimneset	60°23.7'	05°16.1'	49F51	P	
I241	Nordnes	60°24.1'	05°18.2'	49F51	P	
I243	Hagreneset	60°24.9'	05°18.3'	49F51	P	
<b>SUNNDALSFJORDEN</b>						
I912	Honnhammer	62°51.2'	08°09.7'	54F81	P	
I911	Horvika	62°44.1'	08°31.4'	54F85	P	
I913	Fjøseid	62°49.0'	08°16.48'	54F85	P	
I914	Flåøya, southeast	62°45.35'	08°26.70'	54F85	P	
I915	Flåøya, northwest	62°45.48'	08°26.39'	54F85	P	
<b>[TRONDHEIM AREA - not related to INDEX investigation]</b>						
*	80A Østmerknes	63°27.5'	10°27.5'	56G04	P	
<b>ORKDALSFJORD AREA, supplementary area (cf. Walday <i>et al.</i> 1995)</b>						
JAMP 82A	Flakk	63°27.1'	10°12.6'	56G01	P	
JAMP 84A	Trossavika	63°20.8'	09°57.8'	55F97	P	
JAMP 87A	Ingdalsbukta	63°27.8'	09°54.8'	55F97	P	
<b>INNER RANFJORD</b>						
I969	Bjørnbærviken (B9)	66°16.8'	14°02.1'	61G42	P	
I965	Moholmen (B5)	66°19.0'	14°07.5'	61G42	P	
I964	Toraneskaien	66°19.3'	14°08.0'	61G42	P	
I962	Koksverkkaien (B2)	66°19.4'	14°08.0'	61G42	P	3
<b>OUTER RANFJORD, Helgeland area</b>						
*	96A Breiviken	66°17.6'	12°50.5'	61G28	R	1
<b>LOFOTEN AREA</b>						
JAMP 98A	Husvågen (1997)	68°15.4'	14°40.6'	65G46	R	5
JAMP 98A	Husvågen (1998)	68°16.9'	14°40.1'	65G46	R	
<b>FINNSNES-SKJERVØY AREA</b>						
JAMP 41A	Fensneset, Grytøya	68°56.9'	16°38.5'	66G64	R	3
<b>HAMMERFEST-HONNINGSVÅG AREA</b>						
JAMP 44A	Elenheimsundet	70°30.8'	22°14.8'	70H23	R	1, 6
JAMP 46A	Småneset in Altesula	70°58.4'	25°48.1'	70H57	R	3, 6
<b>VARANGER PENINSULA AREA</b>						
JAMP 48A	Trollfjorden i Tanafjord	70°41.6'	28°33.3'	70H85	R	
JAMP 10A	Skagoodden	70°04.2'	30°09.8'	69J03	R	
JAMP 11A	Sildkroneneset, Bøkfjorden	69°47.0'	30°11.1'	68J02	R	
JAMP 11X	Brashavn	69°53.9'	29°44.7'	68H97	R	2

notes:

- \* - JAMP station but not sampled in accordance to JAMP guidelines, see Appendix text.
- \*\* - Sufficient mussel-sample not found in 1996.
- 1 - mussels collected from buoy and/or buoy anchor lines
- 2 - mussels collected from sand/gravel bottom
- 3 - mussels collected from iron/cement pilings
- 4 - mussels collected from metal navigation buoys
- 5 - mussels collected from floating dock
- 6 - mussels collected from wooden docks
- 7 - mussels collected from tire on jetty

## Appendix L2

### INDEX - Sampling and analyses for 1995-2003

**Appendix L2.** Blue mussel samples planned or used in INDEX and other purposes besides JAMP 1995-2003, where P = "Pollution Index" and R = "Reference Index" (contaminated and assumed "background" stations, respectively). + indicates JAMP sampling and analyses (i.e. equivalent to analysis code A). The number indicates the number samples analysed. Codes for analysis (A, B etc.) are defined in Appendix L3. See Walday *et al.* (1995) for discussion of selection of stations and analyses.

JAMP st.	STATION	INDEX	ANALYSIS CODE									
			+	A	B	C	D	E	F	G	H	I
<b>HVALER/SINGLEFJORD AREA</b>												
021	Kjøkø, south	P	.	.	.	.	.	3	.	.	.	.
024	Kirøy, north west	P	.	.	.	.	.	3	.	.	.	.
022	West Damholmen	P	.	.	.	.	.	3	.	.	.	.
023	Singlekalven, south	P	.	.	.	.	.	3	.	.	.	.
<b>IDDEFJORD</b>												
01A	Sponvikskansen	P	.	.	.	.	.	3	.	.	.	.
011	Krækenebchet	P	.	.	.	.	.	3	.	.	.	.
<b>OSLOFJORD</b> , inner												
30A	Gressholmen	P	.	.	.	.	+	.	.	.	3	.
301	Akershuskaia	P	.	.	.	.	.	3	.	.	.	2
304	Gåsøya	P	.	.	.	.	.	3	.	.	.	.
307	Ramtonholmen	P	.	.	.	.	.	3	.	.	.	.
306	Håøya	P	.	.	.	.	.	3	.	.	.	.
<b>OSLOFJORD</b> , mid and outer												
31A	Solbergstrand	R	.	.	.	.	+	.	.	.	.	.
35A	Mølen	R	.	.	.	.	+	.	.	.	.	.
36A	Færder	R	.	.	.	.	+	.	.	.	.	2
<b>FRIERFJORD AREA</b> , west of outer Oslofjord												
712	Gjermundsholmen	P	.	.	.	.	.	.	3	.	.	2
713	Strømtangen	P	.	.	.	.	.	.	3	.	.	1
71A	Bjørkøya	P	.	.	.	+	.	.	.	.	.	2
76A	Risøy	R	.	.	.	+	.	.	.	.	.	1
<b>INNER KRISTRIANSANDSFJORD</b>												
132	Fiskåtangen	P	.	.	.	.	.	.	3	.	.	2
133	Odderø, west	P	.	.	.	.	.	.	3	.	.	2
<b>LISTA AREA</b>												
15A	Gåsøya	R	.	.	.	.	+	.	.	.	.	2
131	Lastad	R	.	.	.	.	.	3.	.	.	.	.
<b>SAUDAFJORD</b>												
201	Ekkjegrunn (G1)	P	.	.	.	.	.	.	.	3	.	.
205	Bølsnes (G5)	P	.	.	.	.	.	.	3	.	.	.
<b>[HAUGESUND AREA not related to INDEX investigation]</b>												
227	Melandsholmen	O	.	.	.	.	+	.	.	.	.	1
<b>BØMLO-SOTRA AREA</b>												
22A	Espevær, west	R	.	.	.	.	+	.	.	.	.	2
<b>SØRFJORD</b>												
51A	Byrkjeneset	P	.	.	.	.	3	.	.	.	.	.
52A	Eirtrheimneset	P	.	.	.	.	+	.	.	.	.	.

\*) indicates Toxaphene included

**Appendix L2 (cont'd)**

STATION JAMP st.	INDEX	ANALYSIS CODE										
		+	A	B	C	D	E	F	G	H	I	J
<b>BYFJORDEN, BERGEN</b>												
242 Valheimsneset	P	.	.	.	.	.	.	.	.	3	.	.
241 Nordnes	P	.	.	.	.	.	.	.	3	.	.	.
243 Hagreneset	P	.	.	.	.	.	.	.	3	.	.	.
<b>SUNNDALSFJORD</b>												
912 Honnhammer	P	.	.	.	.	.	.	.	.	3	.	.
913 Fjøseid	P	.	.	.	.	.	.	.	.	3	.	.
914 Flåøya, southeast	P	.	.	.	.	.	.	.	.	3	.	.
915 Flåøya, northwest	P	.	.	.	.	.	.	.	.	3	.	.
<b>[TRONDHEIM AREA - not related to index investigation]</b>												
80A Østmarknes	-	.	.	.	.	.	.	.	.	3	.	.
<b>ORKDALSFJORD AREA (not suggested in Walday et al. 1995)</b>												
82A Flakk	P	.	.	.	.	+	.	.	.	.	.	.
84A Trossavika	P	.	.	.	.	+	.	.	.	.	.	.
87A Ingdalsbukta	P	.	.	.	.	+	.	.	.	.	.	.
<b>INNER RANFJORD</b>												
964 Toraneskaien	P	.	.	.	.	.	.	.	.	3	.	.
965 Moholmen (B5)	P	.	.	.	.	.	.	.	.	3	.	.
969 Bjørnbærviken (B9)	P	.	.	.	.	.	.	.	.	3	.	.
<b>OUTER RANFJORD, HELGELAND AREA</b>												
96A Breivika, Tomma	R	.	.	.	.	3	.	.	.	.	.	.
<b>LOFOTEN AREA</b>												
98A Husvågen	R	.	.	.	.	+	.	.	.	.	.	2
<b>FINNSNES-SKJERVØY AREA</b>												
41A Fensneset, Grytøya	R	.	.	.	.	3	.	.	.	.	.	.
<b>HAMMERFEST-HONNINGSVÅG AREA</b>												
44A Elenheimsundet	R	.	.	.	.	3	.	.	.	.	.	.
46A Smineset in Altesula	R	.	.	.	.	3	.	.	.	.	.	.
<b>VARANGER PENINSULA AREA</b>												
48A Trollfjorden i Tanafjord	R	.	.	.	.	3	.	.	.	.	.	.
10A Skagoodden	R	.	.	.	.	+	3	.	.	.	.	.
11X Brashavn	R	.	.	.	.	+	3	.	.	.	.	2

\*) indicates Toxaphene included

### Appendix L3

## INDEX - Key to analysis codes and sample counts

(Used in Appendix L2)

**ANALYSIS CODES<sup>1)</sup>** See Walday *et al.* (1995) for discussion of selection of analyses.

Contaminant	Analysis code										
	A	B	C	D	E	F	G	H	I	J	K
Lead	.	.	.	.	.	X	X	.	.	X	.
Cadmium	.	.	.	.	.	X	X	X	.	X	.
Copper <sup>2)</sup>	.	.	.	.	.	X	X	X	.	.	.
Mercury	.	.	.	.	.	X	X	X	.	.	.
Zinc <sup>2)</sup>	.	.	.	.	.	X	X	X	.	X	.
EPOCI	.	.	.	.	.	.	.	X	.	.	.
PAHs	.	.	.	.	.	.	X	.	X	X	.
PCBs	.	.	.	.	.	X	.	X	X	.	X
"Dioxin" <sup>3)</sup>	.	.	.	.	.	.	.	.	.	.	X
TBT <sup>4)</sup>	.	.	.	.	.	.	.	.	.	.	.

<sup>1)</sup> Concerns MUSSEL - 1 size group (3-5 cm), 3 replicate samples each a bulk of 20 individuals (see text)

<sup>2)</sup> Concerns MUSSEL - discontinued since 1996

<sup>3)</sup> Concerns MUSSEL - discontinued since 1995, but reinstated 2002 for st.30A, 71A, I711, I712, I713, 76A, I132 and I133

<sup>4)</sup> Concerns MUSSEL – not included in Walday *et al.* (1995).



**Appendix L4**  
**INDEX - SFT Environmental quality classes**  
(Molvær *et al.* 1997)

<b>As</b>	arsenic
<b>Pb</b>	lead
<b>F</b>	fluoride
<b>Cd</b>	cadmium
<b>Cu</b>	copper
<b>Cr</b>	chromium
<b>Hg</b>	mercury
<b>Ni</b>	nickel
<b>Zn</b>	zinc
<b>Ag</b>	silver
<b>TBT</b>	tributyltin
<b>PAH_S</b>	total PAH excluding dicyclic (=PAH_Σ)*
<b>BAP</b>	benzo[a]pyrene
<b>DDTSS</b>	DDTPP+DDEPP+TDEPP (=DDTΣΣ)*
<b>HCB</b>	hexachlorobenzene
<b>HCHSS</b>	HCHG+HCHA+HCHB (=HCHΣΣ)*
<b>CBSSe</b>	sum of CB: 28+52+101+118+138+153+180 *
<b>TCDDN</b>	Sum of TCDD-toxicity equivalents *

\* ) See also Appendix B for definitions.

Basis: D = dry weight, W = wet weight

Units: M = ppm (mg/kg), U = ppb (μg/kg), P = ppp (ng/kg)



**SFT's Environmental quality classes for blue mussel (Molvær *et al.* 1997).**

Contaminant	basis	unit	Class I	Class II	Class III	Class IV	Class V
As	D	M	<10	10-30	30-100	100-200	>200
Pb	D	M	<3	3-15	15-40	40-100	>100
F	D	M	<15	15-50	50-150	150-300	>300
Cd	D	M	<2	2-5	5-20	20-40	>40
Cu	D	M	<10	10-30	30-100	100-200	>200
Cr	D	M	<3	3-10	10-30	30-60	>60
Hg	D	M	<0.2	0.2-0.5	0.5-1.5	1.5-4	>4
Ni	D	M	<5	5-20	20-50	50-100	>100
Zn	D	M	<200	200-400	400-1000	1000-2500	>2500
Ag	D	M	<0.3	0.3-1	1-2	2-5	>5
TBT	D	M	<0.1	0.1-0.5	0.5-2	2-5	>5
PAH_S	W	U	<50	50-200	200-2000	2000-5000	>5000
BAP	W	U	<1	1-3	3-10	10-30	>30
DDTSS	W	U	<2	2-5	5-10	10-30	>30
HCB	W	U	<0.1	0.1-0.3	0.3-1	1-5	>5
HCHSS	W	U	<1	1-3	3-10	10-30	>30
CBSSe	W	U	<4	4-15	15-40	40-100	>100
TCDDN	W	P	<0.2	0.2-0.5	0.5-1.5	1.5-3	>3



**Appendix L5  
INDEX - Summary table “Pollution index”  
2003-2004**



**Pollution index 2003-new (with supplementary analyses and stations)**

Average of Max E.C is 3.6

Index areaname (Pollution area) 2003	n	N	As ppm d.w.	Pb ppm d.w.	F ppm d.w.	Cd ppm d.w.	Cu ppm d.w.	Cr ppm d.w.	Hg ppm d.w.	Ni ppm d.w.	Zn ppm d.w.	Ag ppm d.w.	PAH_S ppb w.w.	BAP ppb w.w.	DDTSS ppb w.w.	HCB ppb w.w.	HCHSS ppb w.w.	CBSSe ppb w.w.	TCDDN ppb w.w.	TBT ppm w.w.	Max E.C I:V
Hvaler/Singlefjorden	3	4	i	1.73	i	2.03	i	i	0.23	i	i	i	i	<0.54	<0.05	<0.10	<2.06	i	i	II	
Iddefjord	0	2	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	miss	
Inner Oslofjord	3	5	i	i	i	1.56	i	i	0.07	i	i	i	<44.72	<0.50	<1.17	<0.05	<0.10	11.15	<0.10	2.17	IV
Frierfjorden	3	4	i	i	i	i	i	i	i	i	i	i	i	0.7	0.48	<0.10	<2.48	<4.26	0.91	V	
Inner Kristiansfjord	2	2	i	i	i	i	i	i	i	i	i	i	<412.47	13	<0.37	0.29	<0.32	<1.93	<0.62	0.44	IV
Saudafjord	2	2	i	3.4	i	1.25	i	i	i	i	i	i	<102.90	1.2	i	i	i	i	i	II	
Sørfjord	2	3	i	108.46	i	14.68	i	i	1.47	i	i	i	i	3.73	<0.05	<0.10	<2.08	i	i	V	
Byfjorden	3	3	i	i	i	i	i	i	i	i	i	i	i	7.15	<0.05	<0.10	53.75	i	i	IV	
Sunndalsfjord	3	4	i	i	i	i	i	i	i	i	i	i	<40.97	0.64	i	i	i	i	i	I	
Orkdalsfjord area	0	3	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	miss	
Inner Ranfjord	3	4	i	6.45	i	0.81	i	i	i	i	i	i	1069.9	44	i	i	i	i	i	V	



**Pollution index 2004-new (with supplementary analyses and stations)**

Average of Max E.C is 3.4

Index areaname (Pollution area) 2004	n	N	As ppm d.wt	Pb ppm d.wt	F ppm d.wt	Cd ppm d.wt	Cu ppm d.wt	Cr ppm d.wt	Hg ppm d.wt	Ni ppm d.wt	Zn ppm d.wt	Ag ppm d.wt	PAH_S ppb w.wt	BAP ppb w.wt	DDTSS ppb w.wt	HCB ppb w.wt	HCHSS ppb w.wt	CBSSe ppb w.wt	TCDDN ppb w.wt	TBT ppm d.wt	Max E.C I:V
Hvaler/Singlefjorden	3	4	i	3.51	i	1.57	i	i	0.21	i	i	i	i	i	<0.36	0.06	<0.10	1.89	i	ii	
Iddefjord	0	2	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	miss		
Inner Oslofjord	5	5	i	i	i	2.16	i	i	0.15	i	i	i	626.62	4	1.83	0.11	<0.05	6.6	<0.12	2.83	
Frierfjorden	3	4	i	i	i	i	i	i	i	i	i	i	i	i	0.52	0.47	<0.05	<2.49	3.25	0.3	
Inner Kristiansfjord	2	2	i	i	i	i	i	i	i	i	i	i	375.54	15	<0.18	0.44	<0.11	<1.41	<0.38	0.46	
Saudafjord	2	2	i	3.33	i	1.02	i	i	i	i	i	i	378.27	6.8	i	i	i	i	i	iii	
Sørfjord	2	2	i	42.23	i	5.37	i	i	0.3	i	i	i	i	i	3.38	0.07	<0.05	<1.57	i	iv	
Byfjorden	3	3	i	i	i	i	i	i	i	i	i	i	i	i	1.89	0.13	<0.05	39.76	i	iii	
Sunndalsfjord	3	4	i	i	i	i	i	i	i	i	i	i	<37.19	0.39	i	i	i	i	i	i	
Orkdalsfjord area	0	3	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	i	miss		
Inner Ranfjord	3	4	i	13.12	i	1.9	i	i	i	i	i	i	1706.59	44	i	i	i	i	i	v	

I	24
II	13
III	10
IV	3
V	2

**Appendix L6  
INDEX - Summary table “Reference Index”  
2003-2004**

**Reference index 2003-new (with supplementary analyses and stations)****Average of Max E.C is 1.8**

Index areaname (Reference area) 2003	n	N	As ppm	Pb ppm	F ppm	Cd ppm	Cu ppm	Cr ppm	Hg ppm	Ni ppm	Zn ppm	Ag ppm	PAH_S ppb	BAP ppb	DDTSS ppb	HCB ppb	HCHSS ppb	CBSSe ppb	TCDDN ppb	TBT ppm	Max E.C	I:V
			d.w.	d.w.	d.w.	d.w.	d.w.	d.w.	d.w.	d.w.	d.w.	w.w.	w.w.	w.w.	w.w.	w.w.	w.w.	w.w.	d.w.			
Mid and outer Oslofjord	3	3	w	0.81	w	1.33	i	w	0.06	w	i	w	w	w	<0.93	<0.05	<0.10	<1.54	w	0.1	II	
Lista area	2	2	w	0.76	w	0.86	i	w	0.07	w	i	w	<35.62	<0.50	<0.33	<0.05	<0.10	<1.09	w	0.06	I	
Bømlo-Sotra area	1	1	w	1.46	w	1.04	i	w	0.13	w	i	w	w	w	<0.52	<0.05	<0.10	<1.31	w	0.59	III	
Outer Ranfjord, Helgeland area	0	2	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	miss		
Lofoten area	0	1	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	miss		
Finnsnes- Skjervøy area	0	1	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	miss		
Hammerfest-Honningsvåg area	0	2	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	miss		
Varanger peninsula area	1	3	w	1.65	w	1.98	i	w	0.05	w	i	w	w	w	<0.31	<0.05	<0.10	<0.37	w	<0.01	I	

I	32
II	1
III	1
IV	0
V	0

**Reference index 2004-new (with supplementary analyses and stations)**

Max(median). Statistics for alle areas: (n = Index-station measured, N = Station programmed for index)

**Average of Max E.C is 1.3**

Index areaname (Reference area) <b>2004</b>	n	N	As	Pb	F	Cd	Cu	Cr	Hg	Ni	Zn	Ag	PAH_S	BAP	DDTSS	HCB	HCHSS	CBSSe	TCDDN	TBT	I
			ppm	ppb	ppb	ppb	ppb	ppb	ppb	ppb	ppb	ppb									
			d.wt	w.wt	w.wt	w.wt	w.wt	w.wt	w.wt	w.wt	w.wt	d.wt									
Mid and outer Oslofjord	3	3	w	0.96	w	1.4	i	w	0.11	w	i	w	w	w	0.86	0.05	<0.05	<0.75	w	0.06	
Lista area	2	2	w	0.86	w	1.34	i	w	0.1	w	i	w	<191.16	1.2	<0.32	0.29	<0.05	<0.79	w	0.06	
Bømlo-Sotra area	1	1	w	1.79	w	1.18	i	w	0.16	w	i	w	w	w	<0.23	0.03	<0.05	<0.73	w	0.29	
Outer Ranfjord, Helgeland area	1	2	w	0.49	w	0.72	i	w	0.04	w	i	w	w	w	<0.15	<0.03	<0.05	<0.05	w	w	
Lofoten area	1	1	w	1.09	w	1.07	i	w	0.08	w	i	w	w	w	<0.18	0.09	<0.05	<0.35	w	0.05	
Finnsnes- Skjervøy area	0	1	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	r	
Hammerfest-Honningsvåg area	0	2	w	w	w	w	i	w	w	w	i	w	w	w	w	w	w	w	w	r	
Varanger peninsula area	1	3	w	1.02	w	1.59	i	w	0.05	w	i	w	w	w	<0.15	0.05	<0.05	<0.18	w	w	

I	44
II	4
III	0
IV	0
V	0



