




The toxicity of MEA and amine waste water samples using standardised freshwater bioassays



Main Office

Gaustadalléen 21
 NO-0349 Oslo, Norway
 Phone (47) 22 18 51 00
 Telefax (47) 22 18 52 00
 Internet: www.niva.no

Regional Office, Sørlandet

Jon Lilletuns vei 3
 NO-4879 Grimstad, Norway
 Phone (47) 22 18 51 00
 Telefax (47) 37 04 45 13

Regional Office, Østlandet

Sandvikaveien 59
 NO-2312 Ottestad, Norway
 Phone (47) 22 18 51 00
 Telefax (47) 62 57 66 53

Regional Office, Vestlandet

Thormøhlens gate 53 D
 NO-5006 Bergen Norway
 Phone (47) 22 18 51 00
 Telefax (47) 55 31 22 14

Regional Office Central

Pirsenteret, Havnegata 9
 P.O.Box 1266
 NO-7462 Trondheim
 Phone (47) 22 18 51 00
 Telefax (47) 73 54 63 87

Title The toxicity of MEA and amine waste water samples using standardised freshwater bioassays	Report No.. 6477-2013	Date January 2013
	Project No. 12319	Pages 18
Author(s) Steven Brooks Harald Hasle Heiaas Adam Lillicrap	Topic group Ecotoxicology and Risk Assessment	Distribution
	Geographical area Oslo	Printed NIVA

Client(s) Tel-Tek AS: Jon Hovland	Client ref. Jon Hovland Tel-Tek AS
--------------------------------------	--

Abstract

Water samples provided by Tel-Tek AS were assessed for their toxicity to freshwater organisms from three trophic groups. The water samples included pure monoethanolamine (MEA) and two amine waste water mixtures described as *Amine Reactor Waste (ARW)* and treated amine waste water (TW). The toxicity of these three test solutions to the unicellular algae *Pseudokirchneriella subcapitata*, the freshwater crustacean *Daphnia magna*, and the embryos of the zebra fish *Danio rerio* were performed in accordance with standard protocols. Of the three taxonomic groups tested, the algae were found to be the most sensitive to both MEA and ARW followed by daphnids, with the zebra fish embryo the least sensitive. For the algae, EC₅₀ concentrations of 151 mg/L MEA, 0.019% ARW and 2.4% TW where similar to that recorded in previous tests performed at NIVA in 2009. The pattern in toxicity for TW was different with the lowest EC₅₀ concentration calculated in the zebra fish followed by the algae and daphnids. However, the EC₅₀ concentrations for TW were very close ranging from 1.91% TW in the zebra fish to 3.4% TW in daphnids.

4 keywords, Norwegian	4 keywords, English
1. Monoetanolamin (MEA)	1. Monoethanolamine (MEA)
2. Enkeltcellet grønnsalger	2. Unicellular algae
3. Daphnia	3. Daphnids
4. Sebrafisk embryo	4. Zebra fish embryo


 Steven Brooks
 Project Manager


 Kevin Thomas
 Research Manager


 Kristoffer Næs
 Research Director

**The toxicity of MEA and amine waste water samples
using standardised freshwater bioassays**

Contents

Summary	5
1. Scope of the work	6
2. Objective	6
3. Materials and Methods	6
3.1 Physicochemical parameters of test mixtures	6
3.2 Freshwater algal growth test	7
3.3 <i>Daphnia magna</i> immobilisation test	8
3.4 Zebra fish embryo test	8
4. Results	9
4.1 Physicochemical parameters of the test mixtures	9
4.2 Toxicity to freshwater algae	10
4.3 Toxicity to <i>Daphnia magna</i>	13
4.4 Toxicity to the embryo-larvae of the zebra fish	15
5. Conclusion	17
6. References	18

Summary

Three water samples provided by Tel-Tek AS were assessed for their toxicity to freshwater organisms from three trophic groups. The water samples included two mixtures of amine waste water, which were by-products of the CO₂ capturing methodology using amines. The mixtures included an untreated waste water sample described as *Amine Reactor Waste* (ARW) and a treated waste water sample (TW). In addition, pure monoethanolamine (MEA) an important amine used in the CO₂ capturing process was provided for the toxicity assessment.

The toxicity test performed on the three test solutions included: 1) the unicellular freshwater algae *Pseudokirchneriella subcapitata*; 2) the freshwater crustacean *Daphnia magna*; and 3) the embryos of the freshwater zebra fish *Danio rerio*. All tests were performed in accordance with their standard protocols.

For both MEA and ARW, the unicellular algae were the most sensitive followed by daphnids and then the zebra fish. However for TW, the zebra fish was the most sensitive closely followed by the unicellular algae and the daphnids. The effect concentration (EC) endpoints are summarised in the table below.

Test chemical	Trophic group	EC ₁₀	EC ₅₀
MEA (mg/L)	Unicellular algae	30	151
	Daphnids	128	209
	Zebra fish	164.6	617.5-
ARW (%)	Unicellular algae	0.0089	0.019
	Daphnids	0.060	0.081
	Zebra fish	0.034	0.194
TW (%)	Unicellular algae	0.74	2.4
	Daphnids	2.2	3.4
	Zebra fish	-	1.91

1. Scope of the work

The following report describes the results of three freshwater toxicity tests performed on three test solutions. The test solutions were supplied by Tel-Tek AS and represent pure monoethanolamine (MEA); and mixtures of treated and the untreated amine waste water from a CO₂ capturing facility.

The organisms used within the standardised bioassays represent three major freshwater phyla and include the freshwater algae, (*Pseudokirchneriella subcapitata*) the crustacean (*Daphnia magna*) and zebra fish (*Danio rerio*) embryos.

2. Objective

The main objective of the work was to conduct standard freshwater toxicity tests on the three environmental samples provided by Tel-Tek AS, which included: 1) Pure MEA; untreated amine waste water (amine reclaimed waste, ARW); and treated waste water (TW).

3. Materials and Methods

The three test solutions were supplied by Tel-Tek AS and delivered on ice to the NIVA Oslo laboratory where they were stored until analysis. The MEA and ARW were stored in the dark at 4°C, whilst the TW arrived frozen in approximately fifteen 200 ml plastic containers with screw lids that were stored at -20°C until required for testing.

3.1 Physicochemical parameters of test mixtures

Some important physicochemical parameters of the two test mixtures ARW and the TW were assessed prior to analysis. In addition a third test mixture, identified as 'Feed to bioreactor' was measured for the same parameters. These parameters included chemical oxygen demand (COD), ammonium (NH₄), sulphate (SO₄), chloride (Cl), potassium (K) and sodium (Na).

3.2 Freshwater algal growth test

Growth inhibition of the algae *Pseudokirchneriella subcapitata* (NIVA-CHL 1) was performed in accordance with OECD guideline 201 (OECD, 2011). The specifics of the test are found in Table 1.

Table 1. Information for the algal growth inhibition test (OECD 201), which was performed on the three test solutions.

Test method:	OECD 201: Algae growth inhibition test (2011)
Organism:	<i>Pseudokirchneriella subcapitata</i> (NIVA-CHL1)
Test parameter:	Growth rate 72 hours
Stock culture:	Semi-continuous in 10 % Z8 growth medium (Staub, 1961)
Test date:	03.12.2012 - 06.12.2012 (MEA, TW, ARW)
	11.12.2012 – 14.12.2012 (TW 2 nd test)
Pretreatment of sample:	MEA: 1M HCl was added to the test concentrations to adjust pH to 8.0 ± 0.2
	TW: Test compound was filtered with GF/F filter followed by 0.45 μm filter, and added ISO 8692 stock solutions.
	ARW: Test compound was filtered with 0.45 μm syringe filter, and 0.2M HCl were added to test concentrations to adjust pH to 8.0 ± 0.2
Test concentrations:	MEA: 10, 18, 32, 56, 100, 180 mg/L
	TW: 1 st test: 3.2, 5.6, 10, 18, 32 % of original concentration; 2 nd test: 0.18, 0.32, 0.56, 1.0, 1.8, 3.2 % of original concentration
	ARW: 0.0018, 0.0032, 0.0056, 0.010, 0.018, 0.032, 0.056, 0.1 % of original concentration
Preparation of test concentrations:	By dilution of a stock solution 03.12.2012 and 11.12.2012
Test medium:	ISO 8692
Replicates:	3 in each test concentration, 6 in controls
Incubation:	Incubator with orbital shaking
Test containers:	30 ml glass vials with ca. 12 ml sample
Light:	87 $\mu\text{mol m}^{-2} \text{s}^{-1}$, continuous from daylight type fluorescent tubes
Temperature:	Max: 21.1 Min: 19.1 (3-6.12.2012) Max: 20.9 Min: 20.0 (11-14.12.2012)
Inoculum:	5×10^6 cells/ L (nominal concentration) of an exponentially growing culture.
Registration of growth:	Particle count with Beckman Coulter Multisizer 3 after 24, 48 and 72 \pm 2 hours.
Calculation of growth rate:	Logarithmic increase in density from start to 72 hours.
Calculation of EC_x ¹	Non linear regression with Excel macro Regtox 7.0.5 (Hill 1910, Vinidiman et al 1983)

¹ EC_x : The concentration which results in x % reduction in growth rate compared to the control

3.3 *Daphnia magna* immobilisation test

Acute immobilisation of *Daphnia magna* was performed in accordance with OECD guideline 202 (OECD, 2004). Test vessels were examined under microscope once daily for the duration of the test, and immobilised or dead animals were recorded. The specifics of the test are found in Table 2.

Table 2. Information for the *D. magna* immobilisation test (OECD 202), which was performed on the three test solutions.

Test method	The method is in accordance with the OECD Guideline 202; "Daphnia sp. acute immobilization test"
Test organism	<i>Daphnia magna</i> , clone A (Baird, 1991). Maintained semistatically in Elendt M7 (Elendt, 1990) and fed <i>Pseudokirchneriella subcapitata</i> grown in 10 % Z8 nutrient salt solution and <i>Chlamydomonas reinhardtii</i> grown in 20 % Z8 nutrient salt solution (Staub, 1961). Age at start of test < 24 hours.
Test period	MEA: 10.12.2012 – 12.12.2012 TW: 12.12.2012 – 14.12.2012 ARW: 18.12.2012 – 20.12.2012
Pretreatment of sample:	MEA: Stock solution (10 g/L) was adjusted to pH 7.9 with 18.5 % HCl TW: Filtered with 0.45µm membrane filter ARW: Filtered (0.45µm membrane capsule filter) and adjusted pH with 1M HCl in each test concentration
Dilution medium:	ISO 6341 (ISO, 1996)
Test concentrations	MEA: 18, 31, 55, 98, 176, 312 mg/L TW: 0.56, 1.0, 1.8, 3.2, 5.6 % v/v ARW: 0.009, 0.016, 0.030, 0.056, 0.1 % v/v
Replicates	4 vessels for each concentration, with 5-7 animals per vessel
Test containers	50 ml polystyrene cups with ca. 40 ml medium
Observations:	Every 24 ± 2 hours with a microscope
Temperature	Max: 20.7 Min: 19.3
pH in control (Start – End)	MEA: 7.96 – 7.87 TW: 8.20 – 8.05 ARW: 8.14 – 7.97
pH in highest conc. (Start – End)	MEA: 7.87 – 7.97 TW: 8.34 – 8.48 ARW: 8.10 – 7.96
O₂ saturation, 48 t Highest conc.	MEA: 9.77 mg/L Control: 9.76 mg/L TW: 9.43 mg/L Control: 9.49 mg/L ARW: 9.02 mg/L Control: 8.43 mg/L
Calculation of EC_x[*]	Post hoc logarithmic regression between values of highest concentration without immobilized animals and lowest concentration with 100 % immobilized animals. Performed using Microsoft Office 2010 Excel.

3.4 Zebra fish embryo test

Embryos of the zebra fish, *Danio rerio* were obtained from the Norwegian Veterinary Institute, Oslo. The test method was based on the OECD draft guideline 'Zebra fish Embryo Toxicity Test'. The specifics of the tests are provided in **Table 3**.

The test was initiated immediately after fertilization and continued for 96 hours in duration. Lethal effects were recorded every 24 hours and were based on four apical observations (coagulation of the

* EC_x = The concentration which gives x % immobilization of the test animals.

embryo, non-detachment of the tail, non-formation of somites, and non-detection of the heartbeat). Observations of any one of these four malformations were indicative of lethality. This was compared to the occurrence in the dilution water control to provide sufficient information to calculate lethal concentration (LC) toxicity endpoints.

Table 3. Information on the zebra fish embryo toxicity test, which was performed on the three test solutions.

Test method	OECD draft guidelines ' <i>Zebra fish Embryo Toxicity Test</i> '
Test organism	Zebra fish (<i>Danio rerio</i>) embryos, obtained from the Norwegian Veterinary Institute, Oslo.
Test period	12.12.12 – 15.12.12 (96 h)
Pretreatment of sample:	pH adjusted with 10M HCl
Dilution medium:	Reconstituted freshwater from the Norwegian Veterinary Institute
Test concentrations	MEA: 10, 32, 100, 320 and 1000 mg/L ARW: 0.001, 0.01, 0.1, 1.0 and 10% v/v of original concentration TW: 0.1, 0.32, 1.0, 3.2 and 10% v/v of original concentration
Replicates	20 embryos per test concentration
Test containers	24-well plate
Temperature	27 ± 1°C (incubator controlled temperature)
pH	8.27 – 8.70

4. Results

4.1 Physicochemical parameters of the test mixtures

Due to the high chemical oxygen demand (COD), the waste water mixtures required dilution to 10,000 times its original concentration before analysis. This dilution resulted in some of the parameters falling below the detection limit and in these cases data was not reported.

Table 4. Measured parameters of the amine waste water mixtures

Description (g/L)	ARW (dark liquid)	Feed to bioreactor	Treated waste
COD	74.6	5.4	
NH ₄	8.87	1.01	2.39
K	-	0.23	0.19
Na	68.8	1.1	0.94
pH	11.1	9.82	7.86
conductivity	11.26	6.22	15.22
salinity	4.8	2.8	8.5

4.2 Toxicity to freshwater algae

The increase in cell numbers of the control group during the test was almost exponential. The variation between the control replicates were within the acceptable test criteria (**Table 5**).

MEA:

There was no observed difference in algal growth after 24 h of exposure to MEA up to a concentration of 177 mg/L (**Figure 1A**). However, after 48 h, toxic effects of the MEA on algal growth were evident. There were no differences in the effects seen from exposure to the two highest MEA concentrations with a plateau in the observed toxicity. Effect concentrations were calculated using a nonlinear regression with adjustable minimum effect and are shown in **Figure 2A**.

ARW:

There was no apparent effect of the lower concentrations of ARW (<0.01%). As with MEA, the two top concentrations had higher growth rates compared with the third highest test concentration (**Figure 1B**). Effect concentrations were calculated by nonlinear regression with adjustable minimum effect and are shown in (**Figure 2B**).

TW:

Since inhibition of algal growth was found in the lowest concentration of TW (3.2%) (**Figure 1C**), the test was repeated with a weaker concentration series (**Figure 1D**). However, the second test exhibited less inhibition at 3.2% TW as shown in the first test. Differences in age and sample storage are likely factors, since the first test was performed on a recently thawed sample that was stored at -20°C, whilst the second test was performed on the same sample that was stored in the dark at 4°C for 8 days. Effect concentrations were calculated by non-linear regression (**Figure 2C&D**).

Table 5. Validity criteria for the algal growth inhibition test (OECD, 201).

Criteria	Observed 3.12.12	Observed 11.12.12
Coefficient of variation in control less than 7 %	1.9 %	3.1 %
Coefficient of variation in section by section growth rate less than 35 %	16.1 %	34.6 %
More than 16 times increase in cell concentration from the start to the end of experiment	49 times	53 times
pH increase in the control less than 1 unit	-0.1	0.1




Figure 1. Average cell density over time when exposed to concentrations of the test media: A) MEA (mg/L); B) ARW (%); C) TW (%) 1st test; D) TW (%) 2nd test.




Figure 2. Calculation of effect concentrations of A) MEA, B) ARW C) TW 1st test and D) TW 2nd test.

All test substances were found to inhibit the growth of the algae *P. subcapitata*, although MEA and ARW did not show full inhibition at the highest concentrations tested, but instead the effect appeared to reach a plateau. This was also observed in a previous test with MEA, where test concentrations were even higher (Brooks et al., 2009).

The inhibiting effect of TW on algal growth was lower in the aged (8 days) sample than a freshly thawed sample. The toxic substances in the aged sample might have evaporated or degraded during that time, resulting in lower toxicity. The effect concentrations are summarized in **Table 6**.

Table 6. Effect concentrations of the tested substances on the growth of *P. subcapitata*. For TW the EC₅₀ is calculated from the first test whilst the EC₁₀ is calculated from the repeated test.

Substance	EC ₅₀ (95% confidence interval)	EC ₁₀ (95% confidence interval)
MEA (mg/L)	151	30
TW (%)	2.4 (1.9 – 2.9)	0.74 (0.58 – 0.98)
ARW (%)	0.019	0.0089

4.3 Toxicity to *Daphnia magna*

The test achieved the validity criteria described in OECD 202 “*Daphnia* sp. acute immobilization test” (Table 1). In all tests a monotone concentration-response was observed (table 2). Due to only one partial response in each test EC50 at 48 h was determined by logarithmic regression (figure 1).

Table 7. Validity criteria for the *Daphnia* sp. acute immobilisation test

Criteria	Observed
Less than 10 % immobilization in control	MEA: 0 % TW: 0 % ARW: 3.1 %
≥3 mg/L dissolved oxygen by the end of test in highest test concentration	MEA: 9.77 mg/L TW: 9.43 mg/L ARW: 9.02 mg/L

Table 8. Observed immobilised *D. magna* after 24 and 48 hours in control and test media

Test substance and concentration		# Daphnia	Immobilized 24 h	Immobilized 48 h	pH start	pH 48 h	O ₂ 48 h (mg/L)
MEA	Control	33	0	0	7.96	7.87	9.76
	18	20	0	0	-	7.87	-
	31	21	0	0	7.95	7.95	-
	55	21	0	0	7.82	7.95	-
	98	21	0	0	7.70	7.94	-
	176	24	3	4	7.87	7.96	-
	312	21	4	20	7.87	7.97	9.77
TW	Control	34	0	0	8.20	8.05	9.49
	0.56	20	0	0	8.29	8.07	-
	1	21	0	0	8.28	8.19	-
	1.8	21	0	0	8.29	8.29	-
	3.2	21	0	7	8.34	8.46	-
	5.6	20	20	20	8.34	8.48	9.43
ARW	Control	32	0	1	8.14	7.97	8.43
	0.009	22	0	0	7.89	8.23	-
	0.016	21	0	0	7.98	8.03	-
	0.03	22	0	0	8.01	8.01	-
	0.056	22	0	0	8.03	8.00	-
	0.10	22	3	17	8.10	7.96	9.02




Figure 3. Calculated concentration response curves for A) MEA, B) TW and C) ARW with respect to 24 h and 48 h survival in *D. magna*.

All test substances had a toxic effect on *Daphnia magna* (**Table 9**). ARW was almost 50 times more toxic than TW. EC₅₀ above 100 mg/L is not considered toxic to the aquatic environment

Table 9. Summary of the ecotoxicity data for MEA, TW and ARW on the immobilisation of *D. magna*.

Test substance	Unit	48 hours		
		EC ₅₀	95% conf. int.	EC ₁₀
MEA	mg/L	209		128
TW	%	3.4		2.2
ARW	%	0.081		0.060

4.4 Toxicity to the embryo-larvae of the zebra fish

Table 10. The numbers of successfully hatched and surviving embryos of the zebra fish (*Danio rerio*) following 96 h exposure to the test media indicated.

Treatment	Concentration	Hatched survival after 96h (n=20)	% hatched survival after 96h
Untreated waste (ARW)	control	17	85
	0.001%	18	90
	0.01%	16	80
	0.10%	12	60
	1%	2	10
	10%	0	0
Treated waste (TW)	control	17	85
	0.10%	17	85
	0.32%	18	90
	1%	18	90
	3.20%	1	5
	10%	0	0
MEA	control	17	85
	10 mg/L	18	90
	32 mg/L	17	85
	100 mg/L	18	90
	320 mg/L	12	60
	1000 mg/L	6	30

All test substances were found to be toxic to the developing larvae of the zebra fish. Of the two mixtures, ARW was the most toxic with a calculated EC₅₀ of 0.194 % of its original concentration compared to 1.91% for the TW (**Table 11**). Pure MEA had an EC₅₀ of 617.5 mg/L.




Figure 4. The percentage of surviving hatched larvae following 96 h exposure to MEA and to amine waste water mixtures, ARW and Treated amine waste.

Table 11. Summary of the ecotoxicity data for the zebra fish larvae after 96 h exposure. Calculated with ToxCalc scientific.

	96 h exposure			
	NOEC	LOEC	EC ₁₀	EC ₅₀
MEA (mg/L)	320	1000	164.6	617.5-
ARW (%)	0.1	1.0	0.034	0.194
TW (%)	1.0	3.2	-	1.91

5. Conclusion

Overall, of the three taxonomic groups tested, the algae was found to be the most sensitive to both MEA and ARW followed by daphnids, with the zebra fish embryo the least sensitive. For the algae, EC₅₀ concentrations of 151 mg/L MEA, 0.019% ARW and 2.4% TW, where similar to that recorded in previous tests performed at NIVA in 2009 (i.e. 127 mg/L MEA, 0.014% Untreated waste, 12.1% TW, Brooks et al., 2009).

For the TW a different pattern in toxicity was found with the lowest EC₅₀ concentration calculated in the zebra fish followed by the algae and daphnids. However, the EC₅₀ concentrations for TW were very close ranging from 1.91% TW in the zebra fish to 3.4% TW in daphnids. When compared to previous testing, the TW toxicity was found to be approximately 10 fold lower to the algae than that previously reported (Brooks et al., 2009).

In the previous study (Brooks et al. 2009) the TW was not toxic to the zebra fish after 48 h at the maximum concentration tested 10% TW. This was also the case in the present study, although toxicity was observed after the 96 h exposure duration.

Table 12. Summary of ecotoxicity endpoints for the three test chemicals for the three trophic groups

Test chemical	Trophic group	EC ₁₀	EC ₅₀
MEA (mg/L)	Algae	30	151
	Daphnids	128	209
	Zebra fish	164.6	617.5-
ARW (%)	FW algae	0.0089	0.019
	Daphnids	0.060	0.081
	Zebra fish	0.034	0.194
TW (%)	FW algae	0.74	2.4
	Daphnids	2.2	3.4
	Zebra fish	-	1.91

6. References

- Baird, D. J. et al, 1991, *A Comparative Study of Genotype Sensitivity to Acute Toxic Stress Using Clones of Daphnia magna Strauss*, Ecotoxicology and Environmental Safety, 21, 257 - 265.
- Brooks, S., Lillicrap, A., Heiaas, H., 2009. Ecotoxicity Testing of an amine waste water samples. NIVA, NIVA, p. 12.
- Elendt, B.-P. 1990, *Selenium deficiency in Crustacea; An ultrastructural approach to antennal damage in Daphnia magna Strauss*. Protoplasma, 154, 25-33.
- Hill. A.V. 1910: The possible effects of aggregation of the molecules of hemoglobin on its dissociation curves. J. Physiol. (Lond.) 40. IV-VII.
- ISO/DIS 8692: 1996. Water quality - Algal growth inhibition test
- OECD 2011: Guidelines for testing of chemicals. no. 201; Alga. growth inhibition test. OECD. Paris
- Staub, R., 1961, *Ernährungsphysiologische Untersuchungen an der planktischen Blaualge Oscillatoria rubescens*, D. C., Schweiz, Z., Hydrol, 23, 82-198.
- Vindimian. E. Robaut. R. and Fillion. G. 1983: A method for co-operative and non-cooperative binding studies using non-linear regression analysis on a microcomputer. J. Appl. Biochem. 5: 261-268.

NIVA: Norway's leading centre of competence in aquatic environments

NIVA provides government, business and the public with a basis for preferred water management through its contracted research, reports and development work. A characteristic of NIVA is its broad scope of professional disciplines and extensive contact network in Norway and abroad. Our solid professionalism, interdisciplinary working methods and holistic approach are key elements that make us an excellent advisor for government and society.



Norwegian Institute for Water Research

Gaustadalléen 21 • NO-0349 Oslo, Norway
Telephone: +47 22 18 51 00 • Fax: 22 18 52 00
www.niva.no • post@niva.no