





Norwegian State Pollution Monitoring Programme

Report 716/97

Client State Pollution Control Authority

Contractor NIVA

Joint Assessment and Monitoring Programme (JAMP) National Comments to the Norwegian Data for 1996



Main Office

P.O. Box 173, Kjelsås
N-0411 Oslo
Norway
Phone (47) 22 18 51 00
Telefax (47) 22 18 52 00

Regional Office, Sørlandet

Televeien 1
N-4890 Grimstad
Norway
Phone (47) 37 29 50 55
Telefax (47) 37 04 45 13

Regional Office, Østlandet

Sandvikaveien 41
N-2312 Ottestad
Norway
Phone (47) 62 57 64 00
Telefax (47) 62 57 66 53

Regional Office, Vestlandet

Nordnesboder 5
N-5008 Bergen
Norway
Phone (47) 55 30 22 50
Telefax (47) 55 30 22 51

Akvaplan-NIVA A/S

Søndre Tollbugate 3
N-9000 Tromsø
Norway
Phone (47) 77 68 52 80
Telefax (47) 77 68 05 09


Title Joint Assessment and Monitoring Programme (JAMP). National Comments to the Norwegian Data for 1996 (Norwegian State Pollution Monitoring Programme Report no. 716/97. TA-no. 1489/1997)	Serial No. 3730-97	Date 1997.10.24
	Report No. Sub-No. O-80106	Pages Price 129
Author(s) Norman W. Green	Topic group Marine ecology	Distribution
	Geographical area Oslofjord to Varangerfjord	Printed NIVA

Client(s) Norwegian State Pollution Control Authority (SFT)	Client ref.
---	--------------------


Abstract

This report is part of the Norwegian contribution to the SIME 1997(2) meeting administrated by OSPARCOM. JAMP 1996 included the monitoring of micropollutants in blue mussels (30 stations) and fish (13 stations) at along the entire coast of Norway. The results indicated elevated levels of contaminants (i.e. over provisional "high background") in: Oslofjord proper (PCBs and mercury in fish) and Langesundsfjord (HCB in mussels) and Sør fjord and Hardangerfjord (cadmium, lead, mercury and ppDDE in mussels). Significant downward trends were found for cadmium in mussels from the Sør fjord and Hardangerfjord for the period 1987-1996. A mussel station in a harbour at Lofoten had elevated levels of mercury and PCBs. The results from the remainder of stations showed primarily low levels contamination. Introductory studies of PAH and "dioxins" in mussels were conducted. Overconcentrations of PAH were found in the inner Oslofjord and three stations in the North of Norway. No change in "pollution" or "reference" index classification was found from 1995 to 1996.

4 keywords, Norwegian 1. Miljøgifter 2. Organismer 3. Marin 4. Norge	4 keywords, English 1. Micropollutants 2. Organisms 3. Marine 4. Norway
---	--


 Norman W. Green
 Project manager

ISBN 82-577-3299-0


 Bjørn Braaten
 Head of research department

OSLO AND PARIS CONVENTIONS FOR THE PREVENTION OF MARINE POLLUTION
WORKING GROUP ON CONCENTRATIONS, TRENDS AND EFFECTS OF SUBSTANCES
IN THE MARINE ENVIRONMENT (SIME)

BONN 17-21 NOVEMBER 1997

O-80106

**JOINT ASSESSMENT AND MONITORING PROGRAMME (JAMP)
NATIONAL COMMENTS TO THE NORWEGIAN DATA FOR 1996**

Oslo, 24. October 1997

Submitted by: NORWAY
Norman W. Green
Norwegian Institute for Water Research (NIVA)
PB 173 Kjelsås
N-0411 Oslo

Submitted to: Oslo and Paris Commissions (OSPARCOM)
att. Carolyn Symon
New Court
48 Carey Street
London WC2A 2JQ

Copy to: Norwegian State Pollution Control Authority (SFT)
att. Per Erik Iversen
PB 8100 Dep.
N-0032 Oslo

International Council for the Exploration of the Seas (ICES)
att. Janet Pawlak
Palægade 2-4
DK-1261 Copenhagen

Foreword

This report presents the Norwegian national comments on the 1996 investigations for the Joint Assessment and Monitoring Programme (JAMP). JAMP is administered by the Oslo and Paris Commissions (OSPARCOM) and their Environmental Assessment and Monitoring Committee (ASMO). JAMP receives guidance from the International Council for the Exploration of the Sea (ICES). ASMO has delegated implementation of part of the programme to the Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME). The Norwegian 1996 investigations are directed to particular JAMP issues relating to contaminants and implemented by SIME. Some JAMP issues to be addressed lack adequate guidelines, in such cases guidelines used by the Joint Monitoring Programme (JMP) were applied.

The Norwegian JAMP for 1996 was carried out by the Norwegian Institute for Water Research (NIVA) by contract from the Norwegian State Pollution Control Authority (SFT), (NIVA contract 80106).

The Norwegian contribution to the JAMP was initiated by SFT in 1981 as part of the national monitoring programme. It now comprises three areas: the Oslofjord and adjacent areas (Hvaler-Singlefjord area and Langesundfjord, 1981-), Sør fjord/Hardangerfjord (1983-84, 1987-) and Orkdalsfjord area (1984-89, 1991-93, 1995-96).

Since the North Sea Task Force Monitoring Master Plan was implemented 1990 additional areas have also been monitored. These include: Arendal, Lista and Bømlo-Sotra areas. On the initiative of SFT and NIVA "reference" or merely diffusely contaminated areas from Bergen to Lofoten have been monitored since 1992 and from Lofoten to Norwegian-Russian border from 1994.

These comments are considered as preliminary notes on the 1996 results and are not to be viewed as a final assessment.

The comments are presented in accordance with the agreed standardised format (SIME 1997, Annex 12).

Thanks are due to many colleagues at NIVA, especially: Rita Amundsen, Unni Efraimsen, Frank Kjellberg, Tom Tellefsen for field work, sample preparations, data entry; Einar Brevik and his colleagues for organic analyses; Arne Godal, Marit Villø and Bente Hiort Lauritzen and their colleagues for metal analyses; Audun Rønningen and Gunnar Severinsen for data programme management; Jon Knutzen for constructive criticism. Thanks go also to Risøy Underwater Engineering and their crew aboard 'Risøy' for assisting in the field work and to the numerous fishermen and their boat crews we have had the pleasure working with.

Oslo, 24 October 1997.

*Norman W. Green
Project co-ordinator*

Contents

1. General Details	1
1.1 Executive Summary	1
1.2 Introduction	2
1.3 Information on measurements	2
1.3.1 Oslofjord area	7
1.3.2 Sør fjord and Hardangerfjord	12
1.3.3 Lista areas	17
1.3.4 Bømlo-Sotra area	17
1.3.5 Orkdalsfjord area	17
1.3.6 Open coast areas from Bergen to Lofoten	18
1.3.7 Open coast areas from Lofoten to Russian border	18
1.3.8 PAH and “dioxins”	18
1.3.9 Norwegian Pollution and Reference Indexes	19
1.4 Overall conclusions	20
2. Technical Details	21
2.1 Compliance with guidelines/procedures	21
2.1.1 JAMP programme	21
2.1.2 Overconcentrations and classification of environmental quality	21
2.1.3 Comparison with previous data	24
2.2 Information on Quality Assurance	24
2.3 Description of the Programme	25
3. References	26
Appendix A. Quality assurance programme	29
Appendix B. Abbreviations	33
Appendix C. Participation in intercalibration exercises	43
Appendix D. Analytical overview	47
Appendix E. Overview of localities	59
Appendix F. Overview of materials and analyses 1996	65
Appendix G. Temporal trend analyses of contaminants in biota 1981-96	71
Appendix H. Geographical distributions of contaminants in biota 1995-96	83
Appendix I. Results from INDEX determinations 1996	111

1. General Details

1.1 Executive Summary

The Norwegian JAMP 1996 included the monitoring of micropollutants (contaminants) in blue mussels (30 stations) and fish (13 stations) along the entire coast of Norway. The results indicated elevated levels of contaminants (i.e. over provisional “high background”) in:

- JAMP area 26: Oslofjord proper (PCBs and mercury) and Langesundsfjord (HCB). Lower concentrations of PCB (CB153) were found in liver and fillet of cod from the outer Oslofjord in 1996 compared to 1995.
- JAMP areas 63 and 62: Sjørfjord and Hardangerfjord (especially cadmium and lead and to a lesser degree mercury, zinc, ppDDE and PCBs). Significant downward trends were found for cadmium in mussels from the Sjørfjord and Hardangerfjord for the period 1987-1996.
- JAMP area 65: Orkdalsfjord. No overconcentrations were found in blue mussels.

The remainder of stations are mostly from presumed “reference” areas where only diffuse contamination is expected. The results indicate primarily low levels contamination. One exception is a mussel station in a harbour at Lofoten where elevated levels of mercury and PCBs were detected.

PAH and “dioxins” were measured at 6-12 mussel stations. Overconcentrations of PAH were found in the inner Oslofjord and three stations in the north of Norway. Overconcentrations of “dioxin” (TCDD) were found in the inner Oslofjord and in Langesundsfjord.

A new index was tested for the second year to assess the levels of contamination of mussels in “polluted” and “reference” areas. The results for the “pollution” index indicated a “bad” condition and the “reference” index indicated a “fair”. There was no change in classification from the 1995 results.

1.2 Introduction

The Norwegian contribution to the “Joint Assessment and Monitoring Programme (JAMP) was initiated by the Norwegian State Pollution Control Authority (SFT) and is integrated with SFT’s State Pollution Monitoring Programme. The procedures and practice of JAMP has also provided a basis for other investigations of interest to SFT but not necessarily requested by JAMP (e.g. SFT’s Pollution and Reference Indexes, Chapter 1.3.9).

Data are submitted to ICES under three categories: for Purpose A (Health assessment) on a voluntary basis, Purpose C (spatial distribution) on a voluntary basis and Purpose D (temporal trend assessment) on a mandatory basis. Where practical, data collection was in accordance to agreed procedures (OSPARCOM 1990). Data were screened and submitted to ICES in accordance with procedures outlined by ICES (1996).

This report focuses on issues and situations in Norway concerning contaminants and considered of interest to the implementation of JAMP (Table 1).

Table 1. JAMP issues to which the Norwegian investigations for 1996 can be addressed (cf., ASMO 1997, Annex 30).

Issue	Subject	Description
1.2	Hg, Cd, and Pb	What are the concentrations and fluxes in sediments and biota?
1.7	PCBs	Do high concentrations posed a risk to the marine ecosystem?
1.8	PCBs	Do high concentrations of non-ortho and mono-ortho CBs in seafood pose a risk to human health?
1.10	PAHs	What are the concentrations in the maritime area?
1.15	Chlorinated dioxins and dibenzofurans	What concentrations occur and have the policy goals (for the relevant parts of the maritime area) been met?

This report is structured according to agreed format (ASMO 1997, Annex 12) which *inter alia* presents results before methodology.

1.3 Information on measurements

An overview of JAMP stations in Norway is shown in maps in Figure 1-Figure 4 and Appendix E. The stations and sample counts relevant to the 1996 investigations are noted in Appendix E and F, respectively.

Blue mussels were sampled at 30 stations and fish from 13 stations from the border to Sweden in the south to the border to Russia in the north. Generally, mussels are not in abundance on the exposed coastline from Lista (south Norway) to the North of Norway. A number of samples were collected from dock areas, buoys or anchor lines (see footnotes in Appendix E).




Figure 1. JAMP sampling stations along the southern coast of Norway from the Swedish border to Bergen.




Figure 2. JAMP sampling stations along the western coast of Norway from Bergen to Namsos.




Figure 3. JAMP sampling stations along the Northwest coast of Norway from the region of Ranfjord to Lofoten.




Figure 4. JAMP sampling stations along the north coast of Norway from the region of Lofoten to the Russian border.

1.3.1 Oslofjord area

Moderate overconcentrations of CB153 in mussels were found in the inner Oslofjord (st.30A, up to 5 times provisional "high background" - see Chapter 2.1.2) (Figure 5, Appendix G).

Overconcentrations (see Chapter 2.1.2) were also found in cod liver from the inner Oslofjord (st.30B, up to 4 times "high background"; Figure 6). Further, overconcentrations were also for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in mussels and cod liver from these stations (st.30A and 30B), about 5 times "high background" (Appendix H). There has been no consistent change (see Chapter 2.1.3) for CB153 for mussel from four stations (30A, 31A, 35A and 36A) from 1987 to 1996 or for cod (30B and 36B) from 1990 to 1996.

The median concentration of CB153 in cod liver from the outer Oslofjord was roughly half that found in 1995 but higher than previous years (1990-1994, Figure 6, Appendix I). Similarly for cod fillet, the median concentration for 1996 was lower than for 1995 (by about a quarter) but higher than prior years. Flood conditions from Glomma resulted in high concentrations were found in cod 1995 (cf., Green 1997).

Power analyses (see Chapter 2.1.3) indicated that a hypothetical trend of 10% change per year in CB153 concentration in the blue mussel or cod liver from the inner Oslofjord would take up to 12 years to be detected with 90% significance (Appendix G).

Moderate overconcentrations of mercury (less than twice "high background") were found in the fillet of "large" cod (st.30B) from the inner Oslofjord (Figure 7).

The power, indicated as number of years, to detect a change in mercury in cod fillet from the inner Oslofjord was slightly better for "small" fish (10 years) than "large" fish (12 years) (cf., Appendix G).





Figure 5. Median CB153 concentration in blue mussel (*Mytilus edulis*) from inner (st.30A) to outer (st.36A) Oslofjord. (cf., Figure 1 and Figure 14).


A

CB153 *Gadus morhua*, liver, st.30B




B

CB153 *Gadus morhua*, liver, st.36B



C

CB153 *Gadus morhua*, fillet, st.30B



D

CB153 *Gadus morhua*, fillet, st.36B





Figure 6. Median CB-153 concentration in liver and fillet of cod (*Gadus morhua*) from the inner (st.30B) to outer (st.36B) Oslofjord. (cf., Figure 1 and Figure 14). **Note: for some years the upper confidence interval line is off-scale in Figure D.**


A

HG Gadus morhua Small, fillet, st.30B




B

HG Gadus morhua Large, fillet, st.30B



C

HG Gadus morhua Small, fillet, st.36B



D

HG Gadus morhua Large, fillet, st.36B




Figure 7. Median mercury (Hg) concentration in fillet of cod (*Gadus morhua*): for the inner Oslofjord (st.30B) “small” (A) and “large” (B) fish, and for the outer Oslofjord (st.36B) “small” (C) and “large” (D) fish. (cf., Figure 1 and Figure 14).

Mussels from Langesundsfjord (st.71A) had in 1996 marked overconcentrations of HCB (over 9 times “high background”, Appendix G), and was four times higher than median for 1995. Concentrations have varied greatly during the investigation period (since 1983) but median value have decreased distinctly since 1989 (Figure 8) due to about a 99% reduction in discharge of HCB and other organochlorines from a magnesium factory (cf., Knutzen *et al.* 1996).

Note that the variability in the data is much less after 1989. The relatively large variability found in this series prior to 1990 accounts for the poor power. The power of the monitoring program (expressed in years) for the period 1990-96 is 13 years and better than the power for the entire period which is over 25 years (cf., Appendix G for entire period). The separate analysis for the 1990-96 data also indicated a significant *downward* trend for this period. Moderate overconcentrations of mercury and CB153 (less than twice “high background”) were also found.




Figure 8. Median HCB concentration in blue mussel (*Mytilus edulis*) from Langesundsfjorden (west of Oslofjord). (cf., Figure 1 and Figure 14). **NB: log-scale.**

1.3.2 Sør fjord and Hardanger fjord

The development of the contaminant conditions in these connected fjords and the main remedial actions that have been taken have been outlined in the national comments for 1989 (Green 1991). The results from JAMP 1996 are coupled to other studies in this area (cf., Knutzen *et al.*, in press) and confirm that the fjords continue to be contaminated especially with cadmium (Figure 9 and Figure 10) and lead and to a lesser degree ppDDE (Figure 11 and Figure 12) and mercury.

Results for mussels collected from Sør fjorden (st. 51A, 52A, 56A and 57A) indicated severe overconcentrations of cadmium (up to 12 times provisional "high background", Appendix G) and lead (up to 9 times "high background"), more moderately for mercury and zinc (up to 4 times). Overconcentrations of cadmium and lead could be traced to Ranaskjær (st.63A) in the Hardanger fjord. A significant *downward* trend was found for cadmium at st.57A and 63A from 1987 to 1996 (Appendix G).

Overconcentrations of ppDDE (up to 3 times, "high background"; up 6 times for Σ DDT) were also found in mussels from the three stations in the Sør fjorden, the highest at st.56A, midway along the fjord (Figure 11 and Figure 12, Appendix G). Concentrations in cod liver from Sør fjord and Hardanger fjord (st.53B and 67B, respectively) were up to two times "high background" and higher than median values from 1995 (Figure 13). No trend was evident for the period 1990-96.

The source of ppDDE is uncertain but the Sør fjord and Hardanger fjord area has a considerable number of fruit orchards. Earlier use and persistence of DDT and leaching from contaminated soil is probably the main reason for the elevated levels found. DDT products has been prohibited in Norway since 1970 (excepting the dipping of spruce seedling until 1987). One possible source may be DDT-contaminated material buried in the vicinity of st.56A (Knutzen *et al.*, in press).

Overconcentrations of cadmium were found in flounder liver from Sør fjord and Hardanger fjord, up to eight times "high background". The median concentration in flounder liver from Sør fjorden (st.53B) was over 10 times the median concentration found in 1995. Slight overconcentrations (less than two times "high background") were also found in cod liver from the inner Sør fjord and from Hardanger fjord.

Unexpected and high discharge from industrial areas at the head of the Sør fjord during the period August to December resulted in abnormally high concentrations of cadmium and zinc in sea water in the vicinity (Skei 1997). Preliminary results from the VIC programme (cf., SIME 1996, 1997) indicated no significant local differences for these two metals for flounder (three sites) or cod (two sites). However the concentrations of these metals in cod liver was significantly higher in samples collected in August compared to December.

Moderate overconcentrations (less than two times "high background") were also found for zinc in mussels, mercury in fish, CB153 in cod-liver and lead in flounder. The variability in concentrations of CB153 in the 25 cod from the inner Sør fjord was large (cf., Appendix H) and indicate that exposure and/or net uptake of this contaminant varies considerably from fish to fish.

The power of the sampling strategies for mussels was relatively poor for samples collected from Odda, the head of the Sør fjord (st.52A). For example for lead in mussels, it is estimated that it would take over 25 years to detect hypothetical trend of 10% per year with 90% significance (Appendix G). This reflects the large variability found in the data series from this area. The variability is largely due to the irregular/accidental input of contaminated discharges. The power improved with distance from Odda.




Figure 9. Median cadmium (Cd) concentration in blue mussel (*Mytilus edulis*) from inner (st.52A) to outer (st.57A) Sør fjord. NB: (cf., Figure 1 and Figure 14). **Note: for some years the upper confidence interval line is off-scale in figures A and B.**




Figure 10. Median cadmium (Cd) concentration in blue mussel (*Mytilus edulis*) from Hardangerfjord (st. 63A, 65A and 69A). (cf., 1 and Figure 14). **Note difference in scale from Figure 9 and that for some years the upper confidence interval line is off-scale in Figure A.**




Figure 11. Median ppDDE (DDEPP) concentration in blue mussel (*Mytilus edulis*) from inner (st.52A) to outer (st.57A) Sør fjord. (cf., Figure 1 and Figure 14).




Figure 12. Median ppDDE (DDEPP) concentrations in blue mussel (*Mytilus edulis*) from Hardangerfjord (st. 63A, 65A and 69A). (cf., Figure 1 and Figure 14). **Note difference in scale compared to Figure 11.**




Figure 13. Median ppDDE (DDEPP) concentrations in cod (*Gadus morhua*) from Sør fjord (st.53B) and Hardanger fjord (st.67B) (cf., Figure 1 and Figure 14). **Note that for some years the upper confidence interval line is off-scale in Figures A and B.**

1.3.3 Lista areas

No significant overconcentrations of metals or chlorinated hydrocarbons were found in mussels, cod, or dab (st.15A/B/F, Figure 1, Appendices G and H).

1.3.4 Bømlo-Sotra area

No significant overconcentrations of metals or chlorinated hydrocarbons were found in mussels, cod, or plaice from this area (st. 22A/F and 23B, Figure 1, Appendices G and H). Sufficient samples of dab were not recovered to investigate for elevated levels of mercury (Appendix H, cf., Green 1997).

1.3.5 Orkdalsfjord area

No significant overconcentration of metals or chlorinated hydrocarbons were found in Orkdalsfjord (st. 84A, 82A, and 87A, Figure 2, Appendices G and H).

1.3.6 Open coast areas from Bergen to Lofoten

Only two mussel stations (st.92A and 98X) were investigated in the open coast areas from Bergen to Lofoten, covering 7° of latitude to 68°N (Figure 2 and Figure 3). Unable to obtain mussels from st.98A, as was done in 1993, mussels were collected from nearby Skrova harbour (st.98X). Only cod was collected in the Froan area (st.92B), whereas cod and dab were collected from the Lofoten area (st.98B/F).

Mussels from 98X had moderate overconcentrations (up to 2 times “high background”) of CB153 and slight overconcentrations (less than 2 times) of mercury and lead on a dry weight basis (Appendix G, see also Appendix H). Overconcentration of ppDDE was not found, whereas in 1995 the median value was 3 times higher than the 1996 value and slightly over “high background”. The enhanced concentrations found at this station may be related to fish and sea mammal processing activity in the harbour where the mussels were collected. Overconcentrations of PCBs and Σ DDT are not uncommon in harbours of northern Norway (Knutzen *et al.* 1995, Konieczny & Juliussen 1995, Konieczny 1996).

Moderate overconcentrations (up to 3 times “high background”) of cadmium were found in the livers of cod from st.98B. There were also slight overconcentrations of HCB in cod liver and mercury in dab liver. Slight overconcentration of mercury was found in cod liver from st.92B.

1.3.7 Open coast areas from Lofoten to Russian border

Six mussel stations were investigated in the open coast areas from Lofoten to the Russian border, north of 68°N and a longitude from 17 to 29°E (Figure 3 and Figure 4). In addition, cod was sampled in Kvænangen (st.43B) and Varangerfjorden (st.10B), and lemon sole and dab was sampled in Kvænangen (st.43F).

Slight overconcentrations (less than 2 times “high background”) of cadmium were found at two mussel stations (st. 46A and 10A) (Appendix G and H). Moderate overconcentration (over 4 times “high background”) of cadmium was also found in the homogenate lemon sole liver (st.43F). There was also moderate overconcentration of lead in mussels from st.44A. The relatively large variability in concentrations PCBs (CB153) found in cod from st.10B 1995 was not confirmed in 1996 (cf., Appendix H).

1.3.8 PAH and “dioxins”

Concentrations of PAH were measured at six blue mussel stations: one in the inner Oslofjord (st.30A) and five in the North of Norway (Appendix H). Moderate overconcentrations (up to 4 times “high background”) of Σ PAH, excluding the dicyclic, were found at st.44A and 30A. Highest concentrations were found at st.44A where the median value was slightly higher than the median for 1995. The dicyclics represented 18-86% of Σ PAH, highest at st.10A. This may indicate a significant local influence from small boat traffic. Benzo[a]pyrene (BAP) was below “high background” at all six mussel stations.

“Dioxins” (PCDD and PCDF) were investigated in blue mussels at twelve stations spread along the entire coast (cf. Appendix H). The results expressed as sum TE_{PCDF/D} (PCDF/D toxicity equivalents, JAMP code TCDDN) after Nordic standard (Ahlborg 1989) were below “high background” of 0.2 ng/kg (Appendix H) with two exceptions: st.30A and 71A. The TCDDN concentration at st.71A was over 60 times “high background” and is located near the mouth of Frierfjord with a known source of dioxin (cf., Knutzen *et al.* 1996). Moderate overconcentrations (over 2 times “high background”) were found at st.30A in the inner Oslofjord. The highest non-ortho and mono-ortho PCBs, expressed as sum TE_{CB} (JAMP code TECBW) after WHO standard (Ahlborg, *et al.* 1994) was at st.30A (Appendix H).

1.3.9 Norwegian Pollution and Reference Indexes

The Norwegian State Pollution Control Authority (SFT) is interested in obtaining a select and small group of indices to assess the quality of the environment with respect to contaminants. One index is based on the levels and trends of contaminant concentrations in the blue mussel collected annually from 10-11 of the more contaminated fjords in Norway (Appendix I). SFT has also requested the testing of this index against “reference” stations from selected areas and fjords.

The Index scale varies from 1, in which no overconcentrations were found at any station, to 5, in which at least one sample from each area or fjord could be classified as “very bad” in SFT's system.

The Pollution Index was recalculated for 1995 because of the revisions made in SFT's classification system, which involved lower limits for *inter alia* BAP and HCB. The Index for 1995 is 3.2 compared to 2.8 based on the old system (Appendices I5). The Index for 1996 is 3.1. A value between 3 and 4 would be classified by the SFT system as “Bad”.

The recalculated Reference Index for 1995 is 1.6 and unchanged compared to the old system. The Index for 1996 is 1.8 (Appendices I6). The calculation included (low) suspect dioxin values (TCDDN). A value between 1 and 2 would be classified as “Fair”.

1.4 Overall conclusions

In regards to JMP/JAMP Purpose A (health assessment), attention should be called to the list from Norwegian Food Control Authority (SNT) which names the of restrictions and recommendations concerning the sale and consumption of seafood's in Norway (Table 2).

Table 2. Summary of action taken by the Norwegian Food Control Authority (SNT) concerning the consumption and sale of fish products along the Norwegian Coast (SNT, pers. comm. 1997). **Contact SNT for more detailed information.**

Area of concern	Last year of issue/ evaluation	Main parameters of concern	Main fish/shellfish product of concerned	Recommendations or restrictions of concern:
Inner Oslofjord	1994	PCB	fish liver	Consumption
Drammensfjorden	1992	Dioxins/PCB	cod liver	Consumption and Sale
Sandefjordfjorden	1993	PCB	round fish liver	Consumption and Sale
Grenlandsfjordene, Langesundsfjord	1997	Dioxins	fish, shellfish	Consumption and Sale
Kristiansandsfjorden	1994	Dioxins/PCB	fish, shellfish	Consumption and Sale
Fedafjorden	1995	PAH	flat fish, shellfish	Consumption
Saudafjorden	1992	PAH	fish liver, mussels	Consumption
Sørfjorden and Hardangerfjorden	1997	Cd Pb Hg,	mussels	Consumption
Bergen area including Herdlefjorden, Byfjorden, Hjeltefjorden, Grimstadfjorden and Raunefjorden	1996	PCB	fish, shellfish	Consumption and Sale
Årdalsfjorden	1995	PAH	mussels	Consumption
Sunnalsfjorden	1993	PAH	fish liver, mussels	Consumption
Hommelvik (Trondheimsfjorden)	1985	PAH	mussels	Consumption
Ranfjorden	1997	PAH Pb Hg	mussels	Consumption
Vefsnfjorden	1992	PAH	mussels	Consumption

In regards to JMP/JAMP Purpose C (spatial distribution assessment), the concentrations found in 1996 are indicated in the bar graphs shown in Appendix H. Provisional "high background" levels were used to identify elevated concentrations. This initial assessment revealed no new areas that are not currently under surveillance.

In regards to JMP/JAMP Purpose D (temporal trend assessment) there is evidence that the median concentrations of cadmium in mussels from the Sørfjord has decreased since 1987. A separate analysis of HCB concentrations in mussels from the Langesundsfjorden 1990-1996 have decreased significantly.

The power of temporal trend monitoring expressed as an estimate of the number of years it would take to detect a hypothetical trend of 10% per year with 90% significance (power analysis) is a useful tool in assessing existing sampling strategies. However, in some cases the results can be misleading (cf., HCB-results for mussels from Langesund, page 11) and indicate the need to develop statistical methods more suited to local conditions. Furthermore, there is a general need to obtain a better understanding of the source of variation of concentrations as a basis for assessing/improving sampling strategies and remedial action.

2. Technical Details

2.1 Compliance with guidelines/procedures

2.1.1 JAMP programme

Samples were collected and analysed where practical, according to OSPARCOM guidelines (OSPARCOM 1990) and screened and submitted to ICES by agreed procedures (ICES 1996).

2.1.2 Overconcentrations and classification of environmental quality

This report focuses on the principle cases where *median* concentrations exceeded provisional "high background" (normal). The median concentration can be derived from the tables in Appendix G or figures in Appendix H, depending on the year and concentration basis in question. The provisional "high background" limits are summarised in Table 3. The factor by which concentrations exceeded "high background" is termed **overconcentration**. "High background" limits have not been set for all contaminants and species. It should be noted that there is in general a need for periodical review and supplement of this list of limits in the light of results from reference localities and introduction of new analytical methods, and/or units. Because of changes in the limits, assessments of overconcentrations for years prior to 1996 made in this report may not correspond to figures and assessments made in previous national comments.

In addition to the use of "high background", the Norwegian State Pollution Control Authority's (SFT's) system for **classification of environmental quality** has been applied (Table 4).

No attempt has been made to compensate for differences in size groups or number of individuals of mussels or fish. The exception was with mercury in fish fillet where seven of seventeen data sets in this study showed significant differences between "small" and "large" fish (Appendix G). In regards to mussels, there is some evidence that concentrations do not vary significantly among the three size groups employed for this study (i.e. 2-3, 3-4 and 4-5 cm) (WGSAEM 1993).

The National Comments since 1994 have include two additional analyses. The first is that the upper 95% confidence interval for the last three sampling years is linearly projected for the next three years. This is in line with a proposal submitted earlier (Nicholson, *et al.* 1994) and is used to assess the likelihood of overconcentrations. This estimate is based on the results for the temporal trend analyses. The estimate was made for series with at least 6 years of data.

The second is an estimate of the power of the temporal trend series expressed as the number of years to detect a 10% change per year with a 90% power (cf., Nicholson, *et al.*, 1996b in press). The fewer the years the greater the power needed to detect a change. The power is based on the percentage relative standard deviation (RLSD) estimated using the robust method described ASMO (1994) and Nicholson *et al.* (1996a). The estimate was made for series with at least 3 years of data and covers the *entire* period monitored. This fixed means of treating all the datasets may give misleading results especially where non-linear temporal changes are known to occur, such as for HCB in blue mussels from Langesund (Figure 8).

With respect to Purpose A (health risk assessment), the Norwegian Food Control Authority (SNT) is responsible for official commentary as to possible health risk due to consumption of seafoods. Hence, the results of the JAMP pertaining to this purpose are presented only as a partial basis for evaluation.

Table 3. Provisional "high background levels" of selected contaminants, in **ppm (mg/kg) dry weight** (blue mussel) and **ppm (mg/kg wet weight)** (blue mussel and fish). The respective "high background" limits are from Knutzen & Skei (1990) with mostly minor adjustments (Knutzen & Green 1995; Molvær *et al.* 1997), except for dab where the suggested limit is based on JAMP-data (Knutzen & Green 1995). Especially uncertain values are marked with "?".

Cont.	Blue mussel ¹		Cod ¹		Flounder ¹		Dab ¹	
			liver	fillet	liver	fillet	liver	fillet
	ppm d.w.	ppm w.w.	ppm w.w.	ppm w.w.	ppm w.w.	ppm w.w.	ppm w.w.	ppm w.w.
Lead	3.0 ²⁾	0.6 ³⁾	0.1		0.3 ?		0.3 ?	
Cadmium	2.0 ²⁾	0.4 ³⁾	0.1		0.3 ?		0.3 ?	
Copper	10 ²⁾	2 ³⁾	20		30 ?		10 ?	
Mercury	0.2 ²⁾	0.04 ³⁾		0.1 ²⁾		0.1		0.1 ?
Zinc	200 ²⁾	40 ³⁾	30		60 ?		50 ?	
ΣPCB-7 ⁸⁾	0.020 ³⁾	0.004 ²⁾	0.5 ²⁾	0.005	0.10 ?	0.005 ? ²⁾	0.5 ?	0.010 ?
CB-153	0.005 ³⁾	0.001 ⁴⁾	0.2 ? ⁵⁾		0.05 ? ⁷⁾		0.20 ? ⁷⁾	
ppDDE	0.010 ³⁾	0.002 ⁶⁾	0.2 ²⁾		0.03 ? ⁶⁾		0.1 ? ⁶⁾	
γHCH	0.005 ³⁾	0.001 ⁶⁾	0.05 ^{2,6)}		0.01 ? ⁶⁾		0.03 ? ⁶⁾	
HCB	0.0005 ³⁾	0.0001 ²⁾	0.02 ²⁾		0.005 ?		0.01 ?	
TCDDN	0.000001 ³⁾	0.0000002 ²⁾						

- ¹⁾ Respectively: *Mytilus edulis*, *Gadus morhua*, *Platichthys flesus* and *Limanda limanda*.
- ²⁾ From the Norwegian State Pollution Control Authority Environmental Class I ("good") (Molvær *et al.* 1997).
- ³⁾ Conversion assuming 20% dry weight.
- ⁴⁾ Approximately 25% of ΣPCB-7 (Knutzen & Green 1995)
- ⁵⁾ 1.5-2 times 75% quartile (cf., Annex B in Knutzen & Green 1995)
- ⁶⁾ Assumed equal to limit for ΣHCH or ΣDDT, respectively, from the Norwegian State Pollution Control Authority Environmental Class I ("good") (Molvær *et al.* 1997). Hence, limits for ppDDE and γHCH are probably too high (lacking sufficient and reliable reference values)
- ⁷⁾ Mean plus 2 times standard deviation (cf., Annex B in Knutzen & Green 1995)
- ⁸⁾ Estimated as sum of 7 individual PCB compounds (CB-28, -52, -101, -118, -138, -153 and -180) and assumed to be ca. 50% and 70 % of total PB for blue mussel and cod/flatfish, respectively.

Table 4. Extracts of the Norwegian State Pollution Control Authority partly revised environmental classification system of contaminants in blue mussels and fish (from Molvær *et al.* 1997).

Contaminant		Classification (upper limit for classes I-IV)				
		I "good"	II "fair"	III "poor"	IV "bad"	V "very bad"
BLUE MUSSEL						
Lead	ppm d.w.	3	15	40	100	>100
Cadmium	ppm d.w.	2	5	20	40	>40
Copper	ppm d.w.	10	30	100	200	>200
Mercury	ppm d.w.	0.2	0.5	1.5	4	>4
Zinc	ppm d.w.	200	400	1000	2500	>2500
ΣPCB-7	ppb w.w.	4	15	40	100	>100
ΣDDT	ppb w.w.	2	5	10	30	>30
ΣHCH	ppb w.w.	1	3	10	30	>30
HCB	ppb w.w.	0.1	0.3	1	5	>5
TE_{PCDF/D} ¹⁾	ppp w.w.	0.2	0.5	1.5	3	>3
COD, fillet						
Mercury	ppm w.w.	0.1	0.3	0.5	1	>1
Cod, liver						
ΣPCB-7	ppb w.w.	500	1500	4000	10000	>10000
ΣDDT	ppb w.w.	200	500	1500	3000	>3000
ΣHCH	ppb w.w.	50	200	500	1000	>1000
HCB	ppb w.w.	20	50	200	400	>400

¹⁾ TCDDN (cf., Appendix B)

2.1.3 Comparison with previous data

A simple 3-model approach has been developed to study time trends for contaminants in biota based on *median* concentrations (ASMO 1994). A variation of this method was applied to mercury in fish fillet to distinguish trends in "large" and "small" individuals. The method was first used on a large-scale basis by the Ad Hoc Working Group on Monitoring that met in Copenhagen 8-12. November 1993 (MON 1993). At this meeting it was agreed to apply the method on contaminants in fish muscle and liver on a wet weight basis and contaminants in soft tissue of mussels on a dry weight basis. The results for this assessment are presented earlier (cf., ASMO 1994). The method has been applied to Norwegian data and results are shown in Appendix G. The results can be presented as in Figure 14.




Figure 14. Example presentation of variation in contaminant concentration with time. The figure shows median concentrations, running mean of median values, 95% confidence interval. The provisional "high background level" is marked with a horizontal line and corresponds to values listed in Table 3 (see text).

The statistical analysis was carried out on temporal trend data series for cadmium, copper, mercury, lead, zinc, the PCB congener CB153, ppDDE (ICES code DDEPP), γ -HCH (ICES code HCHG) and HCB. Assessment focused on individual compounds instead of "sum variables". CB153 was chosen because it is persistent and may act as an indicator for other congeners (Atuma *et al.* 1996). Furthermore, there is some evidence that CB153 may correlate with TCDD-equivalents (Boer *et al.* 1993).

2.2 Information on Quality Assurance

NIVA has participated in all the QUASIMEME international intercalibration exercises, including Round 6 (1996). These exercises have included nearly all the contaminants analysed for JAMP. Quality assurance programme for NIVA is similar to the 1995 programme (cf. Green 1995). In addition, NIVA was accredited in 1993 in accordance with the EN45000 standard by the Norwegian Accreditation (reference P009). A summary of the quality assurance programme at NIVA is given in Appendix A. A summary of the intercalibrations exercises that NIVA has participated in is given in Appendix C.

Four JAMP mussel samples were analysed for "dioxins" (PCDDs and PCDFs) and non-ortho chlorobiphenyls by the Norwegian Institute for Air Research (NILU). NILU has participated in seven relevant international intercalibration exercises from 1990 to 1996 (NILU, pers. comm.)

2.3 Description of the Programme

The sampling for 1996 involved sampling of blue mussel at thirty stations and at least one flatfish species and/or cod was sampled at thirteen stations. The Norwegian JAMP has been expanded since 1989 to include monitoring in more diffusely polluted areas. Though new stations are initially intended for temporal trends, there has not always been sufficient funds to do this in following years. Hence, sample/station reduction measures have taken to reduce costs. Furthermore, sufficient samples have not always been practical to obtain. When this applies to mussels a new site in the vicinity is often chosen. As for fish, the quota of 25 individuals ($\pm 10\%$), indicated in Appendix F as either 25 individuals or 5 bulked samples consisting of 5 fish per bulked sample, was met for all stations except for flatfish at st.67B (flounder) and 43F (dab and lemon sole). Extra sampling may occur where a bi-catch of fish provides 5 or more fish of a priority species or co-ordination with a another programme such as VIC SIME 1996, 1997). Initiation of the VIC programme allowed supplementary sampling at st.30B (cod), 33B (flounder), 53B (cod and flounder) and 67B (cod). Appendix E and F gives an overview of the planned and realised sampling.

Concentrations of metal, chlorinated hydrocarbons (including pesticides) and polycyclic aromatic hydrocarbon in mussels and fish were determined at the Norwegian Institute for Water Research (JAMP code NIVA). An overview of the methods applied up to and including 1992 sample material has been presented by Green (1993, also JMG document 19/7 info 3). Only minor modifications have been made since. An overview of analyses applied from 1981 to 1996 ('97) for biological material is given in Appendix D. Parameter abbreviations are given in Appendix B.

A description of the components of variability will be provided with the completion of the Norwegian contribution to the VIC programme.

3. References

Titles translated to English in square brackets [] are not official.

- Ahlborg, U.G., 1989. Nordic risk assessment of PCDDs and PCDFs. *Chemosphere* 19:603-608.
- Ahlborg, U.G., Becking G.B., Birnbaum, L.S., Brouwer, A., Derks, H.J.G.M., Feely, M., Golor, G., Hanberg, A., Larsen, J.C., Liem, A.K.G., Safe, S.H., Schlatter, C., Wärn, F., Younes, M., Yrjänheikki, E., 1994. Toxic equivalency factors for dioxin-like PCBs. Report on a WHO-ECEH and IPSC consultation, December 1993. *Chemosphere* 28:1049-1067.
- ASMO, 1994. Draft assessment of temporal trends monitoring data for 1983-91: Trace metals and organic contaminants in biota. Environmental Assessment and Monitoring Committee (ASMO). Document ASMO(2) 94/6/1.
- ASMO, 1997. Summary Record. Environmental Assessment and Monitoring Committee (ASMO). Document ASMO 97/9/1.
- Atuma, S.S., Linder, C-E, Andersson, Ö., Bergh, A., Hansson, L., Wicklund-Glynn, A., 1996. CB153 as indicator for congener specific determination of PCBs in diverse fish species from Swedish waters. *Chemosphere* 33(8):1459-1464.
- Boer, J. de, Stronck, C.J.N., Traag, W.A., Meer, J van der, 1993. Non-ortho and mono-ortho substituted chlorobiphenyls and chlorinated dibenzo-p-dioxins and dibenzofurans in marine and freshwater fish and shellfish from the Netherlands. *Chemosphere* 26(10):1823-1842.
- Green, N.W., 1987. "Joint Monitoring Group" (JMG). Felles monitoring program i Norge: Oslofjord-området, Sørfjorden og Hardangerfjorden, og Orkdalsfjorden. Programforslag for 1988. 4.Dec.1987. NIVA-project 80106, 12 pp..
- Green, N.W., 1991. Joint Monitoring Programme. National Comments to the Norwegian Data for 1989. Norwegian Institute for Water Research (NIVA) memmo 27pp.. JMG 16/info 13-E
- Green, N.W., 1993. Joint Monitoring Programme - JMP. Overview of analytical methods employed by JMP in Norway 1981-1992. Norwegian Institute for Water Research. Project O-80106 report number 2971, 41 pp.. ISBN number 82-577-2390-8. (Also as document JMG 19/7 info.3-E.)
- Green, N.W., 1997. Joint Assessment and Monitoring Programme (JAMP) National comments to the Norwegian Data for 1995. Norwegian State Pollution Control Authority, Monitoring report no. 685/97 TA no. 3597/1997. NIVA project O-80106, report number 3597-97, 124 pp.. ISBN number 82-577-3152-8. NIVA. (Also as document SIME 97/5/5).
- Helland, A., 1996. Tilførsel av partikulært materiale til Glommaestuariet og områdene utenfor i forbindelse med flommen i Glomma 1995. [Discharge of particulate matter to the Glomma estuary and vicinity in connection with the flood of Glomma 1995]. Norwegian State Pollution Control Authority, Monitoring report no. 664/96 TA no. 1350/1996. NIVA project O-900342, (report number 3503-96) 50 pp.. ISBN number 82-577-3045-9.
- ICES, 1996. ICES Environmental Data Reporting Formats. Version 2.2, revision 2 - July 1996.
- Knutzen, J., Skei, J., 1990. Kvalitetskriterier for miljøgifter i vann, sedimenter og organismer samt foreløpige forslag til klassifikasjon av miljøkvalitet. [Quality criteria for water, sediments and organisms and preliminary proposals for classification of environmental quality]. NIVA-prosjekt O-8000309 (løpenummer 2540), 139 sider. ISBN.82-577-1855-6.
- Knutzen, J., Rygg, B., Thelín, I., 1993. Klassifisering av miljøkvalitet i fjorder og kystfarvann. Virkning av miljøgifter. SFT-rapport TA-923/1993, 20 s. ISBN 82-7655-103-3.
- Knutzen, J., Green, N.W., 1995. Bakgrunnsnivåer av en del miljøgifter i fisk, blåskjell og reker. Data fra utvalgte norske prøvesteder innen den felles overvåking under Oslo-/Paris-kommisjonene 1990-1993. [Background levels of some micropollutants in fish, the blue mussel and shrimps. Data from selected Norwegian sampling sites within the joint monitoring of the Oslo-/Paris Commissions 1990-1993]. Norwegian State Pollution Control Authority, Monitoring report no. 594/94 TA no. 1173/1994. NIVA project O-80106/E-91412, (report number 3302) 105 pp.. ISBN number 82-577-2678-8.

- Knutzen, J., Berglund, L., Brevik E., 1995. Sonderende undersøkelser i norske havner og utvalgte kystområder. Klororganiske stoffer og tributyltinn (TBT) i blåskjell 1993-1994 (Introductory studies in Norwegian harbours. Organochlorines and tributyltin in the common mussel 1993-1994. Norwegian State Pollution Control Authority, Monitoring report no. 610/95 TA no. 1210/1995. NIVA project O-93255 (report number 3296) 79 pp.. ISBN number 82-577-2786-5.
- Knutzen, J., Biseth, A., Brevik, E., Green, N., Schlaback M., Skåre, J.U., 1996. Overvåking av miljøgift i fisk og skalldyr, Grenlandsfjordene 1995. [Monitoring of contaminants in fish and shellfish, Grenlandsfjordene 1995.] Norwegian State Pollution Control Authority, Monitoring report no. 686/96 TA no. 1396/1995. NIVA project O-800312 (report number 3590) 69 pp.. ISBN number 82-577-3143-9.
- Knutzen, J., Green, N.W., Brevik, E.M., 1997 (in press). Tiltaksorienterte miljøundersøkelser i Sjøfjorden og Hardangerfjorden 1996. Delrapport 2: Miljøgifter i organismer. [Investigation of micropollutants in Sjøfjorden and Hardangerfjord 1996. Report 2]. Norwegian State Pollution Control Authority, Monitoring report no. ??/97 TA no. ??/1997. NIVA project O-800309, (report number ??-97) 34 pp.. ISBN number 82-577-??.
- Konieczny, R., Juliussen, A., 1995. Sonderende undersøkelser i norske havner og utvalgte kystområder. Fase 1: Miljøgifter i sedimenter på strekningen Norvik-Kragerø. [Introductory studies in Norwegian harbours and selected coastal areas. Phase 1: Contaminants in sediments along the coast Norvik-Kragerø]. Norwegian State Pollution Control Authority, Monitoring report no. 587/94 TA no. 1159/1994. NIVA project O-93177 (report number 3275) 185 pp.. ISBN number 82-577-2780-6.
- Konieczny, R., 1996. Sonderende undersøkelser i norske havner og utvalgte kystområder. Fase 3: Miljøgifter i sedimenter på strekningen Ramsund-Kirkenes. [Introductory studies in Norwegian harbours and selected coastal areas. Phase 3: Contaminants in sediments along the coast Ramsund-Kirkenes]. Norwegian State Pollution Control Authority, Monitoring report no. 608/95 TA no. 1215/1995. NIVA project O-93177 (report number 3423-96) 117 pp.. ISBN number 82-577-2957-4.
- JMG, 1992. Oslo and Paris Conventions for the Prevention of Marine Pollution,. Seventeenth Meeting of the Joint Monitoring Group. Uppsala: 20-24 January 1992. Summary Record. JMG 17/15/1-E. 29pp. plus appendices.
- Molvær, J., Knutzen, J., Magnusson, J., Rygg, B., Skei J., Sørensen, J., 1997. Klassifisering av miljøkvalitet i fjorder og kystfarvann. Veiledning. *Classification of environmental quality in fjords and coastal waters. A guide*. State Pollution Control Authority. TA no. TA-1467/1997. 36 pp.
- MON, 1993. Draft Summary record. Eleventh meeting of the Ad Hoc Working Group on Monitoring, Copenhagen: 8-12 November 1993. MON 11/1/7-E.
- Nicholson, M., Fryer, R.J., Green, N.W., 1994. Focusing on key aspects of contaminant trend assessments. Nineteenth meeting of the Joint Monitoring Group 24-29 . January, 1994. Document JMG 19/3/3.
- Nicholson, M., Fryer, R.J., Larsen, J.R., 1996a. Contaminants n marine organisms: A robust method for analysing temporal trends, ICES Techniques in Marine Environmental Sciences, No.20 (in press).
- Nicholson, M., Fryer, R.J., Claire, R., 1996b. Designing monitoring programmes for detecting trends in contaminants in fish and shellfish. *Marine Pollution Bulletin* (in press).
- OSPARCOM, 1990. Oslo and Paris Conventions. Principles and methodology of the Joint Monitoring Programme. [Monitoring manual for participants of the Joint Monitoring Programme (JMP) and North Sea Monitoring Master Plan (NSMMP)]. March 1990.
- SIME, 1996. Voluntary international contaminant-monitoring (VIC) for temporal trends in biota. OSPARCOM Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME) Oslo, 22-26 January 1996. SIME 96/25/1 Annex 33 6 pp.
- SIME, 1997. Voluntary international contaminant-monitoring (VIC) for temporal trends with the aim to test sampling strategies for a co-operative revision of guidelines by 1999- status repot January 1997. OSPARCOM Working Group on Concentrations, Trends and Effects of Substances in the Marine Environment (SIME) Ostend, 3-7 February 1997. SIME 96/6/5 3 pp.
- Skei, J., 1997. Tiltaksorienterte miljøundersøkelser i Sjøfjorden og Hardangerfjorden 1996. Delrapport 1. Vannkjemi. [Investigation of micropollutants in Sjøfjorden and Hardangerfjord 1996. Report 1]. Norwegian State Pollution Control Authority, Monitoring report no. 700/97 TA no. 1455/1997. NIVA project O-800309, (report number 3688-97) 27pp.. ISBN number 82-577-3253-2.

- Walday, M., Green, N., Hylland, K., 1995. Kostholds- og tilstandsindikatorer for miljøgifter i marine områder. Norwegian Institute for Water Research project 93254, report number 3280, 39 pp.. ISBN number 82-577-2691-5.
- WGSAEM, 1993. The length effect on contaminant concentrations in mussels. Section 13.2. in the Report of the Working Group on Statistical Aspects of Environmental Monitoring, Copenhagen 27-30, April 1993. International Council for the Exploration of the Sea. C-M- 1993/ENV:6 Ref.: D and E, 61 pp.

Appendix A.

Quality assurance programme

Accreditation

The laboratories at NIVA, both the chemical, microbiological and the ecotoxicological laboratories, were accredited in 1993 for quality assurance system by the National Measurement Service - Norwegian Accreditation and based on European Standard EN45000. NIVA has reference number P009.

Summary of quality control results

A summary of the results for the analyses of the SRM for biota is shown in Table A1.

As standard reference material (SRM) for the control of the determination of metals, dogfish muscle (DORM-2) or dogfish liver (DOLT-2) was used (see Table A1).

For control of PCB's and PAH analyses in biota SRM 350 (mackerel oil) and SRM 1974 was used, respectively. In addition to SRM 1974, an internal standard was used for quality control.

The results are satisfactory.

See also results from intercalibrations exercises listed in Appendix C.

NIVA has also participated in QUASIMEME exercises up to and including Round 10, the latter would apply to the 1996 samples analyzed in 1997. The results from Round 10 (November 1997) have been submitted to QUASIMEME but at the time of publication for these National Comments P and Z scores, assessment values for comparison with other laboratories, had not been received.

Table A1. Summary of the quality control results for the 1996 biota samples analysed 1996-97. The Standard Reference Materials (SRM) were DORM-2* (dogfish muscle) for mussels and fish fillet, DOLT-2* (dogfish liver) for fish liver, 350** (mackerel oil) for mussels and fish liver and NIST SRM 1974*** (mussel tissue) for mussels. SRM was analysed in series with the JAMP-samples for analyses of metals (mg/kg), organic chlorine's or PAH ($\mu\text{g}/\text{kg}$). Tissue types were: mussel softbody (SB), fish liver (LI) and fish fillet (MU).

Code	Contaminant	Tissue type	SRM type	SRM value \pm confidence interval	W	N	Mean value	Standard deviation
Cd	cadmium	SB	DORM	0.043 ± 0.008	15	5	0.042	0.004
		LI	DOLT	20.40 ± 1.80	9	14	20.43	1.00
Cu	copper	SB	DORM	2.34 ± 0.23	15	5	2.23	0.20
		LI	DOLT	25.10 ± 3.60	9	14	27.44	1.22
Pb	lead	SB	DORM	0.066 ± 0.38	15	5	0.07	0.01
		LI	DOLT	0.24 ± 0.11	9	14	0.20	0.02
Hg	mercury	SB MU	DORM	4.720 ± 0.520	26	19	4.564	0.272
Zn	zinc	SB	DORM	26.4 ± 3.7	15	5	25.86	1.52
		LI	DOLT	95.1 ± 11.3	9	14	94.6	2.87
CB-28	PCB congener CB-28	(all)	350	22.5 ± 4	26	20	18.99	0.91
CB-52	PCB congener CB-52	(all)	350	62 ± 9	26	21	57.43	3.60
CB-101	PCB congener CB-101	(all)	350	164 ± 9	26	21	169.38	8.07
CB-118	PCB congener CB-118	(all)	350	142 ± 20	26	21	151.10	10.90
CB-153	PCB congener CB-153	(all)	350	317 ± 20	26	21	346.14	17.69
CB-180	PCB congener CB-180	(all)	350	73 ± 13	26	21	82.24	4.30
PA	phenanthrene	MU	1974	5.6 ± 1.4	28	4	4.15	0.38
ANT	anthracene	MU	1974	0.75 ± 0.21	28	4	0.72	0.07
FLU	fluoranthene	MU	1974	33.6 ± 5.8	28	4	30.95	2.00
PYR	pyrene	MU	1974	34.1 ± 3.7	28	4	29.58	1.11
BBF	benzo[b]fluoranthene	MU	1974	6.5 ± 1.2	28	4	7.60	1.56
BAP	benzo[a]pyrene	MU	1974	2.29 ± 0.47	28	4	2.34	0.43
PER	perylene	MU	1974	1.05 ± 0.29	28	4	1.00	0.09
ICDP	indeno[1,2,3-cd]pyrene	MU	1974	1.80 ± 0.33	28	4	1.85	0.27
BGHIP	benzo[ghi]perylene	MU	1974	2.47 ± 0.28	28	4	2.74	0.30

- *) National Research Council Canada, Division of Chemistry, Marine Analytical Chemistry Standards
 **) BCR, Community Bureau of Reference, Commission of the European Communities
 ***) National Institute of Standards & Technology (NIST)

Appendix B. Abbreviations

Abbreviation¹	English	Norwegian
ELEMENTS		
Al	aluminium	<i>aluminium</i>
As	arsenic	<i>arsen</i>
Cd	cadmium	<i>kadmium</i>
Co	cobalt	<i>kobolt</i>
Cr	chromium	<i>krom</i>
Cu	copper	<i>kobber</i>
Fe	iron	<i>jern</i>
Hg	mercury	<i>kvikksølv</i>
Li	lithium	<i>litium</i>
Mn	manganese	<i>mangan</i>
Ni	nickel	<i>nikkel</i>
Pb	lead	<i>bly</i>
Pb210	lead-210	<i>bly-210</i>
Se	selenium	<i>selen</i>
Ti	titanium	<i>titan</i>
Zn	zinc	<i>sink</i>
PAHs		
PAH	polycyclic aromatic hydrocarbons	<i>polysykliske aromatiske hydrokarboner</i>
ACNE	acenaphthene	<i>acenaften</i>
ACNLE	acenaphthylene	<i>acenaftilen</i>
ANT	anthracene	<i>antracen</i>
BAA³	benz[<i>a</i>]anthracene	<i>benz[<i>a</i>]antracen</i>
BAP³	benzo[<i>a</i>]pyrene	<i>benzo[<i>a</i>]pyren</i>
BBF³	benzo[<i>b</i>]fluoranthene	<i>benzo[<i>b</i>]fluoranten</i>
BBJKF	benzo[<i>b,j,k</i>]fluoranthene	<i>benzo[<i>b,j,k</i>]fluoranten</i>
BBKF	benzo[<i>b+k</i>]fluoranthene	<i>benzo[<i>b+k</i>]fluoranten</i>
BEP	benzo[<i>e</i>]pyrene	<i>benzo[<i>e</i>]pyren</i>
BGHIP	benzo[<i>ghi</i>]perylene	<i>benzo[<i>ghi</i>]perylen</i>
BIPN²	biphenyl	<i>bifenyl</i>
BBJKF³	benzo[<i>b+j,k</i>]fluoranthene	<i>benzo[<i>b+j,k</i>]fluorantren</i>
BJKF³	benzo[<i>j,k</i>]fluoranthene	<i>benzo[<i>j,k</i>]fluorantren</i>
CHR	chrysene	<i>chrysen</i>
CHRTR	chrysene+triphenyl	<i>chrysen+trifenylen</i>
COR	coronene	<i>coronen</i>
DBAHA³	dibenz[<i>a,h</i>]anthracene	<i>dibenz[<i>a,h</i>]antracen</i>
DBA3A³	dibenz[<i>a,c/a,h</i>]anthracene	<i>dibenz[<i>a,c/a,h</i>]antracen</i>
DBP³	dibenzopyrenes	<i>dibenzopyren</i>
DBT	dibenzothiophene	<i>dibenzotiofen</i>
DBTC1	C ₁ -dibenzothiophenes	<i>C₁-dibenzotiofen</i>
DBTC2	C ₂ -dibenzothiophenes	<i>C₂-dibenzotiofen</i>
DBTC3	C ₃ -dibenzothiophenes	<i>C₃-dibenzotiofen</i>
FLE	fluorene	<i>fluoren</i>
FLU	fluoranthene	<i>fluoranten</i>

Abbreviation ¹	English	Norwegian
PAHs (cont.)		
ICDP ³	indeno[1,2,3-cd]pyrene	<i>indeno[1,2,3-cd]pyren</i>
NAPTM ²	2,3,5-trimethylnaphthalene	<i>2,3,5-trimetylnaftalen</i>
NAP ²	naphthalene	<i>naftalen</i>
NAPC1 ²	C ₁ -naphthalenes	<i>C₁-naftalen</i>
NAPC2 ²	C ₂ -naphthalenes	<i>C₂-naftalen</i>
NAPC3 ²	C ₃ -naphthalenes	<i>C₃-naftalen</i>
NAP1M ²	1-methylnaphthalene	<i>1-metylnaftalen</i>
NAP2M ²	2-methylnaphthalene	<i>2-metylnaftalen</i>
NAPDI ²	2,6-dimethylnaphthalene	<i>2,6-dimetylnaftalen</i>
PA	phenanthrene	<i>fenantren</i>
PAC1	C ₁ -phenanthrenes	<i>C₁-fenantren</i>
PAC2	C ₂ -phenanthrenes	<i>C₂-fenantren</i>
PAM1	1-methylphenanthrene	<i>1-metylfenantren</i>
PER	perylene	<i>perylen</i>
PYR	pyrene	<i>pyren</i>
DI-Σn	sum of "n" dicyclic "PAH"s (footnote 2)	<i>sum "n" disykliske "PAH" (fotnote 2)</i>
P-Σn	sum "n" PAH	<i>sum "n" PAH</i>
PK-Σn	sum carcinogen PAH's (footnote 3)	<i>sum kreftfremkallende PAH (fotnote 3)</i>
PAHΣΣ	DI-Σn + P-Σn etc.	<i>DI-Σn + P-Σn mm..</i>
SPAH	"total" PAH, specific compounds not quantified (outdated analytical method)	<i>"total" PAH, spesifikke forbindelser ikke kvantifisert (foreldret metode)</i>
PCBs		
PCB	polychlorinated biphenyls	<i>polyklorete bifenyler</i>
CB	individual chlorobiphenyls (CB)	<i>enkelte klorobifenyl</i>
CB28	CB28 (IUPAC)	<i>CB28 (IUPAC)</i>
CB31	CB31 (IUPAC)	<i>CB31 (IUPAC)</i>
CB44	CB44 (IUPAC)	<i>CB44 (IUPAC)</i>
CB52	CB52 (IUPAC)	<i>CB52 (IUPAC)</i>
CB77 ⁴	CB77 (IUPAC)	<i>CB77 (IUPAC)</i>
CB81 ⁴	CB81 (IUPAC)	<i>CB81 (IUPAC)</i>
CB95	CB95 (IUPAC)	<i>CB95 (IUPAC)</i>
CB101	CB101 (IUPAC)	<i>CB101 (IUPAC)</i>
CB105	CB105 (IUPAC)	<i>CB105 (IUPAC)</i>
CB110	CB110 (IUPAC)	<i>CB110 (IUPAC)</i>
CB118	CB118 (IUPAC)	<i>CB118 (IUPAC)</i>
CB126 ⁴	CB126 (IUPAC)	<i>CB126 (IUPAC)</i>
CB128	CB128 (IUPAC)	<i>CB128 (IUPAC)</i>
CB138	CB138 (IUPAC)	<i>CB138 (IUPAC)</i>
CB149	CB149 (IUPAC)	<i>CB149 (IUPAC)</i>
CB153	CB153 (IUPAC)	<i>CB153 (IUPAC)</i>
CB156	CB156 (IUPAC)	<i>CB156 (IUPAC)</i>
CB169 ⁴	CB169 (IUPAC)	<i>CB169 (IUPAC)</i>

Abbreviations (cont'd.)

Abbreviation ¹	English	Norwegian
PCBs (cont.)		
CB170	CB170 (IUPAC)	<i>CB170 (IUPAC)</i>
CB180	CB180 (IUPAC)	<i>CB180 (IUPAC)</i>
CB194	CB194 (IUPAC)	<i>CB194 (IUPAC)</i>
CB209	CB209 (IUPAC)	<i>CB209 (IUPAC)</i>
CB-Σ7	CB: 28+52+101+118+138+153+180	<i>CB: 28+52+101+118+138+153+180</i>
CB-ΣΣ	sum of CBs, includes CB-Σ7	<i>sum CBer, inkluderer CB-Σ7</i>
TECBW	Sum of CB-toxicity equivalents after WHO model, see TEQ	<i>Sum CB- toksitets ekvivalenter etter WHO modell, se TEQ</i>
TECBS	Sum of CB-toxicity equivalents after SAFE model, see TEQ	<i>Sum CB-toksitets ekvivalenter etter SAFE modell, se TEQ</i>
DIOXINS		
TCDD	2, 3, 7, 8-tetrachloro-dibenzo dioxin	<i>2, 3, 7, 8-tetrakloro-dibenzo dioksin</i>
CDDST	Sum of tetrachloro-dibenzo dioxins	<i>Sum tetrakloro-dibenzo dioksiner</i>
CDD1N	1, 2, 3, 7, 8-pentachloro-dibenzo dioxin	<i>1, 2, 3, 7, 8-pentakloro-dibenzo dioksin</i>
CDDSN	Sum of pentachloro-dibenzo dioxins	<i>Sum pentakloro-dibenzo dioksiner</i>
CDD4X	1, 2, 3, 4, 7, 8-hexachloro-dibenzo dioxin	<i>1, 2, 3, 4, 7, 8-heksakloro-dibenzo dioksin</i>
CDD6X	1, 2, 3, 6, 7, 8-hexachloro-dibenzo dioxin	<i>1, 2, 3, 6, 7, 8-heksakloro-dibenzo dioksin</i>
CDD9X	1, 2, 3, 7, 8, 9-hexachloro-dibenzo dioxin	<i>1, 2, 3, 7, 8, 9-heksakloro-dibenzo dioksin</i>
CDDSX	Sum of hexachloro-dibenzo dioxins	<i>Sum heksakloro-dibenzo dioksiner</i>
CDD6P	1, 2, 3, 4, 6, 7, 8-heptachloro-dibenzo dioxin	<i>1, 2, 3, 4, 6, 7, 8-heptakloro-dibenzo dioksin</i>
CDDSH	Sum of heptachloro-dibenzo dioxins	<i>Sum heptakloro-dibenzo dioksiner</i>
CDDO	Octachloro-dibenzo dioxin	<i>Oktakloro-dibenzo dioksin</i>
PCDD	Sum of polychlorinated dibenzo-p-dioxins	<i>Sum polyklorinaterte-dibenzo-p-dioksiner</i>
CDF2T	2, 3, 7, 8-tetrachloro-dibenzofuran	<i>2, 3, 7, 8-tetrakloro-dibenzofuran</i>
CDFST	Sum of tetrachloro-dibenzofurans	<i>Sum tetrakloro-dibenzofuraner</i>
CDFDN	1, 2, 3, 7, 8/1, 2, 3, 4, 8-pentachloro-dibenzofuran	<i>1, 2, 3, 7, 8/1, 2, 3, 4, 8-pentakloro-dibenzofuran</i>
CDF2N	2, 3, 4, 7, 8-pentachloro-dibenzofurans	<i>2, 3, 4, 7, 8-pentakloro-dibenzofuran</i>
CDFSN	Sum of pentachloro-dibenzofurans	<i>Sum pentakloro-dibenzofuraner</i>
CDFDX	1, 2, 3, 4, 7, 8/1, 2, 3, 4, 7, 9-hexachloro-dibenzofuran	<i>1, 2, 3, 4, 7, 8/1, 2, 3, 4, 7, 9-heksakloro-dibenzofuran</i>
CDF6X	1, 2, 3, 6, 7, 8-hexachloro-dibenzofuran	<i>1, 2, 3, 6, 7, 8-heksakloro-dibenzofuran</i>
CDF9X	1, 2, 3, 7, 8, 9-hexachloro-dibenzofuran	<i>1, 2, 3, 7, 8, 9-heksakloro-dibenzofuran</i>
CDF4X	2, 3, 4, 6, 7, 8-hexachloro-dibenzofuran	<i>2, 3, 4, 6, 7, 8-heksakloro-dibenzofuran</i>
CDFSX	Sum of hexachloro-dibenzofurans	<i>Sum heksakloro-dibenzofuraner</i>
CDF6P	1, 2, 3, 4, 6, 7, 8-heptachloro-dibenzofuran	<i>1, 2, 3, 4, 6, 7, 8-heptakloro-dibenzofuran</i>
CDF9P	1, 2, 3, 4, 7, 8, 9-heptachloro-dibenzofuran	<i>1, 2, 3, 4, 7, 8, 9-heptakloro-dibenzofuran</i>
CDFSP	Sum of heptachloro-dibenzofurans	<i>Sum heptakloro-dibenzofuraner</i>
CDFO	Octachloro-dibenzofurans	<i>Oktakloro-dibenzofuran</i>
PCDF	Sum of polychlorinated dibenzo-furans	<i>Sum polyklorinerted dibenzo-furaner</i>
CDDFS	Sum of PCDD and PCDF	<i>Sum PCDD og PCDF</i>

Abbreviations (cont'd.)

Abbreviation ¹	English	Norwegian
DIOXINS (cont.)		
TCDDN	Sum of TCDD-toxicity equivalents after Nordic model, see TEQ	<i>Sum TCDD- toksitets ekvivalenter etter Nordisk modell, se TEQ</i>
TCDDI	Sum of TCDD-toxicity equivalents after international model, see TEQ	<i>Sum TCDD-toksitets ekvivalenter etter internasjonale modell, se TEQ</i>
PESTICIDES		
ALD	aldrin	<i>aldrin</i>
DIELD	dieldrin	<i>dieldrin</i>
ENDA	endrin	<i>endrin</i>
CCDAN	cis-chlordane (=α-chlordane)	<i>cis-chlordan (=α-chlordan)</i>
TCDAN	trans-chlordane (=γ-chlordane)	<i>trans-chlordan (=γ-chlordan)</i>
OCDAN	oxy-chlordane	<i>oxy-chlordan</i>
TNONC	trans-nonachlor	<i>trans-nonaklor</i>
TCDAN	trans-chlordane	<i>trans-chlordan</i>
OCS	octachlorostyrene	<i>octaklorstyren</i>
QCB	pentachlorobenzene	<i>pentaklorbenzen</i>
DDD	dichlorodiphenyldichloroethane 1,1-dichloro-2,2-bis-(4-chlorophenyl)ethane	<i>diklordifenyldikloreten</i> <i>1,1-dikloro-2,2-bis-(4-klorofenyl)etan</i>
DDE	dichlorodiphenyldichloroethylene (principle metabolite of DDT) 1,1-dichloro-2,2-bis-(4-chlorophenyl)ethylene*	<i>diklordifenyldikloretylen</i> <i>(hovedmetabolitt av DDT)</i> <i>1,1-dikloro-2,2-bis-(4-klorofenyl)etylen</i>
DDT	dichlorodiphenyltrichloroethane 1,1,1-trichloro-2,2-bis-(4-chlorophenyl)ethane	<i>diklordifenyltrikloreten</i> <i>1,1,1-trikloro-2,2-bis-(4-klorofenyl)etan</i>
DDEOP	o,p'-DDE	<i>o,p'-DDE</i>
DDEPP	p,p'-DDE	<i>p,p'-DDE</i>
DDTOP	o,p'-DDT	<i>o,p'-DDT</i>
DDTPP	p,p'-DDT	<i>p,p'-DDT</i>
TDEPP	p,p'-DDD	<i>p,p'-DDD</i>
DDTEP	p,p'-DDE + p,p'-DDT	<i>p,p'-DDE + p,p'-DDT</i>
DD-nΣ	sum of DDT and metabolites, n = number of compounds	<i>sum DDT og metabolitter,</i> <i>n = antall forbindelser</i>

Abbreviations (cont'd.)

Abbreviation ¹	English	Norwegian
HCB	hexachlorobenzene	<i>heksaklorbenzen</i>
HCHG	lindane γ HCH = gamma hexachlorocyclohexane (γ BHC = gamma benzenehexachloride, outdated synonym)	<i>lindan</i> γ HCH = <i>gamma heksaklorsykloheksan</i> (γ BHC = <i>gamma benzenheksaklorid</i> , <i>foreldret betegnelse</i>)
HCHA	α HCH = alpha HCH	<i>α HCH = alpha HCH</i>
HCHB	β HCH = beta HCH	<i>β HCH = beta HCH</i>
HC-nΣ	sum of HCHs, n = count	<i>sum av HCHs, n = antall</i>
EOCI	extractable organically bound chlorine	<i>ekstraherbart organisk bundet klor</i>
EPOCI	extractable persistent organically bound chlorine	<i>ekstraherbart persistent organisk bundet klor</i>
NTOT	total organic nitrogen	<i>total organisk nitrogen</i>
CTOT	total organic carbon	<i>total organisk karbon</i>
CORG	organic carbon	<i>organisk karbon</i>
GSAMT	grain size	<i>kornfordeling</i>
MOCON	moisture content	<i>vanninnhold</i>

Abbreviations (cont'd.)

Abbreviation ¹	English	Norwegian
INSTITUTES		
FIER	Institute for Nutrition, Fisheries Directorate	<i>Fiskeridirektoratets Ernæringsinstitutt</i>
FORC	FORCE Institutes, Div. for Isotope Technique and Analysis [DK]	<i>FORCE Institutterne, Div. for Isotopteknik og Analyse [DK]</i>
IMRN	Institute of Marine Research (IMR)	<i>Havforskningsinstituttet</i>
NACE	Nordic Analytical Center	<i>Nordisk Analyse Center</i>
NILU	Norwegian Institute for Air Research	<i>Norsk institutt for luftforskning</i>
NIVA	Norwegian Institute for Water Research	<i>Norsk institutt for vannforskning</i>
SERI	Swedish Environmental Research Institute	<i>Institutionen för vatten- och luftvårdsforskning</i>
VETN	Norwegian Veterinary Institute	<i>Veterinærinstituttet</i>
SIIF	Fondation for Scientific and Industrial Research at the Norwegian Institute of Technology - SINTEF (a division, previously: Center for Industrial Research SI)	<i>Stiftelsen for industriell og teknisk forskning ved Norges tekniske høgskole- SINTEF (en avdeling, tidligere: Senter for industriforskning SI)</i>

- 1) After: ICES Environmental Data Reporting Formats. International Council for the Exploration of the Sea. July 1996 and supplementary codes related to non-ortho and mono-ortho PCB's and "dioxins" (ICES pers. comm.)
- 2) Indicates "PAH" compounds that are dicyclic and not truly PAH's typically identified during the analyses of PAH, include naphthalenes and "biphenyls".
- 3) Indicates PAH compounds potentially cancerogenic for humans according to IARC (1987), i.e., categories 2A+2B (possibly and probably carcinogenic).
- *) The Pesticide Index, second edition. The Royal Society of Chemistry, 1991.

Other abbreviations andre forkortelser

	English	Norwegian
TEQ	"Toxicity equivalency factors" for the most toxic compounds within the following groups:	" <i>Toxisitetskvivalentfaktorer</i> " for de giftigste forbindelsene innen følgende grupper.
	<ul style="list-style-type: none"> polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDD/PCDFs). Equivalents calculated after Nordic model (Ahlborg <i>et al.</i>, 1989)¹ or international model (Int./EPA, cf. Ahlborg <i>et al.</i>, 1992)² non-ortho and mono-ortho substituted chlorobiphenyls after WHO model (Ahlborg <i>et al.</i>, 1994)³ or Safe (1994, cf., NILU pers. comm.) 	<ul style="list-style-type: none"> <i>polyklorete dibenzo-p-dioksiner og dibenzofuraner (PCDD/PCDF)</i>. <i>Ekvivalentberegning etter nordisk modell (Ahlborg et al., 1989)¹ eller etter internasjonal modell (Int./EPA, cf. Ahlborg et al., 1992)²</i> <i>non-orto og mono-orto substituerte klorobifenylar etter WHO modell (Ahlborg et al., 1994)³ eller Safe (1994, cf., NILU pers. medd.)</i>
ppm	parts per million, mg/kg	<i>deler pr. milliondeler, mg/kg</i>
ppb	parts per billion, µg/kg	<i>deler pr. milliarddeler, µg/kg</i>
ppp	parts per trillion, ng/kg	<i>deler pr. tusen-milliarddeler, ng/kg</i>
d.w.	dry weight basis	<i>tørrvekt basis</i>
w.w.	wet weight or fresh weight basis	<i>våttvekt eller friskvekt basis</i>

¹) Ahlborg, U.G., 1989. Nordic risk assessment of PCDDs and PCDFs. *Chemosphere* 19:603-608.

²) Ahlborg, U.G., Brouwer, A., Fingerhut, M.A., Jacobson, J.L., Jacobson, S.W.; Kennedy, S.W., Kettrup, A.F., Koeman, J.H., Poiger, H., Rappe, C., Safe, S.H., Schlatter, C., Seegal, R.F., Tuomisto, J., van den Berg, M., 1992. Impact of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls on human and environmental health, with special emphasis on application of the toxic equivalency factor concept *European Journal of Pharmacology . Environmental Toxicology and Pharmacology Section* 228 (1992) 179-199

³) Ahlborg, U.G., Becking G.B., Birnbaum, L.S., Brouwer, A., Derks, H.J.G.M., Feely, M., Golor, G., Hanberg, A., Larsen, J.C., J.C., Liem, A.K.G., Safe, S.H., Schlatter, C., Wärn, F., Younes, M., Yrjänheikki, E., 1994. Toxic equivalency factors for dioxin-like PCBs. Report on a WHO-ECEH and IPSC consultation , December 1993. *Chemosphere* 28:1049-1067.

Appendix C.

Participation in intercalibration exercises

Participation in intercalibration exercises

General: The main contributor to JAMP in 1996 has been NIVA which has participated in all QUASIMEME exercises relevant to the parameter and tissues monitored

Sea water:

- 4H ICES/JMG Fifth Round Intercalibration on Trace Metals in Sea Water - Section 4, analysis for Hg - 1983 - (5/TM/SW:4).
- 4I JMG Sixth Intercalibration on Trace Metals in Estuarine Waters - 1986 - (6/TM/SW).
- 4Z Intercalibration exercise for SIIF/SERI (Cd) and NIVA/IAMK (IAMK=Chalmers Inst., Göteborg) - 1985.

Seabed sediment:

- 7E ICES, First Intercalibration Exercise on Trace metals in Marine Sediments - 1984 - (1/TM/MS).
- 8B ICES/OSPARCOM, First Intercomparison Exercise on Organochlorines (individual chlorobiphenyl congeners) in Marine Sediments - Phase 1, analysis of standard solutions - 1989 - (1/OC/MS:1).
- 8C ICES/OSPARCOM, First Intercomparison Exercise on Organochlorines (individual chlorobiphenyl congeners) in Marine Sediments - Phase 2, analysis of standard solutions - 1991 - (1/OC/MS:2).
- 8B ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 1 - (analysis of standard solutions) - 1989 - (1/OC/MS-1).
- 8C ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 2 - 1990 - (1/OC/MS-2).
- 8D ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3a (1/OC/MS-3a) 1991.
- 8E ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3b - (1/OC/MS-3b) 1992.
- 8F ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 4 - (1/OC/MS-4) 1993.

Marine biota:

- 1E ICES, Fifth Intercalibration Exercise on Trace Metals in Biological Tissues - 1978 - (5/TM/BT).
- 1F ICES, Sixth Intercalibration Exercise on Trace Metals (Cadmium and Lead only) in Biological Tissues - 1979 - (6/TM/BT).
- 1G ICES, Seventh Intercalibration Exercise on Trace Metals in Biological Tissues - Part A - 1983 - (7/TM/BT).

- 1H ICES, Seventh Intercalibration Exercise on Trace Metals in Biological Tissues - Part B - 1985 - (7/TM/BT) (preliminary report 1987).
- 1Z VETN Interlabcalibration exercise with VETN and SIIF 1983, mercury and cadmium in cod filet and liver.
- 1Z NIVA Interlabcalibration exercise with VETN, NACE and NIVA 1986 (Hg, Cd, Cu, Pb and Zn in 6 samples).
- 2D ICES Fourth Intercalibration Exercise on Organochlorines (mainly PCBs) in Biological Tissues (Sample No.5) - 1979 - (4/OC/BT).
- 2E ICES Fifth Intercalibration Exercise on Organochlorines (PCBs only) in Biological Tissues - 1982 - (5/OC/BT).
- 2G ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 1 - (analysis of standard solutions) - 1989 - (7/OC/BT-1).
- 2H ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 2 - 1990 - (7/OC/BT-2).
- 2I ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3a - (7/OC/BT-3a) 1991.
- 2J ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 3b - (7/OC/BT-3b) 1992.
- 2K ICES/IOC/OSPARCOM Intercomparison Programme on the Analysis of Chlorobiphenyls in Marine Media - Step 4 - (7/OC/BT-4) 1993.
- 2Z VETN Interlabcalibration exercise with VETN among others, 1983, PCB and HCB in cod liver.
- 2Z NACE Interlabcalibration exercise with NACE, VETN and SIIF 1986 (PCB (all labs), DDE, OCS, HCB and DCB (NACE and VETN)).

Appendix D. Analytical overview

sorted in three ways:

- Method, laboratory**
- Laboratory, method**
- Contaminant, year, laboratory, intercalibration**

abbreviations are defined in Appendices B and C

Filename : I:\TBX\JMG\BIO\TAB-4TIS.TB1

Niva 24/10-1997

Analytical overview BIOTA; Sorted by METHOD, LAB. and TISSUE.

Method	Lab.	Tissue	Monitoring Year	Contaminants
110	SIIF	Fish fillet	1981	PCB
		Fish liver	1981	PCB
		Mussel	1981	PCB
111	SIIF	Mussel	1982-1991	PCB
		Mussel	1983-1991	DDTEP, HCB
		Mussel	1986-1987, 1989-1991	HCHG
		Mussel	1987-1991	CB101, CB180, CB52
		Mussel	1988-1991	CB138, CB153, CB28
		Mussel	1989-1991	CB118
		Shrimp tail	1982, 1984, 1986, 1988, 1990	PCB
		Shrimp tail	1984, 1986, 1988, 1990	DDTEP, HCB
		Shrimp tail	1986, 1990	HCHG
		Shrimp tail	1988, 1990	CB101, CB138, CB153, CB180, CB28, CB52
		Shrimp tail	1990	CB118
		Other	1988	CB101, CB138, CB153, CB180, CB28, CB52, DDTEP, HCB, PCB
120	SIIF	Fish fillet	1981	HG
		Fish liver	1981	HG
		Mussel	1981-1985	HG
		Shrimp tail	1982, 1984	HG
130	SIIF	Fish fillet	1981	CD
		Fish liver	1981	CD
		Mussel	1981-1985	CD
		Mussel	1983	NI
		Mussel	1983-1984	CU
		Mussel	1983-1985	PB
		Shrimp tail	1982, 1984	CD
		Shrimp tail	1984	CU, PB
131	SIIF	Mussel	1983	ZN
132	SIIF	Mussel	1984-1985	MN, ZN
		Shrimp tail	1984	MN, ZN
210	VETN	Fish fillet	1983	DDEPP, HCB
		Fish liver	1982-1985	DDEPP, PCB
		Fish liver	1983-1985	HCB
211	VETN	Fish fillet	1982-1985	PCB
		Fish fillet	1983	DDEPP, HCB
220	VETN	Fish fillet	1982-1985	HG
		Fish liver	1982	HG
230	VETN	Fish liver	1982-1985	CD
240	VETN	Fish fillet	1982	SE
		Fish liver	1982	SE
309	NIVA	Fish fillet	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, COR, DBA3A, DBP, FLE, FLU, ICDP, NAP, NAP1M, NAP2M, NAPDI, NAPTM, PA, PAM1, PER, PYR
		Fish liver	1987	PAH
		Fish liver	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, COR, DBA3A, DBP, FLE, FLU, ICDP, NAP, NAP1M, NAP2M, NAPDI, NAPTM, PA, PAM1, PER, PYR
		Mussel	1992	COR, DBP
		Mussel	1992, 1995	NAP1M, NAP2M, NAPDI, NAPTM, PAM1
		Mussel	1992, 1995-1996	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, DBA3A, FLE, FLU, ICDP, NAP, PA, PER, PYR
		Mussel	1995	CHRTR
		Mussel	1995-1996	BBJKF, DBTC1, DBTC2, DBTC3, NAPC1, NAPC2, NAPC3, PAC1, PAC2
		Shrimp tail	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, COR, DBA3A, DBP, FLE, FLU, ICDP, NAP, NAP1M, NAP2M, NAPDI, NAPTM, PA, PAM1, PER, PYR
310	NIVA	Fish fillet	1986-1996	HG
		Mussel	1986-1996	HG
		Shrimp tail	1986, 1988, 1990, 1992, 1995	HG
		Other	1988	HG
311	NIVA	Fish liver	1986-1996	CU, ZN
		Mussel	1986-1996	CU, ZN
		Shrimp tail	1986, 1988, 1990, 1992, 1995	CU, ZN
		Other	1988	CU, ZN
312	NIVA	Fish liver	1986-1996	CD, PB
		Mussel	1986-1996	CD, PB
		Mussel	1992	CR, NI
		Shrimp tail	1986, 1988, 1990, 1992, 1995	CD, PB
		Other	1988	CD, PB
340	NIVA	Fish liver	1987	PCB
		Fish liver	1990-1996	CB101, CB118, CB138, CB153, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP
		Fish liver	1991-1996	CB105, CB156
		Fish liver	1996	DDTTPP
		Mussel	1995-1996	DDTTPP
341	NIVA	Fish fillet	1990-1996	CB101, CB118, CB138, CB153, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP
		Fish fillet	1991-1996	CB105, CB156
		Mussel	1992-1996	CB101, CB105, CB118, CB138, CB153, CB156, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP
		Shrimp tail	1992, 1995	CB101, CB105, CB118, CB138, CB153, CB156, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP
401	FIER	Fish fillet	1984, 1987	HG
402	FIER	Fish liver	1984, 1987	CD
403	FIER	Fish liver	1987	PB
404	FIER	Fish liver	1987	CU
405	FIER	Fish liver	1987	ZN
510	NACE	Fish liver	1986-1989	DDEPP, DDTTPP, HCB, HCHG, PCB
		Fish liver	1989	CB101, CB118, CB138, CB153, CB180, CB28, CB52

Tab.Length cont'd.

Method	Lab.	Tissue	Monitoring Year	Contaminants
511	NACE	Fish fillet	1986-1989	PCB
		Fish liver	1986	PCB
605	SIIF	Mussel	1986-1991	EPOCL
		Mussel	1989	EOCL
		Shrimp tail	1986, 1988, 1990	EPOCL
		Other	1988	EPOCL
610	NACE	Fish liver	1986-1989	EPOCL
615	NIVA	Fish liver	1990-1991	EPOCL
841	NILU	Mussel	1995-1996	CB126, CB169, CB77, CB81, CDD1N, CDD4X, CDD6P, CDD6X, CDD9X, CDDO, CDDSN, CDDSP, CDDST, CDDSX, CDF2N, CDF2T, CDF4X, CDF6P, CDF6X, CDF9P, CDF9X, CDFDN, CDFDX, CDFO, CDFSN, CDFSP, CDFST, CDFSX, PCDD, PCDF, TCDD, PCC26, PCC32, PCC50, PCC62
842	NILU	Mussel	1996	AG, AS, CO, CR, NI, V
999	NIVA	Mussel	1996	AG, AS, CO, CR, NI, V

Analytical overview BIOTA; sorted by TISSUE, METHOD and LAB.

Tissue	Method	Lab.	Monitoring Year	Contaminants
Fish fillet	110	SIIF	1981	PCB
	120	SIIF	1981	HG
	130	SIIF	1981	CD
	210	VETN	1983	DDEPP, HCB
	211	VETN	1982-1985	PCB
	211	VETN	1983	DDEPP, HCB
	220	VETN	1982-1985	HG
	240	VETN	1982	SE
	309	NIVA	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, COR, DBA3A, DBP, FLE, FLU, ICDP, NAP, NAP1M, NAP2M, NAPDI, NAPTM, PA, PAM1, PER, PYR
	310	NIVA	1986-1996	HG
	341	NIVA	1990-1996	CB101, CB118, CB138, CB153, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP
	341	NIVA	1991-1996	CB105, CB156
	401	FIER	1984, 1987	HG
511	NACE	1986-1989	PCB	
Fish liver	110	SIIF	1981	PCB
	120	SIIF	1981	HG
	130	SIIF	1981	CD
	210	VETN	1982-1985	DDEPP, PCB
	210	VETN	1983-1985	HCB
	220	VETN	1982	HG
	230	VETN	1982-1985	CD
	240	VETN	1982	SE
	309	NIVA	1987	PAH
	309	NIVA	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, COR, DBA3A, DBP, FLE, FLU, ICDP, NAP, NAP1M, NAP2M, NAPDI, NAPTM, PA, PAM1, PER, PYR
	311	NIVA	1986-1996	CU, ZN
	312	NIVA	1986-1996	CD, PB
	340	NIVA	1987	PCB
340	NIVA	1990-1996	CB101, CB118, CB138, CB153, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP	
340	NIVA	1991-1996	CB105, CB156	
340	NIVA	1996	DDTPP	
402	FIER	1984, 1987	CD	
403	FIER	1987	PB	
404	FIER	1987	CU	
405	FIER	1987	ZN	
510	NACE	1986-1989	DDEPP, DDTTP, HCB, HCHG, PCB	
510	NACE	1989	CB101, CB118, CB138, CB153, CB180, CB28, CB52	
511	NACE	1986	PCB	
610	NACE	1986-1989	EPOCL	
615	NIVA	1990-1991	EPOCL	
Mussel	110	SIIF	1981	PCB
	111	SIIF	1982-1991	PCB
	111	SIIF	1983-1991	DDTEP, HCB
	111	SIIF	1986-1987, 1989-1991	HCHG
	111	SIIF	1987-1991	CB101, CB180, CB52
	111	SIIF	1988-1991	CB138, CB153, CB28
	111	SIIF	1989-1991	CB118
	120	SIIF	1981-1985	HG
	130	SIIF	1981-1985	CD
	130	SIIF	1983	NI
	130	SIIF	1983-1984	CU
	130	SIIF	1983-1985	PB
	131	SIIF	1983	ZN
132	SIIF	1984-1985	MN, ZN	
309	NIVA	1992	COR, DBP	
309	NIVA	1992, 1995	NAP1M, NAP2M, NAPDI, NAPTM, PAM1	
309	NIVA	1992, 1995-1996	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN, BJKF, CHR, DBA3A, FLE, FLU, ICDP, NAP, PA, PER, PYR	
309	NIVA	1995	CHRTR	
309	NIVA	1995-1996	BBJKF, DBTC1, DBTC2, DBTC3, NAPC1, NAPC2, NAPC3, PAC1, PAC2	
310	NIVA	1986-1996	HG	
311	NIVA	1986-1996	CU, ZN	
312	NIVA	1986-1996	CD, PB	
312	NIVA	1992	CR, NI	
340	NIVA	1995-1996	DDTPP	
341	NIVA	1992-1996	CB101, CB105, CB118, CB138, CB153, CB156, CB180, CB209, CB28, CB52, DDEPP, HCB, HCHA, HCHG, OCS, QCB, TDEPP	
605	SIIF	1986-1991	EPOCL	
605	SIIF	1989	EOCL	
841	NILU	1995-1996	CB126, CB169, CB77, CB81, CDD1N, CDD4X, CDD6P, CDD6X, CDD9X, CDDO, CDDSN, CDDSP, CDDST, CDDSX, CDF2N, CDF2T, CDF4X, CDF6P, CDF6X, CDF9P, CDF9X, CDFDN, CDFDX, CDFO, CDFSN, CDFSP, CDFST, CDFSX, PCDD, PCDF, TCDD	
842	NILU	1996	PCC26, PCC32, PCC50, PCC62	
999	NIVA	1996	AG, AS, CO, CR, NI, V	
Shrimp tail	111	SIIF	1982, 1984, 1986, 1988, 1990	PCB
	111	SIIF	1984, 1986, 1988, 1990	DDTEP, HCB
	111	SIIF	1986, 1990	HCHG
	111	SIIF	1988, 1990	CB101, CB138, CB153, CB180, CB28, CB52
	111	SIIF	1990	CB118
	120	SIIF	1982, 1984	HG
	130	SIIF	1982, 1984	CD
	130	SIIF	1984	CU, PB
	132	SIIF	1984	MN, ZN
	309	NIVA	1992	ACNE, ACNLE, ANT, BAA, BAP, BBF, BEP, BGHIP, BIPN,

Tab.length cont'd.

Tissue	Method	Lab.	Monitoring Year	Contaminants
	309	NIVA		BJKF , CHR , COR , DBA3A, DBP , FLE , FLU , ICDP , NAP , NAP1M, NAP2M, NAPDI, NAPTM, PA , PAM1 , PER , PYR
	310	NIVA	1986,1988,1990,1992,1995	HG
	311	NIVA	1986,1988,1990,1992,1995	CU , ZN
	312	NIVA	1986,1988,1990,1992,1995	CD , PB
	341	NIVA	1992,1995	CB101, CB105, CB118, CB138, CB153, CB156, CB180, CB209, CB28 , CB52 , DDEPP, HCB , HCHA , HCHG , OCS , QCB , TDEPP
Other	605	SIIF	1986,1988,1990	EPOCL
	111	SIIF	1988	CB101, CB138, CB153, CB180, CB28 , CB52 , DDTEP, HCB , PCB
	310	NIVA	1988	HG
	311	NIVA	1988	CU , ZN
	312	NIVA	1988	CD , PB
	605	SIIF	1988	EPOCL

Tab. length cont'd.

Tissue		Fish liver					Fish fillet, Shrimptail, Mussel, Other					
Contam.	Mon. Year	Lab. Inter-calibr.+Basis	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim
CB156	1990-SIIF	2G W						111	0.30	41		
	1991-NIVA	2H W	340	1.00	179			341	0.05	62		
	1991-SIIF	2H W						111	0.50	35		1
	1992-NIVA	2J W	340	5.00	192			341	0.10	140		
	1993-NIVA	2K W	340	4.00	212	3		341	0.10	133		
	1994-NIVA	2Z W	340	3.00	300			341	0.05	165	9	
	1995-NIVA	W	340	3.00	318	1		341	0.05	216		
	1996-NIVA	W	340	3.00	332	1		341	0.05	228		
	1991-NIVA	2H W	340	1.00	87		15	341	0.05	47		5
	1992-NIVA	W	340	5.00	192	3		341	0.10	140		
	1993-NIVA	QM W	340	4.00	212	31		341	0.10	133		
	1994-NIVA	2Z W	340	3.00	300	24	1	341	0.05	162	70	
1995-NIVA	W	340	3.00	317	27		341	0.05	216	67		
1996-NIVA	W	340	3.00	332	48		341	0.05	228	62		
CB169	1995-NILU	W					841	.20E-04	6			
	1996-NILU	W					841	.10E-03	18	2		
CB180	1987-SIIF	W					111	0.20	21	6		
	1988-SIIF	D					111	0.10	6			
	1988-SIIF	W					111	0.10	22			
	1989-NACE	W	510	20.00	93	1						
	1989-SIIF	W					111	0.10	36			
	1990-NIVA	2G W	340	1.00	169		341	0.05	58			
	1990-SIIF	2G W					111	0.20	41	8		
	1991-NIVA	2H W	340	1.00	179		341	0.05	62			
	1991-SIIF	2H W					111	0.20	35			
	1992-NIVA	2J W	340	5.00	192	3	341	0.10	140			
	1993-NIVA	2K W	340	4.00	212	15	341	0.10	133			
	1994-NIVA	2Z W	340	3.00	300	3	341	0.05	162	49		
	1995-NIVA	W	340	3.00	318	5	341	0.05	216	22		
	1996-NIVA	W	340	3.00	332	14	341	0.05	228	25		
CB209	1990-NIVA	W	340	2.00	169	24	341	0.05	58			
	1991-NIVA	W	340	2.00	179	11	341	0.05	62	5		
	1992-NIVA	W	340	5.00	192	3	341	0.10	140		7	
	1993-NIVA	W	340	4.00	212	46	341	0.10	133		1	
	1994-NIVA	W	340	3.00	300	29	341	0.05	165	91		
	1995-NIVA	W	340	3.00	318	36	341	0.05	216	92		
	1996-NIVA	W	340	3.00	332	255	341	0.05	228	107		
CB28	1988-SIIF	D					111	0.10	6			
	1988-SIIF	W					111	0.10	22			
	1989-NACE	W	510	20.00	93							
	1989-SIIF	W					111	0.10	36		1	
	1990-NIVA	2G W	340	1.00	169	2	341	0.05	58			
	1990-SIIF	2G W					111	0.20	41	7		
	1991-NIVA	2H W	340	1.00	179	2	341	0.05	62	5	1	
	1991-SIIF	2H W					111	0.30	35			
	1992-NIVA	2J W	340	5.00	192	3	341	0.10	137			
	1993-NIVA	2K W	340	4.00	212	44	341	0.10	133			
	1994-NIVA	2Z W	340	3.00	282	18	341	0.05	163	73		
	1995-NIVA	W	340	3.00	313	27	341	0.05	216	75		
	1996-NIVA	W	340	3.00	332	107	341	0.05	227	70		
CB52	1987-SIIF	W					111	0.20	20	1		
	1988-SIIF	D					111	0.10	6			
	1988-SIIF	W					111	0.10	22			
	1989-NACE	W	510	20.00	93							
	1989-SIIF	W					111	0.10	36			
	1990-NIVA	2G W	340	1.00	169	2	341	0.05	58			
	1990-SIIF	2G W					111	0.40	41	7		
	1991-NIVA	2H W	340	1.00	179	1	341	0.05	62	5	1	
	1991-SIIF	2H W					111	0.30	35			
	1992-NIVA	2J W	340	5.00	192	3	341	0.10	137			
	1993-NIVA	2K W	340	4.00	212	40	341	0.10	133			
	1994-NIVA	2Z W	340	3.00	300	9	341	0.05	165	64		
	1995-NIVA	W	340	3.00	312	19	341	0.05	205	28		
	1996-NIVA	W	340	3.00	332	49	341	0.05	226	31		
CB77	1995-NILU	W					841	.20E-04	6			
	1996-NILU	W					841	.10E-03	18			
CB81	1995-NILU	W					841	.20E-04	6			
	1996-NILU	W					841	.10E-03	18			
CD	1981-SIIF	1E W	130	10.00	28		130	5.00	27			
	1981-SIIF	1F W					130	10.00	7			
	1982-SIIF	1F W					130	10.00	18			
	1982-VETN	W	230	10.00	54							
	1983-SIIF	1F W					130	10.00	17			
	1983-VETN	1Z W	230	10.00	46							
	1984-FIER	1H W	402	1.00	23							
	1984-SIIF	1G W					130	10.00	27			
	1984-VETN	1Z W	230	10.00	66							
	1985-SIIF	1G D					130	10.00	35			
	1985-VETN	1Z W	230	10.00	45							
	1986-NIVA	1H D	312	30.00	56	1	312	30.00	20			
	1987-FIER	1G W	402	1.00	37							
	1987-NIVA	1H D	312	30.00	57		312	30.00	37			
	1988-NIVA	1H D	312	30.00	61	11	312	30.00	55			
	1989-NIVA	1H D	312	30.00	135	11						
	1989-NIVA	1H W					312	30.00	36			
	1990-NIVA	1H W	312	10.00	189	9	312	30.00	77	5		
	1991-NIVA	1H W	312	10.00	190	29	312	10.00	67			
	1992-NIVA	1H W	312	10.00	191	4	312	10.00	111			

Tab.length cont'd.

Tissue				Fish liver					Fish fillet, Shrimptail, Mussel, Other				
Contam.	Mon. Year	Lab. Inter-calibr. +Basis	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	
	1993-NIVA	1H W	312	50.00	221	98		312	50.00	79			
	1994-NIVA	1Z W	312	50.00	302	134		312	50.00	81			
	1995-NIVA	W	312	50.00	318	129		312	50.00	139	2		
	1996-NIVA	V1 W						312	50.00	125			
	1996-NIVA	V2 W	312	50.00	368	128							
CDD1N	1995-NILU	W						841	.20E-04	6	1	1	
	1996-NILU	W						841	.10E-04	18		2	
CDD4X	1995-NILU	W						841	.20E-04	6	3	1	
	1996-NILU	W						841	.200E+08	18	18		
CDD6P	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.40E-04	18			
CDD6X	1995-NILU	W						841	.20E-04	6		1	
	1996-NILU	W						841	.20E-04	18		1	
CDD9X	1995-NILU	W						841	.20E-04	6	2	1	
	1996-NILU	W						841	.20E-04	18		1	
CDDO	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-03	18			
CDDSN	1995-NILU	W						841	.20E-04	5			
	1996-NILU	W						841	.10E-04	18		3	
CDDSP	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.40E-04	18			
CDDST	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18			
CDDSX	1995-NILU	W						841	.20E-04	5			
	1996-NILU	W						841	.20E-04	18		2	
CDF2N	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18		1	
CDF2T	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18			
CDF4X	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.20E-04	18		1	
CDF6P	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.40E-04	18	2	1	
CDF6X	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.20E-04	18		1	
CDF9P	1995-NILU	W						841	.20E-04	6	2	1	
	1996-NILU	W						841	.80E-04	17	3	1	
CDF9X	1995-NILU	W						841	.20E-04	6	3	1	
	1996-NILU	W						841	.20E-04	18		1	
CDFDN	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18		1	
CDFDX	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.20E-04	18		1	
CDFO	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-03	18	3	1	
CDFSN	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18		1	
CDFSP	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.80E-04	18	6	1	
CDFST	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.10E-04	18			
CDFSX	1995-NILU	W						841	.20E-04	6			
	1996-NILU	W						841	.20E-04	18		1	
CHR	1992-NIVA	W	309	0.20	8			309	0.20	44			
	1995-NIVA	W						309	0.20	56			
	1996-NIVA	W						309	0.20	65		3	
CHRTR	1995-NIVA	W						309	0.20	15		2	
CO	1996-NIVA	W						999	miss	3			
COR	1992-NIVA	W	309	0.20	8			309	0.20	46			
CR	1992-NIVA	W						312	10.00	6			
	1996-NIVA	W						999	miss	3			
CU	1983-SIIF	1G W						130	10.00	12			
	1984-SIIF	1G W						130	10.00	27			
	1986-NIVA	1H D	311	150.00	56			311	150.00	20			
	1987-FIER	1G W	404	50.00	37								
	1987-NIVA	1H D	311	150.00	57			311	150.00	37			
	1988-NIVA	1H D	311	150.00	61			311	150.00	55			
	1989-NIVA	1H D	311	150.00	135								
	1989-NIVA	1H W						311	150.00	36			
	1990-NIVA	1H W	311	150.00	189			311	150.00	77			
	1991-NIVA	1H W	311	50.00	193	2		311	50.00	67			
	1992-NIVA	1H W	311	10.00	191			311	10.00	111			
	1993-NIVA	1H W	311	10.00	221			311	10.00	79			
	1994-NIVA	1Z W	311	10.00	302			311	10.00	81			
	1995-NIVA	W	311	10.00	318			311	10.00	124			
	1996-NIVA	V1 W						311	10.00	113			
	1996-NIVA	V2 W	311	10.00	368								
DBA3A	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	71		48	
	1996-NIVA	W						309	0.20	65		53	
DBP	1992-NIVA	W	309	0.20	8			309	0.20	46			
DBTC1	1995-NIVA	W						309	0.20	57		14	
	1996-NIVA	W						309	0.20	65		9	
DBTC2	1995-NIVA	W						309	0.20	56		9	
	1996-NIVA	W						309	0.20	62		11	
DBTC3	1995-NIVA	W						309	0.20	57		4	
	1996-NIVA	W						309	0.20	65		5	
DDEPP	1982-VETN	W	210	50.00	53								

JAMP National Comments 1996 - Norway

Tab.length cont'd.

Tissue		Fish liver							Fish fillet, Shrimptail, Mussel, Other				
Contam.	Mon. Year	Lab.	Inter-calibr.+Basis	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim
	1983-VETN		2E W		210	50.00	48		211a	50.00	48		
	1984-VETN		2E W		210	50.00	66						
	1985-VETN		2E W		210	50.00	45						
	1986-NACE		2Z W		510	20.00	56						
	1987-NACE		2Z W		510	40.00	53						
	1988-NACE		2Z W		510	40.00	61						
	1989-NACE		2Z W		510	20.00	93						
	1990-NIVA		W		340	1.00	169		341	0.05	58		
	1991-NIVA		W		340	1.00	179		341	0.05	62		
	1992-NIVA		W		340	5.00	192	2	341	0.10	140		
	1993-NIVA		W		340	4.00	212	3	341	0.10	133		
	1994-NIVA		2Z W		340	4.00	300		341	0.10	165	27	
	1995-NIVA		W		340	4.00	318	2	341	0.10	216	30	
	1996-NIVA		W		340	4.00	332	2	341	0.10	228	47	
DDTEP	1983-SIIF		W						111	0.50	12		
	1984-SIIF		W						111	0.50	24		1
	1985-SIIF		W						111	0.50	27	1	5
	1986-SIIF		W						111	0.50	21		
	1987-SIIF		W						111	0.50	21	1	
	1988-SIIF		D						111	0.50	6		
	1988-SIIF		W						111	0.50	22	1	
	1989-SIIF		W						111	0.50	36	1	
	1990-SIIF		W						111	0.20	41	1	
	1991-SIIF		W						111	0.30	35		
DDTPP	1986-NACE		W		510	40.00	56						
	1987-NACE		W		510	40.00	53						
	1988-NACE		W		510	40.00	61						
	1989-NACE		W		510	20.00	93						
	1995-NIVA		W						340	0.05	60		
	1996-NIVA		W		340	0.05	54	4	340	0.05	36		
EOCL	1989-SIIF		W						605	170.00	5		
EPOCL	1986-NACE		W		610	800.00	56						
	1986-SIIF		W						605	5000.00	21	21	
	1987-NACE		W		610	800.00	53						
	1987-SIIF		W						605	40.00	20		
	1988-NACE		W		610	800.00	60						
	1988-SIIF		W						605	40.00	27		
	1989-NACE		W		610	800.00	89	1					
	1989-SIIF		W						605	40.00	35		
	1990-NIVA		W		615	40.00	117	3					
	1990-SIIF		W						605	40.00	41		
	1991-NIVA		W		615	40.00	116	12					
	1991-SIIF		W						605	130.00	35		
FLE	1992-NIVA		W		309	0.20	8		309	0.20	45		
	1995-NIVA		W						309	0.20	72		22
	1996-NIVA		W						309	0.20	65		6
FLU	1992-NIVA		W		309	0.20	8		309	0.20	44		
	1995-NIVA		W						309	0.20	72		
	1996-NIVA		W						309	0.20	65		
HCB	1983-SIIF		W						111	0.50	12		
	1983-VETN		2Z W		210	10.00	48		211a	10.00	48		
	1984-SIIF		W						111	0.20	24		1
	1984-VETN		2Z W		210	10.00	66						
	1985-SIIF		W						111	0.20	30	6	2
	1985-VETN		2Z W		210	10.00	45	4					
	1986-NACE		2Z W		510	10.00	56						
	1986-SIIF		2Z W						111	0.20	21	3	
	1987-NACE		2Z W		510	40.00	53						
	1987-SIIF		2Z W						111	0.20	21	4	
	1988-NACE		2Z W		510	40.00	61						
	1988-SIIF		2Z D						111	0.20	6		
	1988-SIIF		2Z W						111	0.20	22	2	
	1989-NACE		2Z W		510	20.00	93						
	1989-SIIF		2Z W						111	0.05	36		
	1990-NIVA		W		340	1.00	169	2	341	0.05	58		
	1990-SIIF		2Z W						111	0.05	41	3	
	1991-NIVA		W		340	1.00	179	4	341	0.05	62	5	
	1991-SIIF		2Z W						111	0.10	35		
	1992-NIVA		W		340	5.00	189	3	341	0.10	140		
	1993-NIVA		W		340	4.00	212	31	341	0.10	133		
	1994-NIVA		2Z W		340	3.00	300	24	341	0.05	165	33	
	1995-NIVA		W		340	3.00	317	37	341	0.05	216	30	
	1996-NIVA		W		340	3.00	332	52	341	0.05	228	37	
HCHA	1990-NIVA		W		340	1.00	168		341	0.05	58		
	1991-NIVA		W		340	1.00	179	2	341	0.05	62	5	10
	1992-NIVA		W		340	5.00	192	3	341	0.10	140		
	1993-NIVA		W		340	4.00	212	45	341	0.10	133		
	1994-NIVA		2Z W		340	3.00	296	32	341	0.05	165	85	
	1995-NIVA		W		340	3.00	318	45	341	0.05	216	98	
	1996-NIVA		W		340	3.00	332	111	341	0.05	222	100	
HCHG	1986-NACE		W		510	30.00	56	1					
	1986-SIIF		W						111	3.00	21		
	1987-NACE		W		510	40.00	53						
	1987-SIIF		W						111	5.00	21		1
	1988-NACE		W		510	40.00	61						
	1989-NACE		W		510	20.00	93						
	1989-SIIF		W						111	50.00	36		
	1990-NIVA		W		340	1.00	169	1	341	0.05	58		

Tab.length cont'd.

Tissue				Fish Liver					Fish fillet, Shrimptail, Mussel, Other				
Contam.	Mon. Year	Lab. Inter-calibr.+Basis	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	
HG	1990-SIIF	W						111	0.10	41			
	1991-NIVA	W	340	1.00	179	3	18	341	0.05	62	5	1	
	1991-SIIF	W						111	0.30	35			
	1992-NIVA	W	340	5.00	192	3		341	0.10	140			
	1993-NIVA	W	340	4.00	212	42	17	341	0.10	133			
	1994-NIVA	2Z W	340	3.00	300	24	1	341	0.05	165	46		
	1995-NIVA	W	340	3.00	313	31		341	0.05	204	29		
	1996-NIVA	W	340	3.00	330	68		341	0.05	211	8		
	1981-SIIF	1E W	120	10.00	15		1	120	10.00	35			
	1982-SIIF	1E W						120	10.00	18			
	1982-VETN	W	220	10.00	51			220	10.00	54			
	1983-SIIF	1E W						120	10.00	17			
	1983-VETN	1Z W						220	10.00	48			
	1984-FIER	1G W						401	10.00	39			
	1984-SIIF	1G W						120	10.00	27	6		
	1984-VETN	1Z W						220	10.00	66			
	1985-SIIF	1G D						120	10.00	30			
	1985-VETN	1Z W						220	10.00	90			
	1986-NIVA	1H D						310	10.00	74			
	1987-FIER	1G W						401	10.00	38			
	1987-NIVA	1H D						310	10.00	93		14	
	1988-NIVA	1H D						310	10.00	116			
	1989-NIVA	1H D						310	100.00	134			
	1989-NIVA	1H W						310	10.00	36	5		
	1990-NIVA	1H W						310	10.00	266			
	1991-NIVA	1H W						310	100.00a	264	126		
	1992-NIVA	1H W						310	100.00a	303	122		
	1993-NIVA	1H W						310	5.00	300			
	1994-NIVA	1Z W						310	5.00	381			
	1995-NIVA	W						310	5.00	442	1		
1996-NIVA	V1 W						310	5.00	481				
ICDP	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	72		29	
MN	1996-NIVA	W						309	0.20	65		23	
	1984-SIIF	W						132	40.00	27			
NAP	1985-SIIF	D						132	40.00	35			
	1992-NIVA	W	309	0.20	8			309	0.20	46			
NAP1M	1995-NIVA	W						309	0.20	70		21	
	1996-NIVA	W						309	0.20	61		11	
NAP2M	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	15		13	
NAP3	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	15		13	
NAPC1	1995-NIVA	W						309	0.20	55		6	
	1996-NIVA	W						309	0.20	61			
NAPC2	1995-NIVA	W						309	0.20	57		6	
	1996-NIVA	W						309	0.20	60			
NAPC3	1995-NIVA	W						309	0.20	57		5	
	1996-NIVA	W						309	0.20	60			
NAPDI	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	15		6	
NAPTM	1992-NIVA	W	309	0.20	8			309	0.20	46			
	1995-NIVA	W						309	0.20	15		11	
NI	1983-SIIF	1G W						130	20.00	12			
	1992-NIVA	W						312	10.00	6			
	1996-NIVA	W						999	miss	3			
	1990-NIVA	W	340	2.00	169	31	24	341	0.05	58		1	
OCS	1991-NIVA	W	340	2.00	179	14	81	341	0.05	62	5	8	
	1992-NIVA	W	340	5.00	192	3		341	0.10	140			
	1993-NIVA	W	340	4.00	212	51	16	341	0.10	133			
	1994-NIVA	W	340	3.00	300	39	22	341	0.05	165	96		
	1995-NIVA	W	340	3.00	318	44		341	0.05	216	102		
	1996-NIVA	W	340	3.00	332	287		341	0.05	228	114		
	1992-NIVA	W	309	0.20	8			309	0.20	45			
	1995-NIVA	W						309	0.20	72			
PAC1	1996-NIVA	W						309	0.20	65			
	1995-NIVA	W						309	0.20	57		1	
PAC2	1996-NIVA	W						309	0.20	65			
	1995-NIVA	W						309	0.20	56			
PAH	1996-NIVA	W						309	0.20	65		2	
	1987-NIVA	W	309	0.02	1								
PAM1	1992-NIVA	W	309	0.20	8			309	0.20	45			
	1995-NIVA	W						309	0.20	15		2	
PB	1983-SIIF	1G W						130	20.00	12			
	1984-SIIF	1G W						130	20.00	27		2	
	1985-SIIF	1G D						130	20.00	35			
	1986-NIVA	1Z D	312	150.00	56	4		312	150.00	20			
	1987-FIER	1G W	403	10.00	37	1							
	1987-NIVA	1Z D	312	150.00	57		12	312	150.00	37			
	1988-NIVA	1Z D	312	150.00	61	17	3	312	150.00	55			
	1989-NIVA	1Z D	312	150.00	135	9	9						
1989-NIVA	1Z W						312	150.0					

Tab.length cont'd.

Tissue			Fish liver					Fish fillet, Shrimptail, Mussel, Other				
Contam.	Mon. Year	Lab. Inter-calibr.+Basis	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim	Analys Method Code	Detect Limit (ppb)	Total Value Count	Count Below D.Lim	N (<) Above D.Lim
PCB	1996-NIVA	V1 W					109	312	30.00	110		
	1996-NIVA	V2 W	312	30.00	368							
	1981-SIIF	2D W	110	10.00	27			110	10.00	35		
	1982-SIIF	2D W						111	5.00	17		
	1982-VETN	W	210	50.00	53			211	50.00	54		
	1983-SIIF	2E W						111	5.00	14		
	1983-VETN	2E W						211	50.00	48		
	1983-VETN	2Z W	210	50.00	48							
	1984-SIIF	2E W						111	5.00	24		
	1984-VETN	2E W						211	50.00	66		
	1984-VETN	2Z W	210	50.00	66							
	1985-SIIF	2E W						111	5.00	32		6
	1985-VETN	2E W						211	50.00	90		1
	1985-VETN	2Z W	210	50.00	45							
	1986-NACE	2Z W	511a	40.00a	56			511	20.00	56		
	1986-SIIF	2E W						111	5.00	21		
	1987-NACE	2Z W	510	40.00	53			511	20.00	54		
	1987-NIVA	W	340	0.10	2							
	1987-SIIF	2E W						111	5.00	21		
	1988-NACE	2Z W	510	40.00	61			511	20.00	13		
1988-SIIF	2E D						111	5.00	6			
1988-SIIF	2E W						111	5.00	22	4		
1989-NACE	2Z W	510	20.00	93			511	20.00	17			
1989-SIIF	2E W						111	5.00	36	6		
1990-SIIF	2E W						111	5.00	41			
1991-SIIF	2E W						111	5.00	35			
PCC26	1996-NILU	W					842	.10E-02	6			
PCC32	1996-NILU	W					842	.30E-02	6		4	
PCC50	1996-NILU	W					842	.10E-02	6			
PCC62	1996-NILU	W					842	0.03	6		6	
PCDD	1995-NILU	W					841	.20E-04	6			
	1996-NILU	W					841	.10E-03	18			
PCDF	1995-NILU	W					841	.20E-04	6			
	1996-NILU	W					841	.10E-03	18			
PER	1992-NIVA	W	309	0.20	8			309	0.20	46		
	1995-NIVA	W						309	0.20	72		32
	1996-NIVA	W						309	0.20	65		40
PYR	1992-NIVA	W	309	0.20	8			309	0.20	44		
	1995-NIVA	W						309	0.20	72		4
	1996-NIVA	W						309	0.20	65		1
QCB	1990-NIVA	W	340	2.00	169	33	39	341	0.05	58		
	1991-NIVA	W	340	2.00	178	13	97	341	0.05	57	5	7
	1992-NIVA	W	340	5.00	192	3		341	0.10	125		
	1993-NIVA	W	340	4.00	212	52	24	341	0.10	133		
	1994-NIVA	W	340	3.00	299	38	23	341	0.05	165	93	
	1995-NIVA	W	340	3.00	318	45		341	0.05	216	103	
	1996-NIVA	W	340	3.00	332	306		341	0.05	228	109	
SE	1982-VETN	W	240	10.00	46			240	10.00	54		
TCDD	1995-NILU	W						841	.20E-04	6	1	
	1996-NILU	W						841	.10E-04	18		
TDEPP	1991-NIVA	W	340	1.00	138		1	341	0.05	62		
	1992-NIVA	W	340	5.00	191	3		341	0.10	140		
	1993-NIVA	W	340	4.00	212	24	3	341	0.10	133		
	1994-NIVA	2Z W	340	3.00	300	17	5	341	0.05	165	47	
	1995-NIVA	W	340	3.00	318	36		341	0.05	213	51	
	1996-NIVA	W	340	3.00	332	23		341	0.05	228	16	
V	1996-NIVA	W						999	miss	3		
ZN	1983-SIIF	1G W						131	400.00	12		
	1984-SIIF	1G W						132	400.00	27		
	1985-SIIF	1G D						132	400.00	35		
	1986-NIVA	1H D	311	3000.00	56			311	3000.00	20		
	1987-FIER	1G W	405	20.00	37							
	1987-NIVA	1H D	311	3000.00	57			311	3000.00	37		
	1988-NIVA	1H D	311	3000.00	61			311	3000.00	55		
	1989-NIVA	1H D	311	3000.00	135		1					
	1989-NIVA	1H W						311	3000.00	36		
	1990-NIVA	1H W	311	3000.00	189			311	3000.00	77		
	1991-NIVA	1H W	311	1000.00	193			311	1000.00	67		
	1992-NIVA	1H W	311	1000.00	191			311	1000.00	111		
	1993-NIVA	1H W	311	1000.00	221			311	1000.00	79		
	1994-NIVA	1Z W	311	1000.00	302			311	1000.00	81		
	1995-NIVA	W	311	1000.00	318			311	1000.00	142		
	1996-NIVA	V1 W						311	1000.00	131		
	1996-NIVA	V2 W	311	1000.00	368							
Sum of Counts					40365	3611	1007			32068	2896	966

a(6) > Ambiguous value in cell (Maximum value displayed).

Appendix E. Overview of localities

Station positions are shown on maps (Figure 1 to Figure 4).

JAMP stations and programme 1996

Appendix E1. JMP station positions and sampling overview for 1996. WSBOFR: W=water, S=sediment, B=blue mussel, O=other shellfish, F=flatfish, R=roundfish. second station position indicates previous location. NSTF=North Sea Task Force. Mussels were sampled from rock surfaces unless otherwise noted.

JAMP area	St.	Locality name	North latitude	East longitude	ICES position	1996						notes
						W	S	B	O	F	R	
26 OSLOFJORD AREA EAST, Hvaler and Singlefjorden												
26	01A	Sponvika	59°05.4'	11°12.5'	47G13							
			59°05.1'	11°13.9'	47G13							
26	02A	Fugleskjær	59°06.6'	10°59.3'	47G09							
			59°06.9'	10°59.0'	47G09							
26	03A	Tisler	58°59.0'	10°57.8'	46G07							C
			58°58.8'	10°57.5'	46G07							
26 OSLOFJORD AREA CENTRAL, Oslofjord proper												
26	30A	Gressholmen	59°52.5'	10°43.0'	48G07	+	+					
26	30B	Oslo city area / Håøya	59°49'	10°33'	48G04						+	
			59°44'	10°32'	48G04							
26	30B	Oslo city area / Nesodden	59°52'	10°39'	48G04							
26	30F	Oslo city area / Håøya	59°47'	10°34'	48G04							
26	30X	West of Nesodden	59°48.5'	10°36'	48G04							
26	30G	Steilene area (Spro)	59°45.8'	10°34.5'	48G05							
26	30H	Steilene area (Storegrunn)	59°48.5'	10°33.5'	48G05							
26	40C	Steilene	59°49'	10°33'	48G05							
			59°49'	10°39'	48G05							
26	30S	Steilene	59°49.1'	10°33.8'	48G05							
26	31A	Solbergstrand	59°36.9'	10°39.4'	48G06	+	+					
26	31B	Solbergstrand (Filtvet, 1982)	59°37'	10°39'	47G07							
26	32A	Rødtangen	59°31.5'	10°25.6'	48G06							
26	33X	Sande, west side	59°31.7'	10°20.4'	48G06							
26	33B	Sande, east side	59°31.7'	10°21.0'	48G06					+		
26	35A	Mølen	59°29.2'	10°30.1'	47G04	+	+					
26	35C	Holmenstrand-Mølen	59°29'	10°27'	47G04							
26	35S	Mølen	59°30'	10°35'	47G04							
26	36A	Færder	59°01.6'	10°31.7'	47G06	+	+					
26	36B	Færder area	59°02'	10°27'	47G06						+	
			59°02'	10°32'	47G06							
26	36F	Færder area	59°04'	10°23'	47G06					+		
26	36S	Færder area (NSTF-54)	59°00.4'	10°41.6'	47G09							N
26 OSLOFJORD AREA WEST, outer Sandefjord-Langesundsfjord												
26	73A	Lyngholmen	59°02.6'	10°18.1'	47G03							C
26	74A	Oddeneskjær	58°57.3'	09°52.1'	46F97							C
26	71A	Bjørkøya (Risøyodden)	59°01.4'	09°45.4'	47F99	+	+					
ARENDAL AREA												
	76A	Risøy	58°43.6'	09°17.0'	46F92	+	+					C
	77A	Flostafjord	58°31.5'	08°56.9'	46F89							C
	77B	Borøy area	58°33'	09°01'	46F93							
	77F	Borøy area	58°33'	09°01'	46F93							
	77C	Borøy area	58°29'	09°10'	45F91							

Appendix E (cont'd)

JAMP area	St.	Locality name	North latitude	East longitude	ICES position	1996						notes
						W	S	B	O	F	R	
ARENDALE AREA (cont.)												
	77S	Arendal area (NSTF-57)	58°24.2'	09°01.8'	45F91							N, C
	79A	Gjerdvoldsøyen, east	58°25.0'	08°45.3'	45F87							C
LISTA AREA												
	13A	Langøysund	57°59.8'	07°34.6'	44F74							C
	14A	Aavigen	58°02.2'	07°13.2'	45F73							C
	15A	Gåsøy (Ullerø area)	58°03.1'	06°53.3'	45F69	+	+					
	15B	Ullerø area	58°03'	06°43'	45F69						+	
	15F	Ullerø area	58°03'	06°43'	45F69						+	
	15S	Lista area (NSTF-39)	58°01.0'	06°34.3'	45F66							N, C
BØMLO AREA												
	22A	Espevær, west	59°35.2'	05°00.5'	48F59	+	+					C, 1
	22F	Borøyfjorden	59°43'	05°21'	48F55						+	
	22C	Bømløfjorden	59°34'	05°11'	48F53							
	22S	Bømlø (NSTF-36)	59°25.9'	04°50.2'	47F47							N
	23A	Austvik	59°52.2'	05°06.6'	48F51							
	23B	Karihavet area	59°55'	05°07'	48F51							+
	24A	Vardøy	60°10.2'	05°00.8'	49F52							C
	24S	Sotra	60°15.1'	04°33.3'	49F45							N
62 HARDANGERFJORDEN												
62	69A	Lille Terøy	59°58.8'	05°45.4'	49F59	+	+					
62	69S	Kvinnheradsfjorden	60°01.3'	05°56.1'	49F59							
62	67B	Strandebarm	60°16'	06°02'	49F62						+	+
62	67S	Strandebarm	60°13.5'	06°05.1'	49F62							
62	65A	Vikingneset	60°14.5'	06°09.6'	49F62	+	+					
62	63A	Ranaskjær	60°25.1'	06°24.5'	49F64	+	+					
62	63S	Ranaskjær	60°23.6'	06°27.1'	49F64							
63 SØRFJORDEN												
63	51A	Byrkjenes	60°05.1'	06°33.1'	49F66							
63	52A	Eitrheimsneset	60°05.8'	06°32.2'	49F66	+	+					3
63	52S	Tyssedal	60°06.9'	06°32.9'	49F66							
63	53B	Inner Sørfjord	60°10'	06°34'	49F65						+	+
63	56A	Kvalnes	60°13.4'	06°36.1'	49F65	+	+					
63	56S	Kvalnes	60°13.7'	06°35.6'	49F65							
63	57A	Krossanes	60°23.2'	06°41.2'	49F67	+	+					
63	57S	Krossanes	60°23.1'	06°40.7'	49F67							
ÅLESUND AREA												
	25A	Hinnøy	61°22.2'	04°52.8'	51F47							5
	26A	Hamnen	61°52.7'	05°13.6'	52F51							5
	27A	Grinden	62°12.2'	05°25.4'	53F55							1
	27X	Kvame area	62°12.3'	05°22.2'	53F55							
	27S	Statlandet (east of)	62°09.3'	05°21.3'	53F56							
	28A	Eiksundet	62°14.9'	05°54.5'	53F58							1
		Eiksundet (1992)	62°14.9'	05°54.5'	53F58							1

Appendix E (cont'd)

JAMP area	St.	Locality name	North latitude	East longitude	ICES position	1996						notes
						W	S	B	O	F	R	
65 ORKDALSFJORDEN												
65	80A	Østmerknes	63°27.5'	10°27.5'	56G04							
65	81A	Biologisk station	63°26.5'	10°21.4'	56G04							
65	82A	Flakk	63°27.1'	10°12.6'	56G01	+		+				
65	82S	Flakk	63°27.5'	10°11.8'	56G01							
65	83A	Frøsetskjær	63°25.5'	10°07.8'	56G01							
65	84A	Trossavika	63°20.8'	09°57.8'	55F97	+		+				
65	84B	Trossavika	63°20.8'	09°57.8'	55F97							
65	84S	Trossavika	63°21.7'	09°57.4'	55F97							
		(1987)	63°21.2'	09°57.2'	55F97							
65	89S	Thamshavn (indre Orkdal)	63°19.7'	09°52.3'	55F98							
		(1987)	63°19.8'	09°52.5'	55F98							
65	90S	Outer Orkdalsfjord	63°27.4'	10°03.0'	56G01							
		(1987)	63°27.4'	10°04.3'	56G01							
65	85A	Geitastrand	63°21.9'	09°56.3'	55F97							
65	86A	Geitnes	63°26.6'	09°59.2'	55F97							
65	87A	Ingdalsbukta	63°27.8'	09°54.8'	55F97	+		+				
65	88A	Rødberg	63°27.2'	10°00.0'	55G01							
FROAN AREA												
	91A	Nerdvika	63°21.2'	08°09.6'	55F81							3
		Fosflua (1992)	63°23.8'	08°17.6'	55F81							4
	92A	Stokken	64°04.6'	10°00.7'	57G03	+		+				4
	92B	Stokken area	64°09.9'	09°53.0'	57F99						+	
	92F	Stokken area	64°09.9'	09°53.0'	57F99							
	93S	Raudøya (northeast of)	64°22.7'	10°27.8'	57G04							
	93A	Låven (Sætervik)	64°23.7'	10°29.0'	57G04							4
		Låven (Sætervik, 1992))	64°23.5'	10°28.0'	57G04							4
HELGELAND AREA												
	94A	Landfast	65°38.4'	12°00.5'	60G23							1
	96A	Breiviken	66°17.6'	12°50.5'	61G28							1
	95S	Rødø (east of)	66°41.8'	13°09.9'	62G32							
	95A	Flatskjær	66°42.6'	13°15.8'	62G32							4
LOFOTEN AREA												
	97A	Klakholmen	67°39.9'	14°44.6'	64G49							4
	99A	Brunvær	68°00.3'	15°05.6'	65G53							4
	98B	Lille Molla	68°12.0'	14°48.0'	65G48						+	
	98F	Lille Molla	68°12.0'	14°48.0'	65G48					+		
	98S	Skrova (south of)	68°07.0'	14°41.0'	65G49							
	98A	Skrova	68°09.4'	14°39.3'	65G46	*		*				1
	98X	Skrova	68°10.5'	14°40.2'	65G48	+		+				7
	99S	Lundøy (north of)	68°05.8'	15°10.1'	65G53							

Appendix E (cont'd)

JAMP area	St.	Locality name	North latitude	East longitude	ICES position	1996						notes
						W	S	B	O	F	R	
FINNSNES-SKJERVØY AREA												
	41S	Andfjord	68°56.3'	17°05.2'	66G71							
	41A	Fensneset, Grytøya	68°56.9'	16°38.5'	66G64	+	+					3
	42S	Tromsø area	69°60.4'	18°06.8'	68G83							
	42A	Tennskjær, Malangen	69°28.6'	18°18.0'	67G81	+	+					3
	43S	Kvænangen	70°03.3'	21°07.9'	69H13							
	43A	Lyngneset, Langfjorden	70°06.2'	20°32.8'	69H06	+	+					2
	43B	Kvænangen	70°09.0'	21°22.0'	69H16						+	
	43F	Kvænangen	70°09.0'	21°22.0'	69H16					+		
HAMMERFEST-HONNINGSVÅG AREA												
	44S	Sørøya, south	70°25.9'	22°31.8'	69H24							
	44A	Elenheimsundet	70°30.8'	22°14.8'	70H23	+	+					1, 6
	45S	Hammerfest area	70°42.9'	24°26.6'	70H45							
	45A	Ytre Sauhamneset	70°45.8'	24°19.2'	70H42	+	+					
	46S	Porsangen area	70°52.9'	26°11.9'	70H61							
	46A	Smineset in Altesula	70°58.4'	25°48.1'	70H57	+	+					3, 6
	46B	Hammerfest area	70°50.0'	23°44.0'	70H37						+	
	46F	Honningsvåg area	00°00.0'	00°00.0'					*			
	47S	Laksefjord	70°55.0'	26°55.1'	70H67							
	47A	Kifjordeneset	70°52.9'	27°22.2'	70H74	+	+					
VARANGER PENINSULA AREA												
	48S	Tanafjord	70°52.5'	28°38.5'	70H84							
	48A	Trollfjorden i Tanafjord	70°41.6'	28°33.3'	70H85	+	+					
	49S	Syltefjord	70°33.9'	30°19.9'	70J03							
	49A	Nordfjorden, Syltefjord	70°33.1'	30°05.2'	70J03	+	+					
	10S	Varangerfjord	69°56.1'	30°06.7'	68J01							
	10A	Skagoodden	70°04.2'	30°09.8'	69J03	+	+					2
	10B	Varangerfjorden	69°54.5'	29°30.0'	68H97						+	
	10F	Varangerfjorden	00°00.0'	00°00.0'					*			
	11A	Sildkroneset, Bøkfjorden	69°47.2'	30°11.1'	68J02	+	+					

notes:

- + - samples collected
- * - planned but insufficient material
- N - official NSTF station
- C - at or near SFT's coastal monitoring programme station
- 1 - mussels collected from buoy and/or buoy anchor lines
- 2 - mussels collected from sand/gravel bottom
- 3 - mussels collected from iron/cement pilings
- 4 - mussels collected from metal navigation buoys
- 5 - mussels collected from floating dock
- 6 - mussels collected from wooden docks
- 7 - mussels collected from rocks under ferry terminal

Appendix F. Overview of materials and analyses 1996

Including sampling for VIC (cf., SIME 1996, 1997)

Station positions are shown on maps (Figure 1 to Figure 4)

Appendix F1. Sampling and analyses for 1996, L-liver, F-fillet. (See Appendix F2 for descriptions of codes for analysis (M0, M1, M3, M4, M5, C2, C4, A1, G1), fish (P, F, D, L, M, C) and counts). Analytical overview for liver (-L) or fillet (-F) tissue is distinguished.

JAMP area	STATION	WATER		SEDIMENT			MUSSEL/		OTHER		FISH								
											FLAT- (P,F,D,M)				COD- (C)				
		M0	M1	C4	A1	G1	M3	C2	A1	M3	C2	-L	M4	C2	A1	-L	M4	C2	A1
26	OSLOFJORD AREA CENTRAL, Oslofjord proper																		
26	30A Gressholmen	1	3	3
26	30S Steilene
26	30B Oslo city Area / Håøya
	VIC: Steilene - 15.1.97.	C-L 10	10
	VIC: Steilene - 22.1.97.	C-F 10	2B
	VIC: Steilene - 3.2.97.	C-L 10	10
	VIC: Steilene - 3.2.97.	C-F 10	2B
	VIC: Svestad - 18.1.97.	C-L 10	10
	VIC: Håøya 16.1.97.	C-F 10	2B
	VIC: Håøya 16.1.97.	C-L 10	10
	VIC: Håøya 16.1.97.	C-F 10	2B
26	31A Solbergstrand	1	3	3
26	33B Sande, east side
	VIC: 10.96.	F-L 5B	5B
	VIC: 11.96.	F-F 5B	5B
	VIC: 12.96.	F-L 5B	5B
	VIC: 12.96.	F-F 5B	5B
	VIC: 12.96.	F-F 5B	5B
26	35S Mølen	.	3	2	2
26	35A Mølen	1	3	3
26	36S Færder	.	17	4	4	1
26	36A Færder	1	3	3
26	36B Færder area	C-L 25	25
	VIC: 12.96.	C-F 25	5B
20	36F Færder area	D-L 5B	5B
	VIC: 12.96.	D-F 5B	5B
26	OSLOFJORD AREA WEST, outer Sandefjord-Langesundsfjord																		
26	71A Bjørkøya	1	3	3
	ARENDAL AREA																		
	76A Risøy	1	3	3

Appendix F1 (cont.)

JAMP area	STATION	WATER		SEDIMENT			MUSSEL/			OTHER	FISH							
											FLAT- (P,F,D,M)				COD- (C)			
		M0	M1	C4	A1	G1	M3	C2	A1		-L	M4	C2	A1	-L	M4	C2	A1
									-F	M5	C2	A1	-F	M5	C2	A1		
LISTA AREA																		
	15A Ullerø area	1	3	3	
	15B Ullerø area	C-L 25	25	
		C-F 25	5B	
	15F Ullerø area	D-L 5B	5B	.	.	.	
		D-F 5B	5B	.	.	.	
BØMLO-SOTRA AREA																		
	22A Espevær, west	1	3	3	
	22F Borøyfjorden	P-L 5B	5B	.	.	.	
		P-F 5B	5B	.	.	.	
	23B Karihavet	C-L 25	25	
		C-F 25	5B	
HARDANGERFJORDEN																		
62	69A Lille Terøy	1	3	3	
62	67B Strandebarm	
	VIC: 18.8--.	FL 3B	3B	.	C-L 25	25	
		FF 3B	3B	.	C-F 25	5B	
	VIC: 31.10--.	MF 5B	5B	.	C-L 10	0	
		MF 5B	5B	.	C-F 10	0B	
62	65A Vikingneset	1	3	3	
62	63A Ranaskjær	1	3	3	
SØRFJORDEN																		
63	52A Eitrheimsneset	1	3	3	
63	53B Inner Sørfjord	
	VIC: Tyssedal - 8.96--.	F-L 5B	5B	.	C-L 15	15	
		F-F 5B	5B	.	C-F 15	3B	
	VIC: Tyssedal - 12.96--.	C-L 10	.	
		C-F 10	.	
	VIC: Edna - 8.96--.	F-L 3B	3B	.	C-L 15	15	
		F-F 3B	3B	.	C-F 15	3B	
	VIC: Edna - 12.96--.	C-L 10	.	
		C-F 10	.	
	VIC: Odda - 8.96--.	F-L 3B	3B	.	.	.	
		F-F 3B	3B	.	.	.	
63	56A Kvalnes	1	3	3	
63	57A Krossanes	1	3	3	

Appendix F1 (cont.)

JAMP area	STATION	WATER					SEDIMENT			MUSSEL/ OTHER			FISH							
													FLAT- (P,F,D,M)				COD- (C)			
		M0	M1	C4	A1	G1	M3	C2	A1	-L	M4	C2	A1	-L	M4	C2	A1	-L	M4	C2
ORKDALSFJORD AREA																				
65	82A Flakk	1	3
65	84A Trossavika	1	3	3
65	87A Ingdalsbukta	1	2
FROAN AREA																				
92A	Stokken	1	3	3
92B	Stokken	C-L 25	25	.
92F	Stokken	C-F 25	5B	.
LOFOTEN AREA																				
98X	Skrova	1	3	3
98B	Lille Molla	C-L 25	25	.
98F	Lille Molla	D-L 5B	5B	.	.	C-F 25	5B	.
FINNSNES-SKJERVØY AREA																				
41A	Fensneset, Grytøya	1	3	3
42A	Tennskjær, Malangen	1	3
43A	Lyngneset, Langfjorden	1	3	3
43B	Kvænangen	C-L 25	25	.
43F	Kvænangen	D-L 3B	3B	.	.	C-F 25	5B	.
HAMMERFEST-HONNINGSVÅG AREA																				
44A	Elenheimsundet	1	3
45A	Ytre Sauhamneset	1	3	3
45B	Hammerfest area
45F	Honningsvåg area
46A	Smineset in Altesula	1	3	3	3
47A	Kifjordeneset	1	3
VARANGER PENINSULA AREA																				
48A	Trollfjorden i Tanafjord	1	3	3
49A	Nordfjorden, Syltefjord	1	3
10A	Skagoodden	1	3	3
10B	Varangerfjorden	C-L 25	25	.
10F	Varangerfjorden	C-F 25	5B	.
11A	Sildkroneset, Bøkfjorden	1	3	3

Appendix F2: Key to analysis codes and sample counts used in Appendix F1.

ANALYSIS CODES:

Code	Analyses
M0	suspended matter
M1	Hg, Cd, Cu, Pb, Zn, Li (normalising element) total organic carbon (TOC)
M3	Hg, Cd, Cu, Pb, Zn
M4	Cd Cu Pb Zn (for fish liver)
M5	Hg (for fish fillet)
C1	CB-28,-52,-101,-105,-118,-138,-153,-156,-180, 209, 5-CB, OCS, a+gHCH, HCB, DDT, EPOCI (optional), dry weight percent
C2	CB-28,-52,-101,-105,-118,-138,-153,-156,-180, 209, 5-CB, OCS, a+gHCH, HCB, DDT, EPOCI (optional), fat and dry weight percent
A1	PAH
G1	Sediment core geological dating

SAMPLE COUNT CODES:

Medium	Count	Explanation
SEAWATER	1	sample for suspended matter determination
SEDIMENT	17	17 samples for metal analyses; two cores each with samples from 0-1, 1-2, 2-4, 4-6, 6-10, 10-15, 15-20cm and deepest 5cm slice plus one core with sample from 0-1cm.
	4	4 samples for PCB or PAH analyses; two each cores with samples from 0-1cm and deepest 5cm slice.
	3	3 samples for metal analyses; three cores each with samples from 0-1cm.
MUSSEL	3/6	3 size groups (2-3, 3-4, 4-5cm) each a bulk of ca.50 individuals and/or 1 size group (3-4 or 4-5cm), 3 parallel samples each a bulk of 20 individuals.
	1/2	1 size group (2-3 or 3-4cm), 2 parallel samples each a bulk of 50 individuals.
SHRIMP	2	2 samples of 100 individuals (edible size)
FISH		The number of individual fish or bulk samples of fish (-B) for analyses is shown. Bulk samples of fish consist of 5 fish. The five longest fish make up one bulk sample, the next five longest fish make up the another bulk sample and so on. The letter following the number indicates the fish type: D=dab, F=flounder, L=lemon sole, M=megrim, P=plaice, W=witch and C=cod.

Appendix G.

Temporal trend analyses of contaminants in biota 1981-96

Sorted by contaminant, species and area/station:

Cadmium (Cd)
Copper (Cu)
Mercury (Hg)
Lead (Pb)
Zinc (Zn)
CB-153
DDEPP (ppDDE)
γHCH
HCB

MYTI EDU - Blue Mussel (*Mytilus edulis*)
GADU MOR - Atlantic cod (*Gadus morhua*)
LEPI WHI - Megrin (*Lepidorhombus whiff-iaconis*)
LIMA LIM - Dab (*Limanda limanda*)
PLAT FLE - Flounder (*Platichthys flesus*)

OC	Overconcentration expressed as quotient of median of last year and “high background”)
TRND	trend
D-	Significant linear trend, downward
U-	Significant linear trend, upward
--	No significant trend
-?	No significant linear trend, systematic non-linear trend can not be tested because of insufficient data (<6 years)
-Y	No significant linear trend, but a systematic non-linear trend
DY or UY	Significant linear trend (downward or upward) and a significant non-linear trend. This is considered the same as "-Y"
	SIZE length effect mercury in fillet)
L	Significant difference in concentration levels but pattern of variation same
D	As "L" but pattern of variation significantly different
-	No significant difference between “small” and “large” fish
U95+3	Projected upper 95% confidence interval in three years expressed as quotient of value and “high background”
POWER	Estimated number of years to detect a hypothetical situation of 10% trend a year with a 90% power

Annual Median Concentrations of C D (ppm).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S OC TRND U95+3 POWER	
30A	MYTI EDU				1.065	0.810	1.410	0.600	0.610	0.736	0.769	0.769	1.117	1.257	1.174	0.776	0.800	no	no
31A	MYTI EDU			1.386	1.314	0.890	1.930	0.400	0.430	0.412	0.719	0.727	0.914	0.933	0.781	1.324	0.789	no	no
35A	MYTI EDU			1.347	0.932	1.170	1.300	0.520	0.660	0.647	0.926	1.053	1.350	1.111	0.958	0.894	0.766	no	no
36A	MYTI EDU			0.845	1.191	0.840	1.380	0.590	0.560	0.502	0.407	1.217	1.063	0.899	1.215	1.172	1.602	no	no
71A	MYTI EDU			2.520	1.975	1.419	2.004	0.980	2.110	2.021	0.968	1.088	1.657	1.895	1.974	2.253	1.497	no	no
76A	MYTI EDU										0.658	1.088	1.657	1.895				no	no
15A	MYTI EDU										0.860	1.088	1.657	1.895				no	no
22A	MYTI EDU										0.505	0.831	1.098	1.182	0.794	1.439	1.220	no	no
51A	MYTI EDU										0.532	1.139	1.117	0.844	1.020	1.414	1.145	no	no
52A	MYTI EDU							42.800	58.200									no	no
56A	MYTI EDU							94.400	55.200	94.400	80.083	80.083	43.089	14.651	8.713	36.837	25.272	12.6	?
57A	MYTI EDU							55.900	98.413	45.034	69.444	69.444	51.667	59.681	11.357	19.759	18.377	9.2	?
63A	MYTI EDU							21.100	43.200	36.692	32.803	32.803	32.121	15.397	11.800	30.761	20.000	10.0	128.9
65A	MYTI EDU							47.200	10.300	19.006	30.394	35.140	18.212	7.811	4.228	12.207	8.482	4.2	30.8
69A	MYTI EDU							15.000	5.219	8.291	14.604	23.964	5.088	7.733	3.006	5.367	5.400	2.7	5.9
82A	MYTI EDU												4.261	2.368	2.084	2.910	3.195	1.8	5.6
82A	MYTI EDU							0.990	0.400	1.257	1.200	1.200	1.213	1.147		2.901	3.195	1.6	?
84A	MYTI EDU							2.100	0.960	1.191	1.818	1.818	2.105	1.596		0.981	1.223	no	no
87A	MYTI EDU							0.770	0.690	0.756	0.872	0.872	0.978	0.927		1.639	1.288	no	no
91A	MYTI EDU												1.677	1.275	1.818	1.150	1.270	no	no
92A	MYTI EDU												1.078	0.544	0.939	0.743	0.691	no	no
98X	MYTI EDU														0.712	0.688	0.781	no	no
41A	MYTI EDU														1.891	2.947	1.628	no	no
44A	MYTI EDU														1.691	2.741	1.954	no	no
46A	MYTI EDU														2.696	2.058	2.718	1.4	?
30B	GADU MOR														0.063	0.049	0.045	no	no
36B	GADU MOR														0.021	0.042	0.033	no	no
36B	GADU MOR														0.014	0.016	0.024	no	no
15B	GADU MOR														0.015	0.026	0.014	no	no
23B	GADU MOR														0.015	0.026	0.014	no	no
53B	GADU MOR														0.007	0.007	0.180	1.8	4.7
67B	GADU MOR														0.099	0.033	0.111	1.1	>25
84B	GADU MOR														0.029	0.022	0.066	no	no
92B	GADU MOR														0.025	0.113	0.330	no	no
98B	GADU MOR														0.168	0.183	0.097	3.3	?
43B	GADU MOR														0.230	0.188	0.095	no	no
10B	GADU MOR														0.304	0.259	0.200	no	no
67B	LEPI WHI														0.135	0.147	0.139	no	no
36F	LIMA LIM														0.136	0.125	0.153	no	no
15F	LIMA LIM														0.169	0.125	0.153	no	no
22F	LIMA LIM														0.980	0.182	0.225	no	no
98F	LIMA LIM														0.087	0.091	0.110	no	no
33B	PLAT FLE														0.087	0.091	0.110	no	no
53B	PLAT FLE														0.087	0.135	2.530	8.4	19.1

SB Soft body tissue.
LI Liver tissue.
OC Overconcentration; Median(LastYear)/Background (=?) if missing Background)
U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?) if missing Background or N(years)<=5
POWER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of C U (ppm).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S			
																			OC	TRND	U95+3
30A	MYTI EDU				4.57		7.45	4.96	5.48	5.97	10.26	10.47	5.84	6.67	8.56	6.94	7.69	no	--	1.2	10
31A	MYTI EDU				6.57		8.06	4.44	4.52	4.51	9.04	11.00	5.49	5.67	6.21	7.26	6.61	no	UY	1.2	9
35A	MYTI EDU			7.03	3.62		8.06	4.89	4.58	5.26	8.02	10.06	6.56	6.34	6.61	6.41	6.94	no	--	1.2	11
36A	MYTI EDU			6.32	3.57		6.08	4.47	4.87	4.30	5.50	9.23	5.16	5.76	5.63	7.67	9.06	no	--	2.2	11
71A	MYTI EDU			8.47	5.24			6.08	8.43	6.99	8.33	10.26	7.40	7.88	7.18	8.11	7.66	no	--	1.1	8
76A	MYTI EDU										8.51	10.80	5.65	5.57			6.65	no	--	?	10
15A	MYTI EDU										5.72	7.21		5.50			7.32	no	--	?	8
22A	MYTI EDU										6.35	6.69		5.76			9.29	no	UY	1.5	7
51A	MYTI EDU							7.14	6.14				5.38				10.20	no	1.0	?	9
52A	MYTI EDU										7.49	72.08	9.35	8.45			6.28	no	--	3.3	21
56A	MYTI EDU							8.07	8.04	8.40	5.37	7.54	7.40	9.15			7.03	no	--	no	9
57A	MYTI EDU							8.21	6.33	6.01	6.16	7.27	6.59	6.77			6.54	no	D	no	6
63A	MYTI EDU							9.84	6.16	5.11	6.84	11.07	6.32	6.11			6.56	no	--	1.1	9
65A	MYTI EDU							7.98	4.86	5.19	12.23	8.14	5.51	5.32			5.82	no	--	1.0	11
69A	MYTI EDU												6.19	5.32			7.28	no	--	?	6
82A	MYTI EDU							4.69	5.77	7.49		11.77	6.87	9.61			7.93	no	--	1.2	10
84A	MYTI EDU							56.80	39.30	26.76		17.07	22.24	23.99			7.21	no	--	1.0	18
87A	MYTI EDU							20.10	8.35	5.88		7.28	6.30	6.88			7.52	no	--	2.4	15
91A	MYTI EDU												6.59	6.36				no	--	?	8
92A	MYTI EDU												6.65	6.03				no	--	?	8
98X	MYTI EDU																	no	--	?	6
41A	MYTI EDU																	no	--	?	7
44A	MYTI EDU																	no	--	?	10
46A	MYTI EDU																	no	--	?	<=5
30B	GADU MOR						7.00	28.00	5.12	4.21	4.52	3.74	8.54	6.80			5.11	no	--	no	17
36B	GADU MOR						14.20	20.00	6.98	12.18	10.70	8.74	9.65	7.43			5.11	no	--	no	12
15B	GADU MOR										11.30	1.63	5.83	2.28			9.82	no	--	7.6	22
23B	GADU MOR										8.15	11.10	5.83	7.06			8.75	no	--	1.1	10
53B	GADU MOR						12.27		7.11	5.86	2.40	3.29	4.68	6.01			6.34	no	DY	no	10
67B	GADU MOR							8.00	6.91	13.90	8.94	8.76	5.15	7.98			7.17	no	--	no	13
92B	GADU MOR													7.78			6.98	no	--	?	8
98B	GADU MOR												5.87	8.20			6.17	no	--	?	15
43B	GADU MOR																5.41	no	--	?	<=5
10B	GADU MOR																4.31	no	--	?	11
67B	LEPI WHI							15.30	14.19	18.63	10.70	13.80	13.00	12.40			19.50	no	--	?	9
36F	LIMA LIM										13.40	5.29	8.49	8.09			7.06	no	--	1.7	13
15F	LIMA LIM										9.83	2.77	4.11	4.11			9.12	no	--	?	12
22F	LIMA LIM											6.66	3.38				3.46	no	D?	?	11
98F	LIMA LIM																5.00	no	D?	?	8
33B	PLAT FLE						21.74	34.98	16.26	34.32	20.30	18.80	15.70	18.10			13.20	no	--	no	12
53B	PLAT FLE								13.99	9.02	8.19	18.90	10.50	17.20			14.50	no	--	no	15

SB Soft body tissue.
 LI Liver tissue.
 OC Overconcentration; Median(LastYear)/Background (=?" if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?" if missing Background or N(years)<=5)
 POWER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of H G (ppm).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S OC TRND U95+3 POWER
30A MYTI EDU	SB D.WT						0.147	0.050	0.130	0.044	0.064	0.053	0.051	0.070	0.086	0.057	0.070	no
31A MYTI EDU	SB D.WT			0.076	0.118	0.073	0.147	0.050	0.090	0.022	0.060	0.048	0.051	0.045	0.050	0.062	0.043	no
35A MYTI EDU	SB D.WT			0.093	0.074	0.086	0.120	0.050	0.180	0.050	0.062	0.058	0.058	0.054	0.061	0.037	0.038	no
36A MYTI EDU	SB D.WT			0.052	0.043	0.084	0.140	0.050	0.140	0.034	0.045	0.048	0.039	0.032	0.048	0.033	0.044	no
71A MYTI EDU	SB D.WT			0.393	0.242	0.218	0.247	0.120	0.340	0.249	0.182	0.145	0.178	0.140	0.212	0.201	0.222	2.5
76A MYTI EDU	SB D.WT										0.071	0.068	0.050	0.020	0.057	0.057	0.057	?
15A MYTI EDU	SB D.WT										0.056	0.072	0.078	0.024	0.050	0.022	0.049	no
22A MYTI EDU	SB D.WT						0.240	0.240	0.250		0.053	0.053	0.112	0.048	0.067	0.066	0.072	no
51A MYTI EDU	SB D.WT														1.510	0.901	0.901	4.5
52A MYTI EDU	SB D.WT									2.352	0.321	3.008	0.976	0.372	0.282	1.510	0.901	?
56A MYTI EDU	SB D.WT						0.530	0.355	0.355	1.087	0.710	1.540	0.935	1.223	0.352	0.679	0.178	8.4
57A MYTI EDU	SB D.WT						0.170	0.210	0.269	0.269	0.411	0.758	0.576	0.349	0.350	0.260	0.155	4.8
63A MYTI EDU	SB D.WT						0.310	0.140	0.177	0.394	0.394	0.468	0.294	0.143	0.190	0.252	0.172	no
65A MYTI EDU	SB D.WT						0.100	0.135	0.104	0.513	0.513	0.528	0.124	0.119	0.134	0.148	0.118	no
69A MYTI EDU	SB D.WT												0.106	0.026	0.083	0.070	0.104	no
82A MYTI EDU	SB D.WT							0.080	0.120	0.067	0.067	0.074	0.052	0.026	0.049	0.049	0.049	no
84A MYTI EDU	SB D.WT							0.080	0.240	0.057	0.066	0.066	0.090	0.057	0.054	0.043	0.043	no
87A MYTI EDU	SB D.WT							0.050	0.260	0.046	0.056	0.056	0.054	0.076	0.044	0.044	0.062	no
91A MYTI EDU	SB D.WT											0.052	0.054	0.076				?
92A MYTI EDU	SB D.WT											0.055	0.055	0.033				?
98X MYTI EDU	SB D.WT														0.335	0.340	0.328	1.6
41A MYTI EDU	SB D.WT														0.069	0.063	0.064	?
44A MYTI EDU	SB D.WT														0.055	0.050	0.052	?
46A MYTI EDU	SB D.WT														0.059	0.062	0.056	?
30B GADU MOR(s)	MU W.WT														0.102	0.080	0.108	no
308 GADU MOR(l)	MU W.WT														0.129	0.119	0.142	1.1
368 GADU MOR(s)	MU W.WT														0.053	0.076	0.066	1.3
368 GADU MOR(l)	MU W.WT														0.060	0.095	0.066	1.4
15B GADU MOR(s)	MU W.WT														0.080	0.075	0.066	no
158 GADU MOR(l)	MU W.WT														0.045	0.043	0.058	no
23B GADU MOR(s)	MU W.WT														0.081	0.045	0.087	no
238 GADU MOR(l)	MU W.WT														0.051	0.069	0.059	no
53B GADU MOR(s)	MU W.WT														0.073	0.109	0.057	no
67B GADU MOR(l)	MU W.WT														0.120	0.053	0.229	no
84B GADU MOR(s)	MU W.WT														0.090	0.050	0.277	2.3
84B GADU MOR(l)	MU W.WT														0.071	0.073	0.073	2.8
92B GADU MOR(s)	MU W.WT														0.106	0.072	0.089	no
92B GADU MOR(l)	MU W.WT																	1.6
98B GADU MOR(s)	MU W.WT																	?
98B GADU MOR(l)	MU W.WT																	?
43B GADU MOR(s)	MU W.WT																	?
43B GADU MOR(l)	MU W.WT																	?
108 GADU MOR(s)	MU W.WT																	?
108 GADU MOR(l)	MU W.WT																	?
67B LEPI WHI(s)	MU W.WT																	?
67B LEPI WHI(l)	MU W.WT																	?
36F LIMA LIM(s)	MU W.WT																	?
36F LIMA LIM(l)	MU W.WT																	?
15F LIMA LIM(s)	MU W.WT																	?
15F LIMA LIM(l)	MU W.WT																	?
22F LIMA LIM(s)	MU W.WT																	?
22F LIMA LIM(l)	MU W.WT																	?
33B PLAT FLE(s)	MU W.WT																	?
33B PLAT FLE(l)	MU W.WT																	?
53B PLAT FLE(s)	MU W.WT																	?
53B PLAT FLE(l)	MU W.WT																	?

(S) Small fish
 (L) Large fish
 MU Soft body tissue.
 MU Muscle tissue.
 OC Overconcentration; Median(LastYear)/Background (=?" if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?" if missing Background or N(years)<=5)
 POWER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of P B (ppm).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S		
																			OC	TRND
30A MYTI EDU	SB D.WT										1.859	1.361	3.952	2.270	2.543	1.579	2.120	no	--	1.3
31A MYTI EDU	SB D.WT										1.383	1.212	1.264	1.027	1.370	1.679	1.789	no	UY	1.1
35A MYTI EDU	SB D.WT										1.437	1.071	1.676	1.198	1.284	0.507	0.628	no	--	no
36A MYTI EDU	SB D.WT										1.009	0.847	0.787	1.123	1.394	1.238	2.044	no	U-	1.2
71A MYTI EDU	SB D.WT										1.161	0.745	1.716	1.421	1.923	1.494	2.209	no	--	1.7
76A MYTI EDU	SB D.WT										1.773	0.968	1.505	0.913			0.796	no	--	?
15A MYTI EDU	SB D.WT										1.457	0.777		0.976	1.053	0.522	0.671	no	--	no
22A MYTI EDU	SB D.WT										1.371	1.456	2.778	0.976			0.671	no	--	no
52A MYTI EDU	SB D.WT										12.073	312.500	189.430	65.504	16.354	17.532	1.506	no	Y	16.3
56A MYTI EDU	SB D.WT										20.738	23.413	121.500	109.380	24.691	46.418	9.843	3.3	--	27.2
57A MYTI EDU	SB D.WT										10.548	12.137	33.258	19.194	15.071	13.195	5.596	1.9	--	2.8
63A MYTI EDU	SB D.WT										12.137	10.093	15.430	10.938	7.215	12.086	7.600	2.5	--	6.0
65A MYTI EDU	SB D.WT										5.605	3.784	5.190	6.533	3.277	4.725	2.412	no	--	1.5
69A MYTI EDU	SB D.WT											4.619	3.421		2.800	3.166	4.024	1.3	--	?
82A MYTI EDU	SB D.WT											1.278	0.933	0.916		0.622	0.674	no	D?	?
84A MYTI EDU	SB D.WT										1.010	1.152	1.383	1.383		1.378	0.833	no	--	?
87A MYTI EDU	SB D.WT										0.974	0.870	0.634	0.634		1.402	2.468	no	U?	?
91A MYTI EDU	SB D.WT											0.898	1.465		2.013	1.094	0.654	no	--	?
92A MYTI EDU	SB D.WT											0.933	0.628		1.094	0.664	0.654	no	--	?
98X MYTI EDU	SB D.WT														4.341	3.121	4.115	no	--	?
41A MYTI EDU	SB D.WT														1.287	0.900	0.793	no	--	?
44A MYTI EDU	SB D.WT														2.813	2.568	1.657	no	--	?
46A MYTI EDU	SB D.WT														1.257	1.570	1.385	no	--	?
30B GADU MOR	LI W.WT														0.120	0.110	0.060	no	--	1.4
36B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
15B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
23B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
53B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
67B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
92B GADU MOR	LI W.WT														0.030	0.020	0.030	no	--	no
43B GADU MOR	LI W.WT														0.090	0.040	0.040	no	--	1.2
10B GADU MOR	LI W.WT														0.030	0.030	0.040	no	--	no
67B LEPI UHT	LI W.WT														0.030	0.030	0.040	no	--	?
36F LIMA LIM	LI W.WT														0.030	0.030	0.040	no	U?	?
15F LIMA LIM	LI W.WT														0.030	0.020	0.040	no	--	?
22F LIMA LIM	LI W.WT														0.040	0.070	0.030	no	--	?
98F LIMA LIM	LI W.WT														0.030	0.020	0.030	no	D-	no
33B PLAT FLE	LI W.WT														0.060	0.070	0.030	no	D?	?
53B PLAT FLE	LI W.WT														0.020	0.040	0.030	no	--	?
SB Soft body tissue.															0.030	0.020	0.460	no	DY	no
LI Liver tissue.															0.030	0.020	0.030	no	DY	no
OC Overconcentration; Median(LastYear)/Background (=?" if missing Background)															0.030	0.020	0.460	no	--	1.5
U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?" if missing Background)															0.030	0.020	0.460	no	--	5.8
POWER Number of years to detect a 10% trend/year with 90% power.															0.030	0.020	0.460	no	--	>25

Annual Median Concentrations of Z N (ppm).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	OC	TRND	U95+3	POMER	A N A L Y S E S
30A MYTI EDU	SB D.Wt				137.80	90.50	140.30	120.00	93.10	76.19	160.90	115.98	147.14	104.32	117.30	108.77	114.00	no	--	no	11	
31A MYTI EDU	SB D.Wt	88.14	132.41	76.90	106.00	66.30	67.70	58.05	67.70	58.05	180.84	127.88	124.87	96.43	96.80	151.20	103.21	no	--	1.4	12	
35A MYTI EDU	SB D.Wt	91.89	79.63	75.90	89.80	68.40	81.50	83.19	81.50	83.19	166.05	139.18	131.29	118.54	97.56	82.88	94.26	no	UY	no	8	
36A MYTI EDU	SB D.Wt	66.50	85.78	66.10	57.70	61.50	73.60	65.27	73.60	65.27	125.69	126.98	104.33	83.96	121.15	115.35	137.02	no	UY	1.2	9	
71A MYTI EDU	SB D.Wt	123.61	125.00	77.04	114.93	101.00	169.00	128.18	161.81	143.03	165.68	120.22	157.05	150.00	122.16	112.44	121.44	no	--	no	10	
76A MYTI EDU	SB D.Wt					158.16	127.27	124.35	71.43	88.95	62.61	231.71	1.2	1.2	4.6	16		no	--	4.6	16	
15A MYTI EDU	SB D.Wt					156.56	143.70	171.96	162.03	135.00	115.72	98.02	143.94	220.65	1.1	1.1	2.3	8	UY	2.3	8	
22A MYTI EDU	SB D.Wt					378.00	253.00								1.1	1.1	2.2	13	--	2.4	14	
51A MYTI EDU	SB D.Wt														1.2	1.2	2.7	14	--	2.7	14	
52A MYTI EDU	SB D.Wt														no	no	11	D-	no	2.1	12	
56A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
57A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
63A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
65A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
69A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
82A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
84A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
87A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
91A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
92A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
98X MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
98X MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
41A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
44A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
46A MYTI EDU	SB D.Wt														no	no	11	D-	no	1.6	11	
30B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
368 GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
15B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
23B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
53B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
67B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
92B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
98B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
43B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
10B GADU MOR	LI W.Wt														no	no	11	D-	no	1.6	11	
67B LEPI WHI	LI W.Wt														no	no	11	D-	no	1.6	11	
36F LIMA LIM	LI W.Wt														no	no	11	D-	no	1.6	11	
15F LIMA LIM	LI W.Wt														no	no	11	D-	no	1.6	11	
22F LIMA LIM	LI W.Wt														no	no	11	D-	no	1.6	11	
98F LIMA LIM	LI W.Wt														no	no	11	D-	no	1.6	11	
33B PLAT FLE	LI W.Wt														no	no	11	D-	no	1.6	11	
53B PLAT FLE	LI W.Wt														no	no	11	D-	no	1.6	11	

SB Soft body tissue.
 LI Liver tissue.
 OC Overconcentration; Median(LastYear)/Background (=1?) if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+5years)/Background (=1?) if missing Background or N(years)<=5)
 POMER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of C B 1 5 3 (ppb).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S	OC	TRND	U95+3	POWER	
30A	MYTI EDU											19.53	24.37	10.86	20.87	19.65	26.36	5.3	---	---	5.3	15.3	12
31A	MYTI EDU											3.66	6.59	3.30	11.23	9.41	10.88	2.2	---	---	2.2	10.2	15
32A	MYTI EDU											2.92	5.52	3.03	3.76	3.69	3.07	no	---	---	no	1.2	12
36A	MYTI EDU											4.36	3.54	1.59	1.59	3.62	3.72	no	---	---	no	3.7	13
71A	MYTI EDU											5.87	4.59	3.35	5.45	3.02	7.24	1.4	---	---	1.4	3.9	16
76A	MYTI EDU											4.62	2.49	1.18	2.00	0.90	4.21	no	---	---	no	?	17
15A	MYTI EDU											2.84	0.98	0.98	2.00	0.90	0.79	no	---	---	no	?	17
22A	MYTI EDU											2.96	2.78	1.31	2.33	2.12	1.94	no	---	---	no	?	17
52A	MYTI EDU											3.47	3.77	3.62	5.00	5.32	2.62	no	---	---	no	1.9	17
56A	MYTI EDU											5.57	2.80	2.08	4.00	3.97	3.09	no	---	---	no	2.6	17
57A	MYTI EDU											3.82	1.44	1.59	2.69	2.04	1.81	no	---	---	no	?	17
63A	MYTI EDU											4.85	1.48	1.23	1.77	1.91	1.40	no	---	---	no	?	17
65A	MYTI EDU											4.31	1.17	0.74	1.69	1.30	1.83	no	---	---	no	?	17
69A	MYTI EDU											4.45	1.17	1.58	1.66	1.85	0.93	no	---	---	no	?	17
84A	MYTI EDU											2.53	3.13	1.70	1.82	1.43	2.40	no	---	---	no	?	17
92A	MYTI EDU											1.27	1.27	0.45	27.42	1.17	0.89	no	---	---	no	?	17
98X	MYTI EDU														0.99	25.29	14.64	no	---	---	no	?	17
41A	MYTI EDU														1.68	0.84	0.73	no	---	---	no	?	17
46A	MYTI EDU														1.82	1.28	0.98	no	---	---	no	?	17
30B	GADU MOR											955.00	957.39	856.41	885.00	807.00	871.90	no	---	---	no	?	17
36B	GADU MOR											126.00	147.00	254.00	138.00	138.00	350.00	no	---	---	no	?	17
13B	GADU MOR											164.86	99.00	66.50	97.00	117.73	112.00	no	---	---	no	?	17
23B	GADU MOR											60.00	86.00	79.00	48.00	75.00	73.00	no	---	---	no	?	17
53B	GADU MOR											104.44	86.00	65.00	48.00	61.00	262.44	no	---	---	no	?	17
67B	GADU MOR											156.00	105.76	83.50	128.45	85.00	92.00	no	---	---	no	?	17
92B	GADU MOR											111.00	105.76	51.00	50.99	102.00	129.00	no	---	---	no	?	17
98B	GADU MOR											106.00	65.00	72.97	33.00	55.00	80.00	no	---	---	no	?	17
43B	GADU MOR														110.00	113.00	50.00	no	---	---	no	?	17
10B	GADU MOR														201.00	171.00	66.00	no	---	---	no	?	17
30B	GADU MOR														1.23	0.89	2.54	no	---	---	no	?	17
36B	GADU MOR											3.70	9.20	3.30	1.23	0.89	2.54	no	---	---	no	?	17
15B	GADU MOR											0.63	0.70	1.30	0.15	5.42	1.37	no	---	---	no	?	17
23B	GADU MOR											0.52	0.49	0.20	0.13	0.36	0.43	no	---	---	no	?	17
53B	GADU MOR											0.26	0.20	0.20	0.05	0.27	0.16	no	---	---	no	?	17
67B	GADU MOR											3.36	4.79	0.40	0.47	0.12	6.43	no	---	---	no	?	17
92B	GADU MOR											0.24	0.40	0.14	0.06	0.19	1.23	no	---	---	no	?	17
98B	GADU MOR														0.03	0.08	0.12	no	---	---	no	?	17
43B	GADU MOR											0.20	0.20	0.10	0.14	0.21	0.13	no	---	---	no	?	17
10B	GADU MOR														0.55	0.85	0.11	no	---	---	no	?	17
30B	GADU MOR														87.00	79.00	38.80	no	---	---	no	?	17
36B	GADU MOR											37.00	61.00	45.00	87.00	79.00	38.80	no	---	---	no	?	17
15F	LIMA LIM											0.27	0.60	0.10	0.17	0.63	0.17	no	---	---	no	?	17
22F	LIMA LIM											129.48	138.00	171.00	148.00	161.00	150.80	no	---	---	no	?	17
36F	LIMA LIM											53.00	52.44	21.54	28.93	28.00	22.80	no	---	---	no	?	17
15F	LIMA LIM											65.41	48.96	3.20	24.00	29.60	22.80	no	---	---	no	?	17
22F	LIMA LIM											1.13	2.30	0.24	2.06	3.22	3.33	no	---	---	no	?	17
98F	LIMA LIM											1.62	1.28	0.24	0.12	0.41	0.69	no	---	---	no	?	17
98F	LIMA LIM											2.00	1.28	0.24	0.50	1.66	0.69	no	---	---	no	?	17
33B	PLAT FLE											11.00	30.00	22.00	17.00	17.00	9.40	no	---	---	no	?	17
53B	PLAT FLE											123.00	80.00	10.00	17.00	17.00	9.40	no	---	---	no	?	17
53B	PLAT FLE											0.65	0.50	0.30	0.27	0.44	0.69	no	---	---	no	?	17
53B	PLAT FLE											5.88	3.40	0.40	0.12	0.22	0.93	no	---	---	no	?	17

SB Soft body tissue.
 LI Liver tissue.
 MU Muscle tissue.
 OC Overconcentration; Median(LastYear)/Background (=?) if missing Background
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?) if missing Background or N(years)<=5
 POWER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of DDE P P (ppb).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S OC TRND U95+3 POWER				
30A	MYTI EDU												5.24	3.86	7.08	5.70	2.56	no	-?	?	13	
31A	MYTI EDU												3.30	1.89	3.45	1.84	0.50	no	-?	?	?	17
35A	MYTI EDU												4.91	2.08	3.13	2.84	0.57	no	-?	?	?	19
36A	MYTI EDU												2.76	1.06	1.03	1.76	0.44	no	-?	?	?	19
71A	MYTI EDU												2.61	1.58	3.21	1.29	0.74	no	-?	?	?	15
76A	MYTI EDU												1.40	0.79			0.36	no	-?	?	?	14
15A	MYTI EDU																0.73	no	-?	?	?	15
22A	MYTI EDU												2.22	1.31	1.88	1.45	0.39	no	-?	?	?	16
52A	MYTI EDU												12.26	25.48	19.43	18.48	9.53	no	-?	?	?	13
56A	MYTI EDU												50.00	47.53	114.57	40.84	33.89	3.4	-?	?	?	15
57A	MYTI EDU												25.90	18.25	34.97	25.31	15.80	1.6	-?	?	?	12
63A	MYTI EDU												12.94	9.29	9.68	8.36	5.53	no	-?	?	?	9
65A	MYTI EDU												7.60	5.19	7.79	4.12	5.00	no	-?	?	?	12
69A	MYTI EDU												3.55	3.16	3.54	2.91	0.40	no	-?	?	?	19
84A	MYTI EDU												3.13	2.23		0.99	0.74	no	-?	?	?	10
92A	MYTI EDU												0.68	2.09	1.41	0.77	0.28	no	-?	?	?	17
98X	MYTI EDU														31.65	22.87	5.16	no	-?	?	?	19
41A	MYTI EDU														0.62	0.42	0.29	no	-?	?	?	10
46A	MYTI EDU														1.05	0.76	0.27	no	-?	?	?	15
30B	GADU MOR												182.46	158.97	191.00	194.00	312.50	1.6	--	5.6	11	
36B	GADU MOR												91.90	51.00	55.00	105.00	141.00	no	--	2.4	15	
15B	GADU MOR												50.00	135.99	86.00	33.47	75.00	no	--	no	18	
23B	GADU MOR												68.00	85.38	42.00	31.00	49.00	no	--	no	10	
53B	GADU MOR												637.00	805.84	939.42	85.00	42.00	491.39	2.5	--	7.9	24
67B	GADU MOR												776.00	554.00	471.14	391.80	109.00	460.00	2.3	--	4.1	19
92B	GADU MOR													53.00	50.50	50.00	196.00	no	-?	?	?	17
98B	GADU MOR													83.40	43.00	48.96	138.00	no	-?	?	?	14
43B	GADU MOR												73.00		126.00	69.00	60.00	no	-?	?	?	11
10B	GADU MOR														211.00	71.00	75.00	no	-?	?	?	16
67B	LEPI WHI												294.00	240.00	250.00	163.00	142.60	no	-?	?	?	10
36F	LIMA LIM												27.98	34.41	50.00	40.00	40.00	no	--	no	11	
15F	LIMA LIM												39.00	13.42	23.49	9.00	20.70	no	--	no	?	19
22F	LIMA LIM												68.93	47.96	21.00	9.17	7.00	no	-?	?	?	11
33B	PLAT FLE												13.00	9.10	13.00	7.00	10.20	no	-?	?	?	14
53B	PLAT FLE												94.00	70.14	41.00	8.00	25.00	no	--	no	1.4	19

SB Soft body tissue.
 LI Liver tissue.
 OC Overconcentration; Median(LastYear)/Background (=11?) if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=11?) if missing Background or N(years)<=5)
 POWER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of H C H G (ppb).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S	OC	TRND	U95+3	POMER	
30A	MYTI EDU																						13
	SB D.WT																						no
31A	MYTI EDU																						11
	SB D.WT																						no
35A	MYTI EDU																						10
	SB D.WT																						no
36A	MYTI EDU																						14
	SB D.WT																						no
71A	MYTI EDU																						15
	SB D.WT																						no
76A	MYTI EDU																						18
	SB D.WT																						?
15A	MYTI EDU																						14
	SB D.WT																						?
22A	MYTI EDU																						12
	SB D.WT																						no
52A	MYTI EDU																						1,2
	SB D.WT																						1,2
56A	MYTI EDU																						10
	SB D.WT																						no
57A	MYTI EDU																						12
	SB D.WT																						no
63A	MYTI EDU																						12
	SB D.WT																						no
65A	MYTI EDU																						17
	SB D.WT																						no
69A	MYTI EDU																						10
	SB D.WT																						?
84A	MYTI EDU																						14
	SB D.WT																						?
92A	MYTI EDU																						10
	SB D.WT																						?
98X	MYTI EDU																						13
	SB D.WT																						?
41A	MYTI EDU																						11
	SB D.WT																						?
46A	MYTI EDU																						9
	SB D.WT																						?
30B	GADU MOR																						19
	LI W.WT																						no
36B	GADU MOR																						21
	LI W.WT																						no
15B	GADU MOR																						20
	LI W.WT																						no
23B	GADU MOR																						17
	LI W.WT																						no
53B	GADU MOR																						9
	LI W.WT																						no
67B	GADU MOR																						12
	LI W.WT																						no
92B	GADU MOR																						10
	LI W.WT																						?
98B	GADU MOR																						17
	LI W.WT																						no
43B	GADU MOR																						8
	LI W.WT																						?
10B	GADU MOR																						8
	LI W.WT																						?
67B	LEPI WHI																						18
	LI W.WT																						?
36F	LIMA LIM																						21
	LI W.WT																						no
15F	LIMA LIM																						10
	LI W.WT																						?
22F	LIMA LIM																						18
	LI W.WT																						?
33B	PLAT FLE																						23
	LI W.WT																						no
53B	PLAT FLE																						15
	LI W.WT																						no

SB Soft body tissue.
 LI Liver tissue.
 OC Overconcentration; Median(LastYear)/Background (=?" if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=?" if missing Background or N(years)<=5)
 POMER Number of years to detect a 10% trend/year with 90% power.

Annual Median Concentrations of H C B (ppb).

St. Species	Tissue Base	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	A N A L Y S E S	OC	TRND	U95+3	POWER
30A	MYTI EDU				1.18	0.88	2.06	0.92	1.15	0.87	0.35	0.59	0.95	0.54	0.27	0.24	0.25	no	D-	no	D-	13
31A	MYTI EDU		13.37		1.38	3.83	1.89	0.93	0.89	0.36	0.32	0.61	0.55	0.45	0.24	0.31	0.22	no	D-	no	D-	18
35A	MYTI EDU		12.83		0.95	3.33	0.79	0.98	1.12	0.47	0.42	0.58	0.58	0.51	0.23	0.28	0.22	no	D-	no	D-	20
36A	MYTI EDU		15.02		0.95	3.83	2.90	2.37	0.96	0.43	0.33	0.55	0.39	0.53	0.24	0.33	0.28	no	D-	no	D-	20
71A	MYTI EDU		15.25		10.37	91.37	11.11	206.90	1.84	149.17	8.48	6.92	4.14	3.91	1.47	2.13	4.48	9.0	--	320.0	>25	
76A	MYTI EDU										0.38	0.57	0.50	0.79			0.25	no	-?	?	?	16
15A	MYTI EDU										0.20			0.49	0.25	0.22	0.29	no	-?	?	?	14
22A	MYTI EDU										0.26	0.61	0.56	0.44	0.25	0.25	0.30	no	--	no	--	11
52A	MYTI EDU									0.85	0.38		0.81	0.81	0.28	0.32	0.26	no	--	no	--	15
56A	MYTI EDU						0.20			0.79	0.41	0.79	0.93	1.04	0.31	0.38	0.31	no	--	1.2	16	
57A	MYTI EDU									0.77		0.76	0.72	0.79	0.36	0.30	0.26	no	DY	no	DY	9
63A	MYTI EDU									1.05		0.97	0.74	0.63	0.32	0.33	0.33	no	DY	no	DY	8
65A	MYTI EDU									0.43	0.52	0.86	0.62	0.67	0.28	0.30	0.29	no	DY	no	DY	12
69A	MYTI EDU												0.53	0.53	0.29	0.25	0.29	no	D?	?	?	8
82A	MYTI EDU				2.26	10.73	0.66	0.62	0.80	0.53			0.63	0.53			0.21	1.1	--	14.0	24	
84A	MYTI EDU				3.41	8.79	3.33	2.04	1.23	0.48			0.68	0.42			0.23	no	D-	no	D-	13
92A	MYTI EDU														0.24		0.23	no	D?	?	?	9
98X	MYTI EDU														0.56		0.32	no	--	?	?	11
41A	MYTI EDU														0.29		0.26	no	--	?	?	6
46A	MYTI EDU														0.26		0.29	no	--	?	?	6
30B	GADU MOR										10.00	17.00	7.48	16.00	11.00	11.00	12.00	no	--	1.3	14	
36B	GADU MOR										7.00	9.00	9.00	10.00	9.00	5.00	9.00	no	--	no	--	11
15B	GADU MOR										5.00	20.49	10.00	14.00	14.00	9.00	11.00	no	--	1.3	17	
23B	GADU MOR										6.00	9.49	12.00	9.00	8.00	6.00	10.00	no	--	no	--	10
53B	GADU MOR										10.00	10.00	16.49	7.00	8.00	5.00	7.00	no	--	no	--	12
67B	GADU MOR										14.00	8.00	7.94	8.00	8.49	10.00	8.00	no	--	no	--	9
92B	GADU MOR												20.00	17.00	11.00	14.00	13.00	no	--	?	?	10
98B	GADU MOR													9.95	12.00	18.00	35.00	1.8	-?	?	?	13
43B	GADU MOR														15.00	16.49	13.00	no	--	?	?	8
10B	GADU MOR														13.00	11.00	16.00	no	--	?	?	10
67B	LEPI WHI														5.00	2.00	4.60	?	--	?	?	16
36F	LIMA LIM										9.00	4.00	5.00	4.00	3.00	2.00	2.30	no	--	no	--	14
15F	LIMA LIM										5.48	3.00	5.00	2.00	3.00	2.00	3.00	no	--	no	--	11
22F	LIMA LIM										6.00	3.00	5.00	4.00	4.00	2.00	3.00	no	--	no	--	19
33B	PLAT FLE										1.00	0.50	5.00	2.00	1.00	1.41	0.60	no	--	?	?	11
53B	PLAT FLE										6.00	4.47	5.00	2.00	1.00	1.00	2.00	no	--	no	--	20
																		no	D-	no	D-	13

SB Soft body tissue.
 LI Liver tissue.
 OC Overconcentration; Median(LastYear)/Background (=!?) if missing Background)
 U95+3 Upper 95% ConfidenceInterval(Last+3years)/Background (=!?) if missing Background or N(years)<=5)
 POWER Number of years to detect a 10% trend/year with 90% power.

Appendix H. Geographical distribution of contaminants in biota 1995-96

Sorted by contaminant and species:


Cadmium (Cd)
Copper (Cu)
Mercury (Hg)
Lead (Pb)
Zinc (Zn)
Sum of 7 CBs (CB-28, -52, 101, -118, -138, -153 and -180)
CB-153
DDEPP (ppDDE)
γ-HCH
HCB
PAH
B[a]P
TCDDN
TECBW

MYTI EDU - Blue Mussel (*Mytilus edulis*)
GADU MOR - Atlantic cod (*Gadus morhua*)
PLAT FLE - Flounder (*Platichthys flesus*)
LIMA LIM - Dab (*Limanda limanda*)
PLEU PLA - Plaice (*Pleuronectes platessa*)
MICR KIT - Lemon sole (*Microstomus kitt*)
LEPI WHI - Megril (*Lepidorhombus whiff-iaconis*)
GLYP CYN - Witch (*Glyptocephalus cynoglossus*)
BROS BRO - Torsk (*Brosme brosme*)

Station positions are shown on maps (Figure 1 to Figure 4)

Appendix H
Geographical distribution of contaminants in biota 1995-96
(cont.)

A



B





Figure 15. Median, standard deviation and provisional "high background" concentration for cadmium in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1- Figure 4).

A



B





Figure 16. Median, standard deviation and provisional "high background" concentration for cadmium in fish liver 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 17. Median standard deviation and provisional "high background" concentration for copper in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 18. Median, standard deviation and provisional "high background" concentration for copper in fish liver 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 19. Median, standard deviation and provisional "high background" concentration for mercury in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 20. Median, standard deviation and provisional "high background" concentration for mercury in fish fillet 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 21. Median, standard deviation and provisional "high background" concentration for lead in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 22. Median, standard deviation and provisional "high background" concentration for lead in fish liver 1995 (**A**) and 1996 (**B**), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 23. Median, standard deviation and provisional "high background" concentration for zinc in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 24. Median, standard deviation and provisional "high background" concentration for zinc in fish liver 1995 (A) and 1996 (B), ppm wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 25. Median, standard deviation and provisional "high background" concentration for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 26. Median, standard deviation and provisional "high background" concentration for sum of 7 PCBs (CB-28, -52, 101, -118, -138, -153 and -180) in fish liver 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 27. Median, standard deviation and provisional "high background" concentration for CB-153 in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 28. Median, standard deviation and provisional "high background" concentration for CB-153 in fish liver 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).

A



B





Figure 29. Median, standard deviation and provisional "high background" concentration for ppDDE (DDEPP) in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1- Figure 4). (See also footnote in Table 3.)

A



B





Figure 30. Median, standard deviation and provisional "high background" concentration for ppDDE (DDEPP) in fish liver 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4). (See also footnote in Table 3.)

A



B





Figure 31. Median, standard deviation and provisional "high background" concentration for γ -HCH (Lindan) in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1- Figure 4). (See also footnote in Table 3.)

A



B





Figure 32. Median, standard deviation and provisional "high background" concentration for γ -HCH (Lindan) in fish liver 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4). (See also footnote in Table 3.)

A



B





Figure 33. Median, standard deviation and provisional "high background" concentration for HCB in mussels (*Mytilus edulis*) 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).

A



B




Figure 34. Median, standard deviation and provisional "high background" concentration for HCB in fish liver 1995 (A) and 1996 (B), ppb wet weight (see maps in Figure 1-Figure 4).





Figure 35. Median, standard deviation and provisional "high background" concentration for PAH (excluding dicyclics) in mussels 1995 (**A**) and 1996 (**B**) ppb wet weight (see maps in Figure 1 and Figure 4).

A



B





Figure 36. Median, standard deviation and provisional "high background" concentration for B[a]P (BAP) in mussels 1995 (**A**) and 1996 (**B**) ppb wet weight (see maps in Figure 1 and Figure 4).

A



B





Figure 37. Measured and provisional "high background" concentration for sum of TCDD-toxicity equivalents after Nordic model (TCDDN) in mussel 1995 (**A**) and 1996 (**B**), **NB! ppp (nanogram/kg)** wet weight (see maps in Figure 1 and Figure 4).

A



B




Figure 38. Measured and provisional "high background" concentration for sum of TECB-toxicity equivalents after WHO model (TECBW) in mussel 1995 (**A**) and 1996 (**B**), **NB!** ppp (nanogram/kg) wet weight (see maps in Figure 1 and Figure 4).

Appendix I.

Results from INDEX determinations 1995-96

Introduction

The Norwegian State Pollution Control Authority (SFT) is interested in obtaining a small group of indices to assess the quality of the environment with respect to contaminants. One index is based on the levels and trends of contaminant concentrations in the blue mussel collected annually from 10 of the more contaminated fjords in Norway (Walday *et al.* 1995), herein designated "Pollution Index". JAMP results from Orkdalsfjorden were also included in the calculation of this index (SFT, pers. comm.). This index was to be tested in 1995-96. It was practical to organise sampling within JAMP, which included monitoring of mussels from in or near the 11 fjords. Some JAMP results could be used to calculate the index value. Also, mussels from one station in Trondheim harbour were collected and analysed but were not included in this initial evaluation of the index.

In addition, a "Reference Index" was tested in 1995-96 based on contaminant concentrations in the blue mussel found along the entire coast in areas with presumably low levels of contamination. JAMP stations sampled in 1996 were used for the most part. The importance of "reference" stations for monitoring of contaminants has been discussed earlier (cf., Green 1987). One of the main reasons for this work is to establish points of reference for contaminated fjords that may fall under the criteria for Areas of Special Concern (JMG 1992). This is also of national and international interest.

The target medium for both indices may vary depending on the purpose, however sediment, cod and mussels are considered to be the most likely choices. Blue mussels were selected for the 1995-96 investigation (Appendix I1).

Calculation of the index

A detailed discussion of calculation of the Pollution Index has been given earlier (cf. Walday *et al.* 1995) and only a brief summary will be given here. The relevant contaminants for each of the Pollution Index fjords (including Orksalsfjord) are summarised in Appendices I2 and I3. Their selection is based on earlier investigations. Two to five stations were sampled from each area. Three parallels of 20 individuals from 3-5cm are collected from each station. Each sample was analysed for the contaminants according to the scheme in Appendix I3. Due to expense only one or two samples per fjord were analysed for dioxins where this contaminant group was relevant.

One to three stations were sampled from selected areas for the determination of the Reference Index. Each station included three parallels which were analysed for the usual JAMP contaminants (cf., analysis code A, Appendix I3). Some samples were also analysed for PAHs and dioxins.

Mussel sampling differed from (A) stations that were exclusively to be used for index calculations and (B) those included in the JAMP for 1995-96 in that the former allowed a greater size range and that the mussels were not depurated.

The maximum median for each contaminant for all the stations in an area was determined. These concentrations were classified according to SFT's classification system for contaminants in the marine environment (Appendix I4). The highest class found for any contaminant measured in an area determined the index value for that area.

The classes are roughly and not systematically based on the provisional "high background" levels. This system has been recently revised (Molvær *et al.* 1997); where among other changes the sum of

CB-28, -52, -101, -118, -138, -153, and -180 (CBΣΣe) is now a distinct parameter for classification. The sum of all PAHs (including the dicyclic) (PAHΣΣ) was compared to the system's "sum-PAH". "Dioxins" were assessed based on toxicity equivalency factors (TEQ) according to a Nordic model (Ahlborg 1989). Note that EPOCl is considered a relevant contaminant for one area but is not included in the part of the classification system based on levels in mussels. Likewise, there are contaminants which are included in the classification system but have not been measured in any area.

The maximum class found for any contaminant determined the class (I-V) of the area. The average class for all the contaminated sub areas and all the reference localities determined the Pollution or Reference Index, respectively. The lowest Index value is 1 and means that all median values were in Class I ("Good"). The highest Index value is 5 and means that all median values were in Class V ("Very bad").

Conclusion from first application of the indices

The Pollution Index was recalculated for 1995 because of the revisions made in SFTs classification system, which involved lower limits for *inter alia* BAP and HCB. The Index for 1995 is 3.2 compared to 2.8 based on the old system (Appendices I5). The Index for 1996 is 3.1. A value between 3 and 4 would be classified by the SFT system as "Bad".

The Reference Index for 1995 is 1.6 and unchanged compared to the old system. The Index for 1996 is 1.8 (Appendices I6). The calculation included (low) suspect dioxin values (TCDDN). A value between 1 and 2 would be classified as "Fair".

There may be need to review the selection of areas/stations and contaminants for the Pollution Index (cf. Walday *et al.* 1995) as well as for the Reference Index. The anticipated remedial actions should eventually cause the index to decrease, hence, the initial "Pollution Index" could be closer to the upper end of the scale (5) than where it now (close to and below the middle). With the given analytical scheme, two of the eleven "polluted" areas were classified as "good" (Iddefjord and Orkdalsfjord), and this was the situation for both years 1995-96. Removing these fjords from the calculation would increase the index. Another means of raising the index is by changing the classification system, which was recently done (cf., Molvær *et al.* 1997).

Only two of the eight "reference" areas are classified as "good". The remaining stations are classified as "fair" and show that the majority are more polluted. One reason could be that the stations were selected from the JAMP program and are not necessarily representative of "diffusely" polluted areas along the entire coast of Norway.

Appendix I1.

INDEX - Stations and programme 1995-96

Appendix I1. INDEX station positions and sampling overview for blue mussels 1995-96, where P = "Pollution Index" and R = "Reference Index" (contaminated and assumed "background" stations, respectively). Mussels were sampled from rock surfaces unless otherwise noted. See Walday *et al.* (1995) for discussion of station selection and analyses.

Station	Locality name	North latitude	East longitude	ICES position	INDEX type P/R	notes
HVALER/SINGLEFJORDEN, east of outer OSLOFJORD						
I021	Kjøkkø, south	59°07.8'	10°57.1'	47G13	P	
I024	Kirkø, north west	59°04.9'	10°59.2'	47G09	P	
I022	West Damholmen	59°06.2'	10°57.9'	47G09	P	
I023	Kirkø, north west	59°05.7'	11°08.2'	47G09	P	
IDDEFJORD, east of outer OSLOFJORD						
I001	Sponvikskansen	59°05.4'	11°12.5'	47G09	P	
I011	Kråkenebbet	59°06.1'	11°17.3'	47G09	P	
INNER OSLOFJORD						
JAMP 30A	Gressholmen	59°52.5'	10°43.0'	48G07	P	
I301	Akershuskaia	59°54.2'	10°45.5'	48G07	P	
I304	Gåsøya	59°51.0'	10°35.5'	48G04	P	
I307	Ramtonholmen	59°44.7'	10°31.4'	48G05	P	
I306	Håøya	59°24.7'	10°33.4'	48G04	P	
MID and OUTER OSLOFJORD						
JAMP 31A	Solbergstrand	59°36.9'	10°39.4'	48G06	R	
JAMP 35A	Mølen	59°29.2'	10°30.1'	47G04	R	
JAMP 36A	Færder	59°01.6'	10°31.7'	47G06	R	
FRIERFJORD AREA, west of outer Oslofjord						
I711	Steinholmen	59°31.7'	09°40.7'	48F99	P	
I712	Gjemesholmen	59°21.7'	09°42.6'	47F99	P	
JAMP 71A	Bjerkøya (Risøyodden)	59°01.4'	09°45.4'	47F99	P	
INNER KRISTIANSANDSFJORD						
I132	Fiskåtangen	57°07.7'	07°59.2'	43F79	P	
I133	Odderø, west	57°07.9'	08°00.3'	43F83	P	
LISTA AREA						
JAMP 15A	Gåsøy (Ullerø area)	58°03.1'	06°53.3'	45F69	R	
I131	Lastad	58°03.3'	07°42.4'	45F79	R	
SAUDAFJORD						
I201	Ekkjegrunn (G1)	59°38.7'	06°21.4'	48F66	P	
**	I205 Bølsnes (G5)	59°35.5'	06°18.3'	48F63	P	
BØMLO AREA						
JAMP 22A	Espevær, west	59°35.2'	05°58.5'	48F59	R	C, 1
SØRFJORDEN						
*	51A Byrkjenes	60°05.1'	06°33.1'	49F66	P	
JAMP 52A	Eitrheimsneset	60°05.8'	06°32.2'	49F66	P	3

Appendix I1 (cont'd)

Station	Locality name	North latitude	East longitude	ICES position	INDEX type P/R	notes
BYFJORDEN, Bergen						
I242	Valheimneset	60°23.7'	05°16.1'	49F51	P	
I241	Nordnes	60°24.1'	05°18.2'	49F51	P	
I243	Hagreneset	60°24.9'	05°18.3'	49F51	P	
SUNNDALSFJORDen						
I912	Honnhammer	62°51.2'	08°09.7'	54F81	P	
I911	Horvika	62°44.1'	08°31.4'	54F85	P	
[TRONDHEIM AREA - not related to INDEX investigation]						
* 80A	Østmerknes	63°27.5'	10°27.5'	56G04	P	
ORKDALSFJORD AREA, supplementary area (cf. Walday <i>et al.</i> 1995)						
JAMP 82A	Flakk	63°27.1'	10°12.6'	56G01	P	
JAMP 84A	Trossavika	63°20.8'	09°57.8'	55F97	P	
JAMP 87A	Ingdalsbukta	63°27.8'	09°54.8'	55F97	P	
INNER RANFJORD						
I969	Bjørnbærviken (B9)	66°16.8'	14°02.1'	61G42	P	
I962	Koksverkkaien (B2)	66°19.4'	14°08.0'	61G42	P	3
OUTER RANFJORD, Helgeland area						
* 96A	Breiviken	66°17.6'	12°50.5'	61G28	R	1
LOFOTEN AREA						
JAMP 98X	Skrova	68°10.5'	14°40.2'	65G48	R	
FINNSNES-SKJERVØY AREA						
JAMP 41A	Fensneset, Grytøya	68°56.9'	16°38.5'	66G64	R	3
HAMMERFEST-HONNINGSVÅG AREA						
JAMP 44A	Elenheimsundet	70°30.8'	22°14.8'	70H23	R	1, 6
JAMP 46A	Smineset in Altesula	70°58.4'	25°48.1'	70H57	R	3, 6
VARANGER PENINSULA AREA						
JAMP 48A	Trollfjorden i Tanafjord	70°41.6'	28°33.3'	70H85	R	
JAMP 10A	Skagoodden	70°04.2'	30°09.8'	69J03	R	2
JAMP 11A	Sildkroneset, Bøkfjorden	69°47.0'	30°11.1'	68J02	R	

notes:

* - JAMP station but not sampled in accordance to JAMP guidelines, see appendix text.

** - Sufficient mussel-sample not found in 1996.

1 - mussels collected from buoy and/or buoy anchor lines

2 - mussels collected from sand/gravel bottom

3 - mussels collected from iron/cement pilings

4 - mussels collected from metal navigation buoys

5 - mussels collected from floating dock

6 - mussels collected from wooden docks

Appendix I2.

INDEX - Sampling and analyses for 1995-96

Appendix I2. Blue mussel samples used in INDEX 1995-96, where P = "Pollution Index" and R = "Reference Index" (contaminated and assumed "background" stations, respectively). + indicates JAMP sampling and analyses (i.e. equivalent to analysis code A). The number indicates the number samples analyzed. Codes for analysis (A, B etc.) are defined in Appendix I3. See Walday *et al.* (1995) for discussion of selection of stations and analyses.

JAMP st.	STATION	INDEX	ANALYSIS CODE													
			+	A	B	C	D	E	F	G	H	I	J			
HVALER/SINGLEFJORD AREA																
021	Kjøkkø, south	P	3
024	Kirøy, north west	P	3
022	West Damholmen	P	3
023	Singlekalven, south	P	3
IDDEFJORD																
01A	Sponvikskansen	P	3
011	Kråkenebbet	P	3
OSLOFJORD, inner																
30A	Gressholmen	P	+	3	.	.1
301	Akershuskaia	P	3
304	Gåsøya	P	3
307	Ramtonholmen	P	3
306	Håøya	P	3
OSLOFJORD, mid and outer																
31A	Solbergstrand	R	+	1
35A	Mølen	R	+
36A	Færder	R	+	2*
FRIERFJORD AREA, west of outer Oslofjord																
711	Steinholmen	P	3	1
712	Gjemesholmen	P	3	1
71A	Bjørkøya	P	+	1
INNER KRISTRIANSANDSFJORD																
132	Fiskåtangen	P	3	1
133	Odderø, west	P	3	1
LISTA AREA																
15A	Gåsøya	R	+	1
131	Lastad	R	3.
SAUDAFJORD																
201	Ekkjegrunn (G1)	P	3
205	Bølsnes (G5)	P	3
BØMLO-SOTRA AREA																
22A	Espevær, west	R	+	2*
SØRFJORDEN																
51A	Byrkjeneset	P	3
52A	Eirtheimsneset	P	+

*) indicates Toxaphene included

Appendix I2 (cont'd)

JAMP st.	STATION	INDEX	ANALYSIS CODE											
			+	A	B	C	D	E	F	G	H	I	J	
BYFJORDEN, BERGEN														
242	Valheimsneset	P	3	.	.	.
241	Nordnes	P	3	.	.	.
243	Hagreneset	P	3	.	.	.
SUNNDALSFJORD														
912	Honnhammer	P	3	.	.
911	Horvika	P	3	.	.
[TRONDHEIM AREA - not related to index investigation]														
80A	Østmarknes	-	3	.
ORKDALSFJORD AREA (not suggested in Walday <i>et al.</i> 1993)														
82A	Flakk	P	+
84A	Trossavika	P	+
87A	Ingdalsbukta	P	+
INNER RANFJORD														
962	Koksverkkaien (B2)	P	3	.
969	Bjørnbærvikenn (B9)	P	3	.
OUTER RANFJORD, HELGELAND AREA														
96A	Breivika, Tomma	R	3
LOFOTEN AREA														
98X	Skrova	R	+	1
FINNSNES-SKJERVØY AREA														
41A	Fensneset, Grytøya	R	+	3	1
HAMMERFEST-HONNINGSVÅG AREA														
44A	Elenheimsundet	R	+	3	2*
46A	Smneset in Altesula	R	+	3	1*
VARANGER PENINSULA AREA														
48A	Trollfjorden i Tanafjord	R	+	3	.
10A	Skagoodden	R	+	3	1
11A	Sildkroneset	R	+	3	1

*) indicates Toxaphene included

Appendix I3. INDEX - Key to analysis codes and sample counts

(Used in Appendix I2.)

ANALYSIS CODES¹⁾ See Walday *et al.* (1995) for discussion of selection of analyses.

Contaminant	Analysis code												
	A	B	C	D	E	F	G	H	I	J			
Lead	X	X	.	.	.	X	.
Cadmium	X	X	X	.	.	X	.
Copper	X	X	X
Mercury	X	X	X
Zinc	X	X	X	.	.	X	.
EPOCI	X
PAHs	X	X	X	X	X
PCBs	X	.	X	X	.	.	.
"Dioxin"	X

¹⁾ Concerns MUSSEL 1 size group (3-5cm), 3 parallel samples each a bulk of 20 individuals (see text)

Appendix I4. INDEX - SFT Environment classes

(Molvær *et al.* 1997)

As	arsenic
Åb	lead
F	fluoride
Cd	cadmium
Cu	copper
Cr	chromium
Hg	mercury
Ni	nickel
Zn	zinc
Ag	silver
PAHΣΣ	total PAH including dicyclic *
BAP	benzo[<i>a</i>]pyrene
DDTΣΣ	DDTPP+DDEPP+TDEPP *
HCB	hexachlorobenzene
HCHΣΣ	HCHG+HCHA+HCHB *
CBΣΣe	sum of CB: 28+52+101+118+138+153+180 *
TCDDN	Sum of TCDD-toxicity equivalents *

*) see also Appendix B

15/10-97

LIMIT-CHECK-file; I:\TPX\JMG\LIM\NI970923.ISH

CLASS-limits for M Y T I E D U (Mytilus edulis, GB: Blue mussel, N: Blåskjell).
Tissue : WHOLE SOFT BODY.

Limit Level=> Basis =====> Param. Unit	Class I		Class II		Class III		Class IV		Class V	
	Wet weight	Dry weight	Wet weight	Dry weight	Wet weight	Dry weight	Wet weight	Dry weight	Wet weight	Dry weight
As ppm	.	<10.0	.	<30.0	.	<100.0	.	<200.0	.	>=200.0
Pb ppm	.	<3.0	.	<15.0	.	<40.0	.	<100.0	.	>=100.0
F ppm	.	<15.0	.	<50.0	.	<150.0	.	<300.0	.	>=300.0
Cd ppm	.	<2.0	.	<5.0	.	<20.0	.	<40.0	.	>=40.0
Cu ppm	.	<10.0	.	<30.0	.	<100.0	.	<200.0	.	>=200.0
Cr ppm	.	<3.0	.	<10.0	.	<30.0	.	<60.0	.	>=60.0
Hg ppm	.	<0.2	.	<0.5	.	<1.5	.	<4.0	.	>=4.0
Ni ppm	.	<5.0	.	<20.0	.	<50.0	.	<100.0	.	>=100.0
Zn ppm	.	<200.0	.	<400.0	.	<1000.0	.	<2500.0	.	>=2500.0
Ag ppm	.	<0.3	.	<1.0	.	<2.0	.	<5.0	.	>=5.0
PAHΣ ppb	<50.0	.	<200.0	.	<2000.0	.	<5000.0	.	>=5000.0	.
BAP ppb	<1.0	.	<3.0	.	<10.0	.	<30.0	.	>=30.0	.
DDTΣ ppb	<2.0	.	<5.0	.	<10.0	.	<30.0	.	>=30.0	.
HCB ppb	<0.1	.	<0.3	.	<1.0	.	<5.0	.	>=5.0	.
HCHΣ ppb	<1.0	.	<3.0	.	<10.0	.	<30.0	.	>=30.0	.
CBΣΣe ppb	<4.0	.	<15.0	.	<40.0	.	<100.0	.	>=100.0	.
TCDDN ppp	<0.2	.	<0.5	.	<1.5	.	<3.0	.	>=3.0	.

Appendix I5.
INDEX - Summary table “Pollution index”

1995

Max (Median). Statistics om ALL AREAS: (n = INDEX-Stations measured, N = Station programmed for INDEX)

INDEX_AreaNames (PollutionAreas)	n	N	AS ppm d.wt	Pb ppm w.wt	F ppm d.wt	CD ppm w.wt	CU ppm w.wt	CR ppm d.wt	HG ppm w.wt	NI ppm d.wt	ZN ppm w.wt	AG ppm d.wt	PAH32 ppb w.wt	BAP ppb w.wt	DOT32 ppb w.wt	HCB ppb w.wt	HCH22 ppb w.wt	CB32e ppb w.wt	TCDDN ppb w.wt	Max E.C I:V
-1995																				
Hvaler/Singlefjord	4	4	i	0.20A	i	0.27A	1.71A	i	0.03B	i	17.40A	i	i	i	0.93a	0.13b	0.53a	6.73b	i	II
Iddefjord	2	2	i	0.12A	i	0.17A	1.30A	i	0.01A	i	14.70A	i	i	i	1.95a	<0.05a	0.41a	20.60c	w	I
Inner Oslofjord	5	5	i	i	i	0.20A	1.45A	i	0.01A	i	27.90A	i	<97.90b	0.80a	0.85a	0.60c	0.27a	4.74b	w	III
Frierfjord	3	3	i	i	i	i	i	i	i	i	i	i	<44.80c	15.00d	0.65a	9.55e	s1.01b	5.08b	<1.68d	III
Inner Kristiansandsfjord	2	2	i	i	i	i	i	i	i	i	i	i	<387.80c	15.00d	i	i	i	i	i	V
Saudafjord	2	2	i	0.92B	i	0.16A	i	i	i	i	37.80B	i	4.80c	15.00d	6.01c	0.06a	0.28a	2.67a	i	IV
Serfjorden	2	2	i	14.60E	i	3.61D	1.11B	i	0.15D	i	31.30A	i	162.90b	4.80c	i	i	i	i	i	V
Byfjorden, Bergen	3	3	i	i	i	i	i	i	i	i	25.00A	i	<704.90c	8.00c	i	i	i	i	i	III
Sunnalsfjord	2	2	i	i	i	i	i	i	i	i	23.50A	i	487.90c	31.00e	<0.38a	0.05a	0.28a	<0.73a	i	III
Orkdalsfjord	3	3	i	0.30A	i	0.35A	1.93A	i	0.01A	i	25.00A	i	i	i	i	i	i	i	i	I
Inner Ranfjord	2	2	i	0.56B	i	0.11A	i	i	i	i	23.50A	i	487.90c	31.00e	i	i	i	i	i	V
Count			0	6	0	7	5	0	5	0	7	0	6	6	6	6	6	6	1	11
Min				0.12A		0.11A	1.11A		0.01A		14.70A		<97.90b	0.80a	<0.38a	<0.05a	0.27a	<0.73a	<1.68d	I
Max				14.60E		3.61D	1.93B		0.15D		37.80B		<704.90c	31.00e	6.01c	9.55e	s1.01b	20.60c	<1.68d	V
Mean				2.78C		0.70C	1.50A		0.04B		25.37A		<381.03c	12.43d	<1.80a	<1.74d	s0.46a	<6.76b	<1.68d	3-2

i (11B) ! Value ignored when calculating Environment Class.
 w (2) ! Missing value. Should be included when calculating Environment Class E.C

a/A(52) > Below Class-I Limit.
 b/B(18) > Below Class-II Limit.
 c/C(15) > Below Class-III Limit.
 d/D(12) > Below Class-IV Limit.
 e/E(6) > Below Class-V Limit.

1996

Max(Median). Statistics om ALL AREAS: (n = INDEX-Stations measured, N = Station programmed for INDEX)

INDEX-AreaNames (PollutionAreas)	n	N	AS	Pb	F	CD	CU	CR	HG	NI	ZN	AG	PAHSE	BAP	D0T3E	HOB	HC12E	CB32E	TCDON	Max E-C I:V
			ppm d.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm d.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	
1996																				
Hvaler/Singletfjord	4	4	i	0.30A	i	0.28B	1.64B	i	0.05B	i	20.80A	i	i	i	<0.56a	0.12b	0.27a	4.83b	i	II
Iddefjord	2	2	i	0.15A	i	0.11A	1.47A	i	0.02A	i	15.90A	i	i	3.30c	1.08a	<0.05a	0.30a	20.86c	s<0.56c	I
Inner Oslofjord	5	5	i	i	i	0.15A	1.86B	i	0.01A	i	23.60A	i	<345.70c	0.26a	2.20d	0.19a	0.19a	4.18b	14.30e	III
Frierfjord	3	3	i	i	i	i	i	i	i	i	i	i	<418.40c	0.61a	17.55e	1.32b	6.64b	s<2.25d	V	
Inner Kristiansandsfjord	2	2	i	i	i	i	i	i	i	i	i	i	<850.30c	i	i	i	i	i	V	
Saudafjord	1	2	i	0.75B	i	0.15A	i	i	i	i	35.10B	i	i	17.00d	4.08b	<0.05a	0.60a	1.92a	i	IV
Serfjorden	2	2	i	8.80D	i	3.82D	1.53B	i	0.13C	i	39.70B	i	<38.20a	0.80a	i	i	i	i	i	II
Byfjorden, Bergen	3	3	i	i	i	i	i	i	i	i	i	i	<265.00c	3.80c	i	i	i	i	i	III
Sundalsfjord	2	2	i	i	i	i	i	i	i	i	20.30A	i	<181.60b	6.20c	<0.22a	0.19a	<1.51a	i	i	
Orkdalsfjord	3	3	i	0.50A	i	0.30A	1.73A	i	0.01A	i	44.20B	i	i	i	i	i	i	i	i	I
Inner Ramfjord	2	2	i	1.03B	i	0.12A	i	i	i	i	i	i	i	i	i	i	i	i	i	III
Count			0	6	0	7	5	0	5	0	7	0	6	6	6	6	6	6	3	11
Min			.	0.15A	.	0.11A	1.47A	.	0.01A	.	15.90A	.	<38.20a	0.80a	<0.22a	<0.05a	0.19a	<1.51a	s<0.56c	I
Max			.	8.80D	.	3.82D	1.86B	.	0.13C	.	44.20B	.	<850.30c	35.00e	4.08b	17.55e	1.32b	20.86c	14.30e	V
Mean			.	1.92B	.	0.70B	1.65A	.	0.05B	.	28.51A	.	<349.87c	11.02d	<1.14a	<3.34d	0.48a	<6.66b	s<5.70e	s3.1

i (118) i Value ignored when calculating Environment Class.
a/A/C 49) > Below Class-I limit.
b/B/C 25) > Below Class-III limit.
c/C/C 15) > Below Class-III limit.
d/D/C 9) > Below Class-IV limit.
e/E/C 7) > Below Class-V limit.

Appendix I6.
INDEX - Summary table “Reference index”

1995

Max(Median). Statistics om ALL AREAS: (n = INDEX-Stations measured, N = Station programmed for INDEX)

INDEX-Areas/Names (Reference Areas)	n	N	AS	Pb	F	CD	CU	CR	HG	NI	ZN	AG	PAH2S	BAP	DOT2S	HCB	HCH2S	CB2S2e	TCOON	Max
			ppm d.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm d.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	E-C I-V
Mid and outer Oslofjord	3	3	W	0.27A	W	0.25A	1.62A	W	0.01A	W	24.30A	W	W	W	<0.95a	0.07a	0.40a	7.86b	<0.12a	II
Lista	2	2	W	0.12A	W	0.33A	1.22A	W	0.02A	W	17.50A	W	<17.60a	0.50a	<0.34a	0.05a	0.38a	<1.28a	<0.09a	I
Baino-Sotra	1	1	W	0.22A	W	0.28A	1.39A	W	0.01A	W	28.50A	W	W	W	<0.46a	<0.05a	0.31a	<1.38a	W	I
Outer Rarfjord, Helgeland	1	1	W	0.18A	W	0.15A	0.96A	W	0.01A	W	15.90A	W	<28.20a	<0.50a	0.21a	<0.05a	0.38a	<0.90a	W	I
Lofoten	1	1	W	0.47B	W	0.11A	1.40A	W	0.05B	W	28.50A	W	W	W	4.20b	<0.05a	0.15a	12.31b	<0.23b	II
Firsmes-Skjerøy	1	1	W	0.18A	W	0.56B	1.55A	W	0.01A	W	17.80A	W	W	W	<0.18a	<0.05a	0.16a	<0.81a	W	II
Hammerfest-Honningsvåg	2	2	W	0.47A	W	0.51B	1.75A	W	0.01A	W	19.20A	W	<131.60b	0.70a	<0.23a	<0.05a	<0.15a	<1.34a	W	II
Veranger Peninsula	3	3	W	0.57A	W	0.32A	2.04B	W	0.02A	W	24.70A	W	<6.90a	<0.50a	<0.36a	<0.05a	0.16a	<0.88a	W	II
Count			0	8	0	8	8	0	8	0	8	0	4	4	8	8	8	8	3	8
Min			.	0.12A	.	0.11A	0.96A	.	0.01A	.	15.90A	.	<6.90a	<0.50a	<0.18a	<0.05a	<0.15a	<0.81a	<0.09a	I
Max			.	0.57B	.	0.56B	2.04B	.	0.05B	.	28.50A	.	<131.60b	0.70a	4.20b	0.07a	0.40a	12.31b	<0.23b	II
Mean			.	0.31A	.	0.31A	1.49A	.	0.02A	.	22.05A	.	<46.08a	<0.55a	<0.87a	<0.05a	<0.26a	<3.34a	<0.15a	1.6

w (53) ! Missing value. Should be included when calculating Environment Class E-C
a/A(101) > Below Class-I limit.
b/B(18) > Below Class-II limit.

1996

Max (Median) . Statistics om ALL AREAS: (n = INDEX-stations measured, N = Station programmed for INDEX)

INDEX-Arealknes (ReferenceAreas)	n	N	AS	Pb	F	CD	CU	CR	HG	NI	ZN	AG	PAH22	BAP	D0722	HCB	HCH22	CB22e	TCDDN	Max E.C I-V
			ppm w.wt	ppm w.wt	ppm d.wt	ppm w.wt	ppm w.wt	ppm w.wt	ppm w.wt	ppm w.wt	ppm w.wt	ppm w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	ppb w.wt	
Mid and outer Oslofjord	3	3	1.66A	0.39A	W	0.29A	1.63A	0.11A	0.01A	0.20A	25.50A	0.03A	<19.20a	W	<0.25a	<0.05a	0.25a	13.95b	s<0.17a	II
Lista	2	2	W	0.11A	W	0.20A	1.21A	W	0.01A	W	38.00B	W	<0.20a	<0.50a	<0.20a	0.05a	0.29a	<2.14a	s<0.04a	II
Bemlo-Sotra	1	1	W	0.25A	W	0.19A	1.44A	W	0.01A	W	34.20B	W	<0.11a	W	<0.11a	<0.05a	<0.14a	<0.78a	s<0.07a	II
Outer Ranfjord, Helgeland	1	1	W	0.17A	W	0.16A	1.50A	W	0.01A	W	13.80A	W	W	W	<0.12a	<0.05a	0.21a	<0.62a	W	I
Lofoten	1	1	W	0.79B	W	0.15A	1.44A	W	0.06B	W	27.20A	W	W	W	<1.15a	<0.05a	<0.13a	8.90b	s<0.25b	II
Finnsnes-Skjervei	1	1	W	0.13A	W	0.28A	1.36A	W	0.01A	W	16.30A	W	<10.05a	<0.50a	<0.05a	<0.05a	<0.05a	<0.40a	s<0.03a	I
Hammerfest-Horningsvåg	2	2	W	0.29A	W	0.48B	1.56A	W	0.01A	W	16.40A	W	<111.30b	0.80a	<0.11a	<0.05a	<0.11a	<1.59a	s<0.14a	II
Varanger Peninsula	3	3	W	0.15A	W	0.47B	1.73B	W	0.01A	W	18.20A	W	<36.20a	<0.50a	<0.14a	<0.05a	0.14a	<0.98a	s<0.12a	II
Count			1	8	0	8	8	1	8	1	8	1	4	4	8	8	8	8	7	8
Min			1.66A	0.11A	.	0.15A	1.21A	0.11A	0.01A	0.20A	13.80A	0.03A	<10.05a	<0.50a	<0.05a	<0.05a	<0.05a	<0.40a	s<0.03a	I
Max			1.66A	0.79B	.	0.48B	1.73B	0.11A	0.06B	0.20A	38.00B	0.03A	<111.30b	0.80a	<1.15a	0.05a	0.29a	13.95b	s<0.25b	II
Mean			1.66A	0.29A	.	0.28A	1.48A	0.11A	0.02A	0.20A	23.70A	0.03A	<44.19a	<0.58a	<0.27a	<0.05a	<0.17a	<3.67a	s<0.12a	sI.8

W (.45) ! Missing value. Should be included when calculating Environment Class E.C
a/A(120) > Below Class-I limit.
b/B(19) > Below Class-II limit.