


The model MERLIN applied to Nordmoen, Norway



Miljøverndepartementet
Fagrapport nr. 95



Naturens Tålegrenser

Programmet Naturens Tålegrenser ble satt igang i 1989 i regi av Miljøverndepartementet. Programmet skal blant annet gi innspill til arbeidet med Nordisk Handlingsplan mot Luftforurensninger og til pågående aktiviteter under Konvensjonen for Langtransporterte Grensoverskridende Luftforurensninger (Genevekonvensjonen). I arbeidet under Genevekonvensjonen er det vedtatt at kritiske belastningsgrenser skal legges til grunn ved utarbeidelse av nye avtaler om utslippsbegrensning av svovel, nitrogen og hydrokarboner.

En styringsgruppe i Miljøverndepartementet har det overordnede ansvar for programmet, mens ansvaret for den faglige oppfølgingen er overlatt en arbeidsgruppe bestående av representanter fra Direktoratet for naturforvaltning (DN), Norsk polarinstitutt (NP) og Statens forurensningstilsyn (SFT).

Arbeidsgruppen har for tiden følgende sammensetning:

Gunnar Futsæter - NP
Tor Johannessen - SFT
Else Løbersli - DN
Steinar Sandøy - DN

Styringsgruppen i Miljøverndepartementet består av representanter fra avdelingen for naturvern og kulturminner, avdelingen for vannmiljø, industri- og avfallssaker og avdelingen for internasjonalt samarbeid, luftmiljø og polarsaker.

Henvendelse vedrørende programmet kan rettes til:

Direktoratet for naturforvaltning
Tungasletta 2
7005 Trondheim
Tel: 73 58 05 00

eller
Statens forurensningstilsyn
Postboks 8100 Dep
0032 Oslo 1
Tel: 22 57 34 00

Main Office

P.O. Box 173, Kjelsås
N-0411 Oslo
Norway
Phone (47) 22 18 51 00
Telefax (47) 22 18 52 00

Regional Office Sørlandet

Televeien 1
N-4890 Grimstad
Norway
Phone (47) 37 04 30 33
Telefax (47) 37 04 45 13

Regional Office Østlandet

Rute 866
N-2312 Ottestad
Norway
Phone (47) 62 57 64 00
Telefax (47) 62 57 66 53

Regional Office Vestlandet

Thormøhlensgt 55
N-5008 Bergen
Norway
Phone (47) 55 32 56 40
Telefax (47) 55 32 88 33

Akvaplan-NIVA A/S

Søndre Tollbugate 3
N-9000 Tromsø
Norway
Phone (47) 77 68 52 80
Telefax (47) 77 68 05 09

Title The model MERLIN applied to Nordmoen, Norway	Serial No. 3844-98	Date 25. March 1998
	Report No. Sub-No. O-95173	Pages Price 41
Author(s) Sogn, Trine A. (IJVF, NLH) Wright, Richard F.	Topic group acid precipitation	Distribution
	Geographical area Norway	Printed NIVA

Client(s) Naturens tålegrense (Statens forurensningstilsyn) European Commission	Client ref.
---	-------------

Abstract The process-oriented dynamic model, MERLIN, is been used to predict N leaching and risk of «N saturation» at the experimental field at Nordmoen, Norway. The model was calibrated to the period 1957 to 1987, and then compared to observed data from 4 years of experimental addition of N at 20 kg N ha ⁻² yr ⁻¹ (above ambient 11 kg N ha ⁻² yr ⁻¹). The calibrated model was used to predict future trends in N leaching. MERLIN gave acceptable simulation of the increased N leaching observed in the N addition experiment. The model predicts that with ambient deposition there will be no significant leaching of N within the next 60 years; the forest is not at risk for N saturation. With ambient N load the model predicted the C:N ratio in forest floor to remain above 30. The enhanced deposition, on the other hand, is predicted to lead to significant increased NO ₃ in leachate and a drop in the C:N ratio to below 25 after about 30 years. The critical load for nitrogen for this ecosystem is thus predicted to lie between 11 and 31 kg N ha ⁻² yr ⁻¹ .
--

4 keywords, Norwegian 1. sur nedbør 2. nitrogen 3. modell 4. skog	4 keywords, English 1. acid precipitation 2. nitrogen 3. model 4. forest
---	--



Richard F. Wright
Project manager

ISBN 82-577-3424-1



Bjørn Olav Rosseland
Head of research department

The model MERLIN applied to Nordmoen, Norway

Trine A. Sogn
Department of Soil and Water Sciences
The Agricultural University of Norway
Box 5008
1432 Ås

Richard F. Wright
Norwegian Institute for Water Research
Box 173 Kjelsås
0411 Oslo

Preface

This project came about as a result of interest in applying the new MERLIN model with data from the experimental field at Nordmoen, Norway. The project was supported by Naturens Tålegrense (Statens forurensningstilsyn contract 971800) and European Commission (Contract ENV4-CT95-0030, DYNAMO project).

Oslo, November 1997

Richard F. Wright

Contents

1. Summary	7
2. Introduction	8
3. Site description, methods and data source	8
3.1. Site description	8
3.2. Experimental design	9
3.3. Methods and data for the forest at Nordmoen 1957 and 1987	9
3.3.1. Pools 1987	9
3.3.2. Fluxes 1987	10
3.3.3. Changes in pools and fluxes from 1957 to 1987	11
4. Description of MERLIN	11
5. Calibration procedure	14
6. Results and discussion	15
7. Acknowledgements	19
8. References	19
Appendix A: C and N mass balances used in MERLIN	21
Appendix B: Parameter values for calibration of MERLIN to Nordmoen, Norway	25

1. Summary

Atmospheric N deposition, in excess of plant requirements, is likely to lead to changes in forest ecosystem structure and function. Continuous input of N from the atmosphere may in the long-term produce «N-saturation», which among other things, is characterised by increased leaching of inorganic N below the rooting zone. Increased leaching leads to acidification of soils and waters. MERLIN (Model of Ecosystem Retention and Loss of Inorganic Nitrogen) is a potential tool for modelling the dynamics of N leaching, and can in the future be combined with MAGIC to model the dynamics of acidification of soils and waters and thus predict future exceedence of critical loads. Here MERLIN has been applied to the experimental field at Nordmoen, Norway.

The forest stand at Nordmoen (R1 and R2 plots) is a mixture of Scots pine and Norway spruce, 40 to 50 years old. The site was intensively investigated in the period 1986-1988 as part of the Integrated Forest Study (Johnson and Lindberg 1991). Since 1992 the plots have been added extra doses of N at $20 \text{ kg N ha}^{-2} \text{ yr}^{-1}$ in addition to ambient $11 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (I. Røsberg, pers. comm.).

MERLIN describes C and N cycles, and is comprised of 2 plant and 2 soil compartments (labile (LOM) and refractory (ROM) organic matter). Transfers between compartments occur by means of processes such as plant uptake, litter production, immobilisation, mineralization, nitrification and denitrification. Rates of plant uptake and cycling between pools are governed by the C:N ratios of the plant and soil pools.

The model was calibrated to match the C and N content of the different forest ecosystem compartments in the R1 and R2 plots at Nordmoen. The year 1957 was chosen as the starting point for the hindcast simulation. At this time the N deposition was significantly lower than today. A C and N cycle for the period 1957-1987 was reconstructed. Some historical information of changes in C and N content in different compartments of the young forest at Nordmoen is published, but the paucity of good historical information is a major source of uncertainty in the present model study. The N required to stand development and humus build-up, as well as loss of N in runoff, were assumed to come from deposition of N and «mining» of N from the soil ROM compartment.

The simulated leaching of N was compared to the N leakage measured after 4 years of N addition.

Two different scenarios were simulated. Scenario 1 assumes constant deposition at present-day levels. MERLIN simulated only a small increase in N leaching, and the C:N ratio did not decrease to less than 30 during the 60-year simulation period (1987-2047). For scenario 2 the extra N deposition of $20 \text{ kg ha}^{-1} \text{ yr}^{-1}$ was assumed to continue for 60 years. MERLIN simulated a significant increase in N leaching, and the C:N ratio in forest floor dropped below 25 around the year 2020. With this N load the time needed for the system to switch from being N-limited to N-rich was approximately 30 years.

As emphasised by Kjønaas *et al.* (1997) in the closely related assessment of the MERLIN model to the NITREX field at Gårdsjön, Sweden, the processes which govern the transition of a forest ecosystem from being N-rich to N-limited are still inadequately known. The ability of MERLIN to predict the time needed for a forest system to be N saturated is still uncertain. MERLIN may be an important contribution to the further development of qualitative understanding of the N cycle in forests. In order to aid evaluation of the model an extensive sensitivity analysis of the different model parameters are needed, as well as more data on changes in the C and N content in different compartments during forest succession. More data from long-term N addition experiments is also urgently required in order to properly test the model dynamics and prediction ability.

Coupling of MERLIN with MAGIC may provide a mechanism by which the dynamics of N and S exceedences of critical loads for soil and waters in Norwegian forest ecosystems can be explored.

2. Introduction

During the past years, there has been an increasing number of suggestions that deposition of N compounds from the atmosphere may negatively affect forest ecosystems (e.g. Nihlgård 1985, Skeffington and Wilson 1988). Atmospheric N deposition, in excess of plant requirements, is likely to lead to changes in ecosystem structure and function as well as to increase the loss of inorganic N to runoff. In N limited forest ecosystems typical for the boreal zone, the internal N cycle is tight, and little NO_3^- is usually lost to the soil water below the rooting zone (e.g. Stuanes *et al.* 1995). Even in these N limited forests, continuous input of N from the atmosphere may in the long-term produce «N saturation». True N saturation occurs when N losses are equal to or greater than N inputs. Other definitions put forward have simply viewed N saturation as the availability of NH_4^+ and NO_3^- in excess of the total combined plant and microbial N demand (Aber *et al.* 1989).

Predictions of the amount and number of years of N deposition a given forest ecosystem can tolerate before N saturation occurs, are crucial for the determination of critical loads for N in forest ecosystems. In the present study the process-oriented dynamic model MERLIN (Model of Ecosystem Retention and Loss of Inorganic Nitrogen) (Cosby *et al.* 1997) is used to indicate the time required for a forest ecosystem to move from a N limited to a N saturated state under a given rate of N deposition, assuming N saturation to occur when the N losses are equal or greater than the N inputs. MERLIN describes C and N cycles and is comprised of two plant and two soil compartments (labile and refractory organic matter). Transfer between compartments in the model occurs by means of processes such as plant uptake and cycling by the C:N ratios of the plant and soil pools (Cosby *et al.* 1997).

Intensive investigations of the C and N content in different compartments of a 40 to 50 years forest stand at Nordmoen, south-east Norway, were carried out during the Integrated Forest Study (Johnson and Lindberg 1992). The natural N deposition in the area is $0.079 \text{ mmol N m}^{-2} \text{ yr}^{-1}$. Since 1992 the forest have received extra N, as NH_4NO_3 (I. Røsberg, pers. comm.).

In the present study we describe the calibration of MERLIN to the data from Nordmoen. An attempt to test the model is also carried out. Simulated leaching of N is compared to observed data on soil water N content following enhanced N input. The time for increased N leaching and N saturation at higher N load at Nordmoen is predicted by MERLIN. The potential of MERLIN to predict ecosystem effects following increased N input in forest ecosystems is discussed.

1. Site description, methods and data sources

3.1 Site description

The experimental field station at Nordmoen is located about 60 km north of Oslo at an elevation of 200 m. The R1 and R2 plots, used in the present study were established in 1986 as part of the Integrated Forest Study programme (Johnson and Lindberg 1991) in pristine forest.

The site was located on a flat plain of glaciofluvial deposits partly covered by a sheet of eolian sand. The deposits are about 60 m deep, overlying Precambrian and Permian crystalline bedrocks. The soil in the area was classified as a Typic Udipsamment (Stuanes and Sveistrup 1979).

The mature vegetation in the plots was a mixture of Norway spruce (*Picea abies* (L.) Karst.) and Scots pine (*Pinus sylvestris* L.), 40-50 years old. The dominating vascular plants were *Vaccinium myrtillus* L. and *Deschampsia flexuosa* (L.) Trin. (*Aira flexuosa* L.). The forest floor layer was dominated by moss species such as *Pleurozium schreberi* (Brid.) Mitt., *Hylocomium splendens* (Hedw.) Br Eur., *Ptilium crista-castrensis* (Hedw.) De Not., and *Dicranum* (Hedw.) spp. (Johnson and Lindberg 1991).

3.2 Experimental design

The natural N deposition in this area is $0.079 \text{ mmol N m}^{-2} \text{ yr}^{-1}$ ($11 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). In the period from 1992 to 1997 the R1 and R2 plots have been treated with extra doses of N, as NH_4NO_3 , in doses of 1 and $2 \text{ g N m}^{-2} \text{ yr}^{-1}$ (10 and $20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) (I. Røsberg, pers. comm.). The NH_4NO_3 solution was applied 4 to 6 times during the growing season by use of a hand-driven insects spray pump. Concentrations of N in the soil water below the BC horizon were measured in the late autumn 1995, after 4 years of treatment (I. Røsberg, pers.comm.).

3.3 Methods and data for Normoen forest site 1987 and 1957

The year 1957 was chosen as a starting point for this model application. At this time the N deposition was significantly lower than the present-day deposition, and the forest was 10 to 20 years old. We thus avoid uncertainties involved in modelling the clear-cut and replanting period.

Values for C og N pools and fluxes for the reference year 1987 were obtained from measurements of tree biomass, litterfall, and soil chemistry reported by Johnson and Lindberg (1991).

3.3.1 Pools 1987

Trees:

The total biomass in the R1 and R2 plots is reported in the Table NS-1 in Johnson and Lindberg (1992). The various components were divided into a structural «wood stock» compartment and an active «plants» component. By consulting the closely related work by Kjønåas *et al.* (1997) where the MERLIN model was applied to NITREX Gårdsjön, the active component was decided to be calculated as the sum of foliage, 33.3% of the total branch, understory and 86.5% of the roots. The part of the total branch included in the active compartment was based on the assumption that 33.3% of the branch consists of twigs, which are reconditioned different from the branches. As done by Kjønåas *et al.* (1997) the part of the root biomass considered active was the fine roots less than 5 mm in diameter, assumed to be 86.5% of the total root biomass. The structural «wood stock» component was calculated as the sum of bole, stump, the rest of the branch (total branch-twigs=66.7%) and the rest of the roots (total root biomass-fine roots<5mm=13.5%). Assuming C to be 50% of the organic matter measured, the C and N pools in the active component were 199.5 and 2.11 mol m^{-2} , respectively. In the «wood stock» pool 189 and 1.23 mol m^{-2} , respectively.

Since no functions describing yearly increments in structural and active tree compartments for young Norway spruce forest were found, the yearly increments («pool change») in the structural and active compartments in the 40 to 50 years old stand were estimated using a modified version of the «empirical stand development» curves of Scots pine proposed by Albrektson (1980).

Soil:

The soil organic matter consists of components with quite different decomposition rates, ranging from days to several thousands of years. In the MERLIN programme the organic matter is simplified to include two fractions only. The labile organic matter (LOM) represents a fraction with a rapid turnover time (0 to 10 years), while the refractory organic matter (ROM) represents the organic matter compartments with longer turnover time (10 to 100 years). Although the different types of organic matter are dispersed throughout the soil both vertically and horizontally, in general litter and forest floor consists of the more rapidly decomposed material, whereas the organic matter in the deeper soil horizons usually consists of older material with a slower turnover time. In the present model application, LOM was represented by the organic matter in the O-horizon, while ROM was taken to be the organic matter content of the deeper mineral horizons. The C and N contents in these two compartments were gained by using the data collected during the Integrated Forest Study programme, Table NS-1 in Johnson and Lindberg (1991). This resulted in C:N ratios of the LOM and ROM pools of 39 and 19, respectively.

Yearly changes («pool change») in the LOM and ROM pool were estimated by using data from Sogn *et al.* (In review), where changes in total content of C and N in the mor layer and mineral soil down to 60 cm have been investigated during a forest rotation (0-100 years) at Nordmoen.

3.3.2 Fluxes 1987

Input-output:

Atmospheric input, and the leaching of N at 60 cm soil depth (BC horizon) were measured in the period from October 1986 to October 1988 during the IFS programme (Table NS-2 in Johnson and Lindberg 1991). The mean deposition of N (sum of wet and dry deposition) in this period was 79 mmol m⁻² yr⁻¹. The output of inorganic N at 60 cm soil depth was 1.7 mmol m⁻² yr⁻¹.

Litterfall:

Litterfall was measured during the IFS-programme (Table NS-3 in Johnson and Lindberg 1991). The above-ground litterfall contained 12 mol C m⁻² yr⁻¹. As model input litter flux includes both above and below-ground litterfall. By assuming a below ground yearly litter production equal to the fine roots biomass down to 60 cm depth, a total C litter flux of 32.4 mol m⁻² yr⁻¹ was estimated.

Soil N transformation processes:

A buried-bag incubation technique has been used at Nordmoen to measure the rate and amount of microbiological conversion of organic N to NH₄⁺ (ammonification) and NH₄⁺ to NO₃⁻ (nitrification) (Mahindan 1988). Incubations, made *in situ*, were conducted over 1 and 3 month during the growing season (May-August) and over 1 month in September and October. Based on this study the net N mineralization rate was estimated to be 10.8 mmol m⁻² yr⁻¹, which was only 0.2% of the total N in LOM. In an assembly describing the nutrient cycling at the R1/R2 plots at Nordmoen, Kvindesland and Røsberg (In review) pointed out that a mineralization of all the N in the annual litterfall would still not satisfy the N uptake. Although the deficit could partly be compensated by draining N from the existing large pool of soil N, the study indicated a significantly higher N mineralization rate than found by Mahindan (1988). Using the figure estimated from the study by Mahindan (1988) as a starting value, the net N mineralization percent of the LOM N content was adjusted to be 0.6% (Appendix A) in order to obtain internal consistency in the N cycle (Appendix A) .

3.3.3 Changes in pools and fluxes from 1957 to 1987

The area where the R1 and R2 plots were situated, was after clear cutting replanted in 1937 and 1947. The stand age was in 1987 40 and 50 years old.

Data for variations in bole, active crown and litterfall C and N content within the 40 to 50 years of development are not available for this area. The yearly increments in the structural and active compartments in the 40 to 50 years old stand were estimated using a modified version of the «empirical stand development» curves of Scots pine proposed by Albrektson (1980). The following changes were assumed. As the forest at Nordmoen was growing from age 10 to 20 years in 1957 to age 40 to 50 years in 1987, only about 1 mol N m^{-2} was stored in the boles of the trees, and only about $0.05 \text{ mol N m}^{-2}$ was lost to leachate (Appendix A). During the stand development the active plant compartment was assumed to increase the N storage by 1.8 mol N m^{-2} . Time trends in the C and N content of LOM and ROM were estimated using data from Sogn *et al.* (In review) where changes in total C and N in the mor layer and mineral soil down to 60 cm have been investigated during a forest rotation (0-100 years) at Nordmoen. The LOM pool the C content was estimated to increase by $0.24 \text{ mol m}^{-2} \text{ yr}^{-1}$ with a C:N ratio of the ambient LOM (Appendix A). A «mining» of ROM in order to supply the N necessary for growth of trees and build up of forest floor was assumed. Such «mining» has been reported in several studies (e.g. Johnson 1991, Emmett *et al.* 1996, Sogn *et al.* In review) and is probably a natural feature of boreal forests. Most probably N is translocated from ROM to LOM via the plants, initially by uptake and then litterfall. According to studies by Berg (1986) there is a higher net release of N during the mineralization of older organic matter (ROM) than in younger material. This net released N is then available for plant uptake. During the mineralization of litter in the mor layer, less decomposed litter may cause a net immobilization of N and thus a reimmobilization instead of uptake by roots. In this way a recycling of N within the soil is taken place. The wood pool C content was assumed to increase by $7.7 \text{ mol m}^{-2} \text{ yr}^{-1}$ with a C:N ratio of 154 (Appendix A).

4. Description of MERLIN

MERLIN is a simple process-oriented model focused on simulating and predicting concentrations of inorganic N in soil leachate and runoff in terrestrial ecosystems (Cosby *et al.* 1997). The model links C and N cycles. The ecosystem is simplified to 2 plant compartments (active biomass such as foliage and roots and structural woody biomass) and 2 soil compartments (labile and refractory organic matter) (Figure 1). The C pools and fluxes are set external to the model. N pools and fluxes are then linked to C by C:N ratios in the pools. Transfer between compartments occurs by processes such as plant uptake, litterfall, immobilisation, mineralization, nitrification and denitrification. There is no feedback between N status and C fluxes; this would be the next step in complexity. The key to MERLIN is that the rates of these processes are assumed to be governed by C:N ratios in the various pools. MERLIN is comprised of (1) a book-keeping procedure in which inputs and outputs of C and N to the ecosystem and between the 4 compartments are tallied, and (2) a series of simultaneously-operating processes that describe the transfer of C and N between the compartments and out of the ecosystem. The model is designed to function at annual time steps.




Figure 1. A schematic representation of the MERLIN model. Closed arrows represent fluxes of inorganic C; open arrows represent fluxes of inorganic N. Paired closed/open arrows represent fluxes of organic material containing both C and N (from Cosby *et al.* 1997.).

The model deals with NO_3 , NH_4 and organic N. N inputs are as atmospheric deposition, fertilisers and N fixation. Time series of all inputs of inorganic N must be specified *a priori*. Nitrification, microbially-mediated transformation of NH_4 to NO_3 , is represented in the model by a first-order reaction (1). The rate of loss of NH_4 is the product of a rate constant and the concentration of NH_4 at each time step:

$$\text{FN}_{\text{nit}} = \text{NIT} \times (\text{NH}_4)_{\text{soil}} \times \text{SPV} \quad (1)$$

where FN_{nit} is the nitrification loss of NH_4 ($\text{mmol m}^{-2} \text{yr}^{-1}$), $(\text{NH}_4)_{\text{soil}}$ is the concentration of NH_4 in soil solution (mmol m^{-3}), NIT is the nitrification rate constant (yr^{-1}), and SPV is the pore volume per unit area of the soil (m). Adsorption of NH_4 on the soil matrix is modelled as a non-linear capacity-limited process using a Langmuir isotherm approach. N losses from the system are as inorganic in runoff and as gases (N_2O and N_2) from denitrification.

Plant biomass represents the aggregated pool of C and N present in the active portion of the ecosystem. The time series of net productivity is determined from specified time sequences of plant biomass, litter production and long-term storage such as wood production (Figure 1).

Plant growth requires N uptake from the soil as NH_4 and NO_3 . Uptake is modelled as a non-linear Michaelis-Menten process that depends on the soil water concentration of NH_4 or NO_3 (2).

$$\text{FN}_{\text{Xupt, plt}} = K_{\text{mx, Nx, plt}} \times \frac{(\text{Nx})_{\text{soil}}}{(K_{\text{hlf, Nx, plt}} + (\text{Nx})_{\text{soil}})} \quad (2)$$

where $\text{FN}_{\text{Xupt, plt}}$ is the uptake flux of inorganic N species by the plant compartment ($\text{mmol m}^{-2} \text{yr}^{-1}$), $(\text{Nx})_{\text{soil}}$ is the concentration of inorganic N species in soil solution (mmol m^{-3}). $K_{\text{mx, Nx, plt}}$ is the maximum uptake rate ($\text{mmol m}^{-2} \text{yr}^{-1}$) which is calculated by the formula

$$K_{\text{mx, Nx, plt}} = FC_{\text{pp}} \times M_{\text{Nx, plt}} \quad (3)$$

where FC_{pp} is the net primary production ($\text{mmol C m}^{-2} \text{yr}^{-1}$) and $M_{\text{Nx, plt}}$ is the maximum N element, Nx , uptake per unit of C production (mol N per mol C). $K_{\text{hlf, Nx, plt}}$ is the half-saturation constant (the concentration of inorganic N species at which plant uptake proceeds at half the maximum rate; mmol m^{-3}) calculated by the formula

$$K_{\text{hlf, Nx, plt}} = K_0 \times \exp\left(\frac{C:N_{0, \text{plt}} - C:N_{\text{plt}}}{S_{\text{Nx, plt}}}\right) \quad (4)$$

where K_0 is the upper limit of uptake half saturation parameter as C:N's of biomass decrease (mmol m^{-3}), $C:N_{0, \text{plt}}$ the target C:N ration of the biomass compartment (mol C per mol N), $C:N_{\text{plt}}$ the C or N ratio of the biomass compartment (mol C per mol N) and $S_{\text{Nx, plt}}$ is the steepness of decline in uptake half-saturation parameter as C:N's of biomass increase (unitless). The maximum uptake rate is assumed to be proportional to the net primary productivity of the plants.

Soil organic material is divided into two compartments: labile organic matter (LOM) and refractory organic matter (ROM). Each is an aggregated pool of C and N representing accumulated organic compounds in the ecosystem. These materials provide the energy substrate for soil micro-organisms, which immobilise and mineralise C and N in soils (Figure 1). Similar to plant uptake the microbial immobilisation of inorganic N is modelled as a non-linear Michaelis-Menten process that depends on the concentration of NH_4 or NO_3 using the equations (2), (3) and (4). The maximum immobilisation rate is assumed proportional to microbial secondary productivity.

Litter from the plant compartment enters the LOM pool. C and N leave this pool by decomposition (transformation of organic C to CO_2) and decay (degradation of the quality of the organic material). The decay products are passed on to the next compartment (ROM). Decomposition also occurs in ROM. The model requires input information for litter production, size of the C pool stored in LOM, and decomposition of LOM at each time step. In addition the size of the C pool stored in ROM is required.

These compartments and fluxes are highly aggregated. Conceptually the LOM pool represents that fraction of soil organic matter with relatively short turnover time (0-10 years), whereas ROM represents the fraction of soil organic matter with longer turnover time. The LOM pool may be most readily identified with the forest floor, but organic matter such as decaying roots are also LOM and are found in deeper soil layers. LOM provides a soil organic matter compartment that can respond rather quickly to changing external conditions and inputs. The ROM pool, on the other hand, represents the bulk of C and N present in the rest of the soil profile. The long-term storage losses from the ROM represent losses of organic matter through leaching of dissolved organic compounds in drainage waters or peat formation.

Inputs required for MERLIN are temporal sequences of: 1) C fluxes and pools, 2) hydrologic discharge, and 3) external sources of inorganic N. Initial conditions (amounts of C and N) must be specified for each compartment. A number of «constants» (e.g. uptake parameters) are needed to specify the N dynamics of the organic compartments, and characteristics of the soils must be given (depth, porosity, bulk density, anion/cation exchange characteristics).

Outputs from MERLIN include: 1) concentrations and fluxes of NH_4 and NO_3 in soil water and runoff; 2) total N contents of the various compartments; 3) the C:N ratios of the aggregated plant and soil compartments; and 4) estimates of important processes in the N cycle (i.e. uptake, gross and net mineralization and immobilisation).

5. Calibration procedure

The first step in the calibration of MERLIN to data from the Nordmoen field involved the compilation of C and N pool and flux data for reference year 1987 and for the year 1957 in the past (Appendix A). The necessary data were compiled in spreadsheets (Appendix A) and the complete C and N cycles are calculated and checked for internal consistency.

The next step was to define the time sequences of changes of pools and fluxes over the period 1957-87. In other words, how did the forest grow in terms of C and N. Although the history of the forest in the R1 and R2 plots at Nordmoen is relatively well known, no measurements of changes in C and N storage during the 40 to 50 years of forest growth are available. The following changes was assumed as basis for the calibration procedure. As the forest at Nordmoen was growing from age 10-20 years in 1957 to age 40-50 years in 1987, only about 1 mol N m^{-2} was stored in the boles of the trees, and only about $0.05 \text{ mol N m}^{-2}$ was lost to leachate (Appendix A). During the stand development the active plant compartment was assumed to increase the N storage by 1.8 mol N m^{-2} . We assumed changes in the LOM pool (by an increase in C of $0.24 \text{ mol m}^{-2} \text{ yr}^{-1}$ with a C:N ratio of the ambient LOM, Appendix A) (Sogn *et al.* In review), the wood pool (by an increase in the C content of $7.7 \text{ mol m}^{-2} \text{ yr}^{-1}$ with a C:N ratio of 154, Appendix A), and the N deposition (increasing from 1957 to 1987 following estimates of A. Semb NILU, used by Kjønnaas *et al.* 1997). Changes in ROM pool are calculated by MERLIN at each time step to balance the C budget.

Calibration of MERLIN then involved adjusting the parameters (see eq.(2), (3) and (4)) for the three N uptake/immobilization functions: plants, LOM and ROM. The maximum N uptake per unit of C production ($M_{x,\text{plt}}$, $M_{x,\text{LOM}}$ and $M_{x,\text{ROM}}$, in eq. (3)), the upper limit of uptake half-saturation parameter as C:N's of biomass decrease (K_0 in eq. (4)), the position of the Michaelis-Menten uptake curve (C:N reference, i.e the target C:N ratio of the biomass compartment ($C:N_{0,\text{plt}}$, $C:N_{0,\text{LOM}}$ and $C:N_{0,\text{ROM}}$, in eq. (4))) and the steepness of decline in uptake half-saturation parameter as C:N's of biomass increase ($S_{x,\text{plt}}$, $S_{x,\text{LOM}}$ and $S_{x,\text{ROM}}$, in eq. (4)) (Appendix B). For each trial the model was run from starting values in 1957 to 1987 and the suitability of the calibration was subjectively judged with respect to the best fit for:

- (1) carbon pools in 1987 agree with observed;
- (2) C:N ratios in the pools in 1987 agree with observed;
- (3) Changes in C:N ratios over the 30-year period are acceptable;
- (4) N output (leachate) in 1987 agree with the observed.

Tables of pools and fluxes for the Nordmoen calibration and a print-out of the parameter values are given in Appendix A and B.

The calibrated model was evaluated using results from I. Røsberg's N addition experiments. Finally the model was used to assess the future impact of 2 scenarios of N deposition. Scenario 1 was 60 years of constant N deposition ($11 \text{ kg ha}^{-1} \text{ yr}^{-1}$), while scenario 2 was 60 years of enhanced N deposition ($11+20 = 31 \text{ kg N ha}^{-1} \text{ yr}^{-1}$).

6. Results and discussion

The evaluation

After 4 years with addition of $2 \text{ g N m}^{-2} \text{ yr}^{-1}$, a 4-fold increase in the soil water N concentration below the BC horizon was observed at the R1/R2 plot (I. Røsberg, pers. comm.). MERLIN simulated this increased N leaching (Figure 2). After 4 years with extra N addition MERLIN simulated the flux of N out of the system to be $4.6 \text{ mol N m}^{-2} \text{ yr}^{-1}$. The simulated N flux was about 30 % lower than the observed flux after 4 years with enhanced N input. The discrepancy between simulated and observed is acceptable in that the observed value also is uncertain. This uncertainty stems from the fact that the soil water below the BC-horizon was only sampled occasionally during a limited period the autumn 1995 (I. Røsberg, pers. comm.), and thus seasonal variations are not included. Furthermore the increased N observed in the soil water the autumn 1995 does not necessarily reflect the real N status of the system. The forest system has had a limited time to respond to the sudden enhanced N deposition. On long-term basis, it is the changes in major processes such as the ratio of gross NH_4^+ and NO_3^- immobilization to gross mineralization, and the soil C:N ratio that is expected to determine the leaching level (Aber 1992). Additional years of N treatment and measurements are necessary in order to determine whether the increasing trend is correct, and to fully validate the predictive capacity of MERLIN.

The prediction

The calibrated model was used to assess 2 scenarios for future N deposition at Nordmoen. For scenario 1 (constant deposition) MERLIN simulated a small increase in N leaching from the forest at Nordmoen during the 60-year forecast period (Figure 2). A reduction in the LOM C:N ratio was simulated (Figure 3), but C:N ratio did not decrease below 30 within the 60 year-period. This suggests that this stand at Nordmoen is not at risk of N saturation under present-day levels of N deposition.

For scenario 2 the N deposition was increased by $20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (total $31 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) (Røsberg's experimental dose) for 60 years into the future (Figure 2). The N content in both the active plant pool and LOM were predicted to increase (Figure 3) as did N leaching during the 60-year simulation period (Figure 2). The C:N ratio of the LOM compartment was predicted to drop from around 40 before 1987 down to 25 around 2020, and then below 25 (Figure 3). With this N load the time needed for the system to switch from N-limited, indicated by high C:N ratio in LOM, to N-rich with a C:N ratio below 25, was estimated to be 25-30 years. Since MERLIN simulated the measured N leaching after 4 years of treatment relatively well, some confidence could be put into the results. However, both the rate and capacity of N assimilation, as well as the response of C dynamics to enhanced N deposition are inadequately understood.

The relationship between the C:N ratio and N leaching (Figure 4) corresponded with general results from the European forest stands and NITREX sites, at which significant N leaching appears at sites with low C:N ratios (Gundersen *et al.* 1998) (Figure 4). However, as emphasised by Kjønås *et al.* (1997), the time needed for a system to change from being N limited, indicated by high C:N ratio in LOM, to a N-rich system, with a C:N ratio below 25, is uncertain, and the processes which govern the transition are not fully understood.




Figure 2. Results of MERLIN calibration and prediction for Nordmoen. Deposition and runoff. Solid line represents scenario 1 (constant N deposition at 1987 levels). Dotted line represents scenario 2 (with extra N addition).




Figure 3. Results of MERLIN calibration and prediction for Nordmoen. C:N ratios in the three pools and gross mineralization and immobilisation in LOM and ROM. Solid line represents scenario 1 (constant N deposition at 1987 levels). Dotted line represents scenario 2 (with extra N addition).

General discussion

This application of MERLIN to Nordmoen corroborates applications of the model to NITREX sites at Aber, Wales, UK (Emmett *et al.* 1997) and Gårdsjön, Sweden (Kjønaas *et al.* 1997). In all cases the model calibrations suffer from inadequate data for C and N dynamics in forest ecosystems. In particular the changes in pools and fluxes during a forest rotation cycle are inadequately known. As pointed out by Kjønaas *et al.* (1997) there have been few studies of litter production, build-up of LOM and changes in ROM over a forest life cycle. Traditional forest studies focus mainly on wood production. Studies such as the survey of Sitka spruce stands of different ages in Wales (Emmett *et al.* 1993, Stevens *et al.* 1994) are urgently required for other major tree species such as Norway spruce and Scots pine. More background data regarding variations of C and N during forest succession are needed in order to increase the reliability of MERLIN predictions and those of any model in which N leaching is related to ecosystem N pools.

The application of the MERLIN model to Gårdsjön, Sweden, by Kjønaas *et al.* (1997) showed that MERLIN is a useful tool for bookkeeping of N pools and fluxes. This is also the case for our application to Nordmoen. The model may also be an important contribution to further development of qualitative understanding of the N cycle in forest ecosystems. However, the testing and application of the model is still too limited to generalise on its utility and robustness. Data to fully validate the model are scarce. An extensive sensitivity analysis of the key parameters is also urgently required. However, additional applications of MERLIN to sites such as Nordmoen may indicate whether a relatively narrow range of values for key parameters can satisfactorily explain observed ecosystem effects. Such model applications may also reveal whether major aspects of ecosystem behaviour following changes in N deposition can be accounted for by changes in mineralization and soil N accumulation rates.

This application of MERLIN to Nordmoen can be used to estimate critical load for N in the area. Critical loads for both water (groundwater) and soil at Nordmoen has been calculated previously by Wright et al. (1991) using the dynamic model MAGIC and several static models. These indicate that the critical load for acidity at Nordmoen is zero for both water and soil, largely because the growing forest depletes the soil of base cations faster than these are resupplied by weathering and deposition. More recently the nutrient cycling model NuCM (Liu *et al.* 1992) has been assessed to the Nordmoen site (Kvindesland, In review). While the MAGIC simulations predicted a critical load of $S < 0 \text{ mmol}_c \text{ SO}_4^{2-} \text{ m}^{-2} \text{ yr}^{-1}$ to ensure that the molar Ca:Al ratio in soil solution does not fall below 1, NuCM predicted a load of approximately $70 \text{ mmol}_c \text{ SO}_4^{2-} \text{ m}^{-2} \text{ yr}^{-1}$. Historically and at present most of the acidification at Nordmoen has been due to S deposition. MERLIN indicates that at present-day N deposition there will be no appreciable N leaching and thus no acidity generated by N deposition in the next 60 years. This indicates that the critical load for N is greater than $11 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ but less than $31 \text{ kg N ha}^{-1} \text{ yr}^{-1}$.





Figure 4. N flux out (top panel) and ratio of output to input of inorganic N (bottom panel) versus the C:N ratio in the forest floor (LOM). Solid diamonds: with $20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ addition; grey squares: with no N addition.

6. Acknowledgements

We thank O. Janne Kjønaas, Sheila Kvindesland and Ingvald Røsberg for professional assistance on selecting suitable data from Nordmoen, as well as how to prepare them for MERLIN application.

7. References

- Aber, J.D., Nadelhoffer, K.J., Steudler, P. and Melillo, J.M. 1989. Nitrogen saturation in northern forest ecosystems. *Bioscience* 39: 378-386.
- Aber, J.D. 1992. Nitrogen cycling and nitrogen saturation in temperate forest ecosystems. *Tree* 7: 220-223.
- Albrektson, A. 1980. Total tree production as compared to conventional forestry production. In: Persson, T. (ed.). *Structure and function of northern Coniferous Forests - An Ecosystem Study*. *Ecol. Bull* (Stockholm) 32: 315-327.
- Berg, B. 1986. Nutrient release from litter and humus in coniferous forest soil - a mini review. *Scand. J. For. Res.* 1: 359-369.
- Cosby, B.J., Ferrier, R.C., Jenkins, A., Emmett, B.A., Tietema, A. and Wright, R.F. 1997. Modelling the ecosystem effects of nitrogen deposition at the catchment scale: Model of Ecosystem Retention and Loss of Inorganic Nitrogen (MERLIN). *Hydrol. Earth System Sci.* 1: 137-158.
- Emmett, B.A., Cosby, B.J., Ferrier, R.C., Jenkins, A., Tietema, A., and Wright, R.F. 1997. Modelling the ecosystem effects of nitrogen deposition at the catchment scale: simulation of nitrogen saturation in a Sitka spruce forest, Aber, Wales, UK. *Biogeochemistry* 38: 129-148.
- Emmett, B.A., Reynolds, B., Stevens, P.A., Norris, D.A., Hughes, S., Görres, J. and Lubrecht, I. 1993. Nitrate leaching from afforested Welsh catchments - interactions between stand age and nitrogen deposition. *Ambio* 22: 386-394.
- Emmett, B. A. and Reynolds, B. 1996. Nitrogen critical loads for spruce plantations in Wales: is there too much nitrogen? *Forestry* 69: 205-214.
- Gundersen, P., Emmett, B.A., Kjønaas, O.J., Koopmans, C.J., and Tietema, A. 1998. Impact of nitrogen deposition on nitrogen cycling in forests: a synthesis of NITREX data. *Forest Ecol. Manage.* 101: 37-56.
- Johnson, D.W. 1992. Nitrogen retention in forest soils. *Jour. Environ. Quality* 21: 1-12.
- Johnson, D.W. and Lindberg, S.E. (Eds.) 1991. *Atmospheric Deposition and Forest Nutrient Cycling*. *Ecological Studies* 91. Springer-Verlag, New York, 707pp.
- Kjønaas, O.J., Wright, R.F. and Cosby B.J. 1997. MERLIN applied to NITREX Gårdsjön. *Acid Rain Research Rept.* 46/97, SNO 3696-97, Norwegian Institute for Water research, Oslo, 38pp.
- Kvindesland, S.H.S.B. In review. Testing the Nutrient cycling Model (NuCM) at Nordmoen, a Norway spruce site in south-east Norway. *Ecological Modelling*.

- Kvindesland, S.H.S.B. and Røsberg, I. In review. A comparison of Nutrient cycling in the 30 to 40 years-old Norway spruce plots at Nordmoen, south east Norway. *Water, Air Soil Pollut.*
- Liu, S., Munson, R., Johnson, D.W. *et al.* 1991. The nutrient cycling model (NuCM): overview and application. In D.W. Johnson and S.E. Lindberg (Eds.). *Atmospheric Deposition and Forest Nutrient Cycling*. Ecological Studies 91. Springer-Verlag, New York, pp.583-609.
- Mahindan, M. 1988. Seasonal Changes In Nitrogen Mineralization And Some Other Soil Chemical Parameters In A Forest Ecosystem In Norway. Master thesis, NORAGRIC, The Agricultural University of Norway, Aas, 68pp.
- Nihlgård, B., 1985. The ammonium hypothesis - an additional explanation to the forest dieback in Europe. *Ambio* 14: 2-8.
- Skeffington, R.A. and Wilson, E.J. 1988. Excess nitrogen deposition: Issues for consideration. *Environ. Pollut.* 54: 159-184.
- Sogn, T.A., Stuanes, A.O. and Abrahamsen, G., In review. Accumulation of N in forest soil: Variations due to stand age and N deposition. *Ambio*.
- Stevens, P.A., Norris, D.A., Sparks, T.H. and Hodgson, A.L. 1994. The impacts of atmospheric N inputs on throughfall, soil and stream water interactions for different aged forest and moorland catchments in Wales. *Water Air Soil Pollut.* 73:297-317.
- Stuanes, A.O. and Sveistrup, T.E. 1979. Field experiment with simulated acid rain in forest ecosystems. 2. Description and classification of the soils used in the field, lysimeter and laboratory experiments. SNSF-project, Norway, FR 15/79, 35pp.
- Stuanes, A.O., Kjønnaas, O.J. and van Miegroet, H., 1995. Soil solution response to experimental addition of nitrogen to a forested catchment at Gårdsjön, Sweden. *Forest Ecol. Manage.* 71: 99-110.
- Wright, R.F., Stuanes, A.O. and Frogner, T. 1991. Critical loads for soils in Norway, Normoen. NIVA-report 0-89153, 27p.

Appendix A. C and N mass balances used in MERLIN

Shown are spreadsheets for 2 key years, 1957 and 1987.

Table 1 shows year 1957, the start of the hindcast period at which time the forest was 10/20 years-old.

Table 2 shows the year 1987, the end of the hindcast period. Now the forest is 40/50 years old.

The C and N cycles are calculated from measured (or estimated) values for several parameters. These are shown in **bold** and include estimates for C and N contents of the 2 plant and 2 soil compartments outlined in bold at the centre, the annual C increment to these pools (left-hand column), and the deposition and leaching (right-hand column). In addition the fractions of total mineralisation occurring in LOM and ROM must be specified. All other values are calculated from these, either from mass-balance considerations or from assumptions about C:N ratios. The column at the left shows the annual incremental change in the pools, the next column gives the fluxes between the pools. The columns to the right of the pools show the sinks and sources. The details of the calculations are given in Table 3.

1957

C and N mass balances used in MERLIN		Year: 1957		Nordmoen R1/R2	
Pool change	Fluxes	mol/m2			
Wood increment	Net primary production	Wood stock	Inorganic N		
C 4.1	C 22.10	C 29.9	Sinks		
N 0.030		N 0.22			
C/N 135.9		C/N 135.9	Sources		
delta Plants		Plants (active pool)			
C 3		C 30	Uptake mol	Deposition	
N 0.032		N 0.317	N 0.220	N 0.012	
C/N 94.6		C/N 94.6	Immobilisation LOM	Mineralization % LOM	
delta LOM	Litter above & below	LOM (forest floor)		0.4	
C 0	C 15.00	C 176		Gross/Net calc.LOM	
N 0.000	N 0.159	N 4.47		4.11	
C/N 39.4	C/N 94.6	C/N 39.4		Net Mineralisation LOM	
delta ROM	Decay LOM	ROM (refractory OM)			
C 0.00	C 1.50	C 527	Decomposition LOM		
N 0.000	N 0.038	N 27.2	C 13.50		
C/N 19.4	C/N 39.4	C/N 19.4	N 0.343		
			C/N 39.4		
	Export ROM (orgC,N)		Immobilisation ROM		
	C 0.21		N -0.06		
	N 0.011				
	C/N 19.4		Decomposition ROM		
			C 1.29		
			N 0.066		
			C/N 19.4		
Bold: data inputs			Leaching		
<i>Italics:calculated from C/N, or assumed same C/N as source pool</i>			N 0.0017		
Bold+italics: calculated					
				Gross/Net calc.ROM	
				0.53	
				Net Mineralisation ROM	
				N 0.125	

Balance_1987


C and N mass balances used in MERLIN			mol/m ²		Year: 1987		Nordmoen R1/R2	
Pool change	Fluxes		Pools					
Wood increment	Net primary production	Wood stock	Wood stock		Inorganic N			
C 7.7	C 40.10	C 189	C 189		Sinks			
N 0.050		N 1.23	N 1.23					
C/N 153.7		C/N 153.7	C/N 153.7					
delta Plants	Litter above & below	Plants (active pool)	Plants (active pool)		Uptake		Deposition	
C 0	C 32.40	C 199.5	C 199.5		N 0.393		N 0.079	
N 0.000	N 0.343	N 2.11	N 2.11					
C/N 94.5	C/N 94.5	C/N 94.5	C/N 94.5		Immobilisation LOM		Mineralization % LOM	
delta LOM	Decay LOM	LOM (forest floor)	LOM (forest floor)		N 0.553		0.6	
C 0.24	C 3.00	C 240	C 240				Gross/Net calc.LOM	
N 0.006	N 0.076	N 6.1	N 6.1				3.94	
C/N 39.3	C/N 39.3	C/N 39.3	C/N 39.3					
delta ROM	Export ROM (orgC,N)	ROM (refractory OM)	ROM (refractory OM)		Decomposition LOM		Net Mineralisation LOM	
C -0.94	C 0.03	C 478	C 478		C 29.16			
N -0.048	N 0.002	N 24.6	N 24.6		N 0.741			
C/N 19.4	C/N 19.4	C/N 19.4	C/N 19.4		C/N 39.3			
					Immobilisation ROM		Gross/Net calc.ROM	
					N 0.076		1.60	
					Decomposition ROM		Net Mineralisation ROM	
					C 3.91			
					N 0.201			
					C/N 19.4			
					Leaching			
					N 0.0017			

Bold: data inputs
Italics: calculated from C/N, or assumed same C/N as source pool
Bold+italics: calculated

Balance_1987

Calculations:			
carbon			
C-NPP = C wood increment + C delta plants + C litter			$E7 = B6 + B11 + E13$
C decay LOM = C litter - C delta LOM - C decomp LOM			$E18 = E13 - B16 - K17$
C decomp ROM = C decay LOM - C delta ROM - C export (DOC)			$K25 = E18 - B21 - E23$
nitrogen			
N uptake = N wood increment + N delta plants + N litter			$L11 = B7 + B12 + E14$
N net min LOM = (N uptake - N dep) * % min LOM			$N19 = (L11 - O11) * N14$
N immob LOM = N decomp LOM - N net min LOM			$L14 = K18 - N19$
N net min ROM = (N uptake - N dep) * (1-%min LOM)			$N27 = (L11 - O11) * (1 - N14)$
N immob ROM = N decomp ROM - N net min ROM			$L22 = K26 - N27$


Appendix B. Parameter values for calibration of MERLIN to Nordmoen



FOUR PROCESSES CONTROL INORGANIC NITROGEN SPECIATION AND EXCHANGE BETWEEN BIOMASS POOLS AND THE SOILS.

- UPTAKE: REMOVAL OF NO₃ AND NH₄ FROM SOIL BY AUTOTROPHS (CAN BE FROM BOTH SOILS)
- IMMOBILIZATION: REMOVAL OF NO₃ AND NH₄ BY HETEROTROPHS (FROM UPPER SOIL ONLY)
- MINERALIZATION: RELEASE OF NH₄ INTO SOIL (TO THE UPPER SOIL ONLY)
- NITRIFICATION: CONVERSION OF NH₄ TO NO₃ (CAN OCCUR IN BOTH SOILS)


UPTAKE AND IMMOBILIZATION ARE NONLINEAR FUNCTIONS OF SOIL N CONCENTRATION, C/N OF BIOMASS, AND CARBON PRODUCTIVITY
MINERALIZATION IS A LINEAR FUNCTION OF DECOMPOSITION AND C/N OF BIOMASS



SEVEN FLUXES CONTROL THE CARBON CONTENT OF THE AGGREGATED BIOMASS POOLS.

- (NPP) NET PRIMARY PRODUCTION
- (WP) WOOD PRODUCTION
- (LTR) LITTER PRODUCTION
- (DCY) DECAY OF LABILE ORGANIC MATTER, CONVERTED TO REFRACTORY ORGANIC MATTER
- (DCMP1 & DCMP2) DECOMPOSITION OF LABILE AND REFRACTORY ORGANIC MATTER
- (EXP) ORGANIC MATTER EXPORT FROM SOIL

CARBON FLUXES (MOL/M²/T) AND/OR POOL SIZES (MOL/M²) MUST BE SPECIFIED.
CARBON FLUXES TRANSPORT NITROGEN BY MOVING ORGANIC-N ALONG WITH CARBON IN THE RATIO OF C/N IN THE SOURCE BIOMASS (OR BY USING A SPECIFIED C/N RATIO).



CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 2

----- HINDCAST -----

MERLIN VERSION 3.33 (31 OCT 94)

RUN DATE = 11/17/97

RUN TIME = 16:35:37

CATCHMENT = Nordmoen R1/R2

HINDCAST SIMULATION PERIOD = 30 YEARS

MONTHLY VARIATION = N


START YEAR = 1957

END YEAR = 1987

INTEGRATION TIME STEP = 0.0417 YEARS

ROUTING OF FLOW THROUGH THE CATCHMENT IS:

F1 = % RUNOFF NOT CONTACTING TERRESTRIAL AREAS



	---%---	-----MM-----			
	F1	Q1	Q3	Q4	Q7
ANNUAL AVE	0.	0.	350.	350.	350.

MEAN ANNUAL DISCHARGE = 0.350 M/YR

MEAN ANNUAL PRECIPITATION = 1.000 M/YR

	----- REFERENCE YEAR 1987		----- BACKGROUND YEAR 1957	
	PRECIP CONC (MEQ/M3)	DEPOSITION FACTOR	PRECIP CONC (MEQ/M3)	DEPOSITION FACTOR
NH4	38.0	1.0	5.8	1.0
NO3	41.0	1.0	6.1	1.0

THE FOLLOWING CONDITIONS ARE SET FOR Nordmoen R1/R2

CONSTANT-----	-----STREAM
RETENTION TIME (YR)	0.00
RELATIVE AREA (FRAC)	0.00
MEAN DEPTH (M)	0.00

CONSTANT-----	-----SOIL
SOIL DEPTH (M)	0.45
POROSITY (FRAC)	0.45
BULK DENSITY (KG/M3)	1021.00
NH4 MAX CAP (MEQ/KG)	0.00
NH4 HLF SAT (MEQ/M3)	0.00
NO3 MAX CAP (MEQ/KG)	0.00
NO3 HLF SAT (MEQ/M3)	0.00
PORE VOLUME (M)	0.20
SOIL MASS (KG/M2)	459.45

BIOLOGICAL PARAMETERS FOR NITROGEN ASSIMILATION

AUTOTROPHS (PLANT BIOMASS)		HETEROTROPHS (LAB. ORGANIC)		HETEROTROPHS (REF. ORGANIC)	
NH4 UPT, OPTIMUM C/N	10.000	NH4 UPT, OPTIMUM C/N	1.000	NH4 UPT, OPTIMUM C/N	5.000
NH4 UPT, REFERNC K2	1000.000	NH4 UPT, REFERNC K2	1000.000	NH4 UPT, REFERNC K2	1000.000
NH4 UPT, REFERNC C/N	89.000	NH4 UPT, REFERNC C/N	25.000	NH4 UPT, REFERNC C/N	17.000
NH4 UPT, C/N HFWD K2	1.500	NH4 UPT, C/N HFWD K2	8.000	NH4 UPT, C/N HFWD K2	0.550
NO3 UPT, OPTIMUM C/N	10.000	NO3 UPT, OPTIMUM C/N	1.000	NO3 UPT, OPTIMUM C/N	5.000
NO3 UPT, REFERNC K2	1000.000	NO3 UPT, REFERNC K2	1000.000	NO3 UPT, REFERNC K2	1000.000
NO3 UPT, REFERNC C/N	89.000	NO3 UPT, REFERNC C/N	25.000	NO3 UPT, REFERNC C/N	17.000
NO3 UPT, C/N HFWD K2	1.500	NO3 UPT, C/N HFWD K2	8.000	NO3 UPT, C/N HFWD K2	0.550

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 3 **

C/N RATIOS FOR ORGANIC NITROGEN TRANSFERS

C/N RATIOS (ORGANIC FLUXES)

C/N FOR WOOD PRODUCT	154.000
C/N FOR LITTER	0.000
C/N FOR DECOMP LOM	0.000
C/N FOR DECAY LOM	0.000
C/N FOR DECOMP ROM	0.000
C/N FOR ORGNC EXPORT	0.000

(SETTING THE C/N FOR ANY FLUX = ZERO CAUSES THE PROGRAM TO USE THE C/N OF THE TOTAL SOURCE BIOMASS POOL TO CALCULATE N TRANSFER)

	-----STREAM		-----SOIL	
	SOURCES		SOURCES	
	MEQ/M2/YR	MEQ/M2/YR	MEQ/M2/YR	MEQ/M2/YR
	BACKGROUND	REFERENCE	BACKGROUND	REFERENCE
NH4	0.0	0.0	0.0	0.0
NO3	0.0	0.0	0.0	0.0

	-----STREAM		-----SOIL	
	SINKS		SINKS	
	MEQ/M2/YR	MEQ/M2/YR	MEQ/M2/YR	MEQ/M2/YR
	BACKGROUND	REFERENCE	BACKGROUND	REFERENCE
NH4	0.0	0.0	0.0	0.0
NO3	0.0	0.0	0.0	0.0

(NEGATIVE SINK RATES SET SINKS PROPORTIONAL TO EXTERNAL SOURCES)

	-----STREAM		-----SOIL	
	TEMP DEG C	TEMP DEG C	ROOT DEP %	NITRF RATE
ANNUAL AVE	4.3	4.3	100.0	10000.000

INITIAL CONDITIONS FOR CARBON AND NITROGEN POOLS


INIT PLNT C (MOL/M2)	29.900	INIT LOM C (MOL/M2)	175.200	INIT ROM C (MOL/M2)	526.000
INIT PLNT N (MOL/M2)	0.320	INIT LOM N (MOL/M2)	4.500	INIT ROM N (MOL/M2)	27.200
INIT PLNT C/N	93.438	INIT LOM C/N	38.933	INIT ROM C/N	19.338
INIT SOIL NH4 MEQ/M3	1.000	INIT SOIL NO3 MEQ/M3	1.000		

ANNUAL AVE	-----MOL-C/M2-----		-----MOL-C/M2/YR	
	PLANT BIOMASS	LAB-OM BIOMASS	WOOD PRODUCTION	LITTER PRODUCTION
	1.00000	1.00000	1.00000	1.00000

ANNUAL AVE	-----MOL-C/M2/YR-----		
	SOIL ORGANIC EXPORT	LAB-OM DECOMPOSITION	REF-OM DECOMPOSITION
	0.03000	-90.00000	4.00000

(NEGATIVE DECOMPOSITION RATES SET DECOMPOSITION PROPORTIONAL TO INPUTS TO THE POOL)

HINDCAST WET DEPOSITION USED FOR: NH4 NO3



SEQUENCE NAME: semb

BREAK	YEAR	FACTOR
-	1957	0.000
1	1972	0.670
-	1987	1.000


CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37


** PAGE 4 **

HINDCAST PLANT BIOMASS




SEQUENCE NAME: simple2		
BREAK	YEAR	FACTOR
-	1957	1.000
1	1957	29.900
2	1967	169.600
3	1972	249.400
4	1977	199.500
5	1987	199.500
-	1987	1.000

HINDCAST LAB-OM BIOMASS




SEQUENCE NAME: simple2		
BREAK	YEAR	FACTOR
-	1957	1.000
1	1957	175.200
2	1967	175.200
3	1972	175.200
4	1977	240.000
5	1987	242.400
-	1987	1.000

HINDCAST WOOD PRODUCTION



SEQUENCE NAME: simple4		
BREAK	YEAR	FACTOR
-	1957	1.000
1	1957	5.000
2	1962	4.500
3	1970	4.000
4	1975	1.900
5	1982	5.100
6	1987	7.700
-	1987	1.000

HINDCAST LITTER PRODUCTION



SEQUENCE NAME: simple3		
BREAK	YEAR	FACTOR
-	1957	1.000
1	1957	15.000
2	1967	32.400
3	1972	39.400
4	1977	42.400
5	1987	32.400
-	1987	1.000

THERE ARE NO CHANGES IN HINDCAST DRY-DEP FACTOR FOR ANY IONS
 THERE ARE NO CHANGES IN HINDCAST SOIL SINKS FOR ANY IONS

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 5 **

THERE ARE NO CHANGES IN HINDCAST STREAM SINKS FOR ANY IONS
 THERE ARE NO CHANGES IN HINDCAST SOIL SOURCES FOR ANY IONS
 THERE ARE NO CHANGES IN HINDCAST STREAM SOURCES FOR ANY IONS

THERE ARE NO CHANGES IN HINDCAST SOIL TEMP DEG C
 THERE ARE NO CHANGES IN HINDCAST STREAM TEMP DEG C
 THERE ARE NO CHANGES IN HINDCAST DISCHARGE M/YR
 THERE ARE NO CHANGES IN HINDCAST ROOT DEPTH % SOIL
 THERE ARE NO CHANGES IN HINDCAST NITRIF RATE SOIL

THERE ARE NO CHANGES IN HINDCAST NH4 UPT, OPTIMUM C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, C/N HFWD K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, OPTIMUM C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, REFERNC K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, REFERNC C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, C/N HFWD K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, OPTIMUM C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, C/N HFWD K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, OPTIMUM C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, REFERNC K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, REFERNC C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, C/N HFWD K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, OPTIMUM C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC K2 FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, REFERNC C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NH4 UPT, C/N HFWD K2 FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, OPTIMUM C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN HINDCAST NO3 UPT, REFERNC K2 FOR HETEROTROPHS (REF. ORGANIC)

THERE ARE NO CHANGES IN HINDCAST SOIL ORGANIC EXPORT
 THERE ARE NO CHANGES IN HINDCAST LAB-OM DECOMPOSITION
 THERE ARE NO CHANGES IN HINDCAST REF-OM DECOMPOSITION

----- FORECAST -----

MERLIN VERSION 3.33 (31 OCT 94)

RUN DATE = 11/17/97

RUN TIME = 16:35:37

CATCHMENT = Nordmoen R1/R2


FORECAST SIMULATION PERIOD = 60 YEARS

MONTHLY VARIATION = N

START YEAR = 1988

END YEAR = 2047

INTEGRATION TIME STEP = 0.0417 YEARS



SEQUENCE NAME: ingvald20


BREAK	YEAR	FACTOR
-	1988	1.000
1	1988	3.130
2	2047	3.130
-	2047	0.000

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 7 **



SEQUENCE NAME: simple1

BREAK	YEAR	FACTOR
-	1988	1.000
1	1988	1.000
2	2047	0.600
-	2047	1.000

THERE ARE NO CHANGES IN FORECAST DRY-DEP FACTOR FOR ANY IONS
 THERE ARE NO CHANGES IN FORECAST SOIL SINKS FOR ANY IONS
 THERE ARE NO CHANGES IN FORECAST STREAM SINKS FOR ANY IONS
 THERE ARE NO CHANGES IN FORECAST SOIL SOURCES FOR ANY IONS
 THERE ARE NO CHANGES IN FORECAST STREAM SOURCES FOR ANY IONS

THERE ARE NO CHANGES IN FORECAST SOIL TEMP DEG C
 THERE ARE NO CHANGES IN FORECAST STREAM TEMP DEG C
 THERE ARE NO CHANGES IN FORECAST DISCHARGE M/YR
 THERE ARE NO CHANGES IN FORECAST ROOT DEPTH % SOIL
 THERE ARE NO CHANGES IN FORECAST NITRIF RATE SOIL

THERE ARE NO CHANGES IN FORECAST NH4 UPT, OPTIMUM C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, C/N HFWD K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, OPTIMUM C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, REFERNC K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, REFERNC C/N FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, C/N HFWD K2 FOR AUTOTROPHS (PLANT BIOMASS)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, OPTIMUM C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, C/N HFWD K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, OPTIMUM C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, REFERNC K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, REFERNC C/N FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, C/N HFWD K2 FOR HETEROTROPHS (LAB. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, OPTIMUM C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC K2 FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, REFERNC C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NH4 UPT, C/N HFWD K2 FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, OPTIMUM C/N FOR HETEROTROPHS (REF. ORGANIC)
 THERE ARE NO CHANGES IN FORECAST NO3 UPT, REFERNC K2 FOR HETEROTROPHS (REF. ORGANIC)

THERE ARE NO CHANGES IN FORECAST SOIL ORGANIC EXPORT
 THERE ARE NO CHANGES IN FORECAST LAB-OM DECOMPOSITION

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 8 **

OUTPUT FOR: 1957 ANNUAL AVE

				TOTAL NITROGEN (MEQ/M2) PER UNIT AREA OF COMPARTMENT				
(MEQ/M3)	PRECIP	STREAM	SOIL	INORGANIC-N TOTAL	STREAM	SOIL	ORGANIC-N	
NH4	5.8	0.3	0.3	NH4	0.0	0.1	LAB-OM	4559.6
NO3	6.1	6.4	6.4	NO3	0.0	1.3	REF-OM	27119.9
ΣNx	11.9	6.7	6.7	ΣNx	0.0	1.4	TOT-OM	31679.5
(% SAT)				AQUEOUS			PLANTS	315.0
NH4			0.0	NH4	0.0	0.1	TOTBIO	31994.5
NO3			0.0	NO3	0.0	1.3		
ΣNx			0.0	ΣNx	0.0	1.4		
TEMP DEG C		4.3	4.3	ADSORBED			C/N RATIOS	
ROOT DEP %			100.0	NH4		0.0	PLANTS	94.93
NITRF RATE /YR			10000.0	NO3		0.0	LAB-OM	38.65
NITRF Kup			2025.0	ΣNx		0.0	REF-OM	19.32
NITRF %dN			95.8					

INPUT/OUTPUT FLUXES (MEQ/M2/YR) PER UNIT AREA OF CATCHMENT

							UPTAKE OF NH4	UPTAKE OF N
	WETDEP	DRYDEP	TOTDEP	SOIL	STREAM	NTFLUX	PLANTS Kup	
NH4	5.77	0.00	5.77	0.09	0.09	-5.68	30.3	27
NO3	6.10	0.00	6.10	2.25	2.25	-3.85	44.1	43
ΣNx	11.87	0.00	11.87	2.34	2.34	-9.53	14.8	13
Q CM/YR	100.00	0.00	100.00	35.00	35.00	-65.00	PLANTS %dN	1.4
							LAB-OM	2.1
							REF-OM	0.7
								51
								32
								15

SOURCE/SINK FLUXES (MEQ/M2/YR) PER UNIT AREA OF COMPARTMENT

SOIL	SINKS	SOURCES	NITRIF	NETASS	STREAM	SINKS	SOURCES
NH4	0.00	0.00	-538.21	532.38	NH4	0.00	0.00
NO3	0.00	0.00	538.21	-540.95	NO3	0.00	0.00
ΣNx	0.00	0.00	0.00	-8.56	ΣNx	0.00	0.00

UPTAKE AND MINERALIZATION FLUXES (MEQ/M2/YR) PER UNIT AREA OF SOIL

NET ASSIM	NH4	LAB-OM	REF-OM	SOIL-OM	PLANTS	TOTAL
	337.34	203.09	540.44	-8.05	532.38	
	-277.63	-85.77	-363.40	-177.54	-540.95	
	59.71	117.33	177.04	-185.60	-8.56	
UPTAKE	NH4	11.72	3.94	15.66	8.05	23.72
	NO3	277.63	85.77	363.40	177.54	540.95
	ΣNx	289.36	89.71	379.06	185.60	564.66
MINERALIZ	NH4	349.07	207.03	556.10	0.00	556.10
	NO3	0.00	0.00	0.00	0.00	0.00
	ΣNx	349.07	207.03	556.10	0.00	556.10

ORGANIC CARBON AND NITROGEN POOLS (MOLES/M2) PER UNIT AREA OF SOIL

	PLANTS	LAB-OM	REF-OM	SOIL-OM	WOOD	LITTER	EXPORT
C	29.900	175.200	523.470	698.670	5.0	15.0	0.0
N	0.315	4.560	27.120	31.680	0.0	0.2	0.0
C/N	94.914	38.424	19.302	22.054	154.0	94.9	19.3

ORGANIC CARBON AND NITROGEN FLUXES (MOL/M2/YR) PER UNIT AREA OF SOIL

	WOOD PRD	LITTER	DCMP LOM	DCAY LOM	DCMP ROM	EXPORT	NT PRMPR
C	5.000	15.000	13.500	1.500	4.000	0.030	20.000
N	0.032	0.158	0.349	0.039	0.207	0.002	0.186
C/N	154.000	94.871	38.675	38.675	19.321	19.321	107.759

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 9 **

OUTPUT FOR: 1987 ANNUAL AVE

TOTAL NITROGEN (MEQ/M2) PER UNIT AREA OF COMPARTMENT

				INORGANIC-N			ORGANIC-N	
(MEQ/M3)	PRECIP	STREAM	SOIL	TOTAL	STREAM	SOIL		
NH4	38.0	0.4	0.4	NH4	0.0	0.1	LAB-OM	6067.6
NO3	41.0	4.5	4.5	NO3	0.0	0.9	REF-OM	24481.0
ΣNx	79.0	4.9	4.9	ΣNx	0.0	1.0	TOT-OM	30548.6
(% SAT)				AQUEOUS			PLANTS	
NH4			0.0	NH4	0.0	0.1	TOTBIO	2086.6
NO3			0.0	NO3	0.0	0.9		32635.2
ΣNx			0.0	ΣNx	0.0	1.0		
TEMP DEG C				ADSORBED			C/N RATIOS	
ROOT DEP %		4.3	4.3	NH4			PLANTS	95.65
NITRF RATE /YR			10000.0	NO3		0.0	LAB-OM	40.12
NITRF Kup			2025.0	ΣNx		0.0	REF-OM	19.61
NITRF %dN			90.4					

INPUT/OUTPUT FLUXES (MEQ/M2/YR) PER UNIT AREA OF CATCHMENT

UPTAKE OF NH4 UPTAKE OF N

							PLANTS Kup	86.1	79
	WETDEP	DRYDEP	TOTDEP	SOIL	STREAM	NTFLUX	LAB-OM	108.0	106
NH4	38.00	0.00	38.00	0.15	0.16	-37.84	REF-OM	21.3	19
NO3	41.00	0.00	41.00	1.56	1.57	-39.43	PLANTS %dN	3.8	38
ΣNx	79.00	0.00	79.00	1.71	1.72	-77.28	LAB-OM	4.8	51
Q CM/YR	100.00	0.00	100.00	35.00	35.00	-65.00	REF-OM	0.9	9

SOURCE/SINK FLUXES (MEQ/M2/YR) PER UNIT AREA OF COMPARTMENT

SOIL	SINKS	SOURCES	NITRIF	NETASS	STREAM	SINKS	SOURCES
NH4	0.00	0.00	-875.31	837.46	NH4	0.00	0.00
NO3	0.00	0.00	875.31	-914.70	NO3	0.00	0.00
ΣNx	0.00	0.00	0.00	-77.24	ΣNx	0.00	0.00

UPTAKE AND MINERALIZATION FLUXES (MEQ/M2/YR) PER UNIT AREA OF SOIL

		LAB-OM	REF-OM	SOIL-OM	PLANTS	TOTAL
NET ASSIM	NH4	679.90	194.77	874.67	-37.21	837.46
	NO3	-475.19	-85.75	-560.94	-353.76	-914.70
	ΣNx	204.71	109.02	313.73	-390.97	-77.24
UPTAKE	NH4	46.69	9.19	55.88	37.21	93.10
	NO3	475.19	85.75	560.94	353.76	914.70
	ΣNx	521.88	94.94	616.82	390.97	1007.79
MINERALIZ	NH4	726.59	203.96	930.55	0.00	930.55
	NO3	0.00	0.00	0.00	0.00	0.00
	ΣNx	726.59	203.96	930.55	0.00	930.55

ORGANIC CARBON AND NITROGEN POOLS (MOLES/M2) PER UNIT AREA OF SOIL

	PLANTS	LAB-OM	REF-OM	SOIL-OM	WOOD	LITTER	EXPORT
C	199.499	242.398	479.942	722.339	135.4	1018.7	0.9
N	2.087	6.068	24.481	30.549	0.9	10.5	0.0
C/N	95.609	39.949	19.605	23.646	154.0	96.6	19.5

ORGANIC CARBON AND NITROGEN FLUXES (MOL/M2/YR) PER UNIT AREA OF SOIL

	WOOD PRD	LITTER	DCMP LOM	DCAY LOM	DCMP ROM	EXPORT	NT PRMPR
C	7.700	32.400	29.160	3.000	4.000	0.030	40.100
N	0.050	0.339	0.727	0.075	0.204	0.002	0.391
C/N	154.000	95.656	40.133	40.133	19.612	19.612	102.565

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 10 **

OUTPUT FOR: 2037 ANNUAL AVE

TOTAL NITROGEN (MEQ/M2) PER UNIT AREA OF COMPARTMENT

(MEQ/M3)	PRECIP	STREAM	SOIL	INORGANIC-N			ORGANIC-N	
				INORGANIC-N TOTAL	STREAM	SOIL	LAB-OM	REF-OM
NH4	106.6	0.9	0.9	NH4	0.0	0.2	LAB-OM	13599.6
NO3	115.3	96.6	96.5	NO3	0.0	19.5	REF-OM	27187.5
ΣNx	222.0	97.5	97.4	ΣNx	0.0	19.7	TOT-OM	40787.1
(% SAT)				AQUEOUS			PLANTS	1951.0
NH4			0.0	NH4	0.0	0.2	TOTBIO	42738.1
NO3			0.0	NO3	0.0	19.5		
ΣNx			0.0	ΣNx	0.0	19.7		
TEMP DEG C		4.3	4.3	ADSORBED			C/N RATIOS	
ROOT DEP %			100.0	NH4		0.0	PLANTS	89.55
NITRF RATE /YR			10000.0	NO3		0.0	LAB-OM	18.33
NITRF Kup			2025.0	ΣNx		0.0	REF-OM	17.22
NITRF %dN			98.9					

INPUT/OUTPUT FLUXES (MEQ/M2/YR) PER UNIT AREA OF CATCHMENT

						UPTAKE OF NH4	UPTAKE OF N		
	WETDEP	DRYDEP	TOTDEP	SOIL	STREAM	NTFLUX			
NH4	106.65	0.00	106.65	0.32	0.33	-106.32	PLANTS Kup	4.1	3
NO3	115.34	0.00	115.34	33.78	33.79	-81.54	LAB-OM	16.4	15
ΣNx	221.99	0.00	221.99	34.10	34.12	-187.86	REF-OM	0.7	0
Q CM/YR	100.00	0.00	100.00	35.00	35.00	-65.00	PLANTS %dN	0.2	18
							LAB-OM	0.8	77
							REF-OM	0.0	3

SOURCE/SINK FLUXES (MEQ/M2/YR) PER UNIT AREA OF COMPARTMENT

SOIL	SINKS	SOURCES	NITRIF	NETASS	STREAM	SINKS	SOURCES
NH4	0.00	0.00	-1832.26	1725.93	NH4	0.00	0.00
NO3	0.00	0.00	1832.26	-1913.39	NO3	0.00	0.00
ΣNx	0.00	0.00	0.00	-187.47	ΣNx	0.00	0.00

UPTAKE AND MINERALIZATION FLUXES (MEQ/M2/YR) PER UNIT AREA OF SOIL

NET ASSIM		LAB-OM	REF-OM	SOIL-OM	PLANTS	TOTAL
		NH4	1575.15	154.50	1729.65	-3.73
	NO3	-1499.20	-60.25	-1559.45	-353.94	-1913.39
	ΣNx	75.95	94.25	170.20	-357.67	-187.47
UPTAKE	NH4	14.81	0.64	15.44	3.73	19.17
	NO3	1499.20	60.25	1559.45	353.94	1913.39
	ΣNx	1514.01	60.89	1574.90	357.67	1932.57
MINERALIZ	NH4	1589.96	155.14	1745.10	0.00	1745.10
	NO3	0.00	0.00	0.00	0.00	0.00
	ΣNx	1589.96	155.14	1745.10	0.00	1745.10

ORGANIC CARBON AND NITROGEN POOLS (MOLES/M2) PER UNIT AREA OF SOIL

	PLANTS	LAB-OM	REF-OM	SOIL-OM	WOOD	LITTER	EXPORT
C	174.665	248.376	467.680	716.056	146.0	2638.7	2.4
N	1.951	13.600	27.188	40.787	0.9	28.2	0.1
C/N	89.526	18.263	17.202	17.556	154.0	93.4	18.7

ORGANIC CARBON AND NITROGEN FLUXES (MOL/M2/YR) PER UNIT AREA OF SOIL

	WOOD PRD	LITTER	DCMP LOM	DCAY LOM	DCMP ROM	EXPORT	NT PRMPR
C	0.082	32.400	29.160	3.118	2.671	0.030	31.976
N	0.001	0.362	1.590	0.170	0.155	0.002	0.358
C/N	154.000	89.550	18.340	18.340	17.218	17.218	89.400

CATCHMENT = Nordmoen R1/R2

DATE = 11/17/97

TIME = 16:35:37

** PAGE 11 **

OUTPUT FOR: 2047 ANNUAL AVE

				TOTAL NITROGEN (MEQ/M2) PER UNIT AREA OF COMPARTMENT					
(MEQ/M3)	PRECIP	STREAM	SOIL	INORGANIC-N TOTAL	STREAM	SOIL	ORGANIC-N		
NH4	106.6	1.0	1.0	NH4	0.0	0.2	LAB-OM	14657.8	
NO3	115.3	116.1	116.0	NO3	0.0	23.5	REF-OM	27988.2	
ΣNx	222.0	117.0	117.0	ΣNx	0.0	23.7	TOT-OM	42646.1	
(% SAT)				AQUEOUS			PLANTS	1902.4	
NH4			0.0	NH4	0.0	0.2	TOTBIO	44548.5	
NO3			0.0	NO3	0.0	23.5			
ΣNx			0.0	ΣNx	0.0	23.7			
TEMP DEG C		4.3	4.3	ADSORBED			C/N RATIOS		
ROOT DEP %			100.0	NH4		0.0	PLANTS	89.16	
NITRF RATE /YR			10000.0	NO3		0.0	LAB-OM	17.08	
NITRF Kup			2025.0	ΣNx		0.0	REF-OM	16.92	
NITRF %dN			99.1						

INPUT/OUTPUT FLUXES (MEQ/M2/YR) PER UNIT AREA OF CATCHMENT							UPTAKE OF NH4		UPTAKE OF N	
	WETDEP	DRYDEP	TOTDEP	SOIL	STREAM	NTFLUX	PLANTS Kup			
NH4	106.65	0.00	106.65	0.33	0.35	-106.30	3.5		3	
NO3	115.34	0.00	115.34	40.61	40.62	-74.72	14.7		13	
ΣNx	221.99	0.00	221.99	40.94	40.96	-181.02	0.4		0	
Q CM/YR	100.00	0.00	100.00	35.00	35.00	-65.00	0.2		17	
							LAB-OM		78	
							REF-OM		2	

SOURCE/SINK FLUXES (MEQ/M2/YR) PER UNIT AREA OF COMPARTMENT							
SOIL	SINKS	SOURCES	NITRIF	NETASS	STREAM	SINKS	SOURCES
NH4	0.00	0.00	-1937.32	1831.00	NH4	0.00	0.00
NO3	0.00	0.00	1937.32	-2011.68	NO3	0.00	0.00
ΣNx	0.00	0.00	0.00	-180.68	ΣNx	0.00	0.00

UPTAKE AND MINERALIZATION FLUXES (MEQ/M2/YR) PER UNIT AREA OF SOIL						
NET ASSIM		LAB-OM	REF-OM	SOIL-OM	PLANTS	TOTAL
	NH4	1692.93	141.37	1834.30	-3.30	1831.00
	NO3	-1609.63	-45.89	-1655.52	-356.16	-2011.68
	ΣNx	83.30	95.49	178.78	-359.46	-180.68
UPTAKE	NH4	14.04	0.42	14.46	3.30	17.76
	NO3	1609.63	45.89	1655.52	356.16	2011.68
	ΣNx	1623.68	46.30	1669.98	359.46	2029.44
MINERALIZ	NH4	1706.97	141.79	1848.76	0.00	1848.76
	NO3	0.00	0.00	0.00	0.00	0.00
	ΣNx	1706.97	141.79	1848.76	0.00	1848.76

ORGANIC CARBON AND NITROGEN POOLS (MOLES/M2) PER UNIT AREA OF SOIL							
	PLANTS	LAB-OM	REF-OM	SOIL-OM	WOOD	LITTER	EXPORT
C	169.597	249.595	473.338	722.933	147.0	2962.7	2.7
N	1.902	14.658	27.988	42.646	1.0	31.9	0.1
C/N	89.149	17.028	16.912	16.952	153.6	93.0	18.5

ORGANIC CARBON AND NITROGEN FLUXES (MOL/M2/YR) PER UNIT AREA OF SOIL							
	WOOD PRD	LITTER	DCMP LOM	DCAY LOM	DCMP ROM	EXPORT	NT PRMPR
C	0.100	32.400	29.160	3.118	2.400	0.030	31.993
N	0.001	0.363	1.707	0.183	0.142	0.002	0.359
C/N	92.400	89.166	17.083	17.083	16.926	16.926	89.003

Naturens Tålegrenser - Oversikt over utgitte rapporter

- 1 Nygaard, P. H., 1989. Forurensningers effekt på naturlig vegetasjon en litteraturstudie. Norsk institutt for skogforskning (NISK), Ås.

- Uten nr. Jaworovski, Z., 1989. Pollution of the Norwegian Arctic: A review. Norsk polarinstitutt (NP), rapportserie nr. 55. Oslo.

- 2 Henriksen, A., Lien, L. & Traaen, T.S. 1990. Tålegrenser for overflatevann. Kjemiske kriterier for tilførsler av sterke syrer. Norsk institutt for vannforskning (NIVA), O-89210.

- 3 Lien, L., Henriksen, A., Raddum, G. & Fjellheim, A. 1989. Tålegrenser for overflatevann. Fisk og evertebrater. Foreløpige vurderinger og videre planer. Norsk institutt for vannforskning (NIVA), O-89185.

- 4 Bølviken, B. & medarbeidere, 1990. Jordforsuringsstatus og forsurningsfølsomhet i naturlig jord i Norge. Norges geologiske undersøkelse (NGU), NGU-rapport 90.156. 2 bind (Bind I: Tekst, Bind II: Vedlegg og bilag).

- 5 Pedersen, H. C. & Nybø, S. 1990. Effekter av langtransporterte forurensninger på terrestriske dyr i Norge. En statusrapport med vekt på SO₂, NO_x og tungmetaller. Norsk institutt for naturforskning (NINA), Utredning 005.

- 6 Frisvoll, A. A., 1990. Moseskader i skog i Sør-Norge. Norsk institutt for naturforskning (NINA), Oppdragsmelding 018.

- 7 Muniz, I. P. & Aagaard, K. 1990. Effekter av langtransportert forurensning på ferskvannsdyr i Norge - virkninger av en del sporelementer og aluminium. Norsk institutt for naturforskning (NINA), Utredning 013.

- 8 Hesthagen, T., Berger, H. M. & Kvenild, L. 1992. Fiskestatus i relasjon til forurensning av innsjøer. Norsk institutt for naturforskning (NINA), Forskningsrapport 032.

- 9 Pedersen, U., Walker, S.E. & Kibsgaard, A. 1990. Kart over atmosfærisk avsetning av svovel- og nitrogenforbindelser i Norge. Norsk institutt for luftforskning (NILU), OR 28/90.

- 10 Pedersen, U. 1990. Ozonkonsentrasjoner i Norge. Norsk institutt for luftforskning (NILU), OR 28/90.

- 11 Wright, R. F., Stuanes, A. Reuss, J.O. & Flaten, M.B. 1990. Critical loads for soils in Norway. Preliminary assessment based on data from 9 calibrated catchments. Norsk institutt for vannforskning (NIVA), O-89153.

- 11b Reuss, J. O., 1990. Critical loads for soils in Norway. Analysis of soils data from eight Norwegian catchments. Norsk institutt for vannforskning (NIVA), O-89153.

- 12 Amundsen, C. E., 1990. Bufferprosent som parameter for kartlegging av forsurningsfølsomhet i naturlig jord. Universitetet i Trondheim, AVH (stensil).

- 13 Flatberg, K.I, Foss, B., Løken, A. & Saastad, S.M. 1990. Moseskader i barskog. Direktoratet for naturforvaltning (DN), notat.

-
- 14 Frisvoll, A.A., & Flatberg, K.I., 1990. Moseskader i Sør-Varanger. Norsk institutt for naturforskning (NINA), Oppdragsmelding 55.
 - 15 Flatberg, K.I., Bakken, S., Frisvoll, A.A., & Odasz, A.M. 1990. Moser og luftforurensninger. Norsk institutt for naturforskning (NINA), Oppdragsmelding 69.
 - 16 Mortensen, L.M. 1991. Ozonforurensning og effekter på vegetasjonen i Norge. Norsk landbruksforsk. 5:235-264.
 - 17 Wright, R.F., Stuanes, A.O. & Frogner, T. 1991. Critical Loads for Soils in Norway Nordmoen. Norsk institutt for vannforskning (NIVA), O-89153.
 - 18 Pedersen, H.C., Nygård, T., Myklebust, I. og Sæther, M. 1991. Metallbelastninger i lirype. Norsk institutt for naturforskning (NINA), Oppdragsmelding 71.
 - 19 Lien, L., Raddum, G.G. & Fjellheim, A. 1991. Tålegrenser for overflatevann evertebrater og fisk. Norsk institutt for vannforskning (NIVA), Rapport 0-89185,2.
 - 20 Amundsen, C.E. 1992. Sammenligning av parametre for å bestemme forsuringfølsomhet i jord. NGU-rapport 91.265.
 - 21 Bølviken, B., R. Nilsen, J. Romundstad & O. Wolden. 1992. Surhet, forsuringfølsomhet og lettløselige basekationer i naturlig jord fra Nord-Trøndelag og sammenligning med tilsvarende data fra Sør Norge. NGU-rapport 91.250.
 - 22 Sivertsen, T. & medarbeidere. 1992. Opptak av tungmetaller i dyr i Sør-Varanger. Direktoratet for naturforvaltning, DN-notat 1991-15.
 - 23 Lien, L., Raddum, G.G. & A. Fjellheim. 1992. Critical loads of acidity to freshwater. Fish and invertebrates. Norwegian Institute for Water Research (NIVA), Rapport O-89185,3.
 - 24 Fremstad, E. 1992. Virkninger av nitrogen på heivegetasjon. En litteraturstudie. Norsk institutt for naturforskning (NINA), Oppdragsmelding 124.
 - 25 Fremstad, E. 1992. Heivegetasjon i Norge, utbredelseskart. Norsk institutt for naturforskning (NINA), Oppdragsmelding 188.
 - 26 Flatberg, K.I. & Frisvoll, A. 1992. Undersøkelser av skader hos to sigdmoser i Agder. Norsk institutt for naturforskning (NINA), Oppdragsmelding 134.
 - 27 Lindstrøm, E.A. 1992. Tålegrenser for overflatevann. Fastsittende alger. Norsk institutt for vannforskning (NIVA), O-90137/E-90440, rapport-2.
 - 28 Brettum, P. 1992. Tålegrenser for overflatevann. Planteplankton. Norsk institutt for vannforskning (NIVA), O-90137/E-90440, rapport-3.
 - 29 Brandrud, T.E., Mjelde, M. 1992. Tålegrenser for overflatevann. Makrovegetasjon. Norsk institutt for vannforskning (NIVA), O-90137/E-90440, rapport-1.
 - 30 Mortensen, L.M. & Nilsen, J. 1992. Effects of ozone and temperature on growth of several wild plant species. Norwegian Journal of Agricultural Sciences 6: 195-204.
-

-
- 31 Pedersen, H.C., Myklebust, I., Nygård, T. & Sæther, M. 1992. Akkumulering og effekter av kadmium i lirype. Norsk institutt for naturforskning (NINA), Oppdragsmelding 152.
- 32 Amundsen, C.E. 1992. Sammenligning av relativ forsurningsfølsomhet med tålegrenser beregnet med modeller, i jord. Norges geologiske undersøkelse. NGU-rapport 92.294.
- 33 Frogner, T., Wright, R.F., Cosby, B.J., Esser, J.M., Håøya, A.-O. & Rudi, G. 1992. Map of critical loads for coniferous forest soils in Norway. Norsk institutt for vannforskning (NIVA), O-91147.
- 34 Henriksen, A., Lien, L., Traaen, T.S. & Taubøll, S. 1992. Tålegrenser for overflatevann - Kartlegging av tålegrenser og overskridelser av tålegrenser for tilførsler av sterke syrer. Norsk institutt for vannforskning (NIVA), O-89210.
- 35 Lien, L. Henriksen, A. & Traaen, T.S. 1993. Tålegrenser for sterke syrer på overflatevann - Svalbard. Norsk institutt for vannforskning (NIVA), O-90102.
- 36 Henriksen, A., Hesthagen, T., Berger, H.M., Kvenild, L., Taubøll, S. 1993. Tålegrenser for overflatevann - Sammenheng mellom kjemisk kriterier og fiskestatus. Norsk institutt for vannforskning (NIVA), O-92122.
- 37 Odasz, A.M., Øiesvold, S., & Vange, V. 1993. Nitrate nutrition in *Racomitrium lanuginosum* (Hedw.)Brd., a bioindicator of nitrogen deposition in Norway. Direktoratet for naturforvaltning. Utredning for DN 1993-2.
- 38 Espelien, I.S. 1993. Genetiske effekter av tungmetaller på pattedyr. En kunnskapsoversikt. Norsk institutt for naturforskning (NINA), Utredning 051.
- 39 Økland, J. & Økland, K.A. 1993. Database for bioindikatorer i ferskvann - et forprosjekt . Laboratorium for ferskvannsekologi og innlandsfiske (LFI), Zoologisk Museum, Oslo, Rapport 144, 1993.
- 40 Aamlid, D. & Skogheim, I. 1993. Nikkel, kopper og andre metaller i multer og blåbær fra Sør-Varanger, 1992. Rapport Gkogforsk 14/93. 14/93.
- 41 Kålås, J.A., Ringsby, T.H. & Lierhagen, S. 1993. Metals and radiocesium in wild animals from the Sør-Varanger area, north Norway. Norsk institutt for naturforskning (NINA), Oppdragsmelding 212.
- 42 Fløisand, I. & Løbersli, E. (red.)1993. Tilførsler og virkninger av lufttransporterte forurensninger (TVLF) og Naturens tålegrenser. Sammendrag av foredrag og postere fra møte i Stjørdal, 15.-17.februar 1993. Norsk institutt for luftforskning (NILU), OR 17/93.
- 43 Henriksen, A. & Hesthagen, T. 1993. Critical load exceedance and damage to fish populations. Norsk institutt for vannforskning (NIVA), O-89210.
- 44 Lien, L., Henriksen, A. & Traaen, T.S. 1993. Critical loads of acidity to surface waters, Svalbard. Norsk institutt for vannforskning (NIVA), O-90102.
- 45 Løbersli, E., Johannessen, T. & Olsen, K.V (red.) 1993. Naturens tålegrenser. Referat fra seminar i 1991 og 1992. Direktoratet for naturforvaltning, DN-notat 1993-6.
-

-
- 46 Bakken, S. 1993. Nitrogenforurensning og variasjon i nitrogen, protein og klorofyllinnhold hos barskogsmosen blanksigd (*Dicranum majus*). Direktoratet for naturforvaltning (DN). Utredning for DN 1994-1.
- 47 Krøkje, Å. 1993. Genotoksisk belastning i jord . Effekstudier, med mål å komme fram til akseptable grenser for genotoksisk belastning fra langtransportert luftforurensning. Direktoratet for naturforvaltning (DN). Utredning for DN 1994-2.
- 48 Fremstad, E. 1993. Heigråmose (*Racomitrium lanuginosum*) som indikator på nitrogenbelastning. Norsk institutt for naturforskning (NINA) Oppdragsmelding 239.
- 49 Nygaard, P.H. & Ødegaard, T.H. 1993. Effekter av nitrogengjødsling på vegetasjon og jord i skog. Rapport Skogforsk 26/93.
- 50 Fløisand, I. og Johannessen, T. (red.) 1994. Langtransporterte luftforurensninger. Tilførsler, virkninger og tålegrenser. Sammendrag av foredrag og postere fra møte i Grimstad, 7.-9.3.94. Norsk institutt for luftforskning NILU OR: 17/94
- 51 Kleivane, L. Skåre, J.U. & Wiig, Ø. 1994. Klorerte organiske miljøgifter i isbjørn. Forekomst, nivå og mulige effekter. Norsk Polarinstitutt Meddelelse nr. 132.
- 52 Lydersen, E., Fjeld, E. & Andersen, T. 1994. Fiskestatus og vannkjemi i norske innsjøer. Norsk institutt for vannforskning (NIVA) O-93172
- 53 Schartau, A.K.L. (red.) 1994. Effekter av lavdose kadmium-belastning på littorale ferskvanns-populasjoner og -samfunn. Norsk institutt for naturforskning (NINA) Forskningsrapport 055.
- 54 Mortensen, L. (1994). Variation in ozone sensitivity of *Betula pubescens* Erh. from different sites in South Norway. Direktoratet for naturforvaltning (DN). Utredning for DN, Nr. 1994-6.
- 55 Mortensen, L. (1994). Ozone sensitivity of *Phleum alpinum* L. from different locations in South Norway. Direktoratet for naturforvaltning (DN). Utredning for DN, Nr. 1994-7.
- 56 Frogner, T., Wright, R.F., Cosby, J.B. and Esser, J.M. (1994). Maps of critical loads and exceedance for sulfur and nitrogen to forest soils in Norway. Norsk institutt for vannforskning (NIVA) O-91147.
- 57 Flatberg, K.I. & Frisvoll, A.A. 1994. Moseskader i Agder 1989-92 (1994). Norsk institutt for naturforskning (NINA), Oppdragsmelding 298.
- 58 Hesthagen, T. & Henriksen, A. (1994). En analyse av sammenhengen mellom overskridelser av tålegrenser for overflatevann og skader på fiskebestander. Norsk institutt for naturforskning (NINA), Oppdragsmelding 288.
- 59 Skåre, J.U., Wiig, Ø. & Bernhoft, A. (1994). Klorerte organiske miljøgifter; nivåer og effekter på isbjørn. Norsk Polarinstitutt Rapport nr. 86 - 1994.
- 60 Tørseth, K. & Pedersen, U. 1994. Deposition of sulphur and nitrogen components in Norway. 1988-1992. Norsk institutt for luftforskning (NILU): OR 16/94.
-

-
- 61 Nygaard, P.H. 1994. Virkning av ozon på blåbær (*Vaccinium myrtillus*), etasjehusmose (*Hylocomium splendens*), furumose (*Pleurozium schreberi*) og krussigd (*Dicranum polysetum*). Rapport Skogforsk 9/94.
- 62 Henriksen, A. & Lien, L. 1994. Tålegrenser for overflatevann: Metode og usikkerheter. Norsk institutt for vannforskning (NIVA), O-94122.
- 63 Hilmo, O. & Larssen, H.C. 1994. Morfologi hos epifyttisk lav i områder med ulik luftkvalitet. ALLFORSK Rapport 2.
- 64 Wright, R.F. 1994. Bruk av dynamiske modeller for vurdering av vann- og jordforsuring som følge av redusert tilførsel av sur nedbør. Norsk institutt for vannforskning (NIVA), O-94112.
- 65 Hesthagen, T., A. Henriksen & Kvenild, L. 1994. Overskridelser av tålegrenser for overflatevann og skader på fiskebestander i norske innsjøer med spesiell vekt på Troms og Finnmark. Norsk institutt for naturforskning (NINA), Oppdragsmelding 298.
- 66 Sagmo Solli, I.M, Flatberg, K.I.F., Söderström, L., Bakken S. & Pedersen, B. 1996. Blanksigd og luftforurensningsstudier. NTNU. Vitenskapsmuseet. Rapport botanisk serie 1996-1.
- 67 Stuanes, A. & Abrahamsen, G. 1996. Tålegrenser for nitrogen i skog - en vurdering av kunnskapsgrunnlaget. Aktuelt fra Skogforsk 7-96.
- 68 Ogner, G. 1995. Tålegrenser for skog i Norge med hensyn til ozon. Aktuelt fra Skogforsk 3-95.
- 69 Thomsen, M., Nellemann, C. Frogner, T., Henriksen A., Tomter, S. & Mulder, J. 1995. Tilvekst og vitalitet for granskog sett i relasjon til tålegrenser og forurensning. Rapport fra Skogforsk 22-95.
- 70 Tomter, S. M. & Esser, J. 1995. Kartlegging av tålegrenser for nitrogen basert på en empirisk metode. Norsk institutt for jord- og skogkartlegging (NIJOS). Rapport nr 10/95.
- 71 Pedersen, H.Chr. (red.). 1995. Kadmium og bly i lirype: akkumulering og cellulære effekter. Stiftelsen for naturforskning og kulturminneforskning (NINA-NIKU) Oppdragsmelding 387
- 72 Bakken, S. & Flatberg, K.I.F. 1995. Effekter av økt nitrogendeposisjon på ombrotrof myrvegetasjon. En litteraturstudie. ALLFORSK Rapport 3.
- 73 Sogn, T.A., Stuanes, A.O. & Abrahamsen, G. 1995. Akkumulering av nitrogen - en kritisk parameter for beregning av tålegrenser for nitrogen i skog. Rapport fra Skogforsk 21/95.
- 74 Nygaard, P.H. & Eldhuset, T. 1996. Forholdet mellom basekationer og aluminium i jordløsning som kriterium for tålegrenser i skogsjord. Norsk institutt for skogforskning (NISK). Rapport fra Skogforsk 1/96
- 75 Mortensen, L. 1993. Effects of ozone on growth of several subalpine plant species. *Norw. J. Agric. Sci.* 7:129-138.
- 76 Mortensen, L. 1994. Further studies on the effects of ozone concentration on growth of subalpine plant species. *Norw. J. Agric. Sciences* 8:91-97.
-

-
- 77 Fløisand, I. & Løbersli, E. (red.) 1996. Lufttransporterte forurensninger - tilførsler, virkninger og tålegrenser. Norsk institutt for luftforskning (NILU) OR 2/96.
- 78 Thomsen, M.G., Esser, J., Venn, K. & Aamlid, D. 1996. Sammenheng mellom træs vitalitet og næringsstatus i nåler og humus på skogovervåkingsflater i Sørøst-Norge (in prep).
- 79 Tørseth, K., Mortensen, L. & Hjellbrekke, A.-G. 1996. Kartlegging av bakkenær ozon etter tålegrenser basert på akkumulert dose over 40 ppb. Norsk institutt for luftforskning (NILU) OR 12/96.
- 80 Esser, J.M. & Tomter, S.M. 1996. Reviderte kart for tålegrenser for nitrogen basert på empiriske verdier for ulike vegetasjonstyper. Norsk institutt for jord- og skogkartlegging (NIJOS).
- 81 Henriksen, A., Hindar, A., Styve, H., Fjeld, E. & Lien, L. 1996. Forsuring av overflatevann, beregningsmetodikk, trender og mottiltak. Norsk institutt for vannforskning (NIVA). Rapport LNR 3528-96.
- 82 Henriksen, A., Hesthagen, T. & Fjeld, E. 1996. Overskridelser av tålegrenser for overflatevann og skader på fiskebestander. Norsk institutt for vannforskning (NIVA). Rapport LNR 3565-96.
- 83 Wright, R. F., Raastad, I.A., & Kaste, Ø. 1996. Atmospheric deposition of nitrogen, runoff of organic nitrogen, and critical loads for soils and waters. Norsk institutt for vannforskning (NIVA) Report SNO 3592-97
- 84 Mortensen, L.M. 1995. The influence of ozone pollution on growth of young plants of *Betula pubescens* Ehrh. And *Phleum alpinum* L. Dose-response relations. *Norw. J. Agr. Sci.* 9:249-262
- 85 Mortensen, L.M. 1996. Ozone sensitivity of *Betula pubescens* at different growth stages after budburst in spring. *Norw. J. Agr. Sci.* 10:187-196.
- 86 Tørseth, K., Rosendahl, K.E., Hansen, A.C., Høie, H. & Mortensen, L.M. 1997. Avlingstap som følge av bakkenært ozon. Vurderinger for perioden 1989-1993. SFT-rapport.
- 87 Rognerud, S, Hognve, D. & Fjeld, E. 1997. Naturlige bakgrunnskonsentrasjoner av metaller. Kan atmosfæriske avsetninger påvirke metall-konsentrasjoner slik at det ikke reflekterer berggrunnens geokjemi? Norsk institutt for vannforskning (NIVA) Rapport LNR 3670-97
- 88 Skjelkvåle, B.L., Wright, R.F. & Tjomsland, T. 1997. Vannkjemi, forsuringsstatus og tålegrenser i nasjonalparker; Femundsmarka og Rondane. Norsk institutt for vannforskning (NIVA) Rapport LNR 3646-97
- 89 Nordbakken; J.-F. 1997. Småskalaendringer i ombrotrof myrvegetasjon i SØ-Norge 1990/91-96. Botanisk Hage og Museum, Univ. Oslo Rapp. 1
- 90 Sogn, T.A., Kjønås, J., Stuanes, A.O., & Abrahamsen, G. 1997. Akkumulering av nitrogen - variasjoner avhengig av bestandsutvikling, nitrogentilførsel og simulert snødekke. Norges Landbrukshøgskole, Institutt for jord- og vannfag, Rapport nr. 10/97.
-

- 91 Nygaard, P.H., Ødegård, T. & Flatberg, K.I.F. Vegetasjonsendringer over 60 år i fattig skog- og myrvegetasjon i Karlshaugen skogreservat. Skogforsk (in prep)
- 92 Knutzen, J., Gabrielsen, G.W., Henriksen, O.E., Hylland, K., Källqvist, T. Nygård, T., Pacyna, J.S. Skjegstad, N. & Steinnes, E. 1997. Assessment of the applicability for pollution authorities of the concept "critical load" of long-range transported micropollutants in relation to aquatic and terrestrial ecosystems. Norsk institutt for vannforskning (NIVA) Report SNO 3751-97.
- 93 Tørseth, K. & Semb, A. 1997. Deposition of major inorganic components in Norway 1992-1996. Norsk institutt for luftforskning (NILU), OR 67/97.
- 94 Henriksen, A. 1998. Application of the first order acidity balance (FAB) model to Norwegian surface waters. Norsk institutt for vannforskning (NIVA)
- 95 Sogn, T.A. & Wright, R.F. 1998. The model MERLIN applied to Nordmoen, Norway. Norsk institutt for vannforskning (NIVA) Report SNO 3844-98

Henvendelser vedrørende rapportene rettes til utførende institusjon