

Available NIVA generated ecotoxdata with  
relevance to petroleum related chemical  
compounds



<b>Main Office</b> Gautstadalleen 21 NO-0349 Oslo, Norway Phone (47) 22 18 51 00 Telefax (47) 22 18 52 00 Internet: www.niva.no	<b>Regional Office, Sørlandet</b> Jon Lilletuns vei 3 NO-4879 Grimstad, Norway Phone (47) 22 18 51 00 Telefax (47) 37 04 45 13	<b>Regional Office, Østlandet</b> Sandvikaveien 59 NO-2312 Ottestad, Norway Phone (47) 22 18 51 00 Telefax (47) 62 57 66 53	<b>Regional Office, Vestlandet</b> Thormøhlens gate 53 D NO-5006 Bergen Norway Phone (47) 22 18 51 00 Telefax (47) 55 31 22 14	<b>Regional Office Central</b> Pirsenteret, Havnegata 9 P.O.Box 1266 NO-7462 Trondheim Phone (47) 22 18 51 00 Telefax (47) 73 54 63 87
--	--	---	--	---

Title Available NIVA generated ecotoxdata with relevance to petroleum related chemical compounds	Report No.. 6112-2011	Date 31.01.2011
	Project No. 11061	Pages      Price 22
Author(s) Eivind Farmen	Topic group Ecotoxicology	Distribution Open
	Geographical area Norway	Printed NIVA

Client(s) Akvaplan-NIVA, JoLynn Carroll	Client ref.
--	-------------

**Abstract**  
 This report represents an overview of ecotoxicological data of petroleum-related chemical compounds generated in various NIVA projects. Specific emphasis was put on the sub-Arctic and Arctic marine environment, for applicability to the planned SYMBIOSES model system.

4 keywords, Norwegian 1.      Giftighet 2.      Olje 3.      Produert vann 4.      Arktis	4 keywords, English 1.      Toxicity 2.      Oil 3.      Produced water 4.      Arctic
---	--



*Eivind Farmen*  
Project Manager



*Kevin Thomas*  
Research Manager



*Bjørn Faafeng*  
Senior Adviser

## **Symbioses report**

Available NIVA generated ecotoxdata with relevance to petroleum related chemical compounds

## **Preface**

The work described in this report has been performed on request from Akvaplan-niva, and represents an overview of ecotoxicological data of petroleum-related chemical compounds generated from various NIVA projects. Specific emphasis was put on the sub-Arctic and Arctic marine environment, for applicability to the planned SYMBIOSES model system.

Oslo, 28 January 2011

*Eivind Farmen*

---

# Contents

<b>Summary</b>	<b>5</b>
<b>1. Introduction</b>	<b>6</b>
<b>2. Ecotoxicity testing</b>	<b>7</b>
2.1 Acute toxicity tests	7
2.1.1 Acute toxicity test with the marine algae <i>Skeletonema costatum</i>	7
2.1.2 Acute toxicity test with <i>Daphnia magna</i>	10
2.1.3 Acute toxicity test with fish	10
2.2 Chronic toxicity tests	10
2.2.1 Sediment macroinvertebrate communities	10
2.2.2 Sediment macroinvertebrate <i>Hediste diversicolor</i>	11
2.2.3 Chronic mussel lab exposure: <i>Mytilus edulis</i>	11
2.2.4 Chronic fish lab exposure: <i>Danio rerio</i>	12
2.2.5 Chronic fish lab exposure: <i>Gadhus morhua</i>	12
2.2.6 Chronic mussel field exposure: <i>Mytilus edulis</i>	13
2.2.7 Chronic fish field exposure: <i>Gadhus morhua</i>	14
2.2.8 In vitro systems: Yeast estrogen and androgen screen	14
2.2.9 In vitro systems: primary hepatocytes from <i>Onchorynchus mykiss</i>	17
<b>3. References</b>	<b>20</b>

---

## **Summary**

This report represents an overview of ecotoxicological data of petroleum-related chemical compounds generated in various NIVA projects. Specific emphasis was put on the sub-Arctic and Arctic marine environment, for applicability to the planned SYMBIOSES model system.

## **Sammendrag**

Denne rapporten representerer en oversikt over økotoksikologiske data for petroleum-relaterte kjemiske forbindelser, generert i ulike NIVA prosjekter. Spesiell vekt ble lagt på sub-arktiske og arktiske marine miljø, for innkorporering i det planlagte SYMBIOSES modellsystemet.

# 1. Introduction

This project memo describes an overview of the relevant ecotoxicology data related to petroleum-related chemical compounds generated through projects conducted with NIVA as a partner during the last decade. The memo is as up-dated as possible within the time frame and resources allocated for the work organized by Akvaplan-NIVA within the SYMBIOSES project, and the overview will also be presented at the SYMBIOSES Workshop on Ecotoxicology Meta-data Analysis Exercise on 28 February – 1 March 2011.

The data generated through the various NIVA projects are based on standard experimental systems such as the marine algae *Skeletonema costatum* and the crustacean *Daphnia magna*, but also systems such as fish primary cell culture (*Oncorhynchus mykiss*), in vivo fish experiments with zebrafish *Danio rerio*, Atlantic cod *Gadus morhua*, as well as benthic invertebrate community structure. *Skeletonema costatum* are used as the test organism as they are easily cultured within the laboratory and are recommended for use in marine ecotoxicity assessments (ISO 10253, 2006. Water quality – Marine algal growth inhibition test with *Skeletonema costatum* and *Phaeodactylum tricorutum*). NIVA has worked with *S. costatum* in toxicity studies since 1992. *Daphnia magna* is the one of the most widely used organism for toxicity tests and several guidelines for both acute and chronic testing exists (OECD 202, OECD 211, ISO 6341). It is easy to keep in culture, and it reproduces by parthenogenesis making it ideal as a test organism. NIVA has worked with *D. magna* in toxicity studies since 1992.

Compounds tested vary from dispersants and drilling muds to organic extracts of produced water and synthetic produced water, reflecting the water soluble components of oil.

## 2. Ecotoxicity testing

### 2.1 Acute toxicity tests

#### 2.1.1 Acute toxicity test with the marine algae *Skeletonema costatum*

*Skeletonema costatum* are used as the test organism as they are easily cultured within the laboratory and are recommended for use in marine ecotoxicity assessments (ISO 10253, 2006. Water quality – Marine algal growth inhibition test with *Skeletonema costatum* and *Phaeodactylum tricornutum*).

NIVA has worked with *S. costatum* in toxicity studies since 1992.

A range of tests on dispersants are summarized in NIVA report 3128. Data for comparable tests with *Chlamydomonas reinhardtii*, where the dispersants were tested together with oil, are also present, but not summarized here due to the higher sensitivity and relevance of the marine algae *Skeletonema costatum*. In addition tests of drilling mud from well 7816/12-1 Rendalspasset, Svalbard was also extracted from a different report

Dispersant Arrow Emulsol LW

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	5-100 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 8.4 mg/l, EC50: 29 mg/l
<b>Data format</b>	NIVA Report 3128

Dispersant Corexit 9527

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-30 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 3.2 mg/l, EC50: 13 mg/l
<b>Data format</b>	NIVA Report 3128

Dispersant Dasic Slickgone LTS

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-30 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 11 mg/l, EC50: 15 mg/l
<b>Data format</b>	NIVA Report 3128

Dispersant DisPollene 36S

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-30 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ±



	1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 11 mg/l, EC50: 20 mg/l
<b>Data format</b>	NIVA Report 3128

Dispergant Enersperse 1037

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	1.5-55 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 11 mg/l, EC50: 13 mg/l
<b>Data format</b>	NIVA Report 3128

Dispergant Enersperse 1583

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-20 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 5.8 mg/l, EC50: 8.2 mg/l
<b>Data format</b>	NIVA Report 3128

Dispergant Finasol OSR-5

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	1.5-55 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 10 mg/l, EC50: 13 mg/l
<b>Data format</b>	NIVA Report 3128

Dispergant Finasol OSR-12

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-100 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 10 mg/l, EC50: 24 mg/l
<b>Data format</b>	NIVA Report 3128

Dispergant IKU-9

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-55 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C

<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 18 mg/l, EC50: 25 mg/l
<b>Data format</b>	NIVA Report 3128

## Dispergant Quell Oil C1

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	15-600 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 82 mg/l, EC50: 180 mg/l
<b>Data format</b>	NIVA Report 3128

## Dispergant Shell Dispersant VDC

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-55 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 16 mg/l, EC50: 23 mg/l
<b>Data format</b>	NIVA Report 3128

## Dispergant BP1100 WD

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3-55 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 23 mg/l, EC50: 49 mg/l
<b>Data format</b>	NIVA Report 3128

## Dispergant Ameroid OSD/LT

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	15-1000 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	EC10: 140 mg/l, EC50: 470 mg/l
<b>Data format</b>	NIVA Report 3128

## Dispergant Biosolve

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	1.5-20 mg/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253, 20 ± 1 °C
<b>Target effects</b>	Growth rate

<b>Type of data available</b>	EC10: 2.4 mg/l, EC50: 3.7 mg/l
<b>Data format</b>	NIVA Report 3128

Drilling mud from well 7816/12-1 Rendalspasset, Svalbard was also tested in a different report:

<b>Species and life stage</b>	<i>Skeletonema costatum</i>
<b>Exposure time</b>	3 days (72 h)
<b>Exposure concentration</b>	3.2, 5.6, 10,18, 32, 56, 100 g/l
<b>Experiment description</b>	Marine algal inhibition test ISO DP 10253
<b>Target effects</b>	Growth rate
<b>Type of data available</b>	NOEC: 10 g/l, EC10: 13 g/l, EC50: 49 g/l
<b>Data format</b>	NIVA Report 2593

### 2.1.2 Acute toxicity test with *Daphnia magna*

*Daphnia magna* is the one of the most widely used organism for toxicity tests and several guidelines for both acute and chronic testing exists (OECD 202, OECD 211, ISO 6341). It is easy to keep in culture, and it reproduces by parthenogenesis making it ideal as a test organism. NIVA has worked with *D. magna* in toxicity studies since 1992.

Drilling mud from well 7816/12-1 Rendalspasset, Svalbard

<b>Species and life stage</b>	<i>Daphnia magna</i>
<b>Exposure time</b>	2 days (48 h)
<b>Exposure concentration</b>	3.2, 5.6, 10,18 g/l
<b>Experiment description</b>	<i>Daphnia</i> mobility inhibition test ISO6341
<b>Target effects</b>	Immobility/mortality
<b>Type of data available</b>	LC 20: 14 g/l, LC50: 23 g/l
<b>Data format</b>	NIVA Report 2593

### 2.1.3 Acute toxicity test with fish

Drilling mud from well 7816/12-1 Rendalspasset, Svalbard

<b>Species and life stage</b>	<i>Salmo trutta</i> , 0+, average 3.3 g
<b>Exposure time</b>	4 days (96 h) semi static, renewal every 24 h
<b>Exposure concentration</b>	10 g/l
<b>Experiment description</b>	OECD guidelines for testing of chemicals, No 201: Fish acute toxicity test, 10 ± 1 °C, pH=6.3, conductivity: 3.2 ms, hardness: 11 mg CaCO <sub>3</sub> /l
<b>Target effects</b>	Mortality
<b>Type of data available</b>	No mortality observed
<b>Data format</b>	NIVA Report 2593

## 2.2 Chronic toxicity tests

### 2.2.1 Sediment macroinvertebrate communities

Olefin (Ultidril, Novaplus) and ester (Anco Green, Pterofree) based drilling muds:

<b>Species and life stage</b>	Sediment communities
<b>Exposure time</b>	6 months

<b>Exposure concentration</b>	Approx 500 g slurry, containing 1-1.5 g drilling fluid. Approx 1.5 mm layer
<b>Experiment description</b>	Drill cuttings from drilling sites in the North Sea. Undisturbed communities transferred from Oslofjord, 200 m depth. 7-11 °C, salinity: 32-35 PSU
<b>Target effects</b>	Sediment community effect
<b>Type of data available</b>	Macrofauna community structure
<b>Data format</b>	NIVA rapport 3475-96

Olefin and ester based drilling muds:

<b>Species and life stage</b>	Sediment communities
<b>Exposure time</b>	3 months
<b>Exposure concentration</b>	0.5, 5, 20 mg OP cm <sup>-2</sup>
<b>Experiment description</b>	Undisturbed communities transferred from Roddnessjøen, Porsangen (Finnmark) 120 m depth, -1.5 °C and Oslofjord, 212 m depth. 7°C
<b>Target effects</b>	Sediment community effect
<b>Type of data available</b>	Macrofauna number of species and individuals, diversity
<b>Data format</b>	NIVA rapport 3460-97

Water based drill cuttings:

<b>Species and life stage</b>	Sediment communities
<b>Exposure time</b>	6 months
<b>Exposure concentration</b>	3-24 mm sediment
<b>Experiment description</b>	Undisturbed communities transferred from Oslofjord, 41 m depth. 10°C
<b>Target effects</b>	Sediment community effect
<b>Type of data available</b>	Macrofauna number of species and individuals, diversity
<b>Data format</b>	Schaanning et al., 2009 J Exp Mar Biol Ecol

### 2.2.2 Sediment macroinvertebrate *Hediste diversicolor*

Olefin (Ultidril, Novaplus) and ester (Anco Green, Pterofree) based drilling muds:

<b>Species and life stage</b>	<i>Hediste diversicolor</i>
<b>Exposure time</b>	6 months
<b>Exposure concentration</b>	Approx 500 g slurry, containing 1-1.5 g drilling fluid. Approx 1.5 mm layer
<b>Experiment description</b>	Drill cuttings from drilling sites in the North Sea. Undisturbed communities transferred from Oslofjord, 200 m depth. 7-11 °C, salinity: 32-35 PSU
<b>Target effects</b>	Biomarkers
<b>Type of data available</b>	Catalase activity, glutathione reductase activity
<b>Data format</b>	NIVA rapport 3475-96

### 2.2.3 Chronic mussel lab exposure: *Mytilus edulis*

Produced water from Ormen Lange gas processing plant.

<b>Species and life stage</b>	<i>Mytilus edulis</i> , farmed (from snadderogsnaskum)
<b>Exposure time</b>	5 weeks
<b>Exposure concentration</b>	0.01 %, 0.1 %, 0.5%, 1% PW
<b>Experiment description</b>	Laboratory exposure of mussels exposed to produced water from Ormen Lange gas processing plant.
<b>Target effects</b>	Biomarkers, chemical burdens
<b>Type of data available</b>	Lysosomal stability, micronuclei formation, histochemistry, peroxisomal proliferation, mussel chemistry
<b>Data format</b>	NIVA rapport 5747-2009

#### 2.2.4 Chronic fish lab exposure: *Danio rerio*

Water soluble components of oil, a mixture of low-molecular-weight PAHs (<5 ring) and short-chain APs (C1–C4)

<b>Species and life stage</b>	<i>Danio rerio</i>
<b>Exposure time</b>	1, 7 and 13 weeks
<b>Exposure concentration</b>	sum PAH: 0.54 ppb (low dose), 5.4 ppb (high dose and pulsed dose).
<b>Experiment description</b>	Flow through exposure lasting 1, 7 and 13 weeks, 26 ± 2 °C
<b>Target effects</b>	Biomarkers
<b>Type of data available</b>	Cytochrome P450 1A, Vitellogenin, condition, reproduction, recruitment, F1 deformations, global gene expression
<b>Data format</b>	Holth et al., 2008 Aq Tox

#### 2.2.5 Chronic fish lab exposure: *Gadus morhua*

Water soluble components of oil, low molecular weight PAHs , short chained APs

<b>Species and life stage</b>	<i>Gadus morhua</i>
<b>Exposure time</b>	4, 16, 32, or 44 weeks
<b>Exposure concentration</b>	Low: 0.54 ppb PAH, 1.14 ppb APs, pulsed and high: 5.4 ppb PAH, 11.4 ppb APs
<b>Experiment description</b>	Flow through exposure lasting 4, 16, 32, or 44 weeks
<b>Target effects</b>	Hepatic DNA adduct formation, Hepatic EROD activity
<b>Type of data available</b>	Increased DNA adduct formation following > 16 weeks exposure
<b>Data format</b>	Holth et al., 2009 Environ. Sci. Technol.

Water soluble components of oil, low molecular weight PAHs , short chained APs

<b>Species and life stage</b>	<i>Gadus morhua</i>
<b>Exposure time</b>	4, 16, 32, or 44 weeks
<b>Exposure concentration</b>	Low: 0.54 ppb PAH, 1.14 ppb APs, pulsed and high: 5.4 ppb PAH, 11.4 ppb APs
<b>Experiment description</b>	Flow through exposure lasting 4, 16, 32, or 44 weeks
<b>Target effects</b>	Biomarkers

<b>Type of data available</b>	Condition, gonad maturation, gene expression
<b>Data format</b>	Holth et al., 2010 Can J Fish Aquat Sci

### 2.2.6 Chronic mussel field exposure: *Mytilus edulis*

Produced water from Ormen Lange gas processing plant.

<b>Species and life stage</b>	<i>Mytilus edulis</i> , farmed (from snadderogsnaskum)
<b>Exposure time</b>	6 weeks
<b>Exposure concentration</b>	2 reference station, 4 Exposed stations
<b>Experiment description</b>	Caged mussels placed near produced water discharge point of Ormen Lange gas processing plant. Two reference sites compared to four exposed sites.
<b>Target effects</b>	Biomarkers, chemical burdens
<b>Type of data available</b>	Lysosomal stability, micronuclei formation, histochemistry, peroxisomal proliferation, mussel chemistry
<b>Data format</b>	NIVA rapport 5747-2009

Produced water from Ormen Lange gas processing plant.

<b>Species and life stage</b>	<i>Mytilus edulis</i>
<b>Exposure time</b>	Native mussels (not applicable)
<b>Exposure concentration</b>	Reference station, Exposed station 1, Exposed station 2
<b>Experiment description</b>	Collection of native mussels near produced water discharge point of Ormen Lange gas processing plant. One reference population compared to two exposed populations.
<b>Target effects</b>	Biomarkers, chemical burdens
<b>Type of data available</b>	Lysosomal stability, micronuclei formation, histochemistry, peroxisomal proliferation, mussel chemistry
<b>Data format</b>	NIVA rapport 5747-2009

Produced water discharge in the North Sea.

<b>Species and life stage</b>	<i>Mytilus edulis</i> (farmed), <i>Gadus morhua</i> (farmed)
<b>Exposure time</b>	6 weeks
<b>Exposure concentration</b>	Reference, 500 m, 1000 m, 2000 m, 5000 m, 10 000 m from platform
<b>Experiment description</b>	In situ exposure of Atlantic cod and blue mussel exposed to produced water from oil production platform.
<b>Target effects</b>	Biomarkers
<b>Type of data available</b>	EROD activity, GST activity, Lysosomal stability, DNA adducts, Micronuclei, Histochemistry, Vitellogenin
<b>Data format</b>	Hylland et al., 2008 Mar Poll Bull

### 2.2.7 Chronic fish field exposure: *Gadus morhua*

Produced water discharge in the North Sea.

<b>Species and life stage</b>	<i>Gadus morhua</i> (farmed)
<b>Exposure time</b>	6 weeks
<b>Exposure concentration</b>	Reference, 500 m, 1000 m, 2000 m, 5000 m, 10 000 m from platform
<b>Experiment description</b>	In situ exposure of Atlantic cod and blue mussel exposed to produced water from oil production platform.
<b>Target effects</b>	Biomarkers
<b>Type of data available</b>	EROD activity, GST activity, Lysosomal stability, DNA adducts, Micronuclei, Histochemistry, Vitellogenin
<b>Data format</b>	Hylland et al., 2008 Mar Poll Bull

### 2.2.8 *In vitro* systems: Yeast estrogen and androgen screen

YES/YAS: Samples were analysed for estrogenic activity using the yeast estrogen screen (YES), a yeast-based (*Saccharomyces cerevisiae*) screen developed by Glaxo–Wellcome, plc (Stevenage, Herts, UK), and fully validated by Routledge and Sumpter (1996). The bioassay has been demonstrated to quantitatively detect known estrogens (e.g., 17 $\beta$ -estradiol and estrone) and xenoestrogens (e.g., alkylphenols and Bisphenol A) via a receptor mechanism. The bioassay was carried out using the method of Routledge and Sumpter (1996), with minor variations. Despite not directly relevant to arctic conditions, it could be argued that recombinant yeast screens have importance due to high relevance in determining mechanism of action.. Data exist for two individual PW samples, collected minimum 6 weeks apart. Shown here are results from sampling 1. Both water soluble and particulate fractions of PW were analysed in the yeast screens, but shown here are data from water dissolved fraction only.

Organic extract of water soluble components of produced water from Grane:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extracted with solid phase extraction, followed by static <i>in vitro</i> exposure of recombinant yeast. 32 $\pm$ 1 $^{\circ}$ C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: < 0.03 ng E2/L, Flutamide equivalents: 1578 $\mu$ g Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Gullfaks B:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of

	PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: < 0.03 ng E2/L, Flutamide equivalents: 582 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Heidrun:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: < 0.03 ng E2/L, Flutamide equivalents: 1627 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Oseberg C:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: 2.1 ng E2/L, Flutamide equivalents: 3 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Snorre A:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: <0.03 ng E2/L, Flutamide equivalents: 134 µg Flut/L



<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull
--------------------	--------------------------------------

Organic extract of water soluble components of produced water from Statfjord A:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extracted with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: 3.5 ng E2/L, Flutamide equivalents: 430 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Statfjord B:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extracted with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: - ng E2/L, Flutamide equivalents: - µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Statfjord C:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extracted with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: 4 ng E2/L, Flutamide equivalents: 899 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Troll B:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
-------------------------------	---------------------------------

<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: 0.4 ng E2/L, Flutamide equivalents: 791 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

Organic extract of water soluble components of produced water from Asgard:

<b>Species and life stage</b>	<i>Saccharomyces cerevisiae</i>
<b>Exposure time</b>	3 days (YES), 2 days (YAS)
<b>Exposure concentration</b>	Serial dilutions of PW extract
<b>Experiment description</b>	Yeast Estrogen Screen (YES) and Yeast Androgen Screen (YAS). Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of recombinant yeast. 32 ± 1 °C
<b>Target effects</b>	Estrogenic ad anti-androgenic effects
<b>Type of data available</b>	Estrogen (E2) equivalents: 0.1 ng E2/L, Flutamide equivalents: 8171 µg Flut/L
<b>Data format</b>	Tollefsen et al., 2007 Mar Poll Bull

### 2.2.9 *In vitro* systems: primary hepatocytes from *Onchorynchus mykiss*

ROS/glutathione/cytotoxicity: The bioassays was carried out using the method of Wang and Joseph, 1999 (ROS), Vandeputte et al., 1994 (glutathione), Schreer et al. 2005 (cytotoxicity). Despite maybe not directly relevant to arctic, it could be argued that *in vitro* analyses of biomarkers in rainbow trout hepatocytes have importance due to high relevance in determining mechanism of action. Data exist for both water soluble and particulate fractions of PW, but shown here are data from water dissolved fraction only, as bioavailability of chemicals in the particulate fraction could be discussed

Organic extract of water soluble components of produced water from Grane:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: - , cytotoxicity: 33
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Troll B:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 11, Glutathione: 11 , cytotoxicity: 0.3
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Statfjord C:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: 11 , cytotoxicity: 3
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Statfjord A:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: 11 , cytotoxicity: 1
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Snorre A:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and

	cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: 11 , cytotoxicity: 3
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Gullfaks B:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: 33 , cytotoxicity: 110
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Asgard:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: 3 , cytotoxicity: 1
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Heidrun:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: - , cytotoxicity: 3
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Statfjord B:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)

<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: - , cytotoxicity: 1
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

Organic extract of water soluble components of produced water from Oseberg C:

<b>Species and life stage</b>	<i>Onchorynchus mykiss</i>
<b>Exposure time</b>	1 h (ROS formation), 96 h (glutathione, cytotoxicity)
<b>Exposure concentration</b>	0.3, 1, 3, 10, 30, 100, 300 fold concentrated PW
<b>Experiment description</b>	Organic components of PW extrated with solid phase extraction, followed by static in vitro exposure of hepatocytes.
<b>Target effects</b>	In vitro biomarkers (oxidative stress and cytotoxicity) in primary cell culture
<b>Type of data available</b>	(LOEC): ROS formation: 3, Glutathione: - , cytotoxicity: 1
<b>Data format</b>	Farmen et al., 2010 Mar Poll Bull

### 3. References

Brooks, S., Harman, C. 2009. Biological effects monitoring of the discharge water from the Ormen Lange gas facility, Norway. NIVAreport 5747-2009; ISBN: 978-82-577-5782-2

Hylland K, Tollefsen KE, Ruus A, Jonsson G, Sundt RC, Sanni S, Røe Utvik TI, Johnsen S, Nilssen I, Pinturier L, Balk L, Barsiene J, Marigómez I, Feist SW, Børseth JF. 2008. Water column monitoring near oil installations in the North Sea 2001-2004. Mar Pollut Bull. 2008;56(3):414-29.

Farmen E, Harman C, Hylland K, Tollefsen KE. 2010. Produced water extracts from North Sea oil production platforms result in cellular oxidative stress in a rainbow trout in vitro bioassay. Mar Pollut Bull.;60(7):1092-8

Holth TF, Beylich BA, Skarphédinsdóttir H, Liewenborg B, Grung M, Hylland K. 2009. Genotoxicity of environmentally relevant concentrations of water-soluble oil components in cod (*Gadus morhua*). Environ Sci Technol.;43(9):3329-34

Holth TF, Thorsen, A, Olsvik PA., Hylland K. 2010. Long-term exposure of Atlantic cod (*Gadus morhua*) to components of produced water: condition, gonad maturation, and gene expression. Can J Fish Sci 67: 1685–1698

Holth TF, Nourizadeh-Lillabadi R, Blaesbjerg M, Grung M, Holbech H, Petersen GI, Aleström P, Hylland K. 2008. Differential gene expression and biomarkers in zebrafish (*Danio rerio*) following exposure to produced water components. Aquat Toxicol.;90(4):277-91.

- Källqvist, T. 1991. Biologisk testing av boreslam - Reindalspasset - 1, Svalbard. Norsk institutt for vannforskning report 2593; ISBN: 82-577-1914-5
- Källqvist, T. 1994. Toksisitetstesting av oljedispergeringsmidler - utprøving av en metode med marine alger (*Skeletonema costatum*). Norsk institutt for vannforskning (NIVA) report 3128; ISBN: 82-577-2600-1
- Routledge, E.J., Sumpter, J.P., 1996. Oestrogenic activity of surfactants and some of their degradation products using a recombinant yeast screen. *Environ. Toxicol. Chem.* 15, 241–248.
- Schaanning, M., Hylland, K., Lichtenthaler, R., Rygg, B. 1996. Biodegradation of Anco Green Novaplus Drilling Muds on Cuttings Deposited in Benthic Chambers. Norsk institutt for vannforskning (NIVA) report 3475-96; ISBN: 82-577-3014-9
- Schaanning, M., Lichtenthaler, R., Rygg, B. 1997. Biodegradation of Esters and Olefins in Drilling Mud Deposited on Arctic soft-bottom communities in a low-temperature Mesocosm. Norsk institutt for vannforskning (NIVA) report 3460-97; ISBN: 82-577-3332-6
- Schaanning, M.T, Trannum, H.C, Øxnevad, S, Carroll, J, Bakke, T., 2008. Effects of drill cuttings on biogeochemical fluxes and macrobenthos of marine sediments. *Journal of Experimental Marine Biology and Ecology* 361: 49-57
- Schreer A, Tinson C, Sherry JP, Schirmer K. 2005. Application of Alamar blue/5-carboxyfluorescein diacetate acetoxymethyl ester as a noninvasive cell viability assay in primary hepatocytes from rainbow trout. *Analytical Biochemistry* 344, 76-85.
- Tollefsen KE, Harman C, Smith A, Thomas KV. 2007. Estrogen receptor (ER) agonists and androgen receptor (AR) antagonists in effluents from Norwegian North Sea oil production platforms. *Mar Poll Bull.*;54(3):277-83
- Vandeputte, Guizon I, Genestie-Denis I, Vannier B, Lorenzon G. 1994. A microtiter plate assay for total glutathione and glutathione disulfide contents in cultured/isolated cells: performance study of a new miniaturized protocol. *Cell Biology and Toxicology* 10, 415-421.
- Wang H, Joseph JA. 1999. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radical Biology and Medicine* 27, 612-616.

NIVA: Norway's leading centre of competence in aquatic environments

NIVA provides government, business and the public with a basis for preferred water management through its contracted research, reports and development work. A characteristic of NIVA is its broad scope of professional disciplines and extensive contact network in Norway and abroad. Our solid professionalism, interdisciplinary working methods and holistic approach are key elements that make us an excellent advisor for government and society.



Norwegian Institute for Water Research

Gaustadalléen 21 • NO-0349 Oslo, Norway  
Telephone: +47 22 18 51 00 • Fax: 22 18 52 00  
[www.niva.no](http://www.niva.no) • [post@niva.no](mailto:post@niva.no)