

# The effects of wildfire on the water chemistry of dilute, acidic lakes in southern Norway

Espen Lydersen · Rolf Høgberget ·  
Clara E. Moreno · Øyvind A. Garmo ·  
Per Christian Hagen

Received: 6 July 2013 / Accepted: 31 December 2013 / Published online: 22 January 2014  
© The Author(s) 2014. This article is published with open access at Springerlink.com

**Abstract** Changes in lake water chemistry were studied for >4 years following a large wildfire in a boreal forest area in Mykland, southern Norway, an area characterized by thin and patchy, base-poor and slow-weathering soils and bedrock. Accordingly, the lakes have low acid neutralizing capacity (ANC), calculated as the difference between the total concentration of base cations ( $[\Sigma\text{BC}]$ ) and strong acid anions ( $[\text{SAA}]$ ). During the initial post-fire period, and peaking about two months after the fire, the mobilization of SAA from terrestrial to aquatic systems caused a dramatic drop in ANC. In one of the lakes, ANC dropped from about 20 to  $-80 \mu\text{eq L}^{-1}$ , while  $[\text{H}^+]$  and inorganic aluminum ( $[\text{Al}_i]$ ) increased to

$38 \mu\text{eq L}^{-1}$  (pH 4.42) and  $326 \mu\text{g Al L}^{-1}$  ( $36.2 \mu\text{eq L}^{-1}$  as  $\text{Al}^{3+}$ ), respectively. Sulfate and chloride were the predominant anions responsible for this decline in ANC, as the nitrate increase was small. After the severe chemical episode,  $[\text{SAA}]$  in the lakes declined faster than  $[\Sigma\text{BC}]$ , and within about one year after the wildfire, ANC was back to almost pre-fire values. However, despite the fact that  $[\text{SAA}]$  also continued to decline faster than  $[\Sigma\text{BC}]$  the following years, no further increase in ANC was documented. The strong ionic strength decline and the increase in TOC during the same period have likely counteracted for the potential ANC increase. There were large lake-to-lake variations in water chemistry of the wildfire affected lakes. Hydrology, geology, lake residence time and the catchment area to lake area ratio are important explanatory factors.

Responsible Editor: Jennifer Leah Tank.

E. Lydersen (✉) · C. E. Moreno  
Department of Environmental and Health Studies,  
Telemark University College, 3800 Bø, Norway  
e-mail: espen.lydersen@hit.no

R. Høgberget  
Norwegian Institute for Water Research, 4879 Grimstad,  
Norway

Ø. A. Garmo  
Norwegian Institute for Water Research, 2312 Ottestad,  
Norway

P. C. Hagen  
Department of Business Administration and Computer  
Studies, Telemark University College, 3800 Bø, Norway

**Keywords** Wildfire · Climate change · Lake water chemistry · Acidification · Southern Norway

## Introduction

Wildfire poses a threat to life, property, infrastructure and natural resources in fire-prone areas. Each year wildfires burn large areas of forest land around the world (Smith et al. 2011). Recent projections of future climate changes for the interior of many continents suggest an increase of air temperature by 2–6 °C

coupled with a 10–30 % decrease of summer precipitation (e.g. IPCC 2001; Schär et al. 2004). An increase in drought is directly related to an increase in wildfire occurrence (e.g. Ryan 2000; Flannigan and Wotton 2001). Thus, the question arises where and how much various areas on earth will be affected by these potential changes.

Prolonged drought followed by a strong rain event, may lead to severe water chemical conditions. In Norway Huitfeldt-Kaas (1922) documented mass death of salmon and brown trout during such an event in 1920. Later, many scientists have dealt with this phenomenon (Jeffries et al. 2003; Laudon et al. 2004; Aherne et al. 2006). Drought periods result in lower water table in wetlands and soils, leading to oxidation of previously stored, reduced sulfur (S), and subsequent efflux of oxidized S ( $\text{SO}_4^{2-}$ ) upon re-wetting. In acid sensitive areas with high sulfur input and low acid neutralizing capacity, severe episodic acidification of surface waters may occur as  $\text{H}^+$  and cationic Al ( $\text{Al}^{n+}$ ) appear as predominant counter ions for  $\text{SO}_4^{2-}$  (Dickson 1978; Cronan and Schofield 1979).

Besides drought, the fire itself is also an oxidation process, which under sufficient oxygen conditions, oxidizes organic carbon to  $\text{CO}_2$ , organic S to  $\text{SO}_2$  and organic N to  $\text{NO}_x$ , similar to combustion of biofuels. These gases are emitted to air, and after various physical and chemical transformation processes in the atmosphere, they precipitate as acid rain elsewhere, as carbonic acid, sulfuric acid and nitric acids. The combustion remains, left in the wildfire area are mineralized, alkaline ash products, rich in base cations, phosphate, chlorides, heavy metals (Eriksson et al. 2003; Kelly et al. 2006) and pyrolytic polycyclic aromatic hydrocarbons, PAH (Finkelstein et al. 2005).

Wildfire leads to increased runoff (Schindler et al. 1980; Bayley et al. 1992b), mainly due to reduced evapotranspiration. Particle transport to surface waters also increases, due to both increased runoff and increased erosion (Beatty 1994; Karr et al. 2004). Higher turbidity will temporarily increase the light extinction coefficient and water transparency during flood episodes. These effects are very dependent on time after the wildfire, as the increased particle load to surface water declines rapidly, while total organic carbon (TOC) gradually increases again as new allochthonous material is produced (Carignan et al. 2000). The first period after wildfire, the primary production may also affect these parameters, due to

the increased supply of N, P and base cations to lakes following wildfire (Planas et al. 2000).

Increases in major base cations, strong acid anions, ammonia, total N and total P in surface water have been reported from many wildfire studies (Schindler et al. 1980; Lamontagne et al. 2000; Carignan et al. 2000; Bayley et al. 1992a, b; Eriksson et al. 2003). Studies documenting significant reduction in pH and ANC (Bayley et al. 1992b; Eriksson et al. 2003) are related to areas with natural base-poor soils and low weatherable rocks. Some of the severe acidification reported from a wildfire in Sweden (Eriksson et al. 2003) is attributed to use of brackish water for firefighting, causing an additional sea-salt effect, an acidification phenomena well described earlier (Hindar et al. 1994; Lydersen and Henriksen 1995; Nystroem et al. 1995).

In this paper we report water chemical effects on six lakes in a very acid sensitive area of southern Norway, impacted by a large wildfire in June 2008. Both short-term and long-term chemical effects (>4 years) of the wildfire on lakes were studied.


## Materials and methods

The wildfire area was located in Froland municipality, Aust-Agder county, southern Norway, (Fig. 1). The wildfire started on June 9, 2008 and lasted 1 week. Almost 2,600 ha of forest and wilderness were almost totally damaged, and 120,000  $\text{m}^3$  of forest were lost. Six lakes investigated were within the wildfire area, while three reference lakes were located nearby the impacted area. The control lakes were primarily selected according to the following criteria:

- Proximity to the wildfire affected area
- Negligible influence by land-use and silviculture
- Accessibility by road for use of boat

During the investigated period, powdered limestone was unexpectedly dumped into a stream entering Lake Svarttjenn in September 2009, with significant effects on  $\text{Ca}^{2+}$ , pH and alkalinity both during 2009 (Table 1) and 2010. Another control lake, Lake Melestjenn showed unusually high concentrations of  $\text{Na}^+$  and  $\text{Cl}^-$ , due to sporadic road salting at icy conditions during the winter (Table 1). The remaining control lake, Lake Jordtjenn has not been significantly affected by human activities during the monitoring

**Fig. 1** Map of the wildfire area (red) in Mykland, Aust-Agder County, Norway, showing locations of the studied lakes inside (1–6) and outside (7–9) the wildfire area. Open circles indicate the sampling site in the lakes. 1: Lake Fisketjenn, 2: Lake Hundsvatn, 3: Lake Øyvatt, 4: Lake Grunnetjenn, 5: Lake Heitjenn, 6: Lake Rasvassvatn, 7: Lake Jordtjenn, 8: Lake Svarttjenn, 9: Lake Melestjenn. (Color figure online)



**Table 1** Median water chemistry in the control lakes in 2009 and 2012

Parameter	Unit	Jordtjenn 2009	Jordtjenn 2012	Svarttjenn 2009	Svarttjenn 2012	Melestjenn 2009	Melestjenn 2012
H <sub>25</sub>	mS m <sup>-1</sup>	1.79	1.88	1.61	1.92	1.82	2.87
H <sup>+</sup>	µeq L <sup>-1</sup>	6.0	2.6	3.2	4.6	3.9	5.1
Ca <sup>2+</sup>	µeq L <sup>-1</sup>	47	42	73	46	59	62
Mg <sup>2+</sup>	µeq L <sup>-1</sup>	25	28	26	27	26	29
Na <sup>+</sup>	µeq L <sup>-1</sup>	76	65	69	65	93	122
K <sup>+</sup>	µeq L <sup>-1</sup>	9.5	6.9	9.0	5.9	2.8	5.9
NH <sub>4</sub> <sup>+</sup> -N	µeq L <sup>-1</sup>	3.2	2.3	3.9	2.3	6.9	3.3
SO <sub>4</sub> <sup>2-</sup>	µeq L <sup>-1</sup>	43	39	35	32	15	30
Cl <sup>-</sup>	µeq L <sup>-1</sup>	66	72	65	69	99	141
NO <sub>3</sub> <sup>-</sup> -N	µeq L <sup>-1</sup>	2.2	1.4	2.6	1.1	3.7	1.9
SO <sub>4</sub> *	µeq L <sup>-1</sup>	35	31	27	26	5.0	16
ΣBC*	µeq L <sup>-1</sup>	85	67	102	62	80	68
TOC	mg C L <sup>-1</sup>	5.9	6.0	7.1	7.4	7.4	9.9
ANC	µeq L <sup>-1</sup>	49	29	74	33	67	51
Total-N	µg N L <sup>-1</sup>	309	305	298	356	488	460
Al <sub>a</sub>	µg Al L <sup>-1</sup>	145	111	122	143	120	170
Al <sub>o</sub>	µg Al L <sup>-1</sup>	103	74	95	108	102	127
Al <sub>i</sub>	µg Al L <sup>-1</sup>	34	35	27	39	26	36

**Table 2** Lake water chemistry in the wildfire affected Lake Hundsvatn and Lake Rasvassvatn, during the rain event in August 2008, compared with the nearby long term monitoring site at Birkenes, and the water chemistry in the same wildfire

affected lakes and the least anthropogenic affected control lake, Lake Jordtjenn, in August 2012, &gt;4 years after the wildfire

Parameter	Unit	Hundsvatn	Rasvassvatn August 2008	Birkenes	Hundsvatn	Rasvassvatn August 2012	Jordtjenn
H <sup>+</sup>	μeq L <sup>-1</sup>	38	17	20	3.8	4.2	1.4
Ca <sup>2+</sup>	μeq L <sup>-1</sup>	146	60	28	26	20	42
Mg <sup>2+</sup>	μeq L <sup>-1</sup>	114	63	16	26	30	30
Na <sup>+</sup>	μeq L <sup>-1</sup>	125	84	118	48	48	57
K <sup>+</sup>	μeq L <sup>-1</sup>	70	33	1.0	4.4	9.5	6.1
NH <sub>4</sub> <sup>+</sup>	μeq L <sup>-1</sup>	19	11	0.4	0.7	0.7	2.1
SO <sub>4</sub> <sup>2-</sup>	μeq L <sup>-1</sup>	367	160	55	31	50	31
Cl <sup>-</sup>	μeq L <sup>-1</sup>	165	114	116	65	68	73
NO <sub>3</sub> <sup>-</sup>	μeq L <sup>-1</sup>	2.8	5.4	1.2	0.7	5.7	1.7
ΣBC*	μeq L <sup>-1</sup>	272	115	35	32	32	54
SO <sub>4</sub> *	μeq L <sup>-1</sup>	350	149	43	25	43	24
TOC	mg C L <sup>-1</sup>	5.0	2.6	7.1	6.4	3.6	6.0
ANC	μeq L <sup>-1</sup>	-80	-39	-9	7	-17	29
Tot-N	μg N L <sup>-1</sup>	710	445	235	276	248	259
Al <sub>a</sub>	μg Al L <sup>-1</sup>	368	295	310	77	79	74
Al <sub>o</sub>	μg Al L <sup>-1</sup>	42	22	141	51	42	55
Al <sub>i</sub>	μg Al L <sup>-1</sup>	326	273	169	27	37	19

period, until the winter 2011/2012 when a substantial clear-cut was conducted, but chemical lake effects of this logging have so far not been detected (Table 1).

There are no pre-wildfire water chemistry data from the 6 affected lakes and the 3 control lakes. We thus compared the initial wildfire effects on lakes with a nearby long-term monitoring site, Birkenes, located about 31 km southwest from the wildfire area. At this site, dry and wet deposition has been monitored daily by the Norwegian Institute for Air Research, while freshwater has been monitored weekly by Norwegian Institute for Water Research since the early 1970s.

As the Birkenes site is located closer to the sea (20 km from the coast) compared with the wildfire area (about 50 km from the coast), the comparison of Na<sup>+</sup> and Cl<sup>-</sup> with Birkenes is omitted. In addition, the low acid neutralizing capacity (ANC) at Birkenes, implies generally lower pH and higher concentrations of inorganic Al (Al<sub>i</sub>) at this site compared to the expected pre-fire levels in the wildfire affected area (Table 2). Despite these limitations, it was of interest to compare the chemical effects of wildfire on lakes in Mykland area with the Birkenes site, as this site historically has been the most acid rain impacted

monitoring site in Norway. Thus, we decided to use Lake Jordtjenn and the Birkenes site as the main reference sites for the comparison with the wildfire affected sites.

#### Site description

Since the main lakes in the wildfire area have been artificially limed, we decided to only investigate wildfire affected lakes located upstream the limed lakes. The lakes are located between 229 and 464 m a.s.l., and all lakes drain to the River Tovdal, the cradle of acid rain research in Norway (Drabloes et al. 1980; Sevaldrud et al. 1980). Average annual precipitation in the area is 1,150–1,200 mm and average evapotranspiration 30–36 %. The residence time of the lakes varied from 0.059–3.2 years (Table 3). The catchments of all wildfire affected lakes investigated were totally located within the wildfire area (Fig. 1), except for Lake Rasvassvatn, where ≈ 10 % of the catchment was outside the wildfire area.

Lake Heitjenn, Lake Fisketjenn and Lake Øyvatt are all headwater lakes. Lake Grunnetjenn and Lake Hundsvatn are the first and second downstream lakes

**Table 3** Lake and catchment characteristics of the studied lakes

Lake	Unit	Burnt lakes					Reference lakes				
		Fisketjenn	Hundsvatn	Øyvvatn	Grunnetjenn	Heitjenn	Råsvassvatn	Jordtjenn	Melestjenn	Svarttjenn	
Latitude		N58 36.549	N58 36.413	N58 36.675	N58 36.906	N58 37.073	N58 37.446	N58 38.956	N58 37.598	N58 39.118	
Longitude		E8 17.194	E8 16.770	E8 19.037	E8 16.515	E8 16.370	E8 20.165	E8 18.721	E8 22.063	E8 20.220	
Altitude min	m asl	229	228	255	231	271	173	228	222	191	
Altitude max	m asl	245	320	316	293	288	360	464	243	464	
Lake area (LA)	km <sup>2</sup>	0.56	0.15	0.072	0.068	0.037	0.89	0.024	0.0091	0.021	
Maximum depth	m	5.8	13	3.2	3.2	2.5	15	9.5	6.2	6.7	
Lake volume	1,000 m <sup>3</sup>	109	648	77	72	72	4.6	75	19	46	
Catchment area (CA)	km <sup>2</sup>	0.23	2.6	1.1	0.86	0.14	1.2	2.9	0.13	6.9	
Draining ratio (CA:LA)		4.1	18	15	13	3.8	1.4	122	14	335	
Residence time	yr	0.40	0.21	0.059	0.071	0.19	3.2	0.023	0.12	0.058	
Lake area (LA:CA)	%	25	5.7	6.6	7.8	26	72	0.82	7.0	0.30	
Annual precipitation	mm	1,182	1,184	1,183	1,178	1,165	1,169	1,146	1,184	1,152	
Annual discharge	mm	746	753	802	731	742	745	806	781	760	
Evapotranspiration	%	36.9	36.4	32.2	37.9	36.3	36.2	29.7	34.0	34.0	
Average annual temperature	°C	5.0	4.9	5.0	4.9	4.9	4.9	4.7	5.2	4.8	
Agricultural area	%	0.0	0.0	0.0	0.0	0.0	0.0	1.3	5.2	2.0	
Bog	%	29.8	22.0	11.8	10.7	3.1	7.8	7.8	8.6	13.3	
Lake	%	26.8	12.4	8.1	11.9	26.4	12.5	3.0	15.7	1.8	
Forest	%	37.5	61.0	79.3	70.4	52.3	78.2	85.4	62.4	81.7	
Naked mountain	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Urban area	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Burnt	%	≈ 100	≈ 100	≈ 100	≈ 100	≈ 100	≈ 90				

Areas impacted by fire (named: Burnt lakes) are also to a certain degree impacted by logging after the wild fire. The three control lakes (named: References) are located in the nearby areas from the wildfire area. Except for lake depth, lake volume and residence time data (own calculations), the data are from the Norwegian energy directorate database

**Table 4** Parameters and analytical methods used

Parameter	Unit	Reference
pH	$-\log[\text{H}^+]$	NS 4720, 1979, 2. Edition
Conductivity	$\text{mS m}^{-1}$	NS-ISO 7888, 1993, 1. Edition
Alkalinity	$\text{mmol L}^{-1}$	NS 4754 (described in Ch.2.3)
$\text{Ca}^{2+}$ , $\text{Mg}^{2+}$ , $\text{Na}^+$ , $\text{K}^+$ , $\text{Cl}^-$ , $\text{SO}_4^{2-}$	$\text{mg L}^{-1}$	Ionchromatography (Dionex IC 1100)
$\text{NO}_3^-$ -N, $\text{NH}_4^+$ -N	$\mu\text{g L}^{-1}$	Ionchromatography (Dionex IC 1100)
Total nitrogen	$\mu\text{g N L}^{-1}$	Persulfate oxidation, FIA: Tecator ASN 110-03/92
Total phosphorous	$\mu\text{g P L}^{-1}$	NS 4725, 3. Edition, 1984
Total organic carbon	$\mu\text{g C L}^{-1}$	ISO 1487, TOC analyzer: OI Analytical-1010
Total monomeric Al	$\mu\text{g L}^{-1}$	Barnes/Driscoll (Barnes, 1975; Driscoll, 1984)
Organic monomeric Al	$\mu\text{g L}^{-1}$	Barnes/Driscoll (Barnes, 1975; Driscoll, 1984)

from Lake Heitjenn, while Lake Fisketjenn drains into Lake Hundsvatn.

Lake Melestjenn is a headwater lake. Lake Rasvassvatn (173 m a.s.l.) has a headwater lake about 1 km east from the lake, 332 m a.s.l., with its catchment totally influenced by the wildfire. Lake Jordtjenn has a small headwater lake  $\approx 1$  km north-east from the lake, and Lake Jordtjenn drains into Lake Svarttjenn.

The wildfire area is characterized by slowly weatherable Precambrian rocks (granites and gneisses), thin and patchy soil cover, often with peat marsh and bogs in the bottom of the valleys and in other depressions of the catchments. Locally some more weatherable, base cation richer bedrocks occur as hypabyssal pegmatite rocks. The vegetation is dominated by various heather species with Norwegian pine (*Pinus sylvestris*) and birch (*Betula pubescens*) as predominant trees, with stands of aspen (*Populus tremula*) on south facing hillsides, and Norwegian spruce (*Picea abies*) in more productive areas.

Precipitation data come from a local weather station at Mykland, just outside the wildfire area, while air temperature data were taken from Nelaug, 17 km east from the wildfire area. Both weather stations are operated by the Norwegian Meteorological Institute.

#### Sampling and chemical analysis

For two of the wildfire affected lakes, Lake Hundsvatn and Lake Rasvassvatn, water samples were taken on June 25, 2008 (nine days after the fire), on July 9, 2008, during a rain event on August 13, 2008, and on

October 30, 2008 (Høgberget 2010), after lake turnover. For Lake Øyvvatn, another wildfire affected lake, only one sample was taken in 2008 (on July 16). The control lakes outside the wildfire affected area were not sampled in 2008.

Six lakes within the wildfire area (including Lake Hundsvatn and Lake Rasvassvatn), and three control lakes outside the wildfire area, were monitored monthly from May 19, 2009 to late October 2012. Water samples were collected with a Limnos sampler, transferred to prewashed polyethylene bottles, and stored cold in a darkroom until analyzed. Ion exchange fractionation of aluminum was conducted in field according to Driscoll (1984). Conductivity, pH, alkalinity and Al-fractions were determined within one day after sampled. The analytical methods are given in Table 4.

#### Calculations

Aluminum (Al) was fractionated according to the Barnes/Driscoll method (Barnes 1975; Driscoll 1984). Two Al-fractions were measured: total monomeric Al ( $\text{Al}_a$ ) and organic monomeric Al ( $\text{Al}_o$ ). Based on these fractions, inorganic monomeric Al ( $\text{Al}_i$ ) was calculated as  $\text{Al}_a - \text{Al}_o$ .

Alkalinity ( $\mu\text{eq L}^{-1}$ ) was determined by acid titration (with HCl) down to pH 4.5, and the alkalinity was corrected for pH by the following expression:


$$\text{Alk}_{\text{corrected}} = \text{Alk}_{\text{measured}} - \left( [\text{H}^+]_{\text{pH:4.5}} - [\text{H}^+]_{\text{pH:original}} \right)$$

Acid neutralizing capacity (ANC) was calculated on equivalent basis ( $\text{eq L}^{-1}$ ) according to Reuss and Johnson (1986), i.e.:





**Fig. 3** Concentrations of strong acid anions, base cations and ANC in Lake Hundsvatn *black upper triangle*, Lake Rasvassvatn *white circle*, and the control lake, Lake Jordtjenn *multi sign*, from June 2008 until October 2012. The monitoring started in the control lake in May 2009





than Birkenes. In Lake Rasvassvatn  $[\text{SO}_4^*]$  was  $149 \mu\text{eq L}^{-1}$ , 3.5 times higher than at Birkenes (Table 2). Birkenes, the most acid rain impacted monitoring site in Norway, have never shown such high  $\text{SO}_4^*$  concentrations since the monitoring started in 1973. Only a few events with  $[\text{SO}_4^*] > 200 \mu\text{eq L}^{-1}$  have been registered at Birkenes, all of them during the most severe acidification time period, i.e. before 1986 (Fig. 5).

During the August peak,  $[\Sigma\text{BC}^*]$  in Lake Hundsvatn was  $272 \mu\text{eq L}^{-1}$  (Fig. 5), about eight times

higher than at Birkenes ( $35 \mu\text{eq L}^{-1}$ ), while  $[\Sigma\text{BC}^*]$  in Lake Rasvassvatn was  $115 \mu\text{eq L}^{-1}$ ,  $>3$  times higher than at Birkenes. While the ANC value at Birkenes was  $-8.8 \mu\text{eq L}^{-1}$ , ANC was  $-80$  and  $-39 \mu\text{eq L}^{-1}$  in Lake Hundsvatn and Lake Rasvassvatn, during this episode (Table 2). The strong decreases in ANC in Lake Hundsvatn caused an almost twofold increase in the concentration of  $\text{H}^+$  and inorganic aluminum ( $\text{Al}_i$ ) compared with Birkenes, i.e.  $20 \mu\text{eq H}^+ \text{L}^{-1}$  (pH 4.7) and  $169 \mu\text{g Al}_i \text{L}^{-1}$  ( $19 \mu\text{eq L}^{-1}$  as  $\text{Al}^{3+}$ ) at Birkenes, compared with  $38 \mu\text{eq H}^+ \text{L}^{-1}$  (pH 4.4) and



**Fig. 4** Concentrations of  $H^+$ , TOC, Total-N,  $NH_4^+$ , Total-P and different Al fractions ( $Al_a$ ,  $Al_o$ ,  $Al_i$ ) in Lake Hundsvatn *black upper triangle*, Lake Rasvassvatn *white circle*, and the control lake, Lake Jordtjenn *multi sign*, from June 2008 until October 2012. The monitoring started in the control lake in May 2009



**Fig. 5** Concentrations of non-marine sulfate ( $SO_4^*$ ) in Lake Hundsvatn *black upper triangle*, Lake Rasvassvatn *white circle*, and at the nearby monitoring site, Birkenes *dash*. The left panel

covers the period 1973 (January 1)–2012 (October 23), while the right panel covers the period from January 2008 until October 2012. Dotted lines indicates starts of the wild fire

326  $\mu\text{g Al}_i \text{ L}^{-1}$  (36  $\mu\text{eq L}^{-1}$  as  $\text{Al}^{3+}$ ) in Lake Hundsvatn. The  $\text{H}^+$  concentration in Lake Rasvassvatn (17  $\mu\text{eq L}^{-1}$ , pH 4.8), was very similar to the level at Birkenes, but the  $\text{Al}_i$  concentration was substantially higher in Lake Rasvassvatn, 273  $\mu\text{g Al L}^{-1}$  (30  $\mu\text{eq L}^{-1}$  as  $\text{Al}^{3+}$ ). The concentrations of total monomeric aluminum ( $\text{Al}_a$ ) were about equal at the three sites, i.e. 368, 295 and 310  $\mu\text{g Al L}^{-1}$  in Lake Hundsvatn, Lake Rasvassvatn and Birkenes, respectively (Table 2).

Potassium ( $\text{K}^+$ ) showed the strongest relative increase after the wildfire. The concentration in Lake Hundsvatn was 70  $\mu\text{eq L}^{-1}$ ,  $\approx 70$  times higher than Birkenes (Table 2). However,  $[\text{K}^+]$  had the lowest concentration of the four major base cations.

At the extreme water chemical conditions in August 2008, relatively low levels of both  $[\text{NH}_4^+]$ , 11–19  $\mu\text{eq L}^{-1}$ , and particularly  $[\text{NO}_3^-]$ , 3–5  $\mu\text{eq L}^{-1}$ , were observed in the wildfire affected lakes (Table 2). The concentrations of total phosphorous (Tot-P) in the lakes during the first post-fire year were relatively low, i.e. annual means of  $7.6 \pm 2.1$  and  $5.3 \pm 0.4 \mu\text{g P L}^{-1}$  in Lake Hundsvatn and Lake Rasvassvatn, respectively. As the Regional Kendall test indicated significant decreases in Tot-P in the wildfire affected lakes (slope:  $-0.46 \mu\text{g L}^{-1} \text{ year}^{-1}$ ;  $P = 0.004$ ) and not in the control lakes during the postfire period 2009–2012, we conclude that the wildfire probably caused a minor increase in Tot-P during the first post-fire year.

Similarly, as the Regional Kendall test indicated significant increases in TOC in wildfire affected lakes (slope:  $-0.23 \text{ mg C L}^{-1} \text{ yr}^{-1}$ ;  $p = 0.038$ ) and not in the control lakes during the post-fire period 2009–2012, we assume a decrease in TOC during the first post-fire year, perhaps a consequence of combustion of organic matter with subsequent reduced transport of allochthonous matter into the lakes.

Between the wildfire and start of the main monitoring on May 19, 2009, the lakes went through two turnovers (autumn 2008 and spring 2009) and one snowmelt period (spring 2009). During this period, the concentrations of ions have declined dramatically from the peak observed in August 2008. The electrolytic conductivity ( $\text{K}_{25}$ ) declined by 67 % in Lake Hundsvatn, i.e. from 8.04  $\text{mS m}^{-1}$  in August 2008 to 2.67  $\text{mS m}^{-1}$  in May 2009. The corresponding decrease in Lake Rasvassvatn was 29 %, from 4.21 to 2.98  $\text{mS m}^{-1}$ . The largest percentage decreases were shown by  $\text{H}^+$ ,  $\text{NH}_4^+$ ,  $\text{Al}_i$  and  $\text{SO}_4^{2-}$  (and  $\text{SO}_4^*$ )

in Lake Hundsvatn, all with decreases  $>70$  % from August 2008 until May 2009. Only  $\text{NO}_3^-$  showed an increase in Lake Hundsvatn during the same time period, but the increase from 3 to 12  $\mu\text{eq L}^{-1}$ , is of minor importance for the macro-chemistry in the lake.

As the decrease in  $[\Sigma\text{BC}^*]$  in Lake Hundsvatn from August 2008 to May 2009, was much smaller ( $-177 \mu\text{eq L}^{-1}$ ) compared with the decrease in  $[\text{SO}_4^*]$ ,  $-265 \mu\text{eq L}^{-1}$ , a large increase in ANC occurred in this lake during this period, i.e. from  $-80 \mu\text{eq L}^{-1}$  in August 2008, to  $\approx 0 \mu\text{eq L}^{-1}$  in May 2009.

The largest percentage decreases in Lake Rasvassvatn from August 2008 to May 2009, were observed for  $[\text{NH}_4^+]$  ( $-85$  %),  $[\text{Al}_i]$  ( $-47$  %) and  $[\text{H}^+]$  ( $-44$  %), but the largest absolute decrease in equivalents was documented for  $[\text{SO}_4^*]$ , which decreased by 54  $\mu\text{eq L}^{-1}$ , corresponding to a percentage decrease of 35 %. As in Lake Hundsvatn, the  $\text{NO}_3^-$  levels were generally low, and an increase was recorded from August 2008 to May 2009. The concentration of  $\Sigma\text{BC}^*$  in Lake Rasvassvatn was slightly higher (7.1 %) in May 2009 compared with August 2008. As  $[\text{SO}_4^*]$  decreased significantly during the same time interval, ANC increased from  $-39 \mu\text{eq L}^{-1}$  in August 2008 to 16  $\mu\text{eq L}^{-1}$  in May 2009.

#### Long term trends in water chemistry of the wildfire affected lakes

The most severe water chemical conditions and the most significant water chemical recovery from the fire, occurred within the first post-fire year. In August 2012,  $>4$  years after the fire,  $\Sigma\text{BC}^*$  and  $\text{SO}_4^*$  were reduced by 88 and 93 %, respectively in Lake Hundsvatn, and by 72 and 71 % in Lake Rasvassvatn compared with August 2008.


Based on the Regional Kendall test, no significant chemical trends were found in the control lakes during the period May 2009–October 2012, except for a significant decreases in  $[\text{Ca}^{2+}]$ ,  $[\Sigma\text{BC}^*]$  and ANC, as a direct consequence of the unplanned liming of an inlet stream of Lake Svarttjenn in August 2009.

For the four wildfire affected lakes with the highest ANC values, Lake Øyvatt, Lake Grunnetjenn, Lake Hundsvatn and Lake Rasvassvatn, larger decreasing rates were calculated for  $[\text{SO}_4^*]$  compared  $[\Sigma\text{BC}^*]$  during the period May 2009–October 2012 (Table 5). For the remaining two lakes with the lowest ANC, Lake Fisketjenn and Lake Heitjenn, somewhat

**Table 5** Calculated concentration ( $\mu\text{eq L}^{-1}$ ) and half-life for non-marine base cations ( $\Sigma\text{BC}^*$ ) and non-marine  $\text{SO}_4^*$  in the 6 wild fired affected lakes from May 2009 ( $C_{t=0}$ ) to October 2012  $C_{t=\text{end}}$ , assuming first order reaction kinetic decay patterns

Lake	Parameter	$C_{t=0}$ ( $\mu\text{eq L}^{-1}$ )	$C_{t=\text{end}}$ ( $\mu\text{eq L}^{-1}$ )	slope (k)	SE k	$r^2$	$t_{1/2}$ (days)	$t_{1/2}$ (years)
Heitjenn	BC*	100	17	$-1.4\text{E}-03$	$1.8\text{E}-04$	0.63	495	1.36
Heitjenn	SO4*	75	17	$-1.2\text{E}-03$	$8.6\text{E}-05$	0.84	590	1.62
Fisketjenn	BC*	98	27	$-1.0\text{E}-03$	$1.8\text{E}-04$	0.46	673	1.84
Fisketjenn	SO4*	71	24	$-8.5\text{E}-04$	$9.5\text{E}-05$	0.68	818	2.24
Grunnetjenn	BC*	153	49	$-9.2\text{E}-04$	$1.0\text{E}-04$	0.68	756	2.07
Grunnetjenn	SO4*	89	25	$-1.0\text{E}-03$	$9.6\text{E}-05$	0.76	679	1.86
Hundsvatn	BC*	109	44	$-7.2\text{E}-04$	$9.4\text{E}-05$	0.60	961	2.63
Hundsvatn	SO4*	78	25	$-9.1\text{E}-04$	$7.0\text{E}-05$	0.82	759	2.08
Rasvassvatn	BC*	112	48	$-6.8\text{E}-04$	$9.3\text{E}-05$	0.58	1017	2.79
Rasvassvatn	SO4*	101	35	$-8.4\text{E}-04$	$9.2\text{E}-05$	0.68	828	2.27
Øyvavn	BC*	152	97	$-3.6\text{E}-04$	$7.6\text{E}-05$	0.38	1936	5.30
Øyvavn	SO4*	72	36	$-5.5\text{E}-04$	$1.0\text{E}-04$	0.43	1270	3.48

$k$  slope of the equations,  $SE$   $k$  standard error of  $k$ .  $P < 0.05$  for all regressions



**Fig. 6** Calculated trends in ANC in the wildfire impacted lakes from May 2009, one year after the wild fire, to October 2012, based on the decay constants of BC and SAA presented in Table 5. Only the trend in ANC of Lake Grunnetjenn is significant ( $P = 0.045$ )

stronger decreases were documented in  $[\Sigma\text{BC}^*]$  compared with  $[\text{SO}_4^*]$  during the same time period (Table 5). However, only Lake Grunnetjenn exhibited a significant ( $P = 0.045$ ), change in ANC (decreasing trend) during period May 2009–October 2012 (Fig. 6).

Based on the Regional Kendal test, significant decreases in Tot-N ( $-53 \mu\text{g N L}^{-1} \text{ year}^{-1}$ ,

$P = 0.008$ ) and  $\text{NO}_3^-$  ( $-2 \mu\text{eq N L}^{-1} \text{ year}^{-1}$ ,  $P = 0.0001$ ) occurred from 2009 to 2012, while the decrease in  $\text{NH}_4^+$  was not significant ( $P = 0.071$ ). The highest post-fire concentrations of  $\text{NH}_4^+$  and  $\text{NO}_3^-$ , did not occur during the extreme conditions in August 2008, but during later snow melt episodes. The highest concentrations of  $\text{NH}_4^+$  came during snow-melt in 2010 and 2011 (Fig. 4), with concentrations of  $\approx 20 \mu\text{eq L}^{-1}$  in Lake Hundsvatn and  $\approx 30 \mu\text{eq L}^{-1}$  in Lake Rasvassvatn, while the highest  $\text{NO}_3^-$  concentrations came during the initial snow melt in Mars 2011, with concentrations of  $\approx 35$  and  $\approx 25 \mu\text{eq L}^{-1}$  in Lake Rasvassvatn and Lake Hundsvatn, respectively (Fig. 3).

## Discussion

Post-fire effects on water quality can be highly variable depending on factors such as hydrological regime, topography, geology, fire-size, fire-intensity, lake morphometry, the ratio of catchment to lake area, and water residence time (Ranalli 2004; Carignan et al. 2000). The relatively large differences in chemical response between lakes within the Mykland wildfire area, as well as large chemical variations in surface water responses of wildfires reported elsewhere (e.g. Smith et al. 2011), are primarily explained by these variables.

The wildfire in Mykland occurred in an area dominated by thin and patchy, base-poor and weathering resistant soils and bedrock. Thus, low ANC values in surface water are typical for this area. The long drought period ( $\approx 1$  month) followed by a 1 week-long wildfire, provided excellent conditions for sulfide oxidation within the catchments. Combined with the large mineralization of organic matter and killing of the vegetation, as direct effects of the wildfire, a substantial pool of inorganic cations and anions was ready to be mobilized from terrestrial to aquatic systems during the first rain events. However, the most extreme water chemical conditions did not occur during the first rain events, probably because the soil water first had to be replenished, and runoff was minor. This peak in water chemistry came at the end of a strong rain event, two months after the fire, and was likely the result of a strong flush of wildfire impacted water from the catchments to the lakes.

For the two wildfire affected lakes sampled during this extreme episode, Lake Hundsvatn and Lake Rasvassvatn, the most extreme chemical conditions were seen in Lake Hundsvatn, likely because of the high catchment-to-lake area ratio (17.6), compared with Lake Rasvassvatn (1.38). The significant delay in water chemical response after the fire and the variations in this delay, might be explained by factors such as post-fire hydrologic status of the catchments and lake residence times, factors which have received little attention in earlier studies of wildfire effects on surface waters. Ferreira et al. (2005) studied mechanisms driving hydrological and solute movement processes from terrestrial to aquatic systems as they change with time following forest fire, and Chessman (1986) reported time-variations in peak response of nitrate in various wildfire affected streams in south-eastern Australia.

As in our study, many articles have documented significant peaks in strong acid anions ([SAA]:  $\text{SO}_4^{2-}$ ,  $\text{Cl}^-$  and  $\text{NO}_3^-$ ) and base cations ([ $\Sigma\text{BC}$ ]:  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  and  $\text{K}^+$ ) in lakes and streams during the initial post-fire period (Bayley et al. 1992b; Carignan et al. 2000; Eriksson et al. 2003; Mast and Clow 2008; Smith et al. 2011), but the peak concentrations and the relative increases compared with pre-fire values, particularly in Lake Hundsvatn (Table 2), are among the highest reported in the literature, especially [ $\text{K}^+$ ] (peak:  $70 \mu\text{eq L}^{-1}$ ; 70-fold increase) and [ $\text{SO}_4^{2-}$ ] (peak:  $367 \mu\text{eq L}^{-1}$ ; 8.2-fold increase). This indicates

a very intense wildfire, with significant mineralization of organic matter and killing of vegetation within this acid sensitive area, heavily impacted by acid deposition for decades (Schartau et al. 2012). Bayley et al. (1992b) reported a peak concentration of  $\text{SO}_4^{2-}$  ( $634 \mu\text{eq L}^{-1}$ ) in a stream in Ontario, Canada, following a wildfire in 1980. The background concentration of  $\text{SO}_4^{2-}$  in this stream in 1980 was  $59.7 \mu\text{eq L}^{-1}$ , which is about twice the background levels of that present in the lakes within the Mykland area in 2012. But the very high peak concentration of sulfate in the wildfire affected streams in Canada in 1980, might also be attributed in part to acid deposition. As a result of the significant declines in sulfur emissions in Europe since 1980, the  $\text{SO}_4^{2-}$  concentration in southern Norway has declined by 70–80 % from 1980 to 2012 (Aas et al. 2012).

The concentration peaks in  $\text{Na}^+$  and  $\text{Cl}^-$  ( $[\text{Na}^+] = 125 \mu\text{eq L}^{-1}$ ,  $[\text{Cl}^-] = 165 \mu\text{eq L}^{-1}$ ) in the wildfire affected lakes, were also relatively high compared with many other studies. Both background and post-fire peak concentrations of these ions are very dependent on the distance from the sea, the main source for these constituents. The far most extreme  $\text{Na}^+$  and  $\text{Cl}^-$  conditions reported, are from wildfire affected streams in Tyresta, Sweden, August 1999 (Eriksson et al. 2000), but due to large amounts of brackish water used during the fire-fighting operations (Eriksson et al. 2003), their data are not directly comparable with most other wildfire impacted surface waters.

ANC in Lake Hundsvatn was  $11 \mu\text{eq L}^{-1}$  on June 25, 9 days after the fire. By July 9, ANC had declined to  $-3 \mu\text{eq L}^{-1}$ , and then dropped to  $-80 \mu\text{eq L}^{-1}$  on August 13, during the 3rd post-fire rain event, 2 months after the fire. A decrease in ANC due to faster mobilization of strong acid anions ( $\text{Cl}^-$  and particularly  $\text{SO}_4^{2-}$ ) compared with base cations during the initial post-fire period, has earlier been described in base-poor boreal forest streams in Ontario, Canada (Bayley et al. 1992b). They reported a mean annual ANC decrease of 20 % compared with the mean ANC value of the last pre-fire year, but the annual mean ANC value was still positive. However, even though not commented on in their study, we assume that a severe decline in ANC must have occurred during the extreme peak in  $\text{SO}_4^{2-}$  ( $634 \mu\text{eq L}^{-1}$ ) reported soon after the wildfire.

After the initial post-fire period, [ $\text{SO}_4^{2-}$ ] decreased faster than [ $\Sigma\text{BC}^*$ ] in Lake Hundsvatn water the

**Table 6** Calculated concentration decreases ( $\mu\text{eq L}^{-1}$ ) in non-marine base cations ( $\text{BC}^*$ ) and non-marine sulfate ( $\text{SO}_4^*$ ) from August 2008 to August 2009 ( $n = 6$ ), and from 2009 to August2012 in Lake Hundsvatn, showing different first order reaction decays for the two periods,  $C_{t=0}$  and  $C_{t=\text{end}}$  is the measured concentrations at the start and the end of the calculated periods

Period		$C_{t=0}$	$C_{t=\text{end}}$	k	SE k	$r^2$	p	$t_{1/2}$ (days)	$t_{1/2}$ (years)
Aug08–Aug09	$\text{BC}^*$	272	103	$-2.21\text{E}-03$	$5.02\text{E}-04$	0.83	0.012	314	0.86
	$\text{SO}_4^*$	350	66	$-3.87\text{E}-03$	$6.06\text{E}-04$	0.91	0.003	179	0.49
Aug09–Aug12	$\text{BC}^*$	103	32	$-7.36\text{E}-04$	$1.22\text{E}-04$	0.52	0.000	942	2.58
	$\text{SO}_4^*$	66	25	$-8.62\text{E}-04$	$8.90\text{E}-05$	0.73	0.000	804	2.20

k slope of the equations, SE k standard error of k

following months. Theoretically, assuming first order decay patterns (Table 6),  $[\Sigma\text{BC}^*]$  was equal to  $[\text{SO}_4^*]$  in Lake Hundsvatn, i.e.  $\text{ANC} = 0$ , 150 days ( $\approx 5$  months) after the severe drop in ANC in August 2008. This means that predominantly positive ANC values were re-established in Lake Hundsvatn five months after the extreme water chemical peak, seven months after the fire. In May 2009, almost one year after the wildfire, ANC was back to almost pre-fire values, i.e.  $\text{ANC} \approx 20 \mu\text{eq L}^{-1}$  (Fig. 6).

From May 2009 to October 2012, the concentrations of  $[\text{SO}_4^*]$  decreased faster than  $[\text{BC}^*]$  in four of the six wildfire affected lakes, but without any significant increase in ANC in five of the lakes. Only Lake Grunnetjenn showed a significant declining trend ( $P = 0.045$ ) in ANC during this period. In the two lakes with lowest ANC, Lake Fisketjenn and Lake Heitjenn, the concentration of  $\Sigma\text{BC}^*$  declined somewhat faster than  $[\text{SO}_4^*]$  (Table 5), but in neither of these lakes significant ANC trends were observed from May 2009 to October 2012 (Fig. 6). The most reasonable explanations for no significant ANC changes in lakes after the first post-fire year, are likely an interplay between several factors that decrease ANC as:

- Faster decrease in  $[\Sigma\text{BC}^*]$  in surface water compared with  $[\text{SO}_4^*]$ , after the chemical post-fire peak, as recorded in two of the wildfire affected lakes, implies a gradual ANC decrease by time.
- Significant decrease in ionic strength in surface water after a wildfire, implies a pure dilution effect, which *per se* implies a decrease in ANC by time when the ANC is positive.

and factors that increase ANC as:

- Faster decrease in  $[\text{SO}_4^*]$  in surface water compared with  $[\Sigma\text{BC}^*]$ , after the chemical post-fire

peak, as recorded in 4 of the wildfire affected lakes, implies a gradual ANC increase by time.

- Significant decrease in ionic strength in surface water after a wildfire, implies a pure dilution effect, which *per se* implies an increase in ANC by time when the ANC is negative.
- The significant increase in TOC observed in the wildfire effected lakes (Table 6) after the first post-fire year, should gradually lead to an ANC increase by time, due to increased concentration of weak organic acids and the fact that increase in TOC also implies increase in base cations and subsequent increase in ANC (Lydersen et al. 2004).

In addition, the relative trends in  $[\text{Cl}^-]$  and  $[\text{Na}^*]$  affect trends in ANC.

Despite very low pH and high concentrations of  $\text{Al}_i$ , the toxic, inorganic, cationic forms of Al (Schofield 1977; Dickson 1978) during the extreme event in August 2008, the concentrations of total monomeric aluminum ( $\text{Al}_a$ ) were very similar in Lake Hundsvatn, Lake Rasvassvatn and Birkenes, i.e. 368, 295 and  $310 \mu\text{g Al L}^{-1}$  (Table 2). The most likely reason for the much higher  $\text{Al}_i$  in the wildfire affected lakes compared with Birkenes, is a combination of decrease in organic matter and a substantial increase in sulfuric acid (pH decrease) as direct effects of oxidation of organic and inorganic compounds within the wildfire area.

Little data exists on pH, ANC and aluminum in surface water after wildfires. In the Tyresta wildfire in Sweden, where brackish water was used during the fire-fighting, one of the least affected streams, showed a pH decrease to pH 4.5 and an  $\text{Al}_i$  increase to  $330 \mu\text{g Al L}^{-1}$  (Eriksson et al. 2003), conditions very similar to the levels observed during the post-fire peak in Lake Hundsvatn 2008. The extreme low pH and ANC and the very high concentrations of  $\text{Al}_i$  measured



in Lake Hundsvatn (pH: 4.42; ANC:  $-80 \mu\text{eq L}^{-1}$ ;  $\text{Al}_i$ :  $326 \mu\text{g Al L}^{-1}$ ) and Lake Rasvassvatn (pH: 4.76; ANC:  $-39 \mu\text{eq L}^{-1}$ ;  $\text{Al}_i$ :  $273 \mu\text{g Al L}^{-1}$ ) should be lethal for fish species such as perch and brown trout (Lydersen et al. 2002, 2004), fish species present within the wildfire area. However, no significant fish death has been reported. There might be several reasons for this. Lack of documentation as very few people were in the area at that time, the extreme water chemical episode was too short to have substantial lethal impact and/or fish might have migrated down into the marginal wildfire affected hypolimnion water, as both lakes were temperature stratified at that time of the year.

Large post-fire variations in export of N and P from different forest environments to aquatic systems, are reported in the literature (Smith et al. 2011). In our study, increases in  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and total N were seen during the first extreme water chemical peak in August 2008, but the concentrations were still relatively low. The low initial post-fire increases in N-compounds, followed by higher levels of these compounds during winter and snowmelt in the first post-fire years in our study, are also reported in other wildfire studies (Bayley et al. 1992b; Carignan et al. 2000; Mast and Clow 2008). One exception is the lack of effects of  $\text{NH}_4^+$  in streams after the wildfire in Glacier National Park, Montana, reported by Mast and Clow (2008). As there are no comparable pre-fire values of total phosphorous (Tot-P), we cannot quantify the immediate wildfire effects, but the concentrations of Tot-P in the lakes during the first post-fire year were relatively low. Most wildfire studies document an increase in Tot-P during the initial post-fire period, followed by a gradual decrease again back to pre-fired levels during a few years (Ranalli 2004).

The fact that nitrate concentrations remained elevated in winter, especially related to snowmelt episodes during the three first post-fire winters, and fell back to pre-fire concentrations in the growing season, two years after the wildfire, are time patterns very similar to that observed by Mast and Clow (2008). Thus, two years after the wildfire, the N-uptake by vegetation has increased sufficiently to return concentrations to near background levels during the growing season.

The moderate concentration increases in  $\text{NO}_3^-$  and  $\text{NH}_4^+$  observed after wildfire in Mykland, agree well with earlier studies of wildfire affected streams in

base-poor boreal forest systems in Ontario, Canada (Bayley et al. 1992b) and in northern subalpine rocky mountains areas in Montana, USA (Mast and Clow 2008), and in wildfire affected lakes from boreal subarctic areas in northern Alberta (McEachern et al. 2000). There might be several causes for only moderate increases of nitrogen compounds in wildfire impacted surface waters:

- Very good oxidation conditions during the wildfire imply large amounts of  $\text{NO}_x$  emissions to the atmosphere and subsequent large transport of  $\text{NO}_x$  gases out of the catchments.
- Boreal forests are nutrient poor environments particularly low in N (Foster and Morrison 1976; Gosz 1981).
- Soils are too shallow to allow nutrients to leach below the rooting zone. Accordingly, most of the nutrients are probably retained by the new vegetation (Bayley et al. 1992b)

Regarding TOC, there are no comparable pre-fire data for evaluation of immediate effects of wildfire, but the significant increase in TOC observed from May 2009 to October 2012, indicates an initial TOC decrease, which might be directly related to combustion of organic matter during wildfire, and the subsequent increase in TOC the following years as new vegetation is gradually re-established in the area. Similar post-fire trends in TOC in surface water have also been documented in other studies (McEachern et al. 2000; Carignan et al. 2000).

## Conclusion

The wildfire in Mykland occurred in an area basically consisting of thin and patchy, base-poor soils and slowly weatherable rocks.

The most severe chemical conditions in wildfire affected surface waters occurred during some of the first post-fire rain events, but the amount of rain in relation to the hydrological status of the catchment prior to these events, and lake residence time, are decisive factors regarding when the most extreme water chemical conditions arise in surface waters affected by wildfire.

During the first post-fire hydrological episodes, strong acid anions as  $\text{Cl}^-$  but primarily  $\text{SO}_4^{2-}$ , are mobilized significantly faster from terrestrial to



aquatic systems compared with base cations ( $\Sigma\text{BC}$ ), which means that the acid neutralizing capacity (ANC:  $[\Sigma\text{BC}] - [\text{SAA}]$ ) of lakes decline. For lakes with low pre-fire ANC values, significant drops in ANC down to very low negative values might occur, with subsequent increase in  $[\text{H}^+]$  and toxic inorganic Al  $[\text{Al}_i]$  to levels potentially harmful to aquatic life. In lakes with higher pre-fire ANC values, the drop in ANC during the first post-fire period is too small to cause severe acidification, and therefore harmless to aquatic life. Normal ANC values seem to be reestablished relatively fast in wildfire affected lakes, often within the first post-fire year.

After the extreme, post-fire water chemical peak, the main mobile anions ( $\text{Cl}^-$  and  $\text{SO}_4^{2-}$ ) decreased faster than  $\Sigma\text{BC}$ . This combined with increase in TOC should imply a potential increase in ANC with time after fire in wildfire affected lakes. However, at the same time, the continuous dilution effect back to pre-fire levels, might counteract this potential ANC increase. This might explain why no ANC increase was seen in five of the six wildfire affected lakes, from one year after the fire and the next three post-fire years.

Increases in nutrients, N and P, and decreases in TOC in surface water during the initial post-fire period, followed by a gradual decrease in N and P and increase in TOC during the next post-fire years, have also been seen in surface waters from other wildfire impacted areas.

The water chemistry in the wildfire affected lakes in the Mykland area, had almost returned to pre-fire conditions 4.5 years after the wildfire. Thus, in this area of Norway, with significant reductions in acid deposition during the last decades, with  $[\text{SO}_4^*]$  reduction by 75–88 % in lakes from 1980 to 2011 (Schartau 2012), the most severe water chemical episodes, are again primarily driven by sea-salt events, directly linked to hydrologic episodes, directly as a result of heavy rain, or indirectly during snow melt as the sea-salt episode(s) are “accumulated” in the snowpack.

Hydrological regime, topography, geology, fire intensity, fraction of catchment burnt, lake morphometric conditions such as catchment-to-lake area ratio and lake residence time, are key factors for the variations in water chemical response in wildfire affected lakes.

**Acknowledgments** This study is part of a forest fire Project (Pyrowater) funded by the Norwegian Research Council (NRC) during the period 2009–2012. We greatly appreciate the Norwegian Water Research Institute (NIVA), for funding and initiating water chemical investigations during the very

essential, initial post-fire period. We also want to acknowledge the technical staff at Telemark University College, Bjørn Steen and Karin Brekke Li, who have conducted almost all the water chemical analysis, and Dick Wright (NIVA) for his fruitful scientific and linguistic contribution during the preparation of this manuscript.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- Aas W, Solberg S, Manø S, Yttri KE (2012) Monitoring long range transported air and precipitation. Atmospheric contributions–2011. (In Norwegian with English summary). Report 1126/2012. TA-2940/2012, pp 206
- Aherne J, Larssen T, Cosby BJ, Dillon PJ (2006) Climate variability and forecasting surface water recovery from acidification: modelling drought-induced sulfate release from wetlands. *Sci Total Environ* 365(1–3):186–199
- Barnes RB (1975) Determination of species forms of aluminum in natural water. *Chem Geol* 15(3):177–191
- Bayley SE, Schindler DW, Beaty KG, Parker BR, Stainton MP (1992a) Effects of multiple fires on nutrient yields from streams draining boreal forest and fen watersheds—nitrogen and phosphorus. *Can J Fish Aquat Sci* 49(3):584–596
- Bayley SE, Schindler DW, Parker BR, Stainton MP, Beaty KG (1992b) Effects of forest-fire and drought on acidity of a base-poor boreal forest stream—similarities between climate warming and acidic precipitation. *Biogeochemistry* 17(3):191–204
- Beaty KG (1994) Sediment transport in a small stream following 2 successive forest-fires. *Can J Fish Aquat Sci* 51(12):2723–2733
- Carignan R, D’Arcy P, Lamontagne S (2000) Comparative impacts of fire and forest harvesting on water quality in Boreal Shield lakes. *Can J Fish Aquat Sci* 57:105–117
- Chessman BC (1986) Impact of the 1983 wildfires on river water quality in East Gippsland, Victoria. *Aust J Mar Freshw Res* 37:399–420
- Cronan CS, Schofield CL (1979) Aluminum leaching response to acid precipitation: effects on high elevation watersheds in the Northeast. *Science* 204:304–306
- Dickson W (1978) Some effects of the acidification of Swedish lakes. *Verh Internat Verein Limnol* 20:851–856
- Drabloes D, Sevaldudr I, Timberlid JA (1980) Historical land-use changes related to fish status development in different areas of southern Norway. In: Drabloes D, Tollan A (eds) Ecological impact of acid precipitation. Proceedings of an international conference, Sandefjord, March 11–14 1980: 367–369
- Driscoll CT (1984) A procedure for the fractionation of aqueous aluminum in dilute acidic waters. *Int J Environ Anal Chem* 16(4):267–283
- Eriksson H, Edberg F, Borg H (2000) Water chemical effects of the wildfire and the fire-fighting. In: Pettersson U (ed) The wildfire in Tyresta 1999. Documentation of effects. (In

- Swedish with English summary). The Swedish environmental protection agency, Report 5604:156–174
- Eriksson H, Edberg F, Borg H (2003) Effects of forest fire and fire-fighting operations on water chemistry in Tyresta National Park, Stockholm, Sweden. *J Phys IV* 107:427–430
- Ferreira AJD, Coelho COA, Boulet AK, Lopes FP (2005) Temporal patterns of solute loss following wildfires in central Portugal. *Int J Wildland Fire* 14(4):401–412
- Finkelstein DB, Pratt LM, Curtin TM, Brassell SC (2005) Wildfires and seasonal aridity recorded in Late Cretaceous strata from south-eastern Arizona. *USA Sedimentol* 52(3): 587–599
- Flannigan MD, Wotton BM (2001) Climate, weather, and area burned. In: Johnson EA, Miyanishi K (eds) *Forest fires—behavior and ecological effects*. Academic Press, San Diego, pp 351–373
- Foster NW, Morrison IK (1976) Distribution and cycling of nutrients in a natural *Pinus banksiana* ecosystem. *Ecology* 57:110–120
- Gosz JR (1981) Nitrogen cycling in coniferous ecosystems. In: Clark FE, Rosswall TH (eds) *Nitrogen cycling in terrestrial ecosystems: processes, ecosystems strategies, and management implications*. *Ecol. Bull., Stockholm* 33:415–426
- Helsel DR, Frans LM (2006) Regional Kendall test for trend. *Environ Sci Technol* 40(13):4066–4073
- Hindar A, Henriksen A, Toerseth K, Semb A (1994) Acid water and fish death. *Nature* 372(6504):327–328
- Høgberget R (2010) The wildfire in Mykland 2008. Results after 2 years monitoring of chemical effects in lakes. (In Norwegian with English summary). NIVA-report 5979-2010, pp 44
- Huitfeldt-Kaas H (1922) Causes to mass death of salmon and brown trout in the River Frafjordelva. River Hellelva and River Dirdalselva in Ryfylke, autumn 1920. (In Norwegian). *Norges Jaeger og Fiskerforenings tidsskrift* 51:37–44
- IPCC (2001) *Climate change 2001: the scientific basis*. In: Houghton JT, Ding Y, Griggs DJ et al (eds) *Contribution of working group I to the third assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge
- Jeffries DS, Clair TA, Couture S, Dillon PJ, Dupont J, Keller W, McNicol DK, Turner MA, Vet R, Weeber R (2003) Assessing the recovery of lakes in southeastern Canada from the effects of acidic deposition. *Ambio* 32(3):176–182
- Karr JR, Rhodes JJ, Minshall GW, Hauer FR, Beschta RL, Frissell CA, Perry DA (2004) The effects of postfire salvage logging on aquatic ecosystems in the American West. *Bioscience* 54(11):1029–1033
- Kelly EN, Schindler DW, St Louis VL, Donald DB, Vlacicka KE (2006) Forest fire increases mercury accumulation by fishes via food web restructuring and increased mercury inputs. *Proc Natl Acad Sci USA* 103(51):19380–19385
- Lamontagne S, Carignan R, D'Arcy P, Prairie YT, Pare D (2000) Element export in runoff from eastern Canadian Boreal Shield drainage basins following forest harvesting and wildfires. *Can J Fish Aquat Sci* 57:118–128
- Laudon H, Dillon PJ, Eimers MC, Semkin RG, Jeffries DS (2004) Climate-induced episodic acidification of streams in central Ontario. *Environ Sci Technol* 38(22):6009–6015
- Lydersen E, Henriksen A (1995) Seasalt effects on the acid neutralizing capacity of streamwaters in southern Norway. *Nord Hydrol* 26(4–5):369–388
- Lydersen E, Löfgren S, Arnesen RT (2002) Metals in scandinavian surface waters: effects of acidification, liming and potential re-acidification. *Crit Rev Environ Sci Technol* 32:73–295
- Lydersen E, Larssen T, Fjeld E (2004) The influence of total organic carbon (TOC) on the relationship between acid neutralizing capacity (ANC) and fish status in Norwegian lakes. *Sci Total Environ* 326(1–3):63–69
- Mast MA, Clow DW (2008) Effects of 2003 wildfires on stream chemistry in Glacier National Park. *Mont Hydrol Process* 22(26):5013–5023
- McEachern P, Prepas EE, Gibson JJ, Dinsmore WP (2000) Forest fire induced impacts on phosphorus, nitrogen, and chlorophyll a concentrations in boreal subarctic lakes of northern Alberta. *Can J Fish Aquat Sci* 57(Suppl. 2):73–81
- Nystroem U, Hultberg H, Lind B (1995) Can forest-soil liming mitigate acidification of surface waters in Sweden? *Water Air Soil Pollut* 85:1855–1860
- Planas D, Desrosiers M, Groulx SR, Paquet S, Carignan R (2000) Pelagic and benthic algal responses in eastern Canadian Boreal Shield lakes following harvesting and wildfires. *Can J Fish Aquat Sci* 57:136–145
- Ranalli AJ (2004) A summary of the scientific literature on the effects of fire on the concentration of nutrients in surface waters. *US Geological Survey Open-File Report 2004-1296*: pp 23
- Reuss JO, Johnson DW (1986) Acid deposition and the acidification of soils and waters. *Ecological studies*, vol. 59. Springer, 62–65
- Ryan KC (2000) Global change and wildland fire. In: Brown JK, Smith JK (eds) *Wildland fire in ecosystems: effects of fire on Flora*. Gen. Tech. Rep. RMRS-GTR-42–Vol. 2. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Ogden, 97–120
- Schär C, Vidale PL, Lüthi D et al (2004) The role of increasing temperature variability in European summer heatwaves. *Nature* 427:332–336
- Schartau AK, Fjellheim A, Walseng B et al. (2012) Monitoring long range transported air and precipitation. Annual Report—Effects 2011 (In Norwegian with English summary). TA-2934/2012, pp 160
- Schindler DW, Newbury RW, Beaty KG, Prokopowich J, Ruzsyczynski T, Dalton JA (1980) Effects of a windstorm and forest fire on chemical losses from forested watersheds and on the quality of receiving streams. *Can J Fish Aquat Sci* 37(3):328–334
- Schofield CL (1977) Acid snow-melt effects on water quality and fish survival in the adirondack mountains of New York state, U.S. Research Technical Completion Report A-072-NY (Office of Water Research and Technology, Department of the Interior, Washington D.C., pp 27
- Sevaldud I, Muniz IP, Kalvenes S (1980) Loss of fish populations in southern Norway. Dynamics and magnitude of the problem. In: Drabloes D, Tollan A (eds) *Ecological impact of acid precipitation*. Proceedings of an international conference, Sandefjord, March 11–14, 1980:350–351
- Smith HG, Sheridan GJ, Lane PNJ, Nyman P, Haydon S (2011) Wildfire effects on water quality in forest catchments: a review with implications for water supply. *J Hydrol* 396(1–2):170–192
- Weast RC (1988) *CRC handbook of chemistry and physics*, 1st student edition. CRC Press, Boca Raton, p F-100