

ICP Waters Report 127/2016

Biodiversity of macro-invertebrates in acid-sensitive waters:
trends and relations to water chemistry and climate



CORRIGENDUM

Corrections to the electronic version of the ICP Waters Report 127/2016 Biodiversity of macro-invertebrates in acid-sensitive waters: trends and relations to water chemistry and climate (NIVA REPORT S.NO. 7077-2016)

Page 18: “Figure 9. Trends in quarterly air temperature, precipitation, SO₄ and ANC for European river sites over the site-specific period covered by the biological data + denotes increase and – denotes decrease. Statistical significant trends are embedded in a grey circle.” changed to “Figure 9. Trends in biodiversity in European rivers and lakes over the last 15 to 30 years (see Tables 1 and 2 for exact periods). + Increasing trend, – decreasing trend. Statistical significant changes in diversity are embedded in grey circles.”

Oslo, 16.02.2017

Heleen de Wit

REPORT

Main Office

Gaustadalléen 21
NO-0349 Oslo, Norway
Phone (47) 22 18 51 00
Telefax (47) 22 18 52 00
Internet: www.niva.no

NIVA Region South

Jon Lilletuns vei 3
NO-4879 Grimstad, Norway
Phone (47) 22 18 51 00
Telefax (47) 37 04 45 13

NIVA Region East

Sandvikaveien 59
NO-2312 Ottestad, Norway
Phone (47) 22 18 51 00
Telefax (47) 62 57 66 53

NIVA Region West


Thormøhlensgate 53 D
NO-5006 Bergen Norway
Phone (47) 22 18 51 00
Telefax (47) 55 31 22 14

| | | |
|---|--|------------------------|
| Title Biodiversity of macro-invertebrates in acid-sensitive waters: trends and relations to water chemistry and climate | Serial number 7077-2016 ICP Waters report 127/2016 | Date September 2016 |
| Author(s) Gaute Velle, Shad Mahlum, Don T. Monteith, Heleen de Wit, Jens Arle, Lars Eriksson, Arne Fjellheim, Marina Frolova, Jens Fölster, Natalja Grudule, Godtfred A. Halvorsen, Alan Hildrew, Jakub Hruška, Iveta Indrikson, Lenka Kamasová, Jiří Kopáček, Pavel Krám, Stuart Orton, Takaaki Senoo, Ewan M. Shilland, Evžen Stuchlík, Richard J. Telford, Lenka Ungermanová, Magda-Lena Wiklund, Richard F. Wright | Topic group Acid rain | Distribution |
| | Geographical area Europe | Printed NIVA |

| | |
|--|--------------------|
| Client(s) Norwegian Environment Agency (Miljødirektoratet) United Nations Economic Commission for Europe (UNECE) | Client's reference |
|--|--------------------|

| |
|---|
| Summary This is a study of the international impact of acid emission reductions on vulnerable freshwater biodiversity of benthic macroinvertebrates. In order to properly attribute impacts it is necessary to also incorporate the potential effects of changes in temperature and precipitation. Most sites show an increasing number of species towards the present. This can be attributed to a decrease in acidification, while temperature has a secondary influence on diversity. We expect that the biodiversity will continue to increase in the future as acid deposition decreases. The widespread response of aquatic diversity resulting from emission reductions of acidifying components to the atmosphere demonstrates the potential of international policy for achieving positive effects on the state of the environment. |
|---|

| | |
|--|---|
| Four keywords 1. Biodiversity 2. Acidification 3. Climate 4. Water chemistry | Fire emneord 1. Biologisk mangfold 2. Forsuring 3. Klima 4. Vannkjemi |
|--|---|



Heleen de Wit
Project Manager



Øyvind Kaste
Research Manager

BIODIVERSITY OF MACRO-INVERTEBRATES IN ACID-SENSITIVE WATERS: TRENDS AND RELATIONS TO WATER CHEMISTRY AND CLIMATE

Gaute Velle^{1,2}, Shad Mahlum¹, Don T. Monteith³, Heleen de Wit⁴, Jens Arle⁵, Lars Eriksson⁶, Arne Fjellheim¹, Marina Frolova⁷, Jens Fölster⁶, Natalja Grudule⁸, Godtfred A. Halvorsen¹, Alan Hildrew⁹, Jakub Hruška⁸, Iveta Indriksone⁷, Lenka Kamasová¹⁰, Jiří Kopáček¹¹, Pavel Krám⁸, Stuart Orton⁹, Takaaki Senoo¹⁰, Ewan M. Shilland¹², Evžen Stuchlík¹⁰, Richard J. Telford², Lenka Ungermanová¹⁰, Magda-Lena Wiklund⁶, Richard F. Wright⁴

¹Laboratory for freshwater ecology and inland fisheries, Uni Research, Bergen, Norway

²Department of Biology, University of Bergen, Norway

³Centre for Ecology and Hydrology, Lancaster Environment Centre, Lancaster, United Kingdom

⁴Norwegian Institute for Water Research (NIVA), Oslo, Norway

⁵Federal Environment Agency, Section II 2.4 Inland Waters, Germany

⁶Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁷Latvian Environment, Geology and Meteorology centre, Riga, Latvia

⁸Czech Geological Survey, Prague, Czech Republic

⁹School of Biological and Chemical Sciences, Queen Mary University of London, United Kingdom

¹⁰Institute for Environmental Studies, Faculty of Science, Charles University in Prague, Czech Republic

¹¹Biology Centre of the Academy of Sciences of the Czech Republic, Institute of Hydrobiology, České Budějovice, Czech Republic

¹²ENSIS Ltd, Environmental Change Research Centre, Department of Geography, University College London, United Kingdom

Preface

The International Cooperative Programme on Assessment and Monitoring Effects of Air Pollution on Rivers and Lakes (ICP Waters) was established under the Executive Body of the United Nations Economic Commission for Europe (UNECE) Convention on Long-Range Transboundary Air Pollution (CLRTAP) at its third session in Helsinki in July 1985. UNECE is a catalyst in the international work aiming at reducing transboundary air emissions. Norway provides facilities for the ICP Waters Programme Centre at the Norwegian Institute for Water Research (NIVA). A programme subcentre was established at Uni Research. The Norwegian Environment Agency provides financial support to the work of the Programme Centre and with a representative leading the ICP Waters programme. In the long-term strategy for the Convention, adopted in 2010 by the Executive Body, it is stated that environmental effects of acidifying components, and its potential interaction with climate change and biodiversity, continue to be among the significant remaining problems with regard to air pollution effects on the environment. These can be addressed with the multi-pollutant/multi-effects approach of the Gothenburg Protocol.

The main aim of the ICP Waters Programme is to assess, on a regional basis, the degree and geographical extent of the impact of atmospheric pollution, in particular acidification, on surface waters. ICP Waters uses data from existing surface water biological and water chemical monitoring programmes in the participating countries, and the countries contribute to ICP waters voluntarily. The monitoring sites are generally acid-sensitive and representative of low acid-neutralising capacity (ANC) and low critical load levels. The ICP site network is geographically extensive and 30 countries in Europe and North America have partly or fully participated in the programme since its start. The Programme also promotes international harmonisation of monitoring practices. Harmonization practices include intercalibration of biological- and chemical monitoring. The participating countries submit chemical data to NIVA and biological data to Uni Research. The biological data has shown a general increase in the number of acid-sensitive taxa towards the present, indicating a recovery from acidification, in line with results from the chemical monitoring.

In this report we provide results from an extensive array of analyses on biological diversity with climatic and water chemistry drivers of the observed trends. The analysis includes about 1.8 million benthic macro invertebrates sampled from 89 European rivers and lakes collected between 1982 and 2014 at ICP Waters sites. In addition, data on water chemistry from the same sites and periods are included. Few, if any, comparable studies exist at this scale.

The report was prepared by ICP Waters subcentre in Bergen by lead Gaute Velle. Data was provided from Task Force members, and the results were discussed in several steps with the co-authors. We would like to thank all participating countries for data. In addition, we would like to thank those who contributed comments and suggestions to the draft report, especially Richard Wright, Heleen de Wit, Jens Fölster and Don Monteith.

Gaute Velle
ICP Waters Programme Subcentre Uni Research

Bergen, August 2016

Contents

| | |
|--|-----------|
| Summary | 5 |
| 1 Introduction | 6 |
| 1.1 Acid deposition and effects on freshwater biota | 6 |
| 1.2 Climate change and effects of temperature and precipitation on the biota | 7 |
| 1.3 Aims of study | 7 |
| 2 Methods | 9 |
| 2.1 Study sites | 9 |
| 2.2 Numerical analyses | 12 |
| 3 Results | 14 |
| 3.1 Environmental variables | 14 |
| 3.2 Species diversity | 15 |
| 3.3 Possible drivers of biodiversity | 22 |
| 4 Discussion | 23 |
| 4.1 Recovery from acidification as drivers of diversity | 23 |
| 4.2 Temperature and precipitation as drivers of diversity | 24 |
| 4.3 Stability- and length of the environmental gradients | 25 |
| 4.4 Implications for policy | 25 |
| 4.5 Perspectives | 26 |
| 5 References | 27 |
| 6 Reports and publications from the ICP Waters programme | 34 |

Summary

In this report, we provide results from analyses on the species diversity of benthic invertebrates from freshwater monitoring sites in the Czech Republic, Germany, Latvia, Norway, Sweden and the UK. The data includes about 1.8 million benthic macroinvertebrates from 5010 samples in 55 rivers and 34 lakes. The sampling periods cover between 12 and 29 years. In addition, data on water chemistry, precipitation and air temperature from the same sites and periods are included. The study sites were chosen as part of national monitoring programmes and most represent type-sites of nutrient poor waters that have been influenced by long-range air pollution leading to acidification. Our aim has been to find the impact of acid emission reductions on vulnerable freshwater biodiversity. In order to properly attribute impacts it is necessary to also incorporate the potential effects of changes in temperature and precipitation. To our knowledge, no comparable studies exist at this scale.

Increases in species diversity were found at 90% of the lake sites and 95% of the river sites, and with significant trends in 30% and 21% of the sites, respectively. With almost no exception, the concentration of sulphate decreased significantly. The air temperatures of most sites show increasing trends, albeit not significant for any sites during the period covered by the biological data. The amount of precipitation has increased significantly at 17% of the sites.

Most sites show a strong correlation between increasing diversity and a reduction in acidifying components. The pronounced increase in diversity at sites with the most pronounced chemical recovery and the strong correlation between sulphate and/ or ANC and diversity suggest that a reduction in acidifying components of the water has had a strong influence on species diversity of aquatic invertebrates. Most likely, sulphate and ANC have little effect on aquatic invertebrates, whereas pH and inorganic aluminium are the biological drivers.

Year-to-year variation in temperature was negatively correlated with diversity, suggesting that temperature has a secondary influence on diversity since the diversity is increasing, despite the negative correlation with temperature. The trends in warming for the periods we cover were not significant and therefore, cannot address effects from climate change. Still, the effects from temperature suggest that these communities will be sensitive to long-term climate change. There was a positive correlation between precipitation and species diversity for many river sites.

The strong correlation between acid components of the water and species diversity suggests that biodiversity will continue to increase when acid deposition decreases. However, environmental changes in the near future are expected to become less distinct for deposition and more distinct for climate, suggesting that aquatic invertebrate communities may experience stronger impacts from climate in the year to come.

The study has demonstrated that species diversity is a useful metric to accompany the acidification indices in monitoring of benthic invertebrates of acid sensitive sites, and for detection of temperature effects on biodiversity. The widespread response of aquatic diversity resulting from emission reductions of acidifying components to the atmosphere demonstrates the potential of international policy for achieving positive effects on the state of the environment.

1 Introduction

Biological diversity, or biodiversity, refers to the variety and variability of life on earth. Biodiversity as a term may include many aspects that describes the variability of life-forms, such as the variation of a specific gene within a species, the number of species at a site, the evenness of species within an ecosystem or the diversity of ecosystems in a landscape (Purvis and Hector, 2000; 2005). Earth's biodiversity has changed greatly since the initial colonisation of the land. Natural phenomena have caused five mass extinctions with an estimated 75% or more loss of species during the last 540 million years (Barnosky et al., 2011). In addition, many smaller extinction events have caused a decrease in species diversity (Zhu et al., 2006). Typically, a recovery from such events takes about 10 million years (Erwin, 2001; Kirchner and Weil, 2000). We are now facing the sixth mass extinction with an extinction rate of species that is 100 to 1000 times greater than normal background rates (Rockstrom et al., 2009; UN, 2005; Williams et al., 2015). No doubt, humans are directly or indirectly responsible for much of this most recent decline in species.

The loss of species is alarming since biodiversity has important consequences for ecosystem processes. All organisms perform functions in an ecosystem. A 'healthy' ecosystem can be defined as a system where all ecosystem functions are maintained at a natural rate. The loss of species, and also shifts in relative abundance, may halt or speed up certain ecosystem processes and cause changes that may influence the ecosystem and the ecosystem-services crucial for human well-being (Chapin Iii et al., 2000; Hooper et al., 2005).

Freshwater bodies are especially impacted by human perturbations, and freshwaters may be the most endangered ecosystems on Earth (Dudgeon et al., 2006; Sala et al., 2000). The major types of stressors that may combine to affect freshwater ecosystems adversely include over-exploitation, (water) pollution, habitat degradation, species invasion and flow modification (Dudgeon et al., 2006). Successful management of impacted ecosystems involves separation of natural phenomena from human-induced perturbations. We need to understand how biodiversity and ecosystem function have responded to past and present human influence (Condamine et al., 2013; Stendera et al., 2012). Only then can we mitigate for human influence and make predictions on the impact of future change (e.g., Markovic et al., 2014).

This is a study of the international impact of acid emission reductions on vulnerable freshwater biodiversity of benthic macroinvertebrates. In order to properly attribute impacts it is necessary to also incorporate the potential effects of changes in temperature and precipitation.

1.1 Acid deposition and effects on freshwater biota

Paleolimnological studies indicate that the pH started to decrease in surface waters already in the mid-nineteenth century, both in northern Europe and in North America (Battarbee and Charles, 1986). Surface water acidification has subsequently represented one of the major perceived drivers of biodiversity loss, as a consequence of the toxic effects of elevated hydrogen and aluminium ions particularly of a range of acid sensitive organisms. It is hard to assess the full biological impact of acidification since very few biological records exist to inform on the biological and ecological quality of surface waters prior to anthropogenic acidification. Acidification monitoring programmes were initiated only after belated recognition of the effects of acid deposition and rarely started before the 1980's. At that time, susceptible waters were heavily impacted by acidification, and the primary driver, sulphur deposition, had already peaked in most regions of concern. Declining fish populations in dilute lakes and streams were reported from the beginning of the 20th century in Norway and Sweden, and from the mid-20th century in Canada and the United States (Schofield, 1976). Detrimental effects on benthic invertebrates likely began prior to the effects on fish since some invertebrates are more sensitive than fish (Raddum et al., 1984). The threats to freshwaters accelerated during the second half of the last century, when the influence of acidification initiated biodiversity loss and alterations of ecosystem processes (Schindler, 1988).

During the last 30 years, surface water chemistry and biota have been monitored in national programmes, initiated through international cooperation such as the Convention on Long-range Transboundary Air

pollution (CLRTAP) to assess the environmental and ecological impact of emissions of sulphur and nitrogen species to the atmosphere. Monitoring programmes indicate a reduction in atmospheric pollution since the late 1980's (Garmo et al., 2014; Stoddard et al., 1999) resulting in improved water quality and ecological state in a broad range of geographical areas (Evans et al., 2001; Halvorsen et al., 2003; Hesthagen et al., 2011; Johnson and Angeler, 2010; Lento et al., 2012; Monteith et al., 2005; Stendera and Johnson, 2008). However, evidence for associated recovery in biodiversity has, to date, remained largely equivocal (Angeler and Johnson, 2012; Lento et al., 2012; Monteith et al., 2005; Murphy et al., 2014). The influence of reducing the acidity of acidified waters on biodiversity and ecosystem function is particularly poorly understood (Johnson and Angeler, 2010; Ledger and Hildrew, 2005) and it remains unclear whether chemical recovery will lead to a return of biodiversity and ecological states similar to those of pre-acidified conditions or, whether alternative states will emerge. The role of a changing climate also remains unclear.

1.2 Climate change and effects of temperature and precipitation on the biota

The term "climate" normally refers to prevailing weather conditions for a region averaged over multiple years. Globally averaged combined land and ocean surface temperature have increased by 0.85 °C (0.65 to 1.06) over the period 1880 to 2012, the period over which multiple independently produced datasets have been available (IPCC, 2013). In mainland Europe, the temperature increase varies by region and season. Given a scenario for the next 50-100 years of 2°C global warming, the temperature increase during winter is expected to be largest (about 3-3.5°C) in Northern Fennoscandia available (Vautard et al., 2014). During summer, temperature increases are expected to be largest in southern Europe (about 2.5°C). Large parts of northern and central Europe are expected to experience a substantial increase in precipitation (up to 50%) during winter. During summer, the amount of precipitation is expected to increase in Fennoscandia and to decrease in southern Europe. In northern regions, diurnal temperatures ranges have changed as a consequence of minimum temperatures are increasing about twice as fast as the maximum daily temperatures (Easterling et al., 1997). As a result, periods with sub-zero temperatures are shorter causing decreased snow cover and ice extent.

Past changes in climate have been major drivers of speciation and extinctions (Mayhew et al., 2012; Sahney et al., 2010) and temperature is generally considered to be one of the main mechanisms determining species distribution patterns (Parmesan and Yohe, 2003; Quist et al., 2004). This is because many biological rates are temperature-dependent and temperature has a major impact on ecological processes (Brown et al., 2004). Warmer water temperatures will speed up physiological processes (Buisson et al., 2012; Parmesan and Yohe, 2003), and could cause variations in the number of generations a population of a species produces per year (Knell and Thackeray, 2016; Raddum et al., 2008), species replacements (Sweeney, 1984; Velle et al., 2010b), a shift in competitive advantages of communities (Isaak and Hubert, 2004) and changes in the relative timing of life cycles of species occupying different trophic levels (Thackeray et al., 2016). Thermal regimes also indirectly govern the habitat since temperature is a key factor for processes within lake, stream and terrestrial ecosystems.

In sum, the result will be altered relative abundances of taxa and altered species diversity. However, the direction and rate of change in biodiversity caused by the present climate change is known only to some extent (Heller and Zavaleta, 2009; Mantyka-pringle et al., 2012). We can expect the most pronounced changes in biodiversity to occur where the warming is large, such as in high latitudes on the Northern hemisphere, and the smallest changes in biodiversity to occur where the warming is small, such as the mid-southern hemisphere. It is difficult to assess the influence that precipitation has on biodiversity in freshwaters since the effects are not direct, unless availability of water is a limiting factor or causes flooding. In addition to direct impacts of flow, such as scouring energy, changes in precipitation may affect habitat and water quality, particularly with respect to the frequency and intensity of acid episodes.

1.3 Aims of study

The high quality biological and chemical data available from the national monitoring networks participating in ICP Waters cover a period of major reductions in acid deposition. This provide conditions

for a hemispheric-scale natural experiment with which to assess, for the first time, the impacts of emission reductions on freshwater biodiversity, and the possible confounding or amplifying effects of climate change. In the present study, we have used extensive monitoring data on benthic macro invertebrates (Figure 1.) and water chemistry in streams and lakes sampled from mid-1980s and to 2014. These data are supported by estimates of average monthly modelled air temperature and precipitation for the $0.5^\circ \times 0.5^\circ$ grid squares within which each site occurs, to provide some control for possibly influential long-term change in either water temperature or flow. Participating countries in this study include the Czech Republic, Germany, Latvia, Norway, Sweden and the UK. All countries participate in both hydrochemical- and macro-invertebrate intercalibrations that ensure international comparability of methods and taxonomy.

Our primary aims have been to:

1. Identify significant trends in the species diversity of benthic macroinvertebrates from the beginning of the monitoring period and to the present
2. Test, and quantify, the extent to which such biological changes can be explained by changes in water chemistry, temperature and precipitation

In this study, we have used the species richness of Ephemeroptera, Plecoptera and Trichoptera, so-called EPT-taxa (hereby referred to as species diversity) as a measure of diversity. We have chosen to include EPT-taxa only and not the full species assemblage since the taxonomic resolution is generally to species level for EPT, whereas most other taxa are identified at a coarser taxonomic resolution. We can expect the response curves to environmental variables to be narrower for individual species than for a coarser taxonomic entity, thereby reducing noise. EPT taxa are known to be sensitive to acidification (Raddum et al., 1988; Raddum and Fjellheim, 1995), in addition to other pressures, e.g. organic pollution.




Figure 1. Benthic invertebrates from Norwegian rivers. From left to right the mayfly *Ephemerella aurivillii* and the caddisfly *Atrypodes aterrimus*. Photos by G. Velle (Uni Research).

2 Methods

2.1 Study sites

Our study sites include lakes and rivers from the Czech Republic, Germany, Latvia, Norway, Sweden and UK (Table 1 and 0). The data were collected between 1982 and 2014 and includes about 1.8 million benthic macro-invertebrate individuals from 3671 samples in 53 rivers and 1570 samples in 34 lakes (Table 1, Table 2, Figure 2). In addition, the data include about 26 000 samples of water analysed for chemistry. The sites form part of national biological- and chemical monitoring programmes in running and standing freshwaters (Halvorsen et al., 2002; Horecký et al., 2002; Horecký et al., 2006; Horecký et al., 2013; Johnson and Goedkoop, 2007; Kernan et al., 2010; Schaumburg et al., 2008).

Table 1. Samples from rivers with number of biological samples, number of sub-samples, average sample size, number of paired chemistry samples and sampling period. All biological samples have been taken by kick-sampling. *The Norwegian rivers include a total of 59 sampling stations. Full details on the sites are given in Velle et al. (2013).

| | Number of Rivers | Biological samples | Sub-samples | Average sample size | Chemical samples | Sampling period |
|----------------|------------------|--------------------|-------------|---------------------|------------------|-----------------|
| Czech Republic | 3 | 35 | 1 | 1102 | 117 | 1999-2011 |
| Germany | 28 | 1037 | 1 | NA | 7863 | 1982-2010 |
| Latvia | 2 | 48 | 1 | 150 | 302 | 1997-2014 |
| Norway | 3* | 2181 | 1 | 217 | 2316 | 1987-2014 |
| Sweden | 6 | 104 | 5 or 6 | 236 | 2580 | 1985-2014 |
| UK | 11 | 266 | 3 or 5 | 1312 | 3015 | 1988-2013 |

Table 2. Samples from lakes with number of biological samples, number of sub-samples, average sample size, number of paired chemistry samples and sampling period. Note that the Swedish littoral, sublittoral and profundal samples were taken from the same seven lakes. Grab samples were taken by use of an Ekman grab. Full details on the sites are given in Velle et al. (2013).

| | Lakes | Biological samples | Sub-samples | Sampling method | Average sample size | Chemical samples | Sampling period |
|--------------------|-------|--------------------|-------------|-----------------|---------------------|------------------|-----------------|
| Germany | 1 | 37 | 1 | Kick | NA | 1179 | 1997-2009 |
| Norway | 20 | 473 | 1 | Kick | 282 | 3051 | 1997-2013 |
| Sweden littoral | 7 | 284 | 5 | Kick | 195 | 1905 | 1986-2013 |
| Sweden sublittoral | 7 | 261 | 5 | Grab | 227 | 1650 | 1989-2013 |
| Sweden profundal | 7 | 357 | 5 | Grab | 183 | 1636 | 1986-2013 |
| UK | 6 | 158 | 3-7 | Kick | 442 | 600 | 1988-2013 |




Figure 2. Sampling sites. Red dots denote rivers and blue dots denote lakes. *There are a total of 112 sampling sites in rivers and 48 sampling sites in lakes.*

The chemical sensitivity of catchment bedrock varies among sampling regions, but apart from Latvia, most sites are in acid-sensitive bedrock consisting of gneiss, granite or quartzite. The Latvian sites are situated on claystone including smaller fractions of dolomite and gypsum. The northernmost catchments are characterised by boreal vegetation, while the southernmost sites are situated in cool temperate vegetation. Some of the sites, especially in Germany and Latvia, are in small stands of forest surrounded by farmland.



Figure 3. Sampling for benthic invertebrates in Norwegian rivers. Photo by G.Velle (Uni Research).

Most invertebrates were collected by kick sampling (Figure 3) (Frost et al., 1971), following the ICP Waters manual (Wathne et al., 2010). In this procedure, the substrate is disturbed and collected in a 0.25 mm mesh net. An Ekman grab was used in the Swedish sublittoral- and profundal lake sites. The sampling interval varied among sites and years with an average of about 1.5 samples per sampling station per year from the time most monitoring programmes opened in 1987 and up to the present. Each sample was sorted under a stereo microscope in the laboratory, and macroinvertebrates subsequently identified. The consistency of the taxonomic resolution through the 30-year sampling period was checked and corrected for by methods described in Velle et al. (2013). In short, data from sites with inconsistent taxonomic resolution were reduced to a coarser taxonomic level prior to numerical analysis.

Chemical variables include pH, conductivity, alkalinity, calcium (Ca^{2+}), magnesium (Mg^{2+}), potassium (K^+), sodium (Na^+), chloride (Cl^-), sulphate (SO_4^{2-}), total nitrogen (TotN), nitrate (NO_3^-), total organic carbon (TOC), hydrogen (H^+) and labile aluminium (LAL). Acid-neutralizing capacity (ANC) was calculated as $(\text{Ca} + \text{Mg} + \text{Na} + \text{K} + \text{NH}_4) - (\text{Cl} + \text{SO}_4 + \text{NO}_3)$ with units in $\mu\text{Eq/l}$. For a detailed description of measured water chemistry, see Skjelkvaale and de Wit (2011).

Climate variables include monthly data for total precipitation, mean daily minimum air temperature, mean daily maximum air temperature and mean average daily air temperature. These are modelled gridded data for the 0.5-0.5 degree grid cells within which each site is located, obtained from the Climate Research Unit CRU TS3.21 (Harris et al., 2014). The data covers the period 1901-2014 and is modelled from measurements provided by national meteorological networks. In the absence of direct measurements of water temperature, we use air temperature as a surrogate. The relationship between water and air temperature is, however, by no means straightforward. As a general rule of thumb, water temperatures tend to vary spatially and seasonally in a similar manner to air temperatures, as the energy balance of both media is dominated by a similar set of factors (Layden et al., 2015; Livingstone and Lotter, 1998; Toffolon et al., 2014). However, factors such as riparian shading, effects of ice cover, snowmelt and thermal stratification etc., can also lead to significant differences in both space and time.

2.2 Numerical analyses

We first tested for linear trends in species diversity for the sites and periods covered by the biological data, and then tested for correlations between environmental variables and species diversity. Several environmental variables are presently changing concurrently in European freshwaters. The major potential drivers at the sites in this database are acid deposition (Stoddard et al., 1999; Velle et al., 2013) and climate (IPCC, 2013). The sites were specifically chosen to be free from pollution by nutrients (eutrophication).

As a consequence of the dominant effect of deposition chemistry on these systems, temporal variation in many of the chemical variables measured in water are closely correlated, such as some major ions, and some chemical variables are derivatives of others, such as alkalinity and pH. In addition, not all chemical variables were measured at every site. In order to reduce collinearity and noise, we retained sulphate (SO_4), acid neutralizing capacity (ANC), mean monthly air temperature and precipitation in the analyses of environmental drivers. ANC is the overall buffering capacity against acidification and effects from acid rain, and sulphate is the main acidifying component of acid rain. Both chemical variables have previously been found to correlate with diversity.

Biological sensitivity to acidification or climate varies according to site and region (Moe et al., 2010; Raddum and Skjelkvale, 2001). We can also expect that the response to acidification may override the response to climate for sites with a dominating change along the acidification axis. Hence, in an attempt to reduce noise and ease interpretation of results, the sites were clustered into four groups based on likely sensitivity to acidification. Here, pH 5.5 was chosen as clustering basis since the inflection point in the titration curve of pH versus ANC is at pH 5.4-5.6 (Figure 4). A small change of ANC in this pH region will cause a large change in pH. The clusters of sites are:

- Group 1. All sites
- Group 2. Sites where pH always has remained above 5.5 (least acidified sites)
- Group 3. Sites where pH has crossed 5.5 one or multiple times (transition sites)
- Group 4. Sites where pH always has remained below 5.5 (acidified sites)




Figure 4. The change in pH given changes in ANC in freshwaters. Shown are the theoretical curves for several levels of TOC in the water calculated assuming pCO_2 of four times atmospheric and negligible concentrations of labile aluminium (after Fölster et al., 2007).

Biological data inevitably involves noise and random effects, such as clustering of species and specimens within the habitat and changes in species identification (taxonomy) through time. Also, separating the influence of multiple environmental variables on the biological communities is challenging since the effects can overlap and or/ be cumulative and/ or be masked and distorted by each other or confounding variables. Other numerical pit-falls include (1) auto-correlation, (2) pseudo replication, (3) difference in

time-scales, or sampling interval, (4) un-known delayed response in the biological communities where animals sampled at a certain time have not responded to the chemical and climate variables at this time but rather at a previous unknown time, and (5) multiple testing.

In an attempt to correct for potential pit-falls, we have used Markov Chain Monte Carlo generalized mixed effects model (MCMCglmm; Hadfield, 2010), which is a method for analysing multi-responses in the family of Bayesian statistics. MCMCglmm allows the incorporation of random effects, such as site and country, which are sources of pseudo-replication and local effects (Bolker et al., 2009). In addition, the data used in this study have significant over dispersion (residual deviance/degrees of freedom > 2), which can be accounted for using MCMCglmm (Hadfield, 2010). We used an indiscriminate priori that results in approximations similar to the Laplace approximation used in other mixed modelling packages. The model included fixed and random effects and a Poisson distribution, and assumed linearity, but not necessarily a straight line. Interaction among explanatory variables was not tested directly. However, the analysis quantifies the effect of specific explanatory variables on species diversity individually, while taking into account variation in other potentially important variables.

We chose $p < 0.05$ as the determinate of statistical significance. All numerical analyses were performed in R (R development core team 2010) using several statistical packages (RODBC, vegan, maps, mapdata, mgsv, MASS, MCMCglmm and nlme). Data analyses were done in three steps:

Step 1. Quantify trends and significant changes in species diversity over the sampling period covered by the biological data.

First, we calculated the species richness (N) of Ephemeroptera, Plecoptera and Trichoptera (our biodiversity metric) for all samples and all available seasons at all sites. Then, linear trends in diversity for each were assessed by use of linear mixed effect models (lme) with random intercept and slope, and tested for statistical significance. The species diversity trends (negative or positive) for each site were also plotted on a map.

Step 2. Quantify trends and significant changes in environmental data

Because environmental variables did not always occur at the time of biological sampling, we calculated the mean environmental condition over the quarter in which biological sampling took place (Q1: January – March, Q2: April – June, Q3: July – September, Q4: October – December). Trends in environmental variables were analysed by using the mean quarterly environmental value across the time span covered by the biological data at the site and with lme with random intercept and slope to test for significant changes.

Step 3. Test for correlations between environmental variables and species diversity

In order to investigate the relationships between species diversity and environmental variables, we analysed the correlation between species diversity and ANC, sulphate, temperature and precipitation. The two rivers from Latvia were excluded from this analysis because they are considered non-sensitive towards acidification and have a very high ANC compared to the other sites. In the analyses, we used a MCMCglmm, with: $N_g \sim ANC_g + SO4_g + Precipitation_g + Temperature_g$, where the subscript g denotes the pH-group.

3 Results

3.1 Environmental variables

The analyses were performed on all sites (Group 1) and on sites split into one of three pH classes (Groups 2-4) (Figure 5 and Figure 6). Water pH increased significantly during the sampling period at most sites (Table 3), while pH fluctuated more in rivers than in lakes. Most sites showed significant decreases in the concentration of SO₄ during the last 30 years, while ANC increased at most sites (Figure 7 and Figure 8). Mean quarterly air temperatures did not increase over the period covered by the biological data in any of the grid cells in which sites were located (Figure 7 and Figure 8). Eight of the 38 lake sites and 8 of the 54 river sites were situated in cells where quarterly precipitation increased.




Figure 5. All river sites are divided into four groups based on their pH-range. Group 1 are all sites, Group 2 are sites where pH < 5.5, Group 3 are sites where pH has crossed 5.5 and Group 4 are sites where pH > 5.5. The columns on the left hand side indicate sites with a significant (p<0.05) increase (+), decrease (-) or no change (0) in pH and species diversity (N).




Figure 6. All lakes are divided into four groups based on their pH-range. Group 1 are all sites, Group 2 are sites where pH < 5.5, Group 3 are sites where pH has crossed 5.5 and Group 4 are sites where pH > 5.5. The columns on the left hand side indicate sites with a significant (p<0.05) increase (+), decrease (-) or no change (0) in pH and species diversity (N), and the correlation between pH and N.

3.2 Species diversity

Most sites show a positive trend in the number of EPT taxa (N) over time (Figure 9). The species diversity increased at all river sites, except three, and at all lake sites, except six. The increase was statistically significant in 30% of the river sites and 21% of the lakes. There was a non-significant decrease in diversity in one river and six lakes. A larger fraction of sites within the group deemed to be sensitive to acidification (Group 3) showed a significant increase in diversity compared to the other groups (Table 3, Figure 10). Diversity also increased significantly at some of the sites deemed to be least influenced by acidification (Group 2), but only at sites where there has been a significant increase in pH (Figure 5 and Figure 6). It should be noted that while a relatively smaller proportion of Group 2 rivers and lakes showed significant trends, this still amounted to around two thirds of all sites in these groups.




Figure 7. Trends in quarterly air temperature, precipitation, SO₄ and ANC for European lakes over the site-specific period covered by the biological data. + denotes increase and – denotes decrease. Statistical significant trends are embedded in a grey circle.




Figure 8. Trends in quarterly air temperature, precipitation, SO₄ and ANC for European river sites over the site-specific period covered by the biological data. + denotes increase and – denotes decrease. Statistical significant trends are embedded in a grey circle.




Figure 9. Trends in biodiversity in European rivers and lakes over the last 15 to 30 years (see Tables 1 and 2 for exact periods). + Increasing trend, – decreasing trend. Statistical significant changes in diversity are embedded in grey circles.

Table 3. The proportion of sites with a statistically significant increase in pH and number of EPT species (N) over the last ca 30 years. Numbers in brackets show a significant decrease. Group 1 includes all sites, Group 2 includes sites where pH < 5.5, Group 3 includes sites where pH has crossed 5.5 during the sampling period and Group 4 includes sites with pH > 5.5.

| | Rivers | | | | Lakes | | | |
|------------|-------------|-------------|---------|---------|---------|---------|---------|---------|
| | Group 1 | Group 2 | Group 3 | Group 4 | Group 1 | Group 2 | Group 3 | Group 4 |
| Group size | 54 | 14 | 27 | 13 | 38 | 16 | 15 | 7 |
| pH | 0.74 (0.02) | 0.64 (0.07) | 0.74 | 0.85 | 0.84 | 0.69 | 0.93 | 1.0 |
| N | 0.30 | 0.21 | 0.37 | 0.23 | 0.21 | 0.13 | 0.33 | 0.14 |




Figure 10. The proportion of sites with a statistically significant increase in number of EPT species (N) over the last ca 30 years.

Table 4. Correlation analyses between environmental variables and diversity, as analysed by a Markov Chain Monte Carlo generalized mixed effects model. Model outputs for rivers and lakes throughout Germany, Sweden, Latvia, Czech Republic, United Kingdom, and Norway. Post mean is the mean of the posterior distribution. The upper and lower 95% CI are credible intervals analogous to standard confidence intervals. The effective sample size (Eff. sample) is a measure of the auto-correlation within the parameter sample. Group 1 includes all sites, Group 2 includes sites where pH < 5.5, Group 3 includes sites where pH has crossed 5.5 during the sampling period, and Group 4 includes sites with pH > 5.5. *denotes significance at $p < 0.05$. The correlation of single parameters is corrected for the influence of the other explanatory variables.

| Variable | Post Mean | Lower 95%CI | Upper 95%CI | Eff. sample | p-value |
|----------------------|-----------|-------------|-------------|-------------|---------|
| Rivers | | | | | |
| Group 1 (all sites) | | | | | |
| Temperature | -0.006 | -0.010 | 0.002 | 40.0 | 0.001* |
| Precipitation | 0 | 0 | 0.001 | 37.0 | 0.036* |
| ANC | 0 | 0 | 0 | 31.8 | 0.001* |
| SO4 | -0.013 | -0.015 | -0.009 | 12.3 | 0.001* |
| Group 2 (least acid) | | | | | |
| Temperature | -0.02 | -0.009 | -0.006 | 7.5 | 0.34 |
| Precipitation | 0.001 | 0 | 0.003 | 10.4 | 0.108 |
| ANC | 0 | 0 | 0.001 | 3.6 | 0.932 |
| SO4 | -0.044 | -0.054 | -0.035 | 3.2 | 0.001* |
| Group 3 (sensitive) | | | | | |
| Temperature | -0.002 | -0.008 | 0.005 | 152.3 | 0.606 |
| Precipitation | 0 | 0 | 0.001 | 247.0 | 0.036* |
| ANC | 0 | 0 | 0 | 117.4 | 0.076 |
| SO4 | -0.009 | -0.013 | -0.005 | 33.4 | 0.001* |
| Group 4 (acid) | | | | | |
| Temperature | -0.009 | -0.017 | -0.002 | 11.5 | 0.004* |
| Precipitation | 0 | 0 | 0.001 | 5.7 | 0.984 |
| ANC | 0 | 0 | 0.001 | 12.4 | 0.001* |
| SO4 | -0.018 | -0.026 | -0.014 | 12.1 | 0.001* |
| Lakes | | | | | |
| Group 1 (all sites) | | | | | |
| Temperature | -0.034 | -0.049 | -0.023 | 7.0 | 0.001* |
| Precipitation | 0.001 | 0 | 0.002 | 5.2 | 0.064 |
| ANC | 0.003 | 0.002 | 0.004 | 12.4 | 0.001* |
| SO4 | 0 | -0.002 | 0.016 | 10.9 | 0.95 |
| Group 2 (least acid) | | | | | |
| Temperature | -0.055 | -0.070 | -0.048 | 4.4 | 0.001* |
| Precipitation | 0 | 0 | 0 | 8.2 | 0.006* |
| ANC | -0.004 | -0.005 | -0.003 | 13.3 | 0.001* |
| SO4 | -0.063 | -0.070 | -0.034 | 3.5 | 0.001* |
| Group 3 (sensitive) | | | | | |
| Temperature | -0.040 | -0.045 | -0.018 | 4.6 | 0.001* |
| Precipitation | 0.002 | 0 | 0.003 | 2.3 | 0.001* |
| ANC | 0.001 | 0 | 0.002 | 5.7 | 0.052 |
| SO4 | 0.004 | -0.011 | 0.018 | 11.7 | 0.426 |
| Group 4 (acid) | | | | | |
| Temperature | -0.034 | -0.040 | -0.024 | 21.4 | 0.001* |
| Precipitation | 0 | 0 | 0.002 | 5.8 | 0.344 |
| ANC | 0.004 | 0.002 | 0.0055 | 15.8 | 0.001* |
| SO4 | -0.023 | -0.060 | 0.0338 | 1.67 | 0.354 |




Figure 11. The relationship between environmental conditions and diversity for Group1 sites (upper four are rivers and lower four are lakes), as analysed by MCMC glm. Temp: mean monthly temperature ($^{\circ}\text{C}$), Precip: mean monthly precipitation (mm), SO_4 : sulphate (mg/l) and ANC: acid neutralizing capacity ($\mu\text{Eq/l}$). The models include various parameters where points and lines do not need to agree in order to be significant.

3.3 Possible drivers of biodiversity

According to the MCMCglm-model, (ANC, SO₄; Table 4, Figure 11), the diversity of all rivers and lakes (Group 1) is significantly correlated to ANC and temperature, and SO₄ and precipitation is also significant for the rivers. When it comes to the subsets of rivers, SO₄ is important for all three sub-groups.

According to the relationship between SO₄ and species diversity, lower SO₄ leads to higher diversity where a 10 mg/l reduction in SO₄ is associated with an increase of about two species (Figure 12). ANC is most important for the most acidified rivers, has perhaps a marginal effect in the sensitive rivers, and appears unimportant for the least sensitive. In addition, temperature is significant for the most acid rivers and precipitation for the sensitive rivers. For lakes, ANC appears the more important chemical variable, with changes in SO₄ only important for the least acid sites. Clearly, temperature is an important variable in lakes with a significant negative effect in all groups. Precipitation is significant for the least acid sites and for the sensitive sites.

There is no clear indication that the species diversity of sites with the least impact from acidification is more influenced by climate. For example, the species diversity of the least acidified rivers is correlated to SO₄ (Figure 11) and not to temperature or precipitation.




Figure 12. The relationship between sulphate (SO₄, mg/l) and species diversity (N) for the least-acidified rivers (Group 2) (left) and the reductions in sulphate-concentration needed in order to alter the diversity (right). The right hand figure is the change in diversity across time (T2-T1, T3-T1, T3-T2 etc.) versus the corresponding change in SO₄. Both relationships are analysed by a MCMC glm.

4 Discussion

4.1 Recovery from acidification as drivers of diversity

There are very clear relationships between acid chemistry and diversity amongst rivers and lakes in ICP Waters sites. This is evident when all groups are analysed together and when they are analysed separately. The most pronounced increase in diversity is found at sites with the most pronounced chemical recovery (Table 3, Figure 10), yet all sites with an increase in diversity also show a pattern of concurrent increase in pH (Figure 5 and Figure 6). There is a clear overall significant negative correlation between diversity and sulphate, e.g., for all groups of rivers, and a significant positive correlation between ANC and diversity, e.g., for all lakes (Figure 11, Table 4), which is consistent because reduced SO_4 in acid-sensitive sites often will be associated with an increase in ANC. The correlation is evident even for sites anticipated to be least influenced by acidification. The clear relationships between acid chemistry and diversity suggest that a reduction in acidifying components of the water is having a strong influence on species diversity in remote surface waters across north-western Europe. A similar relationship between diversity and sulphate has also been documented for streams in North Eastern USA (Klemm et al., 2002).

The analysis therefore provides a clear demonstration of a regional scale impact of acid emissions reduction policy on aquatic biodiversity. The regional extent of increasing species is surprising, particularly given that not all were initially considered sufficiently acid-sensitive to have responded chemically to reductions in acid deposition. The communities may have been insensitive to mild acid deposition if the pre-disturbance community consisted of insensitive taxa, by assemblages limited by biotic interactions or other environmental variables than acidification, or if the acidification eradicated fish populations (Hesthagen, 1986), allowing invertebrates that are sensitive to fish predation to flourish (Appelberg et al., 1993; Hildrew, 2009; Layer et al., 2011). We lack information on the reference state of biodiversity for our sites and our understanding is therefore largely based on space-for-time approaches, whereby communities in acid-sensitive but un-acidified areas provide our best guide as to the likely composition of sites prior to acidification. From this approach, it is likely that acidification was accompanied by a gradual decrease in biodiversity.

Sulphate was chosen as an explanatory parameter because it has dominated the acid load in the region historically. Furthermore, in catchments with soils that have formed since the last glaciation, the majority of sites in this dataset, the SO_4 anion tends to behave relatively conservatively within catchments, so that changes in sulphate deposition have, for example, been shown to follow those in surface waters at an annual time-step in UK waters (Cooper, 2005). Changes in sulphate concentrations can therefore be assigned to changes in acid deposition in sites with no local sources of sulphate and in sites with soils that have little capacity to sorb SO_4 , which is the case in formerly glaciated and acid-sensitive sites. Sulphate *per se*, however, is generally non-toxic at the concentrations measured at our sites and has little direct effect on aquatic invertebrates. Rates of change in the deposition of SO_4 tend to correlate quite closely to rates of change in ANC (Monteith et al., 2014), while influencing specific acid characteristics of surface waters in different ways depending on the history of acidification, base cation buffering potential and the availability of organic anions provided by dissolved organic matter. As ANC increases, changes in inorganic aluminium will be most marked in highly acidified waters, while pH changes are most sensitive in waters around pH 5.5.

Both pH and inorganic aluminium have been shown to exert direct toxic effects on biota. pH can influence the survival of aquatic insects, e.g., through the increase in hydrogen ion that interferes with the uptake and regulation of sodium and other ions (Havas and Rosseland, 1995; Paradise and Dunson, 1997) and by increasing aluminium toxicity (e.g., Havas and Rosseland, 1995; Sparling and Lowe, 1996). Many organisms cannot cope with high concentrations of H^+ and inorganic aluminium (Raddum and Fjellheim, 1995; Raddum and Skjelkvale, 2001). When toxicity is too great for acid-sensitive taxa, local survival is only possible in refuges, perhaps downstream or in neighbouring catchments, which are less impacted by acidification. As conditions improve, the sensitive taxa reappear from the source populations. Such taxa form the basis for acidification indices and scores from these indices are now beginning to hint at

improved conditions at sites in The Czech Republic, Germany, Norway, Sweden, UK and elsewhere (Angeler and Johnson, 2012; Hesthagen et al., 2011; Monteith et al., 2005; Murphy et al., 2014).

A continuous biotic recovery of the least impacted sites would not be evident when monitoring the sites by indices previously applied to assess trends in macroinvertebrates across the ICP Waters network. Acidification indices generally depend on the fraction of acid sensitive taxa in the species assemblage, whereas species diversity is the sum of all species in the assemblage. Many acidification indices are insensitive to pH above pH >5.5 (Fjellheim and Raddum, 1990; Raddum et al., 1988). The median tolerable pH for most sensitive species of invertebrates is between 5.2 and 6.1 (Lacoul et al., 2011), however, it has been demonstrated that a few macroinvertebrate species are sensitive even at pH slightly below 7 (Bell, 1971; Moe et al., 2010). This suggests that species diversity should accompany the acidification indices in monitoring of acid sensitive sites. Site-specific time series of species diversity can then act as a reference for the most recent developments in diversity.

4.2 Temperature and precipitation as drivers of diversity

There was no significant change in air temperature within the grid cells representing the ICP Waters sites over the period covered by the biological data. This may be due to small absolute changes and the large year-to-year fluctuations. Nevertheless, we observed significant, and invariably negative, correlations between temperature and species diversity for both lake and river sites. As temperature trends were lacking these correlations must therefore reflect inter-annual sensitivity of invertebrate populations to variations in temperature around the time of sampling. Whether such negative effects on temperature would be sustained over a period of ramped warming remains to be seen, but these results point to a potential threat to these surface waters from forecast changes in temperature in the region. It should also be noted that we cannot exclude the possibility that the warmer years result in earlier emergence of adults and thus lower larval catches.

In contrast to air temperatures, the amount of precipitation for grid cells representing the ICP Waters sites increased significantly for about 10% and 20% for river and lake sites, respectively. We found a positive correlation between precipitation and species diversity in rivers for all groups except the most acid sites (Group 4). Increased precipitation may act on species diversity through at least one of four different processes: 1. by decreasing the frequency of drought events and desiccation, 2. by enhancing catchment erosion and thereby increases the level of nutrients in the ecosystem, 3. by increasing the frequency of extreme flooding events that flushed out the animals, and 4. by increasing the frequency and intensity of acid episodes. Both 1 and 2 can be expected to have a positive influence on species diversity, while 3 and 4 can be expected to have a negative effect on species diversity. The lack of correlation between precipitation and species diversity at the most acid sites may either suggest that changes in precipitation have not been sufficiently large to impact on these communities or that multiple hydraulic processes with conflicting effects (including the effect of acid episodes) may be operating.

Surface water organisms are potentially sensitive to variation in various climatological characteristics. Monthly or annual average temperature or precipitation may change, extreme values can become more intense or frequent, seasonality may be altered and the timing of specific weather events can shift (Garcia et al., 2014; Kusch, 2015). Extreme weather may influence the species diversity, e.g., through sea salt sea-salt episodes (Hindar et al., 2004). Many benthic invertebrates have an optimal water temperature and cannot survive if the temperatures are too warm or too cold (Velle et al., 2011). In cold regions, their development requires more accumulated degree-days than are available. Some species might benefit from a future climate in which the length of the growing season increases and the winters become less harsh (Iacarella et al., 2015). Increased temperatures have already resulted in range expansions, pole-ward shifts and altered species phenologies (Dingemanse and Kalkman, 2008; Hickling et al., 2006; Parmesan et al., 1999; Walther et al., 2002). In addition, the biota may be negatively influenced by river bed instability during enhanced precipitation (Lods-Crozet et al., 2001).

A continuous warming can be expected to influence species diversity negatively, e.g. as a result of a decrease in the amount of suitable areas subsequent to climate change (Domisch et al., 2013), it can have a neutral effect on species diversity (Burgmer et al., 2007), or the response can vary according to taxonomic group (Kusch, 2015; Li et al., 2014). On a millennium time scale, we might also expect warmer

temperatures to lead to more diverse communities, in accordance with patterns of species richness that often are correlated with latitudinal gradients (Hawkins et al., 2003; Wright et al., 1993). Rohde (1999) concluded that latitudinal gradients in species diversity result from effective evolutionary time modulated by several factors, such as temperature and energy input. Still, the mechanism responsible for the strong relationship between temperature and freshwater biota is not well understood (Eggermont and Heiri, 2012; Velle et al., 2010a). Complex and multidimensional ecological processes – other than simply temperature preferences – are determining the degree and direction of species distributions in aquatic systems (Velle et al., 2010a). Shifts in species response to temperature are likely to vary both spatially and temporally, depending on the nature of climate change and the setting of the aquatic environment in question (Dingemanse and Kalkman, 2008).

4.3 Stability- and length of the environmental gradients

We expected a stronger correlation between diversity and temperature for the least-acidified sites (Group 2). However, water chemistry has an overriding influence on species diversity for all groups of sites. With changes in water chemistry taken into account, temperature was found to exert a net negative effect on species diversity. It is not likely that temperature had a dominating influence on species diversity given that the diversity has increased for most sites. This can also be expected since the gradient of change along the temperature axis is short (not significant long-term change at any site) compared to the pronounced change along the ANC- and sulphate gradients (significant at most sites). The longer chemistry gradients in rivers may explain why the relationship between diversity and ANC- and sulphate is stronger for rivers than for lakes.

A differing response between rivers and lakes can also result from the dissimilar stability of the two environments. The lake thermal environment is more stable and to a lesser extent affected by shading or by short-term variations in hydrology, implying that the air temperature metric provides a more robust measure of variation in lake temperature than it does for stream sites. The same may potentially apply to the chemistry, in as much as point sampled ANC for a river is highly sensitive to short term variation in hydrology, while the ANC of a lake will be more representative of average recent chemistry. SO₄ takes up more of the variance for the rivers, because SO₄ concentration in rivers tends to be generally more stable and robust predictor of recent average chemistry than ANC. More homogenous conditions in lakes than in rivers may also suggest that the biota of lakes to a smaller is influenced by acidification. For example, lakes are less susceptible to episodic shocks events, such as acid peaks during snow-melt or sea-salt episodes. This can partly explain the observation of a more pronounced increase in species diversity of rivers than lakes. In addition, the biological resolution of rivers is higher than for lakes since the number of EPT-taxa naturally is higher in rivers. In lakes, these insects are less abundant and non-biting midges (Chironomidae) form the most abundant benthic macroinvertebrate (Armitage et al., 1995). Non-biting midges are rarely identified below family-level for monitoring purpose.

4.4 Implications for policy

As a response to global threats to biodiversity, the partners to the Convention on Biological Diversity committed themselves in 2002 to achieve, by 2010, a significant reduction in the current rate of loss of biodiversity at the global, regional and national level. This target was not met (Spyropoulou et al., 2010) and the loss of biodiversity has continued. There is now a Strategic Plan for Biodiversity 2011-2020 that aims to halt and eventually reverse the loss of biodiversity of the planet by 2020. To build support and momentum for this task, the United Nations General Assembly at its 65th session declared the period 2011-2020 to be “The United Nations Decade on Biodiversity”. This is a demanding task given that worldwide, the biodiversity of freshwaters and elsewhere is declining (Barnosky et al., 2011; Sala et al., 2000).

In the context of this grim news, our evidence of an apparent hemispheric-scale response of aquatic diversity to internationally agreed acid emission reductions provides an encouraging demonstration of the potential for international action to achieve positive results in this field. Based on the correlations we have established between water acidity and diversity, even for the least-acidified sites (Figure 12), we can expect a further increase in species diversity providing levels of deposition continue to decrease. In addition, we can expect a further increase in diversity since biological recovery often is lagging improvements in

chemistry (Murphy et al., 2014). A number of hypotheses have been proposed to explain this behaviour (Monteith et al., 2005), and of these the most likely surround biological inertia, in which acid sensitive taxa struggle to re-establish functional niches that have become occupied by acid tolerant generalists (Ledger and Hildrew, 2005) and/ or the impoverished buffer capacity of the catchments that makes the assemblages vulnerable to short term acid episodes. Conversely, the effect of climate warming on species diversity may counteract the effects of reduced acid deposition in the future. The trends in warming for the periods for which biological records in the ICP Waters database exist, were not significant and therefore, this analysis cannot address whether long-term climate change affects aquatic biodiversity in acid-sensitive ecosystems. Still, temperature had effects on variation in biodiversity, suggestion that these communities will be sensitive to long-term temperature change. Continued monitoring of biological communities will be extremely valuable for detection of climate effects on biodiversity.

4.5 Perspectives

For the basic unit of biological classification, the species, α -biological diversity is expressed as a function of the number of species and their frequency, or simply the number of species (Chapin Iii et al., 2000; Tuomisto, 2010), as we have done. Nevertheless, biodiversity metrics reduce complex information on structure and abundance of communities to simple numbers. There are two main limitations to the concept of biodiversity: (1) the term is artificial implying that biodiversity not is an intrinsic property in nature and (2) biodiversity is a simplification of nature and it is necessary to consider that information is lost when complex processes are reduced to a single number (Hurlbert, 1971).

A natural next step in the analysis of long-term trends in biological communities and their relationship to environmental drivers is to analyse changes in the function of the communities. Such analyses will complement the analysis of species diversity. Function reflects the biological complexity of the ecosystem expressed by important biological traits (Heino, 2005, 2008; Schleuter et al., 2010). Such traits could include feeding mode, food source, mode of mobility, size at maturity, life cycle length etc. Changes in these traits over space or time gives direct information about ecological processes, and also are highly relevant for finding effects on ecosystem-services.

5 References

- Angeler DG, Johnson RK (2012) Temporal scales and patterns of invertebrate biodiversity dynamics in boreal lakes recovering from acidification. *Ecological Applications* 22: 1172-1186.
- Appelberg M, Henrikson BI, Henrikson L, Svedang M (1993) Biotic Interactions within the Littoral Community of Swedish Forest Lakes during Acidification. *Ambio* 22: 290-297.
- Armitage PD, Cranston PS, Pinder LCV (1995) *The Chironomidae: Biology and ecology of non-biting midges*. London: Chapman & Hall.
- Barnosky AD, Matzke N, Tomiya S, Wogan GOU, Swartz B, Quental TB, Marshall C, McGuire JL, Lindsey EL, Maguire KC, Mersey B, Ferrer EA (2011) Has the Earth's sixth mass extinction already arrived? *Nature* 471: 51-57.
- Battarbee RW, Charles DF (1986) Diatom-Based Ph Reconstruction Studies of Acid Lakes in Europe and North-America - a Synthesis. *Water, Air, and Soil Pollution* 30: 347-354.
- Bell HL (1971) Effect of low pH on the survival and emergence of aquatic insects. *Water Research* 5: 313-319.
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White J-SS (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution* 24: 127-135.
- Brown JH, Gillooly JF, Allen AP, Savage VM, West GB (2004) Toward a metabolic theory of ecology. *Ecology* 85: 1771-1789.
- Buisson L, Grenouillet G, Villéger S, Canal J, Laffaille P (2012) Toward A Loss Of Functional Diversity In Stream Fish Assemblages Under Climate Change. *Global Change Biology*: n/a-n/a.
- Burgmer T, Hillebrand H, Pfenninger M (2007) Effects of climate-driven temperature changes on the diversity of freshwater macroinvertebrates. *Oecologia* 151: 93-103.
- Chapin Iii FS, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, Hooper DU, Lavorel S, Sala OE, Hobbie SE, Mack MC, Diaz S (2000) Consequences of changing biodiversity. *Nature* 405: 234-242.
- Condamine FL, Rolland J, Morlon H (2013) Macroevolutionary perspectives to environmental change. *Ecology Letters* 16: 72-85.
- Cooper DM (2005) Evidence of sulphur and nitrogen deposition signals at the United Kingdom Acid Waters Monitoring Network sites. *Environmental Pollution* 137: 41-54.
- Dingemanse NJ, Kalkman VJ (2008) Changing temperature regimes have advanced the phenology of Odonata in the Netherlands. *Ecological Entomology* 33: 394-402.
- Domisch S, Araújo MB, Bonada N, Pauls SU, Jähnig SC, Haase P (2013) Modelling distribution in European stream macroinvertebrates under future climates. *Global Change Biology* 19: 752-762.
- Dudgeon D, Arthington AH, Gessner MO, Kawabata ZI, Knowler DJ, Leveque C, Naiman RJ, Prieur-Richard AH, Soto D, Stiassny MLJ, Sullivan CA (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews* 81: 163-182.

Easterling DR, Horton B, Jones PD, Peterson TC, Karl TR, Parker DE, Salinger MJ, Razuvayev V, Plummer N, Jamason P, Folland CK (1997) Maximum and Minimum Temperature Trends for the Globe. *Science* 277: 364-367.

Eggermont H, Heiri O (2012) The chironomid-temperature relationship: expression in nature and palaeoenvironmental implications. *Biological Reviews of the Cambridge Philosophical Society* 87: 430-456.

Erwin DH (2001) Lessons from the past: Biotic recoveries from mass extinctions. *Proceedings of the National Academy of Sciences* 98: 5399-5403.

Evans CD, Cullen JM, Alewell C, Kopacek J, Marchetto A, Moldan F, Prechtel A, Rogora M, Vesely J, Wright R (2001) Recovery from acidification in European surface waters. *Hydrology and Earth System Sciences* 5: 283-297.

Fjellheim A, Raddum GG (1990) Acid Precipitation - Biological Monitoring of Streams and Lakes. *Science of the Total Environment* 96: 57-66.

Frost S, Huni A, Kershaw WE (1971) Evaluation of a Kicking Technique for Sampling Stream Bottom Fauna. *Canadian Journal of Zoology* 49: 167-173.

Fölster J, Andrén C, Bishop K, Buffam I, Cory N, Goedkoop W, Holmgren K, Johnson R, Laudon H, Wilander A (2007) A Novel Environmental Quality Criterion for Acidification in Swedish Lakes—An Application of Studies on the Relationship Between Biota and Water Chemistry. In Brimblecombe, P, Hara, H, Houle, D, Novak, M, editors, *Acid Rain - Deposition to Recovery*, Dordrecht: Springer Netherlands, 331-338.

Garcia RA, Cabeza M, Rahbek C, Araújo MB (2014) Multiple Dimensions of Climate Change and Their Implications for Biodiversity. *Science* 344

Garmo OA, Skjelkvale BL, de Wit HA, Colombo L, Curtis C, Folster J, Hoffmann A, Hruska J, Hogasen T, Jeffries DS, Keller WB, Kram P, Majer V, Monteith DT, Paterson AM, Rogora M, Rzychon D, Steingruber S, Stoddard JL, Vuorenmaa J, Worsztynowicz A (2014) Trends in Surface Water Chemistry in Acidified Areas in Europe and North America from 1990 to 2008. *Water Air and Soil Pollution* 225: 14.

Hadfield JD (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. *Journal of Statistical Software* 33: 1-22.

Halvorsen GA, Heegaard E, Raddum G (2002) Tracing recovery from acidification - a multivariate approach. *ICP Waters report: NIVA*, 39.

Halvorsen GA, Heegaard E, Fjellheim A, Raddum GG (2003) Tracing recovery from acidification in the western Norwegian Nausta watershed. *Ambio* 32: 235-239.

Harris I, Jones PD, Osborn TJ, Lister DH (2014) Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology* 34: 623-642.

Havas M, Rosseland BO (1995) Response of zooplankton, benthos, and fish to acidification: An overview. *Water Air and Soil Pollution* 85: 51-62.

Hawkins BA, Field R, Cornell HV, Currie DJ, Guégan J-F, Kaufman DM, Kerr JT, Mittelbach GG, Oberdorff T, O'Brien EM, Porter EE, Turner JRG (2003) Energy, Water, and Broad-Scale Geographic Patterns of Species Richness. *Ecology* 84: 3105-3117.

Heino J (2005) Functional biodiversity of macroinvertebrate assemblages along major ecological gradients of boreal headwater streams. *Freshwater Biology* 50: 1578-1587.

- Heino J (2008) Patterns of functional biodiversity and function-environment relationships in lake littoral macroinvertebrates. *Limnology and Oceanography* 53: 1446-1455.
- Heller NE, Zavaleta ES (2009) Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* 142: 14-32.
- Hesthagen T (1986) Fish kills of Atlantic salmon (*Salmo salar*) and brown trout (*Salmo trutta*) in an acidified river of SW Norway. *Water, Air, and Soil Pollution* 30: 619-628.
- Hesthagen T, Fjellheim A, Schartau AK, Wright RF, Saksgard R, Rosseland BO (2011) Chemical and biological recovery of Lake Saudlandsvatn, a formerly highly acidified lake in southernmost Norway, in response to decreased acid deposition. *Sci Total Environ* 409: 2908-2916.
- Hickling R, Roy DB, Hill JK, Fox R, Thomas CD (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biology* 12: 450-455.
- Hildrew AG (2009) Sustained Research on Stream Communities: A Model System and The Comparative Approach. *Advances in Ecological Research, Vol 41* 41: 175-312.
- Hindar A, Torseth K, Henriksen A, Orsolini Y (2004) The significance of the North Atlantic Oscillation (NAO) for sea-salt episodes and acidification-related effects in Norwegian rivers. *Environ Sci Technol* 38: 26-33.
- Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA (2005) Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs* 75: 3-35.
- Horecký J, Stuchlík E, Chvojka P, Bitušík P, Liška M, Pšenáková P, Špaček J (2002) Effects of acid atmospheric deposition on chemistry and benthic macroinvertebrates of forest streams in the Brdy Mts (Czech Republic). *Acta Soc. Zool. Bohem.* 66: 189-203.
- Horecký J, Stuchlík E, Chvojka P, Hardekopf DW, Mihaljevic M, Špaček J (2006) Macroinvertebrate community and chemistry of the most atmospherically acidified streams in the Czech Republic. *Water Air and Soil Pollution* 173: 261-272.
- Horecký J, Rucki J, Kram P, Krecek J, Bitušík P, Špaček J, Stuchlík E (2013) Differences in benthic macroinvertebrate structure of headwater streams with extreme hydrochemistry. *Biologia* 68: 303-313.
- Hurlbert SH (1971) The Nonconcept of Species Diversity: A Critique and Alternative Parameters. *Ecology* 52: 577-586.
- Iacarella JC, Dick JTA, Alexander ME, Ricciardi A (2015) Ecological impacts of invasive alien species along temperature gradients: testing the role of environmental matching. *Ecological Applications* 25: 706-716.
- IPCC (2013) Summary for Policymakers. In Stocker, TF, Qin, D, Plattner, G-K, Tignor, M, Allen, SK, Boschung, J, Nauels, A, Xia, Y, Bex, V, Midgley, PM, editors, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- Isaak DJ, Hubert WA (2004) Nonlinear response of trout abundance to summer stream temperatures across a thermally diverse montane landscape. *Transactions of the American Fisheries Society* 133: 1254-1259.
- Johnson RK, Goedkoop W (2007) Bedömningsgrunder för bottenfauna i sjöar och vattendrag – Användarmanual och bakgrundsdokument. Uppsala: Swedish University of Agricultural Sciences, 84 pp.

Johnson RK, Angeler DG (2010) Tracing recovery under changing climate: response of phytoplankton and invertebrate assemblages to decreased acidification. *Journal of the North American Benthological Society* 29: 1472-1490.

Kernan M, Battarbee RW, Curtis CJ, Monteith DT, Shilland EM, editors(2010) *Recovery of lakes and streams in the UK from the effects of acid rain. UK Acid Waters Monitoring Network 20 years Interpretative Report*. London: Environmental Change Research Centre.

Kirchner JW, Weil A (2000) Delayed biological recovery from extinctions throughout the fossil record. *Nature* 404: 177-180.

Klemm DJ, Blocksom KA, Thoeny WT, Fulk FA, Herlihy AT, Kaufmann PR, Cormier SM (2002) Methods Development and use of Macroinvertebrates as Indicators of Ecological Conditions for Streams in the Mid-Atlantic Highlands Region. *Environmental Monitoring and Assessment* 78: 169-212.

Knell RJ, Thackeray SJ (2016) Voltinism and resilience to climate-induced phenological mismatch. *Climatic Change* 137: 525-539.

Kusch J (2015) Interacting influences of climate factors and land cover types on the distribution of caddisflies (Trichoptera) in streams of a central European low mountain range. *Insect Conservation and Diversity* 8: 92-101.

Lacoul P, Freedman B, Clair T (2011) Effects of acidification on aquatic biota in Atlantic Canada. *Environmental Reviews* 19: 429-460.

Layden A, Merchant C, MacCallum S (2015) Global climatology of surface water temperatures of large lakes by remote sensing. *International Journal of Climatology* 35: 4464-4479.

Layer K, Hildrew AG, Jenkins GB, Riede JO, Rossiter SJ, Townsend CR, Woodward G (2011) Long-Term Dynamics of a Well-Characterised Food Web: Four Decades of Acidification and Recovery in the Broadstone Stream Model System. *Advances in Ecological Research, Vol 44* 44: 69-117.

Ledger ME, Hildrew AG (2005) The ecology of acidification and recovery: changes in herbivore-algal food web linkages across a stream pH gradient. *Environmental Pollution* 137: 103-118.

Lento J, Dillon P, Somers K (2012) Evaluating long-term trends in littoral benthic macroinvertebrate communities of lakes recovering from acid deposition. *Environmental Monitoring and Assessment* 184: 7175-7187.

Li F, Kwon YS, Bae MJ, Chung N, Kwon TS, Park YS (2014) Potential Impacts of Global Warming on the Diversity and Distribution of Stream Insects in South Korea. *Conservation Biology* 28: 498-508.

Livingstone DM, Lotter AF (1998) The relationship between air and water temperatures in lakes of the Swiss Plateau: a case study with palaeolimnological implications. *Journal of Paleolimnology* 19: 181-198.

Lods-Crozet B, Lencioni V, Olafsson JS, Snook DL, Velle G, Brittain JE, Castella E, Rossaro B (2001) Chironomid (Diptera : Chironomidae) communities in six European glacier-fed streams. *Freshwater Biology* 46: 1791-1809.

Mantyka-pringle CS, Martin TG, Rhodes JR (2012) Interactions between climate and habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biology* 18: 1239-1252.

Markovic D, Carrizo S, Freyhof J, Cid N, Lengyel S, Scholz M, Kasperdius H, Darwall W (2014) Europe's freshwater biodiversity under climate change: distribution shifts and conservation needs. *Diversity and Distributions* 20: 1097-1107.

- Mayhew PJ, Bell MA, Benton TG, McGowan AJ (2012) Biodiversity tracks temperature over time. *Proceedings of the National Academy of Sciences of the United States of America* 109: 15141-15145.
- Moe SJ, Schartau AK, Baekken T, McFarland B (2010) Assessing macroinvertebrate metrics for classifying acidified rivers across northern Europe. *Freshwater Biology* 55: 1382-1404.
- Monteith DT, Hildrew AG, Flower RJ, Raven PJ, Beaumont WRB, Collen P, Kreiser AM, Shilland EM, Winterbottom JH (2005) Biological responses to the chemical recovery of acidified fresh waters in the UK. *Environmental Pollution* 137: 83-101.
- Monteith DT, Evans CD, Henrys PA, Simpson GL, Malcolm IA (2014) Trends in the hydrochemistry of acid-sensitive surface waters in the UK 1988-2008. *Ecological Indicators* 37: 287-303.
- Murphy JF, Winterbottom JH, Orton S, Simpson GL, Shilland EM, Hildrew AG (2014) Evidence of recovery from acidification in the macroinvertebrate assemblages of UK fresh waters: A 20-year time series. *Ecological Indicators* 37, Part B: 330-340.
- Paradise CJ, Dunson WA (1997) Effects of pH and sulfate on insects and protozoans inhabiting treeholes. *Archives of Environmental Contamination and Toxicology* 33: 182-187.
- Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, Huntley B, Kaila L, Kullberg J, Tammaru T, Tennent WJ, Thomas JA, Warren M (1999) Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* 399: 579-583.
- Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421: 37-42.
- Purvis A, Hector A (2000) Getting the measure of biodiversity. *Nature* 405: 212-219.
- Quist MC, Hubert WA, Isaak DJ (2004) Fish assemblage structure and relations with environmental conditions in a Rocky Mountain watershed. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 82: 1554-1565.
- Raddum G, Fjellheim A, Hesthagen T (1988) Monitoring of acidification through the use of aquatic organisms. *Verhandlungen der internationalen Vereinigung theoretische und angewandte Limnologie* 23: 2291-2297.
- Raddum GG, Hagenlund G, Halvorsen GA (1984) Effects of Lime Treatment on the Benthos of Lake Søndre Boksjø. *Institute of Freshwater Research, Drottningholm Report No. 61*: 167-176.
- Raddum GG, Fjellheim A (1995) Acidification in Norway - Status and trends biological monitoring - Improvements in the invertebrate fauna. *Water Air and Soil Pollution* 85: 647-652.
- Raddum GG, Skjelkvale BL (2001) Critical limit of acidifying compounds to invertebrates in different regions of Europe. *Water Air and Soil Pollution* 130: 825-830.
- Raddum GG, Fjellheim A, Velle G (2008) Increased growth and distribution of *Ephemerella aurivillii* (Ephemeroptera) after hydropower regulation of the Aurland catchment in Western Norway. *River Research and Applications* 24: 688-697.
- Reid WV, Mooney HA, Cropper A, Capistrano D, Carpenter SR, Chopra K, Cropper A, Dasgupta P, Hassan R, Leemans R, May RM, Pingali P, Samper C (2005) *Ecosystems and Human Well-being: Biodiversity Synthesis*. Millennium Ecosystem Assessment - World Resources Institute, Washington, DC.
- Rockstrom J, Steffen W, Noone K, Persson A, Chapin FS, Lambin EF, Lenton TM, Scheffer M, Folke C, Schellnhuber HJ, Nykvist B, de Wit CA, Hughes T, van der Leeuw S, Rodhe H, Sorlin S, Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J, Walker B, Liverman D, Richardson K, Crutzen P, Foley JA (2009) A safe operating space for humanity. *Nature* 461: 472-475.

- Rohde K (1999) Latitudinal gradients in species diversity and Rapoport's rule revisited: a review of recent work and what can parasites teach us about the causes of the gradients? *Ecography* 22: 593-613.
- Sahney S, Benton MJ, Falcon-Lang HJ (2010) Rainforest collapse triggered Carboniferous tetrapod diversification in Euramerica. *Geology* 38: 1079-1082.
- Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DM, Mooney HA, Oesterheld M, Poff NL, Sykes MT, Walker BH, Walker M, Wall DH (2000) Biodiversity - Global biodiversity scenarios for the year 2100. *Science* 287: 1770-1774.
- Schaumburg J, Maetze A, Lehmann R, Kifinger B, Coring E, Baltzer S (2008) Monitoringprogramm für versauerte Gewässer durch Luftschadstoffe in der Bundesrepublik Deutschland im Rahmen der ECE. Bericht der Jahre 2007-2008. Berlin: Bayerisches Landesamt für Umwelt, 134.
- Schindler DW (1988) Effects of acid-rain on fresh-water ecosystems. *Science* 239: 149-157.
- Schleuter D, Daufresne M, Massol F, Argillier C (2010) A user's guide to functional diversity indices. *Ecological Monographs* 80: 469-484.
- Schofield CL (1976) Acid Precipitation: Effects on Fish. *Ambio* 5: 228-230.
- Skjelkvaale BL, de Wit HA (2011) Trends in precipitation chemistry, surface water chemistry and aquatic biota in acidified areas in Europe and North America from 1990 to 2008. ICP Waters Report 106/2011. Oslo.
- Sparling DW, Lowe TP (1996) Environmental hazards of aluminum to plants, invertebrates, fish, and wildlife. *Rev Environ Contam Toxicol* 145: 1-127.
- Spyropoulou R, Martins IP, McInnes G, Uhel R, Uludag A, Herkendell J, Calvo EC, Romao C, Oulton AR, Weber J-L, Biala K, Lund MP, Makarewicz K, Hogeveen Y, Csagoly P, Petersen J-E, Schutyser F, Nieto A, Zisensis M, Richard D, McSharry B, Evans D (2010) *The European environment. State and outlook 2010. Biodiversity*. Luxembourg: Publications Office of the European Union.
- Stendera S, Johnson RK (2008) Tracking recovery trends of boreal lakes: use of multiple indicators and habitats. *Journal of the North American Benthological Society* 27: 529-540.
- Stendera S, Adrian R, Bonada N, Canedo-Arguelles M, Hugueny B, Januschke K, Pletterbauer F, Hering D (2012) Drivers and stressors of freshwater biodiversity patterns across different ecosystems and scales: a review. *Hydrobiologia* 696: 1-28.
- Stoddard JL, Jeffries DS, Lukewille A, Clair TA, Dillon PJ, Driscoll CT, Forsius M, Johannessen M, Kahl JS, Kellogg JH, Kemp A, Mannio J, Monteith DT, Murdoch PS, Patrick S, Rebsdorf A, Skjelkvaale BL, Stainton MP, Traaen T, van Dam H, Webster KE, Wieting J, Wilander A (1999) Regional trends in aquatic recovery from acidification in North America and Europe. *Nature* 401: 575-578.
- Sweeney BW (1984) Factors influencing life-history patterns of aquatic insects. In Resh, VH, Rosenberg, DM, editors, *The Ecology of Aquatic insects*, New York: Praeger, 56-100.
- Team RDC (2010) R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.
- Thackeray SJ, Henrys PA, Hemming D, Bell JR, Botham MS, Burthe S, Helaouet P, Johns DG, Jones ID, Leech DI, Mackay EB, Massimino D, Atkinson S, Bacon PJ, Brereton TM, Carvalho L, Clutton-Brock TH, Duck C, Edwards M, Elliott JM, Hall SJG, Harrington R, Pearce-Higgins JW, Hoyer TT, Kruuk LEB, Pemberton JM, Sparks TH, Thompson PM, White I, Winfield IJ, Wanless S (2016) Phenological sensitivity to climate across taxa and trophic levels. *Nature* 535: 241-U294.

- Toffolon M, Piccolroaz S, Majone B, Soja AM, Peeters F, Schmid M, Wuest A (2014) Prediction of surface temperature in lakes with different morphology using air temperature. *Limnology and Oceanography* 59: 2185-2202.
- Tuomisto H (2010) A diversity of beta diversities: straightening up a concept gone awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity. *Ecography* 33: 2-22.
- UN (2005) *The Millennium Ecosystem Assessment (MA), Ecosystems and Human Well-Being: Synthesis*. Island Press, Washington DC.
- Vautard R, Gobiet A, Sobolowski S, Kjellström E, Stegehuis A, Watkiss P, Mendlik T, Landgren O, Nikulin G, Teichmann C, Jacob D (2014) The European climate under a 2 °C global warming. *Environmental Research Letters* 9: 034006.
- Velle G, Brodersen KP, Birks HJB, Willassen E (2010a) Midges as quantitative temperature indicator species: lessons for palaeoecology. *The Holocene* 20: 989-1002.
- Velle G, Bjune AE, Larsen J, Birks HJB (2010b) Holocene climate and environmental history of Brurskardstjøerni, a lake in the catchment of Øvre Heimdalsvatn, south-central Norway. *Hydrobiologia* 642: 13-34.
- Velle G, Kongshavn K, Birks HJB (2011) Minimizing the edge-effect in environmental reconstructions by trimming the calibration set: Chironomid-inferred temperatures from Spitsbergen. *The Holocene* 21: 417-430.
- Velle G, Telford R, Skjelkvaale BL, C.Curtis, L.Eriksson, A.Fjellheim, J.Fölster, G.A.Halvorsen, Hildrew A, Hoffmann A, Indriksone I, Kamasová L, Kopáček J, Orton S, Krám P, Monteith DT, Senoo T, Shilland EM, Stuchlík E, Wiklund M-L, Wit Hd, Skjelkvaale B-L (2013) 30 years of biodiversity trends in European fresh waters. Bergen: ICP Waters, 66 p.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, Bairlein F (2002) Ecological responses to recent climate change. *Nature* 416: 389-395.
- Wathne BM, Ranneklev SB, Bjørn Olav Rosseland, Braaten HFV, Fjellheim A (2010) ICP Waters Programme Manual 2010. ICP Waters Report 105/2010.
- Williams M, Zalasiewicz J, Haff P, Schwägerl C, Barnosky AD, Ellis EC (2015) The Anthropocene biosphere. *The Anthropocene Review* 2: 196-219.
- Wright DH, Currie DJ, Maurer BA (1993) Species diversity in ecological communities: Historical and geographical perspectives In Ricklefs, RE, Schluter, D, editors, *Energy supply and patterns of species richness on local and regional scales*, Chicago University of Chicago Press, 66-76.
- Zhu M-Y, Babcock LE, Peng S-C (2006) Advances in Cambrian stratigraphy and paleontology: Integrating correlation techniques, paleobiology, taphonomy and paleoenvironmental reconstruction. *Palaeoworld* 15: 217-222.

6 Reports and publications from the ICP Waters programme

All reports from the ICP Waters programme from 2000 up to present are listed below. Reports before year 2000 can be listed on request. All reports are available from the Programme Centre. Reports and recent publications are also accessible through the ICP Waters website; <http://www.icp-waters.no/>

- De Wit, H., Valinia, S. and Steingruber, S. Proceedings of the 31st Task Force meeting of the ICP Waters Programme in Monte Verità, Switzerland 6th –8th October, 2015. **ICP Waters report 126/2015**
- De Wit, H., Hettelingh, J.P. and Harmens, H. 2015. Trends in ecosystem and health responses to long-range transported atmospheric pollutants. **ICP Waters report 125/2015**
- Fjellheim, A., Johannessen, A. and Landås, T.S. 2015. Biological intercalibration: Invertebrates 1915. **ICP Waters report 124/2015**
- Escudero-Oñate, C. 2015 Intercomparison 1529: pH, Conductivity, Alkalinity, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. **ICP Waters report 123/2015**
- de Wit, H., Wathne, B. M. (eds) 2015. Proceedings of the 30th Task Force meeting of the ICP Waters Programme in Grimstad, Norway 14th –16th October, 2014. **ICP Waters report 122/2015**
- Fjellheim, A., Johannessen, A. and Landås, T.S. 2014. Biological intercalibration: Invertebrates 1814. **ICP Waters Report 121/2014**
- Escudero-Oñate. 2014. Intercomparison 1428: pH, Conductivity, Alkalinity, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. **ICP Waters Report 120/2014**
- De Wit, H. A., Garmo Ø. A. and Fjellheim A. Chemical and biological recovery in acid-sensitive waters: trends and prognosis. **ICP Waters Report 119/2014**
- Fjellheim, A., Johannessen, A. and Landås, T.S. 2013. Biological intercalibration: Invertebrates 1713. **ICP Waters Report 118/2014**
- de Wit, H., Bente M. Wathne, B. M. and Hruška, J. (eds) 2014. Proceedings of the 29th Task Force meeting of the ICP Waters Programme in Český Krumlov, Czech Republic 1st –3rd October, 2013. **ICP Waters report 117/2014**
- Escudero-Oñate, C. Intercomparison 1327: pH, Conductivity, Alkalinity, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni and Zn. **ICP Waters Report 116/2013**
- Holen, S., R.F. Wright, I. Seifert. 2013. - Effects of long-range transported air pollution (LTRAP) on freshwater ecosystem services. **ICP Waters Report 115/2013**
- Velle, G., Telford, R.J., Curtis, C., Eriksson, L., Fjellheim, A., Frolova, M., Fölster J., Grudule N., Halvorsen G.A., Hildrew A., Hoffmann A., Indriksone I., Kamasová L., Kopáček J., Orton S., Krám P., Monteith D.T., Senoo T., Shilland E.M., Stuchlík E., Wiklund M.L., de Wit, H., Skjelkvaale B.L. 2013. Biodiversity in freshwaters. Temporal trends and response to water chemistry. **ICP Waters Report 114/2013**
- Fjellheim, A., Johannessen, A. and Landås, T.S. 2013. Biological intercalibration: Invertebrates 1612. **ICP Waters Report 113/2013**
- Skjelkvåle, B.L., Wathne, B.M., de Wit, H. and Michela Rogora (eds.) 2013. Proceedings of the 28th Task Force meeting of the ICP Waters Programme in Verbania Pallanza, Italy, October 8 – 10, 2012. **ICP Waters Report 112/2013**
- Dahl, I. 2012. Intercomparison 1226: pH, Conductivity, Alkalinity, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni and Zn. **ICP Waters report 111/2012**
- Skjelkvåle, B.L., Wathne B. M. and Moiseenko, T. (eds.) 2010. Proceedings of the 27th meeting of the ICP Waters Programme Task Force in Sochi, Russia, October 19 – 21, 2011. **ICP Waters report 110/2012**
- Fjellheim, A., Johannessen, A., Svanevik Landås, T. 2011. Biological intercalibration: Invertebrates 1511. NIVA-report SNO 6264-2011. **ICP Waters report 109/2011**

- Wright, R.F., Helliwell, R., Hruska, J., Larssen, T., Rogora, M., Rzychoń, D., Skjelkvåle, B.L. and Worsztynowicz, A. 2011. Impacts of Air Pollution on Freshwater Acidification under Future Emission Reduction Scenarios; ICP Waters contribution to WGE report. NIVA-report SNO 6243-2011. **ICP Waters report 108/2011**
- Dahl, I and Hagebø, E. 2011. Intercomparison 1125: pH, Cond, HCO₃, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. NIVA-report SNO 6222-2011. **ICP Waters report 107/2011**
- Skjelkvåle B.L. and de Wit, H. (Eds). 2011. Trends in precipitation chemistry, surface water chemistry and aquatic biota in acidified areas in Europe and North America from 1990 to 2008. NIVA-report SNO 6218-2011. **ICP Waters report 106/2011**
- ICP Waters Programme Centre 2010. ICP Waters Programme manual. NIVA SNO 6074-2010. **ICP Waters report 105/2010**. 91 s. ISBN 978-82-577-5953-7,
- Skjelkvåle, B.L., Wathne B. M. and Vuorenmaa J. (eds.) 2010. Proceedings of the 26th meeting of the ICP Waters Programme Task Force in Helsinki, Finland, October 4 – 6, 2010. **ICP Waters report 104/2010**
- Fjellheim, A. 2010. Biological intercalibration: Invertebrates 1410. NIVA-report SNO 6087-2010, **ICP Waters report 103/2010**
- Hovind, H. 2010. Intercomparison 1024: pH, Cond, HCO₃, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. NIVA-report SNO 6029-2010. **ICP Waters report 102/2010**
- De Wit, H. A. and Lindholm M., 2010. Nutrient enrichment effects of atmospheric N deposition on biology in oligotrophic surface waters – a review. NIVA-report SNO 6007 - 2010. **ICP Waters report 101/2010**
- Skjelkvåle, B.L., De Wit, H and and Jeffries, D. (eds.) 2010. Proceedings of presentations of national activities to the 25th meeting of the ICP Waters Programme Task Force in Burlington, Canada, October 19-21 2009. NIVA-report SNO 5995 - 2010. **ICP Waters report 100/2010**
- Fjellheim, A. 2009. Biological intercalibration: Invertebrates 1309. NIVA-report SNO 5883-2009, **ICP Waters report 99/2009**
- Hovind, H. 2009. Intercomparison 0923: pH, Cond, HCO₃, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. NIVA-report SNO 5845-2009. **ICP Waters report 98/2009**
- Ranneklev, S.B., De Wit, H., Jenssen, M. T. S. and Skjelkvåle, B.L., 2009. An assessment of Hg in the freshwater aquatic environment related to long-range transported air pollution in Europe and North America. NIVA-report SNO 5844-2009. **ICP Waters report 97/2009**.
- Skjelkvåle, B.L., Jenssen, M. T. S. and De Wit, H (eds.) 2009. Proceedings of the 24th meeting of the ICP Waters Programme Task Force in Budapest, Hungary, October 6 – 8, 2008. NIVA-report SNO 5770-2009. **ICP Waters report 96/2008**
- Fjellheim, A and Raddum, G.G. 2008. Biological intercalibration: Invertebrates 1208. NIVA-report SNO 5706-2008. **ICP Waters report 95/2008**
- Skjelkvåle, B.L., and De Wit, H. (eds.) 2008. ICP Waters 20 year with monitoring effects of long-range transboundary air pollution on surface waters in Europe and North-America. NIVA-report SNO 5684-2008. **ICP Waters report 94/2008**
- Hovind, H. 2008. Intercomparison 0822: pH, Cond, HCO₃, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. NIVA-report SNO 5660-2008. **ICP Waters report 93/2008**
- De Wit, H. Jenssen, M. T. S. and Skjelkvåle, B.L. (eds.) 2008. Proceedings of the 23rd meeting of the ICP Waters Programme Task Force in Nancy, France, October 8 – 10 , 2007. NIVA-report SNO 5567-2008. **ICP Waters report 92/2008**
- Fjellheim, A and Raddum, G.G. 2008. Biological intercalibration: Invertebrates 1107. NIVA-report SNO 5551 – 2008. **ICP Waters report 91/2008**
- Hovind, H. 2007. Intercomparison 0721: pH, Cond, HCO₃, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. NIVA-report SNO 5486-2007. **ICP Waters report 90/2007**

- Wright, R.F., Posch, M., Cosby, B. J., Forsius, M., and Skjelkvåle, B. L. 2007. Review of the Gothenburg Protocol: Chemical and biological responses in surface waters and soils. NIVA-report SNO 5475-2007. **ICP Waters report 89/2007**
- Skjelkvåle, B.L., Forsius, M., Wright, R.F., de Wit, H., Raddum, G.G., and Sjøeng, A.S.M. 2006. Joint Workshop on Confounding Factors in Recovery from Acid Deposition in Surface Waters, 9-10 October 2006, Bergen, Norway; Summary and Abstracts. NIVA-report SNO 5310-2006. **ICP Waters report 88/2006**
- De Wit, H. and Skjelkvåle, B.L. (eds). 2007. Trends in surface water chemistry and biota; The importance of confounding factors. NIVA-report SNO 5385-2007. **ICP Waters report 87/2007**
- Hovind, H. 2006. Intercomparison 0620. pH, K25, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-report SNO 5285-2006. **ICP Waters report 86/2006**
- Raddum, G.G. and Fjellheim, A. 2006. Biological intercalibration 1006: Invertebrate fauna. NIVA-report SNO 5314-2006. **ICP Waters report 85/2006**
- De Wit, H. and Skjelkvåle, B.L. (eds.) 2006. Proceedings of the 21th meeting of the ICP Waters Programme Task Force in Tallinn, Estonia, October 17-19, 2005. NIVA-report SNO 5204-2006. **ICP Waters report 84/2006**
- Wright, R.F., Cosby, B.J., Høgåsen, T., Larssen, T., Posch, M. 2005. Critical Loads, Target Load Functions and Dynamic Modelling for Surface Waters and ICP Waters Sites. NIVA-report SNO 5166-2005. **ICP Waters report 83/2006**
- Hovind, H. 2005. Intercomparison 0317. pH, K25, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-report SNO 5068-2005. **ICP Waters report 82/2005**
- Raddum, G.G. 2005. Intercalibration 0307: Invertebrate fauna. NIVA-report SNO 5067-2005. **ICP Waters report 81/2005**
- De Wit, H. and Skjelkvåle, B.L. (eds.). 2005. Proceedings of the 20th meeting of the ICP Waters Programme Task Force in Falun, Sweden, October 18-20, 2004. NIVA-report SNO 5018-2005. **ICP Waters report 80/2005**
- Fjeld, E., Le Gall, A.-C. and Skjelkvåle, B.L. 2005. An assessment of POPs related to long-range air pollution in the aquatic environment. NIVA-report SNO 5107-2005. **ICP Waters report 79/2005**
- Skjelkvåle et al 2005. Regional scale evidence for improvements in surface water chemistry 1990-2001. *Environmental Pollution*, 137: 165-176.
- Hovind, H. 2004. Intercomparison 0418. pH, K25, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-report SNO 4875-2004. **ICP Waters report 78/2004**
- Raddum, G.G. 2004. Intercalibration: Invertebrate fauna 09/04. NIVA-report SNO 4863-2004. **ICP Waters report 77/2004**
- Skjelkvåle, B.L. (ed). Proceedings of the 19th meeting of the ICP Waters Programme Task Force in Lugano, Switzerland, October 18-20, 2003. NIVA-report SNO 4858-2004. **ICP Waters report 76/2004**
- Raddum, G.G., *et al.* 2004. Recovery from acidification of invertebrate fauna in ICP Water sites in Europe and North America. NIVA-report SNO 4864-2004. **ICP Waters report 75/2004**
- Hovind, 2003. Intercomparison 0317. pH, K25, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-report SNO 4715-2003. **ICP Waters report 74/2003**
- Skjelkvåle, B.L. (ed). 2003. The 15-year report: Assessment and monitoring of surface waters in Europe and North America; acidification and recovery, dynamic modelling and heavy metals. NIVA-report SNO 4716-2003. **ICP Waters report 73/2003**
- Raddum, G.G. 2003. Intercalibration 0307: Invertebrate fauna. NIVA-report SNO-4659-2003. **ICP Waters report 72/2003**
- Skjelkvåle, B.L. (ed.). 2003. Proceedings of the 18th meeting of the ICP Waters Programme Task Force in Moscow, October 7-9, 2002. NIVA-report SNO 4658-2003. **ICP Waters report 71/2002**

- Wright, R.F and Lie, M.C. 2002. Workshop on models for Biological Recovery from Acidification in a Changing Climate. 9.-11. September 2002 in Grimstad, Norway. Workshop report. NIVA-report 4589-2002.
- Jenkins, A. Larsen, Th., Moldan, F., Posch, M. and Wrigth R.F. 2002. Dynamic Modelling of Surface Waters: Impact of emission reduction - possibilities and limitations. NIVA-report SNO 4598-2002. **ICP Waters report 70/2002.**
- Halvorsen, G.A, Heergaard, E. and Raddum, G.G. 2002. Tracing recovery from acidification - a multivariate approach. NIVA-report SNO 4564-2002. **ICP Waters report 69/2002**
- Hovind, H. 2002. Intercomparison 0216. pH, K₂₅, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-Report SNO 4558-2002. **ICP Waters Report 68/2002**
- Skjelkvåle, B.L. and Ulstein, M. (eds). 2002. Proceedings from the Workshop on Heavy Metals (Pb, Cd and Hg) in Surface Waters; Monitoring and Biological Impact. March 18-20, 2002, Lillehammer, Norway. NIVA-report SNO-4563-2002. **ICP Waters report 67/2002**
- Raddum, G.G. 2002. Intercalibration 0206: Invertebrate fauna. NIVA-report SNO-4494-2002. **ICP Waters report 66/2002**
- Bull, K.R. Achermann, B., Bashkin, V., Chrast, R. Fenech, G., Forsius, M., Gregor H.-D., Guardans, R., Haussmann, T., Hayes, F., Hettelingh, J.-P., Johannessen, T., Kryzanowski, M., Kucera, V., Kvaeven, B., Lorenz, M., Lundin, L., Mills, G., Posch, M., Skjelkvåle, B.L. and Ulstein, M.J. 2001. Coordinated Effects Monitoring and Modelling for Developing and Supporting International Air Pollution Control Agreements. *Water Air Soil Poll.* **130**:119-130.
- Hovind, H. 2001. pH, K₂₅, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-Report SNO 4416-2002. **ICP Waters report 64/2001**
- Lyulko, I. Berg, P. and Skjelkvåle, B.L. (eds.) 2001. National presentations from the 16th meeting of the ICP Waters Programme task Force in Riga, Latvia, October 18-20, 2000. NIVA-report SNO 4411-2001. **ICP Waters report 63/2001**
- Raddum, G.G. 2000. Intercalibration 0005: Invertebrate fauna. NIVA-report SNO4384-2001. **ICP Waters report 62/2001**
- Raddum, G.G. and Skjelkvåle B.L. 2000. Critical Load of Acidifying Compounds to Invertebrates In Different Ecoregions of Europe. *Water Air Soil Poll.* **130**:825-830.
- Stoddard, J. Traaen, T and Skjelkvåle, B.L. 2001. Assessment of Nitrogen leaching at ICP-Waters sites (Europe and North America). *Water Air Soil Poll.* **130**:825-830.
- Skjelkvåle, B.L. Stoddard J.L. and Andersen, T. 2001. Trends in surface waters acidification in Europe and North America (1989-1998). *Water Air Soil Poll.***130**:781-786.
- Kvaeven, B. Ulstein, M.J., Skjelkvåle, B.L., Raddum, G.G. and Hovind, H. 2001. ICP Waters - An international programme for surface water monitoring. *Water Air Soil Poll.***130**:775-780.
- Wright, R.F. 2001. Note on: Effect of year-to-year variations in climate on trends in acidification. NIVA-report SNO 4328-2001. **ICP Waters report 57/2001**
- Hovind, H. 2000. Trends in intercomparisons 8701-9812: pH, K₂₅, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K and aluminium - reactive and nonlabile, TOC, COD-Mn. NIVA-Report SNO 4281-2000. **ICP Waters Report 56/2000**
- Hovind, H. 2000. Intercomparison 0014. pH, K₂₅, HCO₃, NO₃ + NO₂, Cl, SO₄, Ca, Mg, Na, K, total aluminium, aluminium - reactive and nonlabile, TOC, COD-Mn. Fe, Mn, Cd, Pb, Cu, Ni and Zn. NIVA-Report SNO 4281-2000. **ICP Waters Report 55/2000**
- Skjelkvåle, B.L., Olendrzynski, K., Stoddard, J., Traaen, T.S, Tarrason, L., Tørseth, K., Windjusveen, S. and Wright, R.F. 2001. Assessment of trends and leaching in Nitrogen at ICP Waters Sites (Europe And North America). NIVA-report SNO 4383-2001. **ICP Waters report 54/2001**
- Stoddard, J. L., Jeffries, D. S., Lükewille, A., Clair, T. A., Dillon, P. J., Driscoll, C. T., Forsius, M., Johannessen, M., Kahl, J. S., Kellogg, J. H., Kemp, A., Mannio, J., Monteith, D., Murdoch, P. S., Patrick, S., Rebsdorf, A., Skjelkvåle, B. L., Stainton, M. P., Traaen, T. S., van Dam, H., Webster, K.

E., Wieting, J., and Wilander, A. 1999. Regional trends in aquatic recovery from acidification in North America and Europe 1980-95. *Nature* 401:575- 578.

Skjelkvåle, B. L., Andersen, T., Halvorsen, G. A., Raddum, G.G., Heegaard, E., Stoddard, J. L., and Wright, R. F. 2000. The 12-year report; Acidification of Surface Water in Europe and North America; Trends, biological recovery and heavy metals. NIVA-Report SNO 4208/2000.
ICP Waters report 52/2000

Reports before year 2000 can be listed on request.

NIVA: Norway's leading centre of competence in aquatic environments

NIVA provides government, business and the public with a basis for preferred water management through its contracted research, reports and development work. A characteristic of NIVA is its broad scope of professional disciplines and extensive contact network in Norway and abroad. Our solid professionalism, interdisciplinary working methods and holistic approach are key elements that make us an excellent advisor for government and society.



Norwegian Institute for Water Research

Gaustadalléen 21 • NO-0349 Oslo, Norway
Telephone: +47 22 18 51 00 • Fax: 22 18 52 00
www.niva.no • post@niva.no