

Accepted Manuscript

---

This is an Accepted Manuscript of the following article:

Jiří Novák, Branislav Vrana, Tatsiana Rusina, Krzysztof Okonski, Roman Grabic, Peta A. Neale, Beate I. Escher, Miroslava Macová, Selim Ait-Aissa, Nicolas Creusot, Ian Allan, Klára Hilscherová. Effect-based monitoring of the Danube River using mobile passive sampling. *Science of the Total Environment*. Volume 636, 2018, pages 1608-1619, ISSN 0048-9697.

The article has been published in final form by Elsevier at  
<http://dx.doi.org/10.1016/j.scitotenv.2018.02.201>

© 2019. This manuscript version is made available under the

CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

---

# Effect-based monitoring of the Danube River using mobile passive sampling

Jiří Novák<sup>a</sup>, Branislav Vrana<sup>a</sup>, Tatsiana Rusina<sup>a</sup>, Krzysztof Okonski<sup>a</sup>, Roman Grabic<sup>b</sup>, Peta A. Neale<sup>c,d</sup>, Beate I. Escher<sup>d,e,f</sup>, Miroslava Macová<sup>e</sup>, Selim Ait-Aissa<sup>g</sup>, Nicolas Creusot<sup>g</sup>, Ian Allan<sup>h</sup>, Klára Hilscherová<sup>a\*</sup>

<sup>a</sup>Masaryk University, Research Centre for Toxic Compounds in the Environment (RECETOX), Kamenice 753/5, 625 00 Brno, Czech Republic

<sup>b</sup>University of South Bohemia in České Budějovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Zátíší 728/II, CZ-389 25 Vodňany, Czech Republic

<sup>c</sup>Australian Rivers Institute, Griffith School of Environment, Griffith University, Southport QLD 4222, Australia

<sup>d</sup>The University of Queensland, Queensland Alliance for Environmental Health Sciences (QAEHS), Coopers Plains, Qld 4108, Australia

<sup>e</sup>UFZ Helmholtz Centre for Environmental Research – UFZ, Permoserstr. 15, 04318 Leipzig, Germany

<sup>f</sup>Eberhard Karls University Tübingen, Hölderlinstr. 12, 72074 Tübingen, Germany

<sup>g</sup>Institut National de l'Environnement Industriel et des Risques (INERIS), Unité Ecotoxicologie, Parc ALATA - BP2, 60550 Verneuil en Halatte, France

<sup>h</sup>Norwegian Institute for Water Research, Gaustadalle´en 21, NO-0349 Oslo, Norway

## \*Corresponding author:

Klára Hilscherová  
Masaryk University  
Faculty of Science  
RECETOX UKB A29  
Kamenice 753/5,  
625 00 Brno  
Czech Republic

tel: +420 549 493 256

e-mail: [hilscherova@recetox.muni.cz](mailto:hilscherova@recetox.muni.cz)

**Keywords:** passive sampling; river; estrogenicity; androgenicity; dioxin-like activity

## Abstract

Many aquatic pollutants can be present at low concentrations, but their mixtures can still affect health or behavior of exposed organisms. In this study, toxicological and chemical analyses were combined for spatial contamination profiling using an innovative passive sampling approach. A novel Dynamic

40 Passive Sampler (DPS) was employed as a mobile sampler from a ship cruising along 2130 km of the  
41 Danube river during the Joint Danube Survey 3 (JDS3). The sampling was performed in eight  
42 subsequent river stretches with two types of complementary passive samplers: silicone rubber sheets  
43 (SR) used for non-polar chemicals and SDB-RPS Empore™ disks (ED) for more hydrophilic compounds.  
44 Besides extensive chemical analyses, the bioactivity of samples was characterized by a battery of  
45 reporter gene bioassays. Cross-calibration of the employed passive samplers enabled robust  
46 estimation of water concentrations applicable for compounds with a wide range of physicochemical  
47 properties. DPS was suitable for sampling of water contaminants even at  $\text{pg L}^{-1}$  levels, with 209 of 267  
48 analyzed compounds detected in the samples. Biological effects were detected in both ED and SR  
49 extracts across all river stretches by bioassays focused on xenobiotic metabolism mediated by the aryl  
50 hydrocarbon and pregnane X receptors, endocrine disruptive potential mediated by estrogen and  
51 androgen receptors and the oxidative stress response. The bioassay responses expressed as  
52 bioanalytical equivalent concentrations ( $\text{BEQ}_{\text{bio}}$ ) were comparable with data obtained from large  
53 volume active sampling. The extracts of the ED samplers were more biologically active than extracts of  
54 SR samplers. Except of estrogenicity, where the analyzed chemicals explained on average 62 % of the  
55 effects in ED samples, the detected chemicals explained less than 8 % of  $\text{BEQ}_{\text{bio}}$  values. The study shows  
56 the utility of the combination of the innovative passive sampling approach with effect-based tools for  
57 efficient and fast monitoring even in water bodies with relatively low levels of contamination.

58

## 59 **1. Introduction**

60 Contamination of river water with complex mixtures of organic micropollutants poses a challenge for  
61 current pollution monitoring. Monitoring programs are typically driven by a need to protect aquatic  
62 organisms and, in an indirect way, also human populations from potentially toxic effects of  
63 environmental pollutants. This is complicated by the fact that pollutants are mostly present in very  
64 complex mixtures and their biological effects are the outcome of the integrated effects of many  
65 individual chemicals (Escher and Leusch, 2012). Because these mixtures can consist of thousands of  
66 chemicals whose toxic properties are often not known, targeted chemical analysis of individual  
67 compounds can sufficiently describe neither their composition nor toxic properties (Neale et al.,  
68 2015a). It has been previously shown that this problem could be addressed using a bioanalytical  
69 approach such as *in vitro* bioassays (Escher and Leusch, 2012; Giesy et al., 2002). These bioassays  
70 present rapid, sensitive and relatively inexpensive detection tools, which provide complementary  
71 information to data from instrumental chemical analysis. They enable estimation of the overall

72 biological activities of compounds present in environmental mixtures covering potential interactions  
73 among chemicals.

74 In order to cover a wide range of possible adverse effects, the bioassay battery should include  
75 endpoints reflecting a diverse set of possible toxic mechanisms, such as initiation of xenobiotic  
76 metabolism, endocrine disruptive potential and adaptive stress responses (Escher et al., 2014). To  
77 describe the overall bioactive potential of mixtures of chemicals, the concept of bioanalytical  
78 equivalent concentration (BEQ) was developed (Villeneuve et al., 2000; Baston and Denison, 2011). In  
79 this concept, the bioassay-detected biological potential of a complex mixture of chemicals is expressed  
80 as a concentration of a reference compound that would elicit the same effect as the mixture (BEQ<sub>bio</sub>).  
81 To estimate the impact of detected chemicals, a comparable equivalent (BEQ<sub>chem</sub>) can be modeled  
82 based on measured concentrations of individual chemicals using the concentration addition concept.  
83 Thus, by comparison of BEQ<sub>chem</sub> with the equivalents detected in bioassays (BEQ<sub>bio</sub>), it is possible to  
84 identify the chemicals most significantly contributing to the effect detected by bioassays (König et al.,  
85 2016; Neale et al., 2015a, 2017a).

86 Another challenge in aquatic pollutant monitoring is related to the requirement of a representative  
87 sampling in temporally and spatially variable water streams. Since concentrations of pollutants tend  
88 to fluctuate, frequent spot sampling or continuous sampling is required to provide a representative  
89 sample. This type of sampling is often laborious and resource-intensive. Many of the potentially toxic  
90 chemicals are present at very low concentrations and the effective pre-concentration step is thus  
91 needed prior to their instrumental analysis.

92 These problems can be effectively addressed using integrative passive sampling. In this approach,  
93 samplers spontaneously absorb or adsorb chemicals from water. The integrative concentration of  
94 chemicals in a passive sampler throughout the sampling period decreases their limit of detection. It  
95 also allows estimation of their time-weighted average concentrations, which include residues from  
96 episodic pollution events often not detectable even using frequent spot sampling. As the uptake of  
97 pollutants by the passive samplers is affected by environmental variables such as temperature and  
98 flow velocity, accurate in situ assessment of water volume sampled during exposure is required. For  
99 individual compounds, estimation of sampled water volume can be performed through the application  
100 of performance reference compounds (PRCs) and models that relate the sampling rate to properties  
101 that control the compound mass transfer (Booij et al., 2007).

102 Passive sampling as well as effect-based approaches are being considered as potentially suitable tools  
103 that could be employed for monitoring of European water bodies in the implementation strategy of  
104 the EU Water Framework Directive (European Commission, 2015). While both approaches are often

105 employed independently, the utility of their combination has been demonstrated previously in studies  
106 focusing on small water streams and wastewater treatment plant effluents and affected rivers (Creusot  
107 et al., 2013; Jalova et al., 2013; Jarosova et al., 2012). However, these approaches have not been  
108 validated for large rivers with moderate or low levels of contamination.

109 This study investigates the applicability of the newly developed dynamic passive sampling (DPS), which  
110 speeds up the uptake of compounds, for pollution and toxicity profiling of large rivers. The DPS was  
111 employed in the Joint Danube Survey 3 (JDS3) as a mobile sampler from a ship cruising downstream  
112 along 2130 km of the Danube river, one of the largest rivers in Europe. The relatively low contamination  
113 level makes the Danube suitable for testing of sampling techniques for monitoring of less concentrated  
114 pollutants. The sampling was performed in eight subsequent river stretches with two types of  
115 complementary passive samplers: partitioning silicone rubber sheets (SR) focused on non-polar  
116 chemicals, which allow quantification of sampling rate, and adsorption SDB-RPS Empore™ disks (ED)  
117 for non-polar as well as more hydrophilic compounds. Besides extensive chemical analyses, sample  
118 extracts were characterized by a battery of *in vitro* bioassays covering a range of endpoint types  
119 including endocrine disruption, xenobiotic metabolism, and adaptive stress responses. BEQ modeling  
120 was used to estimate the portion of biological effects of the samples that can be explained by detected  
121 chemicals. Since the same bioassays were used also for the assessment of samples from large volume  
122 active sampling during the JDS3 ship cruise (Neale et al., 2015a), it allowed for a mutual comparison of  
123 passive and active sampling approaches for toxicity profiling.

124

## 125 **2. Material and methods**

### 126 **2.1. Passive sampling**

127 Sample collection was performed during the JDS3 (Liska et al., 2015) on a sampling cruise in August  
128 and September 2013 using a dynamic passive sampling (DPS) system described by Vrana et al.  
129 (submitted). Briefly, the DPS device consisted of a stainless-steel chamber equipped with a submersible  
130 pump (approx.  $9 \text{ m}^3 \text{ h}^{-1}$ ) that provided a forced flow of sampled water through the sampling chamber  
131 with a current velocity of  $1\text{--}2 \text{ m s}^{-1}$ . For mobile sampling, the device was immersed in a flow-through  
132 tank located on the frontal deck of the JDS3 expedition ship (Vrana et al., submitted). Each individual  
133 sampling period lasted approximately five days, which resulted in a total of eight samples collected  
134 from each sampler type representing eight stretches of the Danube River (Tab. S1). Two parallel DPS  
135 devices were in operation during each sampler deployment. The samplers exposed in one of the  
136 devices were dedicated to chemical analyses and the samplers from the other device were used for

137 bioanalyses. Besides sampling from the ship, stationary DPS was also performed at a site located at  
138 the Danube river kilometer 1852, approximately 15 km downstream of Bratislava (Fig. 1) and the  
139 sampling dates coincided with time periods when the JDS3 expedition ship moved by the stationary  
140 site.

141 Two types of passive samplers were utilized for bioanalysis: silicone rubber (SR) AlteSil™ sheets, and  
142 SDB-RPS Empore™ disks (ED) based on styrene-divinylbenzene sorbent modified with sulfonic acid  
143 groups.

144 AlteSil™ translucent SR sheets 0.5 mm thick were purchased from Altec, UK. The sampler consisted of  
145 a single AlteSil™ SR sheet with dimensions 14×28 cm. SR samplers in the DPS device dedicated for  
146 chemical analyses were spiked prior to exposure with a set of PRCs. The ED sampler consisted of ten  
147 solid phase Empore™ SDB-RPS extraction disks with 47 mm diameter (Sigma Aldrich, Czech Republic).  
148 ED samplers were not spiked with any PRCs.

149 Besides SR and ED samplers, the DPS devices were equipped with a reference passive sampler that  
150 served to compare the sampling performance of the two DPS devices operating in parallel. It consisted  
151 of a strip of low-density polyethylene (LDPE; 4×28 cm and 70 µm thickness; Brentwood Plastics Inc,  
152 USA). The LDPE samplers mounted in both DPS devices were spiked with another set of PRCs. The LDPE  
153 samplers were located sideways from the SR and ED samplers to minimize cross-contamination of the  
154 samplers by PRCs, for further detailed information on sampling, list of PRCs and sample processing see  
155 SI1 and Vrana et al. (submitted).

156



157

158 **Fig. 1 Map of the sampling cruise with the stretches and stationary exposure site (red dot)**

159

## 160 **2.2. Chemical analysis**

161 SR and ED were processed as described in the SI (1.1.1-1.1.2). SR extracts were analyzed for 81  
162 hydrophobic compounds from the following groups: polycyclic aromatic hydrocarbons (PAHs; 29  
163 individual chemicals), PCBs (7), organochlorine pesticides and their degradation products (OCPs; 12),  
164 brominated diphenyl ethers (PBDEs; 9) and novel brominated flame retardants (15), alkylphenols (3),  
165 alkyl- and aryl- phosphates (14), and synthetic musks (6). ED extracts were analyzed for 204 chemicals  
166 from the following groups: currently used pesticides (CUPs; 40), steroids (5), phytosterols (11),  
167 flavonoids (8), alkylphenols (3), pharmaceuticals (108) and PAHs (29). For a comprehensive list of  
168 analyzed compounds see SI (Tab.S2, S3).

169 PAHs were analyzed by gas chromatography/mass spectrometry (GC/MS) using GC 7890/MS5975  
170 (Agilent, USA) equipped with J&W Scientific fused silica column DB-5MS (0.25 mm I.D., film thickness:  
171 0.25  $\mu\text{m}$ ; Agilent). For PCBs and OCPs, the analysis was performed using GC 7890/MS-MS Triple  
172 Quadrupole 7000B (Agilent), equipped with HT8 SGE Analytical Science column. PBDEs were analyzed  
173 by GC 7890A (Agilent) equipped with 15m  $\times$  0.25mm  $\times$  0.10  $\mu\text{m}$  RTX-1614 column (Restek, USA), HRMS  
174 (AutoSpec Premier) was operated in EI+ mode at the resolution of  $>10,000$ . Alkylphenols were  
175 derivatized with dansyl chloride, separated using an Agilent 1200 Infinity Series liquid chromatograph  
176 equipped with an ACE 5 C18 column (ACE, UK) and detected by MS/MS Agilent 6410 Triple Quadrupole  
177 (Pernica et al., 2015). Novel brominated flame retardants were analyzed using GC 7890A (Agilent)  
178 equipped with 15 m  $\times$  0.25 mm  $\times$  0.10  $\mu\text{m}$  DB5 column coupled to an AutoSpec Premier MS (Waters,  
179 Micromass, UK; Lohmann et al., 2013). Analyses of alkyl phosphates and polycyclic musks were  
180 performed using GC 6890 (Agilent) coupled to MSD 5975 mass spectrometer (Agilent). For more details  
181 see SI 1.2.

182 Currently used pesticides (CUPs) were separated using Agilent 1290 series HPLC (Agilent) and detected  
183 with mass spectrometer AB Sciex Qtrap 5500 (AB Sciex, Canada; Brumovský et al., 2016).

184 Steroid analysis was performed by liquid chromatography (HPLC Agilent 1200 Series) with mass  
185 spectrometry (MS-MS Agilent 6410 Triple Quad) after precolumn derivatization with dansyl chloride  
186 as described previously (Lin et al., 2007; Sadílek et al., 2016).

187 For the detection of flavonoids and phytosterols, the HPLC-MS/MS method previously described by  
188 Bláhová et al. (2016) was employed.

189 Pharmaceuticals were analyzed using a triple stage quadrupole MS/MS TSQ Quantum Ultra mass  
190 spectrometer (Thermo Fisher Scientific, San Jose, CA, USA) coupled with an Accela 1250 LC pump  
191 (Thermo Fisher Scientific) and an HTS XT-CTC autosampler (CTC Analytics AG, Switzerland) as described  
192 previously by Grabic et al., (2012).

193 LDPE extracts were analyzed for PAHs, PCBs, and HCB according to Allan et al. (2013). For further  
194 details on chemical analyses see SI 1.2.

195

## 196 **2.3. Bioanalysis**

197 Specific toxic potentials were assessed using a battery of bioassays (Table 1). A detailed description of  
198 the bioassays can be found in SI 1.3.1 and Neale et al. (2015a). The sample concentrations in bioassays  
199 were expressed as relative enrichment factor (REF), which expresses the water sample enrichment by  
200 passive sampling and sample processing together with the dilution in the bioassays. The effect in the  
201 sample was expressed as either the concentration causing 20% effect ( $EC_{20}$ ), the concentration causing  
202 20% inhibition ( $IC_{20}$ ), or the effect concentration causing an induction ratio (IR) of 1.5 ( $EC_{IR1.5}$ ) (Table 1).  
203 Log-logistic dose-response and linear regression models were used for the calculation of  $EC_{20}$  and  
204  $EC_{IR1.5}$ , respectively. Cell viability was assessed in parallel and cytotoxic sample concentrations were  
205 excluded from further calculations.

206

## 207 **2.4. Data analysis**

### 208 **2.4.1. Estimation of water volume extracted by passive samplers**

209 The calculated concentrations of chemicals in water ( $C_w$ ) derived from the passive sampling correspond  
210 to the freely dissolved concentration  $C_{free}$  (Mayer et al., 2003). Sampling rate  $R_{S,SR}$  of individual  
211 compounds to SR passive samplers was modeled as a function of the molar mass  $M$  by the water  
212 boundary layer-controlled uptake model from Rusina et al. (2010):

$$213 \quad R_{S,SR} = AB \times M^{-0.47} \quad \text{Equation 1}$$

214 with an exposure-specific parameter  $AB$ . The parameter was estimated from the dissipation of PRCs  
215 from samplers during exposure using a nonlinear least squares method by Booij and Smedes (2010),  
216 considering the fractions of individual PRCs that are retained in the sampler after exposure as a  
217 continuous function of their sampler-water partition coefficient  $K_{SR,w}$ . The models applied for SR  
218 samplers are described in detail in Vrana et al. (submitted).



219  $C_w$  of individual compounds were calculated from the mass absorbed by the SR samplers  $N_{SR}$ , the  
 220 degree of equilibrium  $DEQ$  that the compound attained during sampler exposure, the mass of sampler  
 221  $m_{SR}$  and their sampler-water partition coefficients  $K_{SR,w}$  as described in Booij et al. (2007).

$$222 \quad C_w = \frac{N_{SR}}{K_{SR,w} m_{SR} DEQ} \quad \text{Equation 2}$$

223 The  $DEQ$  was calculated as:

$$224 \quad DEQ = \left( 1 - \exp\left( -\frac{R_{S,SR}t}{K_{SR,w}m_{SR}} \right) \right) \quad \text{Equation 3}$$

225 Since the compounds that caused effects detected in passive sampler extracts were unknown they  
 226 were expressed as BEQ per sampler ( $BEQ_{SR,ED}$ ). A conversion to concentration in water ( $BEQ_w$ ) was  
 227 approximated using a sampling rate  $R_{S,SR}$  of a compound with an intermediate molar mass of 300 g.mol<sup>-1</sup>,  
 228 assuming a fully integrative sampling during the entire sampler exposure. In such situations, when  
 229  $DEQ \ll 1$ , Equation 3 can be simplified to  $DEQ = R_{S,SR}t / (K_{SR,w} \times m_{SR})$  and

$$230 \quad C_w = \frac{N_{SR}}{R_{S,SR}t} \quad \text{Equation 4}$$

231 The  $BEQ_w$  for SR was calculated as:

$$232 \quad BEQ_w = \frac{BEQ_{SR}}{R_{S,SR}t} = \frac{BEQ_{SR}}{AB \times 300^{-0.47} t} \quad \text{Equation 5}$$

233 For ED samplers, the PRC approach was not applied since its application in adsorption-based samplers  
 234 is questionable. The sampling rates of ED samplers were derived from a correlation of uptake of PAHs  
 235 and 4-nonylphenol to ED and SR samplers as is described in detail in SI 1.3 and Vrana et al. (submitted).

236 Sampling rate values of the ED samplers  $R_{S,ED}$  were estimated from sampling rates derived for SR  
 237 samplers ( $R_{S,SR}$ ), the calculated overall median  $F_{ED/SR}$  factor for 10 PAH individuals and 4-nonylphenol,  
 238 and the surface areas of both samplers  $A_{ED}$ ,  $A_{SR}$  as has been explained in detail by Vrana et al.  
 239 (submitted):

$$240 \quad R_{S,ED} = F_{ED/SR} \times \frac{A_{ED}}{A_{SR}} \times R_{S,SR} = 0.83 \times \frac{173}{392} \times R_{S,SR} = 0.366 \times R_{S,SR} \quad \text{Equation 6}$$

241 The  $BEQ_w$  for ED was then calculated as:

242 
$$BEQ_w = \frac{BEQ_{ED}}{R_{S,ED}t} = \frac{BEQ_{ED}}{0.366 \times R_{S,SR}t}$$
 Equation 7

243

244 **2.4.1. Bioanalytical Equivalent Concentration**

245 The EC and IC values from the different bioassays (Table 1) were converted to  $BEQ_{bio}$  using Equation 6,  
 246 with the  $EC_{20}$  or  $EC_{IR1.5}$  value of the reference compound (rc) and the matching  $EC_{20}$  or  $EC_{IR1.5}$  value of  
 247 the extract.

248 
$$BEQ_{bio} = \frac{EC_{20}(rc)}{EC_{20}(extract)} \text{ or } \frac{IC_{20}(rc)}{IC_{20}(extract)} \text{ or } \frac{EC_{IR1.5}(rc)}{EC_{IR1.5}(extract)}$$
 Equation 8

249

250 To calculate  $BEQ_{chem}$ , relative effect potency ( $REP_i$ ) of the detected chemicals was calculated from  
 251 measured data, complemented by information from the literature or calculated based on data from  
 252 the US EPA ToxCast database (US EPA, 2015; Tab. S5).  $REP_i$  was calculated using Equation 10, with  $EC_{10}$ ,  
 253  $IC_{20}$  or  $EC_{IR1.5}$  value of the reference compound and the matching  $EC_{10}$ ,  $IC_{20}$  or  $EC_{IR1.5}$  value of detected  
 254 chemical i. REPs derived from the literature data and ToxCast database were based on EC levels  
 255 indicated in Tab. S5. Limit of quantification (LOQ) was calculated using equation 6; the extract effective  
 256 concentration was replaced by the highest non-cytotoxic sample concentration tested.

257 
$$REP_i = \frac{EC_{10}(rc)}{EC_{10}(i)} \text{ or } \frac{IC_{20}(rc)}{IC_{20}(i)} \text{ or } \frac{EC_{IR1.5}(rc)}{EC_{IR1.5}(i)}$$
 Equation 9

258  $BEQ_{chem}$  was calculated from REP and the estimated concentration of each chemical in water in molar  
 259 units ( $C_{w,i}$ ) using Equation 10.

260 
$$BEQ_{chem} = \sum_{i=1}^n REP_i \cdot C_i$$
 Equation 10

261 **3. Results and Discussion**

262 **3.1. Passive sampling**

263 The Danube river watershed covers a significant part of the European continent reaching into the  
 264 territory of nineteen countries of different developmental stages. The river is affected by a range of  
 265 pollution sources discharging a wide spectrum of contaminants. On the other hand, being a large river,  
 266 the Danube has a great dilution capacity and so the pollutant concentrations in the water may be  
 267 relatively low compared to some rivers with a smaller dilution capacity (Keller et al., 2014). To  
 268 representatively characterize the river pollution, the current study employed mobile passive sampling  
 269 during the cruise of the expedition ship along defined stretches of the Danube. The overall sampled

270 river section covered by eight mobile-sampled river stretches spanned through nine countries (Fig. 1).  
271 Employment of two sampler types enabled sampling of chemicals with a wide range of physical-  
272 chemical properties. Partitioning-based silicon rubber (SR) samplers have been shown previously to  
273 effectively sorb non-polar chemicals (Rusina et al., 2010; Smedes and Booij, 2012). In order to facilitate  
274 the collection of more polar chemicals, we employed Empore™ SDB-RPS disc adsorption samplers (ED).  
275 The modification of poly(styrene-divinylbenzene) copolymer with sulfonic acid groups in ED was  
276 selected because it provides improved sorption capacity for polar chemicals with lower log  $K_{ow}$  such as  
277 pharmaceuticals and their metabolites, steroids, pesticides and their metabolites, explosives, or  
278 amine-containing analytes (Vrana et al., submitted).

279 Since the uptake principle is identical in both DPS and classical passive sampling, results may be  
280 evaluated using available passive sampler calibration parameters and models. The sampling rates for  
281 SR were determined using dissipation of PRCs from the samplers during their exposure, while for the  
282 ED samplers, the sampling rates were estimated from the relation of surface specific uptake of PAHs  
283 and 4-nonylphenol between SR and ED. The good correlation for these chemicals, which were  
284 integratively accumulated in SR, provided evidence that sampling rates under water-boundary layer  
285 control in ED are proportional to the sampling rates of SR. The proportionality factor of sampling rates  
286 is roughly given by the ratio of surface areas of the two samplers (details in SI 1.3; Fig. S1; Vrana et al.,  
287 submitted).

288 The increase of the sampling rate achieved by means of DPS was significant: in comparison with data  
289 from the passive samplers with the same dimensions subjected to stationary exposure without forced  
290 water exchange, DPS sampling was approximately five times faster (Vrana et al., submitted). This  
291 allowed us to collect and detect many chemicals with concentrations in the pg/L range during less than  
292 two days of sampler exposure (Tab. S2).


293 The LDPE samplers were used for checking whether the sampled water volume for the DPS device  
294 dedicated to toxicological analyses was equal to the device intended for chemical analysis. PRC release  
295 was monitored for LDPE samplers in both DPS devices per location since they had the same  
296 configuration and were exposed side by side. The comparison was relevant for the situations where  
297 the released PRC fraction  $f_{PRC}$  was quantifiable or lower than 80 % (Fig. S2; i.e. d<sub>10</sub>-FLT and d<sub>12</sub>-CHR).  
298 Their sampling rate ratio was  $1.00 \pm 0.11$  and  $0.93 \pm 0.14$ , respectively, as average across the eight  
299 sampling stretches. The corresponding sampling rates obtained demonstrate that the sampling  
300 performance of the two DPS devices deployed in parallel was equivalent. PRCs were not spiked to SR  
301 and ED samplers for toxicological analysis, but based on the equivalence of LDPE sampling rates in the

302 two devices, sampling rates in SR samplers for chemical analysis can be applied as a good estimate of  
303 sampling rates in SR samplers applied for toxicological analysis (details in SI 1.3.3, Fig. S2).

304

### 305 3.2. Chemical analysis

306 From a total of 267 analyzed chemicals, 209 were detected at least in one sample. A number of  
307 detected chemicals ranged from 52 to 70 and 103 to 131 for the SR and ED samplers across stretches,  
308 respectively (Fig. 2). Among non-polar compounds analyzed in the SR samples, triisobutylphosphate,  
309 4-nonylphenol, naphthalene and phenanthrene, and synthetic musk galaxolide were estimated to be  
310 present in the highest free dissolved concentrations in the river water (4-101 ng L<sup>-1</sup>; Tab. S2). Among  
311 compounds analyzed in the ED samples, the chemicals with the highest estimated concentration in  
312 river water were the pharmaceuticals carbamazepine, irbesartan, and sulphapyridine; atrazine,  
313 bisphenol A, cholesterol and sitosterol (7-17 ng L<sup>-1</sup>; Tab. S3).



314

315 Fig. 2 The sum of freely dissolved molar concentrations of chemicals in water ( $C_w$ ), estimated from  
316 their amounts analyzed in passive samplers (bar graph) and a number of chemicals detected  
317 in the samples (diamond); SR samples (black bar and blue diamond) and ED samples (white  
318 bar and empty diamond). S1-2 samples from stationary exposure of samplers downstream of  
319 Bratislava (Slovakia); 1-8 samples from a mobile sampling of river stretches.

320


### 321 3.3. Bioanalysis

322 In the present study, we employed bioassays for the assessment of xenobiotic metabolism initiation  
323 by the aryl hydrocarbon receptor (AhR) and pregnane X receptor (PXR)-mediated effects. AhR-  
324 activation has been previously described to induce carcinogenicity, immunotoxicity and indirectly also

325 endocrine disruption (Kortenkamp et al., 2012). PXR induces xenobiotic detoxification systems and is  
326 linked, for example, with liver steatosis and is sensitive to a wide range of chemicals (Creusot et al.,  
327 2010). The assessment of endocrine disruptive potential focused on the estrogen receptor (ER) and  
328 androgen receptor (AR)-mediated effects. Effects of xenobiotics on ER and AR signaling are relatively  
329 well characterized and they are reported to affect reproduction, development and play a clear role in  
330 carcinogenicity (Janošek et al., 2006; McLachlan, 2016). While activation and inhibition of ER by  
331 chemicals in water samples has been studied intensively, effects of aquatic samples mediated by AR  
332 have obtained less attention (Brack et al., 2007; Leusch et al., 2017). Exposure to xenobiotics often  
333 causes stress to organisms and it leads to activation of rather non-specific adaptive stress responses  
334 to restore homeostasis such as activation of systems dealing with oxidative stress, genotoxicity or  
335 inflammation. Transcription of many detoxification enzymes is coordinately regulated by antioxidant  
336 response elements (ARE) and their activation can serve as a marker of exposure-related oxidative  
337 stress (Reddy, 2008). The p53-mediated response is triggered by DNA damage and it activates repair  
338 mechanisms or apoptosis. It can thus serve as an indicator of genotoxic chemicals (Duerksen-Hughes  
339 et al., 1999). NF- $\kappa$ B-mediated response plays an important role in inflammatory reactions (Simmons et  
340 al., 2009).

341 Extracts of both SR and ED samplers elicited quantifiable effects in all employed bioassays except for  
342 the assay indicative of the NF- $\kappa$ B response (Table 3; Fig. 3). In the case of p53-mediated response, the  
343 effects were detected only in six ED samplers with higher sampled water volume and even then, they  
344 were very close to the LOQ.

345 All other biological endpoints were detected across all mobile and stationary samples and both  
346 sampler types. When comparing the BEQ<sub>bio</sub> of mobile sampled stretches across the assessed endpoints,  
347 spatial patterns along the river for some endpoints were revealed. For example, there was an increase  
348 in the AhR-mediated response in the ED sample at stretch 8 in the river delta in Romania (Fig. 3). There  
349 was also an increase in anti-androgenic potential from stretch 5 onward (Serbia, Romania, and  
350 Bulgaria). Similarly, an increased estrogenic potential was observed in stretch 5 (mainly Serbia and  
351 Romania), where also the highest levels of steroid estrogens, alkylphenols and musks were observed  
352 (Table 2, Tab. S4). This may be related to the fact that significant amounts of wastewaters are released  
353 to the Danube directly without treatment in this part of Europe, which leads to increased levels of  
354 chemicals with endocrine disruptive potential.




355

356 **Fig. 3 Spatial profiles of biological potentials in Danube river water, derived from passive sampler**  
 357 **data and expressed as BEQ<sub>bio</sub> of a respective reference compound; SR silicone rubber**  
 358 **samples, ED Empore™ disc samples; S1,2 – stationary sampling site; 1-8 sampled river**  
 359 **stretches (mean±SEM).**

360

361 EC values for the SR and ED extracts from both stationary samples and sampled stretches are compared  
 362 in Fig. 4. On average for each bioassay, the ED sample EC values were 3.5 to 12.3 times lower than the  
 363 SR samplers. This indicates that ED samplers were correspondingly more effective in the collection of  
 364 bioactive chemicals across all bioassays which detected significantly more biological potential. Thus,  
 365 the assessed endpoints were most likely elicited mainly by polar chemicals, for which the uptake  
 366 capacity in SR samplers is low. While this is not surprising for chemicals that act as inducers of oxidative  
 367 stress, endocrine disruptors or PXR-activators, the AhR-mediated effect has been often attributed to  
 368 more hydrophobic compounds, such as polychlorinated dibenzo-*p*-dioxins and furans or some PCB  
 369 congeners and PAHs. However, it has been shown that a considerable part of the AhR-mediated

370 activity is elicited by polar chemicals in water and sediment extracts (Jalova et al., 2013; Liu et al.,  
 371 2014). Anyway, the greatest difference between BEQ<sub>bio</sub> of SR and ED samplers was observed for  
 372 estrogenicity where EC values of ED samples were twelve times lower on average. Estrogenicity was  
 373 thus the endpoint where non-polar compounds contributed the least to the overall biological potential  
 374 of river water.



375  
 376 **Fig. 4 Effective concentration (EC) values of samples from dynamic passive sampling expressed**  
 377 **as relative enrichment factors of the samples (REF). EC<sub>20</sub> of PXR, ER, AhR-mediated effects, anti-**  
 378 **androgenic effect (aAR) and EC<sub>IR1.5</sub> of ARE and p53-mediated effects; S1-2 samples from**  
 379 **stationary exposure of the samplers downstream of Bratislava (Slovakia); 1-8 samples from a**  
 380 **mobile sampling of river stretches; SR silicone rubber sampler, ED Empore™ disc sampler.**

381  
 382 BEQ<sub>bio</sub> values detected at samples collected during stationary exposure of the DPS device in a sampling  
 383 site, which was located at the interface of the mobile sampled stretches 1 and 2, were among the  
 384 lowest observed values in the study for all assessed endpoints in the case of ED (Fig. 3). Even when  
 385 comparing data from stationary sampling site and stretch 1, which was directly upstream, the BEQ<sub>bio</sub>  
 386 values of the ED sample were lower for the stationary samples. This might indicate that mobile  
 387 sampling along the river stretches was more effective at sampling bioactive chemicals. On the other  
 388 hand, sampling at the stationary site took more than twice as long as in the mobile sampling and it is  
 389 possible that due to the longer exposure time at least some of bioactive chemicals reached equilibrium  
 390 between the ED samplers and sampled water. Thus, the calculated sampled water volume would be  
 391 overestimated for such chemicals and so the BEQ<sub>bio</sub> could be underestimated in the stationary  
 392 exposure samples. On the other hand, due to the longer sampling period, some non-polar chemicals  
 393 such as benzo(k)fluoranthene, benzo(ghi)perylene or some brominated diphenyl ethers, whose

394 concentrations were below method LOQ in samples from the mobile sampling, were quantified in the  
395 stationary sampling site (Tab. S2, S3).

396 The AhR-mediated BEQ<sub>bio</sub> levels in the samples were 3.8 –14.6 and 11.5–60.1 pg L<sup>-1</sup> for SR and ED,  
397 respectively (Table 3). For this endpoint, van der Oost et al. (2017) has proposed an effect-based trigger  
398 value (EBT; i.e. the BEQ<sub>bio</sub> level with a potential ecological health risk, of 150 ng L<sup>-1</sup> of benzo(a)pyrene  
399 equivalent). When transformed to the corresponding data format using the REP value of  
400 benzo(a)pyrene (Tab. S5), our data are relatively close to the proposed EBT limit and even exceed the  
401 value in ED sample from the stretch 8 (20–76 and 60–300 ng L<sup>-1</sup> of benzo(a)pyrene -equivalent for SR  
402 and ED, respectively).

403 The observed estrogenicity BEQ<sub>bio</sub> ranged 2–15 and 44–320 pg L<sup>-1</sup> in SR and ED samples, respectively.  
404 The concentrations in ED are relatively close to an annual average environmental quality standard of  
405 400 pg L<sup>-1</sup> for 17β-estradiol proposed by the European Commission (European Commission, 2015) as well  
406 as the previously proposed EBT values (Jarošová et al., 2014; van der Oost et al., 2017). Jarošová et al.,  
407 who derived a safe BEQ<sub>bio</sub> level for water burdened by wastewater treatment plants effluents using *in*  
408 *vivo* estrogenic PNECs, proposed the long-time exposure EBT of 200 pg L<sup>-1</sup> specifically for the MELN  
409 bioassay employed in the current study. Van der Oost et al. (2017) derived an EBT of 500 pg L<sup>-1</sup> BEQ<sub>bio</sub>  
410 from LOEC values for the ERα CALUX bioassay. Thus, our results show that estrogenic activity in the  
411 Danube river did not exceed the proposed EBT values, with the exception of stretch 5 in the case of  
412 the BEQ value proposed by Jarošová et al. (2014). On the other hand, even the Danube with its great  
413 dilution capacity contains estrogenic BEQ<sub>bio</sub> levels that are close to or, in case of stretch 5 (Romania,  
414 Serbia), exceed proposed EBT values.

415 The AR-mediated effect was assessed both in agonistic and antagonistic mode (see SI 1.4), but no  
416 androgenic effects were detected (data not shown) by passive nor large volume solid-phase extraction  
417 (LVSPE) sampling during JDS3 (Liska et al., 2015). In fact, androgenic effects in surface water were  
418 described to be associated with wastewater effluents, while anti-androgenic effects mostly prevail in  
419 the less-impacted surface waters (Jalova et al., 2013; König et al., 2016; Neale et al., 2017b).  
420 Antiandrogenic flutamide BEQ<sub>bio</sub> was 63 –432 and 132–2,707 ng L<sup>-1</sup> for SR and ED samples, respectively.  
421 Proposed EBT value for anti-androgenicity based on the AR CALUX assay is 25,000 ng L<sup>-1</sup> (van der Oost  
422 et al., 2017), so the detected levels should be safely below the EBT value, even though there might be  
423 some difference in sensitivity between the assay used for the EBT value estimation and MDA-kb2 assay  
424 employed in the present study.

425




### 3.3.1. Comparison of BEQ<sub>bio</sub> from passive and active sampling

426  
427 Besides passive sampling by DPS, the LVSPe approach for spot sampling was employed during the JDS3  
428 expedition cruise on the Danube River (Fig. S3; Neale et al., 2015a). In this method, suspended particles  
429 were first removed by flow-through centrifugation before extraction of water and so, similarly to  
430 passive sampling, the active sampling method collected mainly the dissolved fraction of chemicals that  
431 can serve as an estimate of  $C_{free}$  for chemicals with log  $K_{ow}$  lower than five (Prokeš et al., 2012). The  
432 LVSPe samples were assessed using a similar bioassay battery as in the current study. Thus, it was  
433 possible to compare BEQ<sub>bio</sub> of samples obtained by both sampling methods. Nevertheless, DPS samples  
434 reflect pollutants integrated along river stretches, while LVSPe was based on spot sampling, which did  
435 not representatively cover the stretches, thus individual samples could not be compared side by side.  
436 Therefore, it was only possible to compare the two sampling approaches using aggregated data from  
437 both studies. According to this comparison, the determined endocrine-disruptive and xenobiotic  
438 metabolism initiating potential of mixtures of compounds from passive sampling corresponded very  
439 well with LVSPe data considering the difference in sampling approaches (Fig. 5). The comparison of  
440 data from passive and active LVSPe sampling independently confirm that the calculation of sampled  
441 water volume by passive samplers was sufficiently accurate and representative to characterize the  
442 toxic potential of most bioactive chemicals contributing to these effects. Overall, BEQ<sub>bio</sub> of samples  
443 from LVSPe correspond much better with ED samplers than SR samples. It is not surprising since the  
444 SR sampler is designed mainly for non-polar chemicals, and the assessed biological endpoints are  
445 triggered rather by polar chemicals. On the other hand, non-polar chemicals tend to bioaccumulate,  
446 which can increase their potential to elicit chronic toxic effects in organisms *in situ*. Anyway, for active  
447 LVSPe and passive ED sampling, the ranges and medians of the BEQ<sub>bio</sub> for estrogenic and PXR-mediated  
448 potentials were in very good agreement and so both sampling methods seem to be similarly efficient  
449 for sampling of chemicals with these modes of action. AhR-mediated potential in SR samples  
450 corresponded very well to LVSPe samples, while it was on average more than two times higher in ED  
451 samples. No such effect was observed for the other endpoints that are more sensitive to polar  
452 chemicals.

453 ARE-mediated oxidative stress potential was on average three-times lower in the ED samples  
454 compared to LVSPe. The difference can be at least partly caused by the fact that LVSPe sample extracts  
455 were assessed for ARE-mediated potential using ARE-bla bioassay, which has differing sensitivity  
456 compared to the AREc32 assay employed in the current study (Stalter et al., 2016). Another adaptive  
457 stress response bioassay detected the p53-mediated effect in six out of ten ED samples with  
458 significantly higher EC<sub>IR1.5</sub> than in the LVSPe samples (median EC<sub>IR1.5</sub> of 609 and 173 REF, respectively),  
459 while no response was detected for the SR extracts. Thus, neither ED nor SR were effective samplers

460 for chemicals affecting the p53 activity. The NF- $\kappa$ B bioassay detected quantifiable effects only in LVSP  
461 samples, but not in the passive samples. While the LVSP water samples needed relatively low pre-  
462 concentration to elicit quantifiable effect, the passive samples were without any effect even at much  
463 higher pre-concentration levels. The low ARE and p53-mediated potentials and non-quantifiable NF $\kappa$ B-  
464 associated effects in the ED extracts indicate the differences in the spectrum of sampled chemicals  
465 between passive and LVSP sampling. Apparently, the applied passive samplers are less suitable for  
466 sampling compounds causing adaptive stress responses. For LVSP sampling, a combination of several  
467 adsorbents was used to quantitatively retain compounds with a very broad range of polarity, as well  
468 as neutral and charged chemicals (Neale et al., 2015a). In contrast, EDs consisted of the SDB-RPS  
469 sorbent material that retains mainly hydrophobic compounds, non-ionized polar compounds and  
470 organic cations. Since the SDB-RPS phase contains sulfonic acid functional groups, secondary sorption  
471 interactions can occur especially with basic compounds that contain amine functional groups. On the  
472 other hand, this sorbent is known to have a limited capacity for anionic compounds or dissociating  
473 compounds present dominantly in anionic form (Kaserzon et al., 2014) at neutral or slightly alkaline pH  
474 such as in the Danube river. The compounds with a low uptake capacity from neutral water include  
475 acidic compounds, such as most carboxylic acids (e.g. acidic herbicides or pharmaceuticals, but also  
476 naturally occurring carboxylic acids), some phenols, sulfinic and sulfonic acids and anions in general.

477



478

479 **Fig. 5 Comparison of bioanalytical equivalent concentrations in river water (BEQ<sub>bio</sub>) calculated**  
 480 **from passive and active sampling throughout all sampling sites and stretches; LVSPE active**  
 481 **sampling (large volume solid phase extraction), SR silicone rubber passive samplers, ED**  
 482 **Empore™ discs passive samplers; n.d.- not detected; data on active sampling taken from Neale**  
 483 **et al. (2015a); the box in the graph consists of two quartiles divided by median, the ends of the**  
 484 **whiskers represent the 10th and the 90th percentile and the dots individual outliers.**

485

### 486 3.3.2. Contribution of detected chemicals to the biological potentials

487 To compare data from chemical and biological analyses, levels of detected chemicals were converted  
 488 to BEQ<sub>chem</sub> using the concentration addition concept. We used REP values either from our own  
 489 experiments, from the peer reviewed literature, or the ToxCast database (Tab. S5). For some  
 490 chemicals, we were not able to find REP values assessed with the same bioassay as in the current study.  
 491 In these cases, use of REP values from an alternative analogous bioassay is indicated in Tab. S5 together  
 492 with EC values that were applied for REP calculation because these values differed among literature  
 493 sources.

494 Unfortunately, information on toxicological properties of many of the detected chemicals and their  
495 bioactivity in the studied endpoints is not available (Fig. S4), which limited the assessment of their  
496 potential contribution to the overall biological response. As soon as possible, the most widespread and  
497 abundant pollutants need to be prioritized for the characterization of their bioactivities to improve the  
498 mass-balance calculations and explanation of the observed effects.

499 Between 0.7 to 7.9 and 0.3 to 2% of BEQ<sub>bio</sub> for SR and ED samples, respectively, was explained by  
500 detected chemicals for the AhR-mediated response (Table 3). The main portion of BEQ<sub>chem</sub> was  
501 contributed by PAHs (namely benzo(*k*)fluoranthene, benzo(*b*)fluoranthene, benzo(*j*)fluoranthene and  
502 chrysene) in the SR samples and benzo(*k*)fluoranthene, terbuthylazine, propiconazole and 4-  
503 nonylphenol in ED (Fig. 6). A higher portion of the BEQ<sub>bio</sub> was explained in samples from the stationary  
504 exposure because of benzo(*k*)fluoranthene, whose concentration was below LOQ in the samples from  
505 the mobile sampling. The detection of this chemical in the stationary samples is likely caused by the  
506 longer integrative sampling at the stationary site which allowed sampling of its quantifiable level.


507 Less than 0.2% of PXR-mediated BEQ<sub>bio</sub> was explained in the SR samples, with 4-nonylphenol as the  
508 main contributor (Fig. S5). In the ED samples, less than 0.4% of the BEQ<sub>bio</sub> was explained, with 4-*t*-  
509 octylphenol, estrone, clotrimazole and metolachlor the main contributors of toxicity. Neale et al.  
510 (2015a) observed a similar level of explicability with samples from the parallel active sampling and also  
511 identified metolachlor as the main driver. Thus, either the more significant PXR activators were not  
512 identified yet or PXR is not very specific and it is activated by a large number of weak agonists. It might  
513 be also possible that synergistic interaction plays a role in the low explicability of this endpoint because  
514 it has been described that the ligand-binding domain of PXR can accommodate more than one weak  
515 agonist at the same time (Delfosse et al., 2015).

516 While estrogenic BEQ<sub>bio</sub> was explained by 0.2–3 % with bisphenol A, 4-nonylphenol, and 4-*t*-  
517 octylphenol as significant contributors in SR, 25 to 104% of estrogenicity was explained in the ED with  
518 estrone, 17 $\beta$ -estradiol and estriol, which showed very limited accumulation in SR (unpublished data).  
519 Other chemicals played a minor role. This supports the importance of steroids as water contaminants  
520 that play the most significant role in estrogenicity of waste and surface waters (Miège et al., 2009).

521 Anti-androgenicity was explained mostly by less than 4% and 6% in the SR and ED sampler extracts,  
522 respectively. The main contributors were tris(1,3-Dichloro-2-propyl)phosphate, bisphenol A, 4-*t*-  
523 octylphenol and 4-nonylphenol in SR extracts. The main identified contributors in ED extracts were  
524 clotrimazole, 4-*t*-octylphenol and ketoconazole. The rest of 19 identified anti-androgens did not  
525 contribute significantly.

526 ARE-mediated oxidative stress  $BEQ_{bio}$  was explained by less than 0.3 % with benzo(*b*)fluoranthene, 4-  
527 nonylphenol and benzo(*a*)pyrene in SR. In ED samples, less than 0.05 % of the  $BEQ_{bio}$  was explained  
528 and the main contributors were bisphenol A, 4-nonylphenol, propiconazole and atrazine in case of  
529 extracts from mobile sampling together with benzo(*b*)fluoranthene and benzo(*a*)pyrene in case of  
530 samples from stationary exposures. Similarly limited fraction of oxidative stress-mediated effects  
531 explained by detected chemicals was reported in the parallel active sampling study using an analogous  
532 ARE-bla bioassay (Neale et al., 2015a). Nevertheless, oxidative stress is probably elicited by a large  
533 number of diverse chemicals so it is not likely that there would be only a few drivers of this effect  
534 explaining a considerable portion of the  $BEQ_{bio}$  in most environmental chemical mixtures. While p53-  
535 response was below LOQ in SR samples and it was quantified in six of ten ED samples (Table 3), the  
536 detected chemicals allowed calculation of  $BEQ_{chem}$  (Fig. S6). The calculated  $BEQ_{chem}$  levels were lower  
537 than 0.004 and 0.015 ng L<sup>-1</sup> for SR and ED samplers, respectively, which was below LOQ of the bioassay  
538 (median value 31 and 59 ng L<sup>-1</sup>, respectively).  $BEQ_{chem}$  explained less than 0.02% of the  $BEQ_{bio}$  levels in  
539 the ED samples with the quantifiable response. This was comparable with data from LVSPE sampling  
540 where  $BEQ_{chem}$  explained 0.004 – 0.07 % of  $BEQ_{bio}$  (Neale et al., 2015a).

541 Our data document that employed passive sampling worked well for capturing chemicals responsible  
542 for the common studied endpoints such as AhR-, ER and AR-mediated activity, which are known to be  
543 associated with numerous known pollutants. On the other hand, in case of adaptive stress endpoints,  
544 namely p53- and NF-kB-mediated response, the effectiveness of the passive sampling was lower  
545 compared to the LVSPE sampling. Nevertheless, the drivers of these responses in surface waters are  
546 mostly unknown. For example, none of the 272 analyzed compounds in LVSPE samples were known to  
547 induce NF-kB according to the ToxCast database (Neale et al., 2015a). This was also documented by  
548 the low explicability of the observed  $BEQ_{bio}$  by the detected chemicals in case of p53-mediated  
549 genotoxic potential and ARE-mediated response despite the relatively large number of chemicals that  
550 were taken into account. Considering the selectivity of the used ED sampler, the chemicals present in  
551 water in an anionic form could contribute to the effects selectively detected in LVSPE samples. To  
552 better capture these types of chemicals, it is possible to broaden the applicability range of Empore™  
553 disk-based passive samplers to increase their capacity for binding anionic compounds. There is a  
554 variant of Empore™ disk, Empore™ Anion-SR available that is, similar to SDB-RPS, also based on  
555 polystyrene divinylbenzene particles, but contains additional strong anion exchange (SAX) functional  
556 groups. This disk variant was specifically designed for the extraction of acidic compounds (e.g.,  
557 carboxylic acids) from water samples. For sampling even broader range of compounds, including the  
558 dissociating acids and bases, SDB-RPS and Anion-SR Empore™ disks could be exposed side by side and  
559 their extracts combined to make a composite sample.



561

562 **Fig. 6 Percentage of the bioanalytical equivalents ( $BEQ_{bio}$ ) of ER and AhR- mediated effects**  
 563 **explained by the detected chemicals ( $BEQ_{chem}$ ) in samples from a dynamic passive sampling**  
 564 **of the Danube river; data from other bioassays are in Fig. S5 and S6; SR silicone rubber**  
 565 **samples, ED Empore™ discs samples; S1,2 – stationary sampling site; 1-8 sampled river**  
 566 **stretches.**

567

## 568 4. Conclusions

569 The DPS system provided a representative picture of the pollution situation at the studied site and  
 570 along the defined river stretches. The system could thus be suitable for sampling of transects of large  
 571 water bodies including rivers, lakes or seas. DPS device has effectively increased the sampling rate of  
 572 the passive samplers and this approach allowed toxicological profiling and detection of many chemicals  
 573 present in the water down to  $\text{pg L}^{-1}$  range in a short sampling time of several days. An integrated  
 574 approach combining passive sampling with chemical and biological analyses was shown suitable for  
 575 spatial profiling of a relatively less polluted river. Our data show that passive sampling should not be  
 576 limited to only one type of sampler. The complementarity of the samplers for hydrophilic and  
 577 hydrophobic compounds has been clearly demonstrated. While the data from bioanalysis, which  
 578 provides information on the sampled pollutant mixtures, indicated that most of the detected biological

579 effects were elicited by hydrophilic chemicals, the hydrophobic compounds have a higher  
580 bioaccumulative potential that could increase their relevancy for chronic impact on the exposed river  
581 ecosystem. However, besides estrogenicity only small portion of the biological effects could be  
582 explained by analyzed chemicals. This finding is also influenced by the lack of data on the biological  
583 potencies of detected chemicals. Comparison of the data from passive and active sampling indicates  
584 that the sampling rate of passive samplers for compounds contributing to endocrine-disruptive and  
585 xenobiotic metabolism potential can be relatively accurately estimated and confirms that both  
586 sampling approaches provide efficient means to monitor these chemicals. While LVSPE provides site-  
587 specific information, DPS enables integrative characterization of the pollution situation over space and  
588 time depending on the design of its employment.

## 589 **Acknowledgements**

590 We acknowledge the SOLUTIONS Project supported by the European Union Seventh Framework  
591 Programme (FP7-ENV-2013-two-stage Collaborative project) under grant agreement 603437 and  
592 RECETOX Research Infrastructure supported by the Czech Ministry of Education, Youth and Sports  
593 (LM2015051) and the European Structural and Investment Funds, Operational Programme Research,  
594 Development, Education (CZ.02.1.01/0.0/0.0/16\_013/0001761). This work was also financially  
595 supported by the French Ministry of Ecology (Programme 190) and Australian National Health and  
596 Medical Research Council (NHMRC) – European Union Collaborative Research Grant  
597 (APP1074775). Authors thank Petra Příbylová, Petr Kukučka, Šimon Vojta, Ondřej Audy, Jiří Kohoutek,  
598 Jitka Bečanová, Marek Pernica, Zdeněk Šimek and Alena Otoupalíková from RECETOX, Masaryk  
599 University for the instrumental analysis of samples. We thank Patrick Balaguer (Inserm, France) for the  
600 provision of the HG5LN-hPXR cell line and Alfhild Kringstad from NIVA for the instrumental analysis of  
601 samples.

602

## 603 **Appendix A. Supplementary data**

604 Supplementary data to this article can be found online at:

## 605 **References**

606 Allan, I.J., Harman, C., Ranneklev, S.B., Thomas, K. V., Grung, M., 2013. Passive sampling for target  
607 and nontarget analyses of moderately polar and nonpolar substances in water. *Environ. Toxicol.*  
608 *Chem.* 32, 1718–1726.

- 609 Balaguer, P., François, F., Comunale, F., Fenet, H., Boussioux, A.M., Pons, M., Nicolas, J.C., Casellas,  
610 C., 1999. Reporter cell lines to study the estrogenic effects of xenoestrogens. *Sci. Total Environ.*  
611 233, 47–56.
- 612 Baston, D.S., Denison, M.S., 2011. Considerations for potency equivalent calculations in the Ah  
613 receptor-based CALUX bioassay: Normalization of superinduction results for improved sample  
614 potency estimation. *Talanta* 83, 1415–1421.
- 615 Bláhová, L., Kohoutek, J., Procházková, T., Prudíková, M., Bláha, L., 2016. Phytoestrogens in milk:  
616 Overestimations caused by contamination of the hydrolytic enzyme used during sample  
617 extraction. *J. Dairy Sci.* 99, 6973–6982.
- 618 Booij, K., Smedes, F., 2010. An improved method for estimating in situ sampling rates of nonpolar  
619 passive samplers. *Environ. Sci. Technol.* 44, 6789–6794.
- 620 Booij, K., Vrana, B., Huckins, J.N., 2007. Theory, modelling and calibration of passive samplers used in  
621 water monitoring, in: Greenwood, R., Mills, G., Vrana, B. (Eds.), *Comprehensive Analytical*  
622 *Chemistry* 48. Passive Sampling Techniques in Environmental Monitoring. Elsevier, Amsterdam,  
623 pp. 141–169.
- 624 Brack, W., Klamer, H.J.C., López de Alda, M., Barceló, D., 2007. Effect-directed analysis of key  
625 toxicants in European river basins - A review. *Environ. Sci. Pollut. Res.* 14, 30–38.
- 626 Brumovský, M., Bečanová, J., Kohoutek, J., Thomas, H., Petersen, W., Sørensen, K., Sáňka, O.,  
627 Nizzetto, L., 2016. Exploring the occurrence and distribution of contaminants of emerging  
628 concern through unmanned sampling from ships of opportunity in the North Sea. *J. Mar. Syst.*  
629 162, 47–56.
- 630 Creusot, N., Kinani, S., Balaguer, P., Tapie, N., LeMenach, K., Maillot-Maréchal, E., Porcher, J.-M.,  
631 Budzinski, H., Aït-Aïssa, S., 2010. Evaluation of an hPXR reporter gene assay for the detection of  
632 aquatic emerging pollutants: screening of chemicals and application to water samples. *Anal.*  
633 *Bioanal. Chem.* 396, 569–583.
- 634 Creusot, N., Tapie, N., Piccini, B., Balaguer, P., Porcher, J.-M., Budzinski, H., Aït-Aïssa, S., 2013.  
635 Distribution of steroid- and dioxin-like activities between sediments, POCIS and SPMD in a  
636 French river subject to mixed pressures. *Environ. Sci. Pollut. Res.* 20, 2784–2794.
- 637 Delfosse, V., Dendele, B., Huet, T., Grimaldi, M., Boulahtouf, A., Gerbal-Chaloin, S., Beucher, B.,  
638 Roecklin, D., Muller, C., Rahmani, R., Cavallès, V., Daujat-Chavanieu, M., Vivat, V., Pascussi, J.-  
639 M., Balaguer, P., Bourguet, W., 2015. Synergistic activation of human pregnane X receptor by  
640 binary cocktails of pharmaceutical and environmental compounds. *Nat. Commun.* 6, 8089.
- 641 Duerksen-Hughes, P.J., Yang, J., Ozcan, O., 1999. p53 induction as a genotoxic test for twenty-five  
642 chemicals undergoing in vivo carcinogenicity testing. *Environ. Health Perspect.*
- 643 Escher, B.I., Allinson, M., Altenburger, R., Bain, P.A., Balaguer, P., Busch, W., Crago, J., Denslow, N.D.,  
644 Dopp, E., Hilscherova, K., Humpage, A.R., Kumar, A., Grimaldi, M., Jayasinghe, B.S., Jarosova, B.,  
645 Jia, A., Makarov, S., Maruya, K.A., Medvedev, A., Mehinto, A.C., Mendez, J.E., Poulsen, A.,  
646 Prochazka, E., Richard, J., Schifferli, A., Schlenk, D., Scholz, S., Shiraishi, F., Snyder, S., Su, G.,  
647 Tang, J.Y.M., Burg, B. van der, Linden, S.C. van der, Werner, I., Westerheide, S.D., Wong, C.K.C.,  
648 Yang, M., Yeung, B.H.Y., Zhang, X., Leusch, F.D.L., 2014. Benchmarking Organic Micropollutants  
649 in Wastewater, Recycled Water and Drinking Water with In Vitro Bioassays. *Environ. Sci.*  
650 *Technol.* 48, 1940–1956.
- 651 Escher, B.I., Dutt, M., Maylin, E., Tang, J.Y.M., Toze, S., Wolf, C.R., Lang, M., 2012. Water quality



652 assessment using the AREc32 reporter gene assay indicative of the oxidative stress response  
653 pathway. *J. Environ. Monit.* 14, 2877–85.

654 Escher, B.I., Leusch, F., 2012. *Bioanalytical Tools in Water Quality Assessment*. IWA Publishing.

655 European Commission, 2015. *The EU Water Framework Directive - integrated river basin management*  
656 *for Europe [WWW Document]*. URL [http://ec.europa.eu/environment/water/water-](http://ec.europa.eu/environment/water/water-framework/)  
657 [framework/](http://ec.europa.eu/environment/water/water-framework/) (accessed 8.24.17).

658 Giesy, J.P., Hilscherova, K., Jones, P.D., Kannan, K., Machala, M., 2002. Cell bioassays for detection of  
659 aryl hydrocarbon (AhR) and estrogen receptor (ER) mediated activity in environmental samples.  
660 *Mar. Pollut. Bull.* 45, 3–16.

661 Grabic, R., Fick, J., Lindberg, R.H., Fedorova, G., Tysklind, M., 2012. Multi-residue method for trace  
662 level determination of pharmaceuticals in environmental samples using liquid chromatography  
663 coupled to triple quadrupole mass spectrometry.

664 Jalova, V., Jarosova, B., Blaha, L., Giesy, J.P.P., Ocelka, T., Grabic, R., Jurcikova, J., Vrana, B.,  
665 Hilscherova, K., Jálková, V., Jarošová, B., Bláha, L., Giesy, J.P.P., Ocelka, T., Grabic, R., Jurčiková, J.,  
666 Vrana, B., Hilscherová, K., 2013. Estrogen-, androgen- and aryl hydrocarbon receptor mediated  
667 activities in passive and composite samples from municipal waste and surface waters. *Environ.*  
668 *Int.* 59, 372–83.

669 Janošek, J., Hilscherová, K., Bláha, L., Holoubek, I., 2006. Environmental xenobiotics and nuclear  
670 receptors--Interactions, effects and in vitro assessment. *Toxicol. Vit.* 20, 18–37.

671 Jarosova, B., Blaha, L., Vrana, B., Randak, T., Grabic, R., Giesy, J.P., Hilscherova, K., 2012. Changes in  
672 concentrations of hydrophilic organic contaminants and of endocrine-disrupting potential  
673 downstream of small communities located adjacent to headwaters. *Environ. Int.* 45, 22–31.

674 Jarošová, B., Bláha, L., Giesy, J.P., Hilscherová, K., 2014. What level of estrogenic activity determined  
675 by in vitro assays in municipal waste waters can be considered as safe? *Environ. Int.* 64, 98–109.

676 Jin, L., Gaus, C., Escher, B.I., 2015. Adaptive Stress Response Pathways Induced by Environmental  
677 Mixtures of Bioaccumulative Chemicals in Dugongs. *Environ. Sci. Technol.* 49, 6963–6973.

678 Kaserzon, S.L., Hawker, D.W., Kennedy, K., Bartkow, M., Carter, S., Booij, K., Mueller, J.F., 2014.  
679 Characterisation and comparison of the uptake of ionizable and polar pesticides,  
680 pharmaceuticals and personal care products by POCIS and Chemcatchers. *Environ. Sci. Process.*  
681 *Impacts* 16, 2517–26.

682 Keller, V.D.J., Williams, R.J., Lofthouse, C., Johnson, A.C., 2014. Worldwide estimation of river  
683 concentrations of any chemical originating from sewage-treatment plants using dilution factors.  
684 *Environ. Toxicol. Chem.* 33, 447–52.

685 König, M., Escher, B.I., Neale, P.A., Krauss, M., Hilscherová, K., Novák, J., Teodorović, I., Schulze, T.,  
686 Seidensticker, S., Kamal Hashmi, M.A., Ahlheim, J., Brack, W., 2016. Impact of untreated  
687 wastewater on a major European river evaluated with a combination of in vitro bioassays and  
688 chemical analysis. *Environ. Pollut.*

689 Kortenkamp, A., Evans, R., Martin, O., Mckinlay, R., Orton, F., Rosivatz, E., 2012. State of the art  
690 assessment of endocrine disrupters Final Report Project Contract Number  
691 070307/2009/550687/SER/D3 [WWW Document]. URL  
692 [http://ec.europa.eu/environment/chemicals/endocrine/pdf/annex1\\_summary\\_state\\_of\\_scienc](http://ec.europa.eu/environment/chemicals/endocrine/pdf/annex1_summary_state_of_scienc)  
693 [e.pdf](http://ec.europa.eu/environment/chemicals/endocrine/pdf/annex1_summary_state_of_scienc) (accessed 11.23.16).

694 Lemaire, G.G., Mnif, W., Pascussi, J.-M., Pillon, A., Rabenoelina, F., Fenet, H.H., Gomez, E., Casellas,  
695 C., Nicolas, J.-C., Cavaillès, V., Duchesne, M.-J.M.-J., Balaguer, P., Cavaillès, V., Duchesne, M.-  
696 J.M.-J., Balaguer, P., 2006. Identification of New Human Pregnane X Receptor Ligands among  
697 Pesticides Using a Stable Reporter Cell System. *Toxicol. Sci.* 91, 501–509.

698 Leusch, F.D.L., Neale, P.A., Hebert, A., Scheurer, M., Schriks, M.C.M., 2017. Analysis of the sensitivity  
699 of in vitro bioassays for androgenic, progestagenic, glucocorticoid, thyroid and estrogenic  
700 activity: Suitability for drinking and environmental waters. *Environ. Int.* 99, 120–130.

701 Lin, Y.-H., Chen, C.-Y., Wang, G.-S., 2007. Analysis of steroid estrogens in water using liquid  
702 chromatography/tandem mass spectrometry with chemical derivatizations. *Rapid Commun.*  
703 *Mass Spectrom.* 21, 1973–1983.

704 Liska, I., Wagner, F., Sengl, M., Deutsch, K., Slobodnik, J. (Eds.), 2015. Joint Danube Survey 3- a  
705 Comprehensive Analysis of Danube Water Quality. ICPDR – International Commission for the  
706 Protection of the Danube River, Vienna.

707 Liu, L., Chen, L., Shao, Y., Zhang, L., Floehr, T., Xiao, H., Yan, Y., Eichbaum, K., Hollert, H., Wu, L., 2014.  
708 Evaluation of the Ecotoxicity of Sediments from Yangtze River Estuary and Contribution of  
709 Priority PAHs to Ah Receptor-Mediated Activities. *PLoS One* 9, e104748.

710 Lohmann, R., Klanova, J., Kukucka, P., Yonis, S., Bollinger, K., 2013. Concentrations, Fluxes, and  
711 Residence Time of PBDEs Across the Tropical Atlantic Ocean. *Environ. Sci. Technol.* 47, 13967–  
712 13975.

713 Mayer, P., Tolls, J., Hermens, J.L.M., Mackay, D., 2003. Peer Reviewed: Equilibrium Sampling Devices.  
714 *Environ. Sci. Technol.* 37, 184A–191A.

715 McLachlan, J.A., 2016. Environmental signaling: from environmental estrogens to endocrine-  
716 disrupting chemicals and beyond. *Andrology* 4, 684–694.

717 Miège, C., Gabet, V., Coquery, M., Karolak, S., Jugan, M.-L., Oziol, L., Levi, Y., Chevreuril, M., 2009.  
718 Evaluation of estrogenic disrupting potency in aquatic environments and urban wastewaters by  
719 combining chemical and biological analysis. *TrAC Trends Anal. Chem.* 28, 186–195.

720 Nagy, S.R., Sanborn, J.R., Hammock, B.D., Denison, M.S., 2002. Development of a Green Fluorescent  
721 Protein-Based Cell Bioassay for the Rapid and Inexpensive Detection and Characterization of Ah  
722 Receptor Agonists. *Toxicol. Sci.* 65, 200–210.

723 Neale, P.A., Ait-Aïssa, S., Brack, W., Creusot, N., Denison, M.S., Deutschmann, B., Hilscherova, K.,  
724 Hollert, H., Krauss, M., Novák, J., Schulze, T., Seiler, T.B., Serra, H., Shao, Y., Escher, B.I., 2015a.  
725 Linking in vitro effects and detected organic micropollutants in surface water using mixture  
726 toxicity modeling. *Environ. Sci. Technol.* 49, 14614–14624.

727 Neale, P.A., Altenburger, R., Aït-Aïssa, S., Brion, F., Busch, W., de Aragão Umbuzeiro, G., Denison,  
728 M.S., Du Pasquier, D., Hilscherová, K., Hollert, H., Morales, D.A., Novák, J., Schlichting, R., Seiler,  
729 T.-B., Serra, H., Shao, Y., Tindall, A.J., Tollefsen, K.E., Williams, T.D., Escher, B.I., 2017a.  
730 Development of a bioanalytical test battery for water quality monitoring: Fingerprinting  
731 identified micropollutants and their contribution to effects in surface water. *Water Res.* 123,  
732 734–750.

733 Neale, P.A., Munz, N.A., Aït-Aïssa, S., Altenburger, R., Brion, F., Busch, W., Escher, B.I., Hilscherová, K.,  
734 Kienle, C., Novák, J., Seiler, T.-B., Shao, Y., Stamm, C., Hollender, J., 2017b. Integrating chemical  
735 analysis and bioanalysis to evaluate the contribution of wastewater effluent on the  
736 micropollutant burden in small streams. *Sci. Total Environ.* 576, 785–795.

- 737 Neale, P.A., Stalter, D., Tang, J.Y.M., Escher, B.I., 2015b. Bioanalytical evidence that chemicals in  
738 tattoo ink can induce adaptive stress responses. *J. Hazard. Mater.* 296, 192–200.
- 739 Pernica, M., Poloucká, P., Seifertová, M., Šimek, Z., 2015. Determination of alkylphenols in water  
740 samples using liquid chromatography–tandem mass spectrometry after pre-column  
741 derivatization with dansyl chloride. *J. Chromatogr. A* 1417, 49–56.
- 742 Prokeš, R., Vrana, B., Klánová, J., 2012. Levels and distribution of dissolved hydrophobic organic  
743 contaminants in the Morava river in Zlín district, Czech Republic as derived from their  
744 accumulation in silicone rubber passive samplers. *Environ. Pollut.* 166, 157–166.
- 745 Reddy, S.P., 2008. The antioxidant response element and oxidative stress modifiers in airway  
746 diseases. *Curr. Mol. Med.* 8, 376–83.
- 747 Rusina, T.P., Smedes, F., Koblížková, M., Klanová, J., 2010. Calibration of Silicone Rubber Passive  
748 Samplers: Experimental and Modeled Relations between Sampling Rate and Compound  
749 Properties. *Environ. Sci. Technol.* 44, 362–367.
- 750 Sadílek, J., Spálovská, P., Vrana, B., Vávrová, M., Maršálek, B., Šimek, Z., 2016. Comparison of  
751 extraction techniques for isolation of steroid oestrogens in environmentally relevant  
752 concentrations from sediment. *Int. J. Environ. Anal. Chem.* 96, 1022–1037.
- 753 Simmons, S.O., Fan, C.Y., Ramabhadran, R., 2009. Cellular stress response pathway system as a  
754 sentinel ensemble in toxicological screening. *Toxicol. Sci.*
- 755 Smedes, F., Booij, K., 2012. Guidelines for passive sampling of hydrophobic contaminants in water  
756 using silicone rubber samplers. *ICES Techniques in Marine Environmental Sciences No. 52.*, ICES  
757 TECHNIQUES IN MARINE ENVIRONMENTAL SCIENCES. International Council for the  
758 Exploration of the Sea, Denmark.
- 759 Stalter, D., O'Malley, E., Von Gunten, U., Escher, B.I., 2016. Fingerprinting the reactive toxicity  
760 pathways of 50 drinking water disinfection by-products. *Water Res.* 91, 19–30.
- 761 US EPA, 2015. US EPA (2015) Interactive Chemical Safety for Sustainability (iCSS) Dashboard v2  
762 [WWW Document]. URL <http://actor.epa.gov/dashboard/> (accessed 2.12.17).
- 763 van der Oost, R., Sileno, G., Suárez-Muñoz, M., Nguyen, M.T., Besselink, H., Brouwer, A., 2017.  
764 SIMONI (Smart Integrated Monitoring) as a novel bioanalytical strategy for water quality  
765 assessment: Part I-model design and effect-based trigger values. *Environ. Toxicol. Chem.* 36,  
766 2385–2399.
- 767 Villeneuve, D.L., Blankenship, A.L., Giesy, J.P., 2000. Derivation and application of relative potency  
768 estimates based on in vitro bioassay results. *Environ. Toxicol. Chem.* 19, 2835–2843.
- 769 Vrana, B., Smedes, F., Allan, I., Rusina, T., Okonski, K., Hilscherová, K., Novák, J., Tarábek, P.,  
770 Slobodník, J., n.d. Dynamic mobile passive sampling of trace organic compounds: evaluation of  
771 sampler performance in the Danube river. *Sci. Total Environ.*
- 772 Wang, X.J., Hayes, J.D., Wolf, C.R., 2006. Generation of a Stable Antioxidant Response Element-  
773 Driven Reporter Gene Cell Line and Its Use to Show Redox-Dependent Activation of Nrf2 by  
774 Cancer Chemotherapeutic Agents. *Cancer Res.* 66, 10983–10994.
- 775 Wilson, V.S., Bobseine, K., Lambright, C.R., Gray, L.E., 2002. A novel cell line, MDA-kb2, that stably  
776 expresses an androgen- and glucocorticoid-responsive reporter for the detection of hormone  
777 receptor agonists and antagonists. *Toxicol. Sci.* 66, 69–81.

778

779

**Table 1 Overview of bioassays in the study**

Bioassay	Endpoint	Positive reference compound	Method reference	Data evaluation method	EC value
CAFLUX-H4G1.1c2	Activation of AhR	2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (TCDD)	(Nagy et al., 2002)	Log-logistic dose-response model	EC <sub>20</sub>
MDA-kb2	Activation / Inhibition of AR	Dihydrotestosterone / Flutamide	(Wilson et al., 2002)	Log-logistic dose-response model	IC <sub>20</sub>
HG5LN-hPXR	Activation of PXR	SR 12813 <sup>1</sup>	(Lemaire et al., 2006)	Log-logistic dose-response model	EC <sub>20</sub>
MELN	Activation of ER	17β-Estradiol	(Balaguer et al., 1999)	Log-logistic dose-response model	EC <sub>20</sub>
AREc32	Oxidative stress response	tert-Butylhydroquinone (tBHQ)	(Escher et al., 2012; Wang et al., 2006)	Linear concentration-effect curve	EC <sub>IR1.5</sub>
p53RE-bla	p53 response	Mitomycin	(Neale et al., 2015b)	Linear concentration-effect curve	EC <sub>IR1.5</sub>
NF-κB-bla	NF-κB response	Tumor necrosis factor alpha (TNFα)	(Jin et al., 2015)	Linear concentration-effect curve	EC <sub>IR1.5</sub>

<sup>1</sup>Tetraethyl 2-(3,5-di-tert-butyl-4-hydroxyphenyl)ethenyl-1,1-bisphosphonate

**Table 2 Sum concentrations (pg L<sup>-1</sup>) of groups of chemicals detected in samples from dynamic passive sampling; 1-8 samples from mobile sampling of river stretches; SR silicone rubber sampler, ED Empore™ disc sampler; No. number of analyzed chemicals in the category; <LOD concentrations below the limit of detection; for molar concentrations or individual chemical levels see SI; CUPs-current use pesticides, PAHs- polycyclic aromatic hydrocarbons, PCBs- polychlorinated biphenyls, OCPs- organochlorinated pesticides, BDEs- brominated diphenyl ethers, NBFRs- novel brominated flame retardants**

Sampler	Chemicals	No.	S1	S2	1	2	3	4	5	6	7	8
ED	ATBs	31	12,670	12,230	11,476	23,922	23,126	17,393	21,699	12,656	33,195	29,262
	Cardiovascular	15	14,183	14,964	11,748	19,693	21,649	11,367	11,730	7,262	10,764	11,026
	Psychoactive	30	25,439	21,421	17,860	28,957	26,290	18,019	24,001	24,855	33,596	28,750
	Antihistamins	8	255	91	275	<LOD	112	348	242	<LOD	583	<LOD
	Antifungals	8	3,260	4,898	4,121	8,291	8,146	8,476	10,560	10,466	10,582	9,297
	Antidiabetics	4	31	4	32	42	98	35	20	21	97	27
	Statins	4	289	318	286	400	1,259	1,031	1,219	418	590	507
	Other pharm. <sup>1</sup>	8	290	248	<LOD	<LOD	122	533	<LOD	295	<LOD	<LOD
	CUPs	40	17,543	28,746	33,152	20,492	32,509	17,067	27,033	28,373	67,140	41,961
	Alkylphenols	3	7,630	7,790	12,225	12,484	15,209	26,311	34,513	20,982	22,399	19,817
	PAHs	29	6,274	7,541	9,633	12,134	22,411	17,384	8,111	8,007	6,981	12,382
	Steroids	5	176	198	201	351	582	626	1,151	398	770	816
	Phytosterols	11	72,010	52,776	29,885	45,964	49,994	34,753	47,458	46,209	48,996	50,809
Flavonoids	8	798	955	954	636	806	2,916	1,990	693	3,211	10,329	
SR	PAHs	29	17,342	23,000	18,215	21,063	51,067	37,296	15,975	12,002	13,965	20,704
	PCBs	7	217	244	171	291	167	172	369	158	307	295
	OCPs	12	188	223	156	244	372	313	358	465	808	1,872
	BDEs	9	4	6	22	2	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
	NBFRs	15	7	13	23	25	15	9	17	22	5	12
	Musks	6	18,582	27,839	11,240	11,894	18,317	17,915	25,395	10,060	13,471	9,732
	Alkylphenols	3	2,932	3,526	5,749	6,270	7,959	10,012	8,511	3,525	7,278	6,236
	Alkylphosphates	14	186,725	213,440	98,815	121,763	111,142	79,046	104,446	147,234	139,681	148,697

<sup>1</sup> furosemide, alfuzosin, naloxone, ranitidine

**Table 3 Toxic potentials of samples assessed with bioassays expressed as bioanalytical equivalents of respective reference compound (BEQ<sub>bio</sub> ± SD) and bioanalytical equivalents predicted from chemical analyses (BEQ<sub>chem</sub>) with percent of BEQ<sub>bio</sub> explained by chemical analyses in brackets. AhR – aryl hydrocarbon receptor-mediated response (tetrachlorodibenzo-*p*-dioxin); ER – estrogenicity (17β-estradiol); anti-AR – antiandrogenicity (flutamide); PXR – pregnane X receptor-mediated response (SR12813).**

		sample	S1	S2	1	2	3	4	5	6	7	8
endpoint	sample	No.										
			Static exposure 1	Static exposure 2	Passau–Bratislava	Bratislava–Budapest	Budapest–Vukovar	Vukovar–Belgrade	Belgrade–Turnu-Severin	Turnu-Severin–Ruse	Ruse–Braila	Braila–Tulcea
AhR [pg L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	9.7 ± 3.5	3.8 ± 1.1	14.6 ± 5	9.7 ± 3.4	6.6 ± 1.9	7.5 ± 1.5	8.2 ± 2.5	7.2 ± 1.9	7.7 ± 0.8	10.6 ± 3.1
		BEQ <sub>chem</sub>	0.32 (3.3 %)	0.3 (7.9 %)	0.1 (0.7 %)	0.13 (1.3 %)	0.16 (2.4 %)	0.07 (0.9 %)	0.11 (1.4 %)	0.11 (1.6 %)	0.06 (0.8 %)	0.08 (0.7 %)
	ED	BEQ <sub>bio</sub>	15.5 ± 1	11.5 ± 1.1	23.2 ± 5.4	30.6 ± 10.2	22 ± 4.9	21.4 ± 0.6	12.9 ± 3.9	28.3 ± 7.7	26.7 ± 7.6	60.1 ± 20.4
		BEQ <sub>chem</sub>	0.249 (1.6 %)	0.265 (2.3 %)	0.141 (0.6 %)	0.131 (0.4 %)	0.267 (1.2 %)	0.11 (0.5 %)	0.135 (1 %)	0.107 (0.4 %)	0.195 (0.7 %)	0.163 (0.3 %)
PXR [ng L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	211 ± 73	148 ± 103	134 ± 23	83 ± 32	123 ± 48	72 ± 41	104 ± 32	261 ± 108	205 ± 68	206 ± 36
		BEQ <sub>chem</sub>	0.04 (0.02 %)	0.07 (0.05 %)	0.09 (0.07 %)	0.09 (0.11 %)	0.12 (0.09 %)	0.14 (0.2 %)	0.12 (0.11 %)	0.05 (0.02 %)	0.11 (0.05 %)	0.1 (0.05 %)
	ED	BEQ <sub>bio</sub>	314 ± 206	200 ± 115	685 ± 281	773 ± 292	237 ± 56	1,118 ± 148	644 ± 130	1,613 ± 670	720 ± 318	1,173 ± 207
		BEQ <sub>chem</sub>	0.39 (0.2 %)	0.46 (0.4 %)	0.49 (0.1 %)	0.61 (0.1 %)	0.85 (0.4 %)	0.98 (0.1 %)	1.52 (0.2 %)	0.84 (0.05 %)	1.23 (0.2 %)	1.03 (0.1 %)
ER [pg L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	3.6 ± 0.2	2.6 ± 0.4	5.7 ± 0.7	2 ± 0.1	10.6 ± 1.9	5.4 ± 0.2	6.8 ± 0.8	11.2 ± 1.5	15.6 ± 2.4	12.6 ± 0.2
		BEQ <sub>chem</sub>	0.12 (3.3 %)	0.12 (4.8 %)	0.08 (1.4 %)	0.08 (3.7 %)	0.13 (1.2 %)	0.15 (2.7 %)	0.11 (1.7 %)	0.03 (0.3 %)	0.08 (0.5 %)	0.09 (0.7 %)
	ED	BEQ <sub>bio</sub>	44.4 ± 24	49.7 ± 9	83.2 ± 22	155.5 ± 24	163.9 ± 37	140.1 ± 26	320.3 ± 67	86 ± 38	131.4 ± 17	150.6 ± 49
		BEQ <sub>chem</sub>	19.7 (44 %)	22.2 (45 %)	22.6 (27 %)	39 (25 %)	105.3 (64 %)	117.9 (84 %)	228.6 (71 %)	47.4 (55 %)	138.3 (105 %)	144.2 (96 %)
anti-AR [ng L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	273 ± 84	74 ± 24	274 ± 100	246 ± 51	258 ± 111	63 ± 11	146 ± 33	202 ± 30	194 ± 36	432 ± 36
		BEQ <sub>chem</sub>	1.96 (0.7 %)	3.29 (4.5 %)	1.79 (0.7 %)	1.13 (0.5 %)	1.65 (0.6 %)	1.17 (1.8 %)	1.04 (0.7 %)	0.85 (0.4 %)	1.23 (0.6 %)	1.1 (0.3 %)
	ED	BEQ <sub>bio</sub>	334 ± 51	132 ± 46	599 ± 189	865 ± 234	592 ± 45	595 ± 148	1,640 ± 542	2,707 ± 289	1,849 ± 531	1,708 ± 495
		BEQ <sub>chem</sub>	7.29 (2.2 %)	8.38 (6.4 %)	9.33 (1.6 %)	15.25 (1.8 %)	17.79 (3 %)	16.41 (2.8 %)	21.29 (1.3 %)	19.42 (0.7 %)	21.48 (1.2 %)	15.23 (0.9 %)

ARE [ng L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	327 ± 230	395 ± 262	782 ± 344	962 ± 520	659 ± 288	827 ± 672	647 ± 351	433 ± 289	1,012 ± 623	1,164 ± 787
		BEQ <sub>chem</sub>	1.324 (0.41 %)	1.25 (0.32 %)	0.792 (0.1 %)	0.83 (0.09 %)	1.09 (0.17 %)	0.62 (0.07 %)	1.15 (0.18 %)	0.7 (0.16 %)	0.51 (0.05 %)	0.58 (0.05 %)
	ED	BEQ <sub>bio</sub>	1,181 ± 511	1,054 ± 214	2,540 ± 577	2,724 ± 971	2,449 ± 622	2,219 ± 193	1,838 ± 339	1,726 ± 378	2,063 ± 439	2,071 ± 361
		BEQ <sub>chem</sub>	0.84 (0.07 %)	1.07 (0.1 %)	0.75 (0.03 %)	0.65 (0.02 %)	0.99 (0.04 %)	0.81 (0.04 %)	1.06 (0.06 %)	1.3 (0.08 %)	1.93 (0.09 %)	1.96 (0.09 %)
p53 [ng L <sup>-1</sup> ]	SR	BEQ <sub>bio</sub>	<17.5	<16.1	<25.3	<50.9	<30.7	<32	<30.7	<33	<54	<59.6
		BEQ <sub>chem</sub>	0.0016	0.0045	0.0005	0.0005	0.0005	0.0004	0.004	0.0003	0.0003	0.0004
	ED	BEQ <sub>bio</sub>	50 ± 1	38 ± 5	83 ± 9	<95	102 ± 16	<60	85 ± 10	86 ± 7	<101	<112
		BEQ <sub>chem</sub>	0.004 (0.008 %)	0.005 (0.012 %)	0.004 (0.005 %)	0.003	0.005 (0.005 %)	0.01	0.014 (0.017 %)	0.007 (0.008 %)	0.008	0.008





## Supporting Information

### Effect-based monitoring of Danube river using mobile passive sampling approach

Jiří Novák<sup>a</sup>, Branislav Vrana<sup>a</sup>, Tatsiana Rusina<sup>a</sup>, Krzysztof Okonski<sup>a</sup>, Roman Grabic<sup>b</sup>, Peta A. Neale<sup>c,d</sup>, Beate I. Escher<sup>d,e,f</sup>, Miroslava Mácová<sup>e</sup>, Selim Ait-Aissa<sup>g</sup>, Nicolas Creusot<sup>g</sup>, Ian Allan<sup>h</sup>, Klára Hilscherová<sup>a\*</sup>

<sup>a</sup>Masaryk University, Research Centre for Toxic Compounds in the Environment (RECETOX), Kamenice 753/5, 625 00 Brno, Czech Republic

<sup>b</sup>University of South Bohemia in České Budějovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Zátíší 728/II, CZ-389 25 Vodňany, Czech Republic

<sup>c</sup>Australian Rivers Institute, Griffith School of Environment, Griffith University, Southport QLD 4222, Australia

<sup>d</sup>The University of Queensland, Queensland Alliance for Environmental Health Sciences (QAEHS), Coopers Plains, Qld 4108, Australia

<sup>e</sup>UFZ Helmholtz Centre for Environmental Research – UFZ, Permoserstr. 15, 04318 Leipzig, Germany

<sup>f</sup>Eberhard Karls University Tübingen, Hölderlinstr. 12, 72074 Tübingen, Germany

<sup>g</sup>Institut National de l'Environnement Industriel et des Risques (INERIS), Unité Ecotoxicologie, Parc ALATA - BP2, 60550 Verneuil en Halatte, France

<sup>h</sup>Norwegian Institute for Water Research, Gaustadalle´en 21, NO-0349 Oslo, Norway

## Table of contents

Supporting Information.....	1
Effect-based monitoring of Danube river using mobile passive sampling approach.....	1
1 Passive sampling.....	2
1.1 Sample processing.....	3
1.1.1 Silicone rubber (SR) sheets.....	3
1.1.2 Empore disks .....	3
1.1.3 Low-density polyethylene (LDPE) sheets .....	4
1.2 Chemical analysis.....	4
1.3 Estimation of sampling rates.....	5
1.3.1 Silicone rubber samplers .....	5
1.3.2 Empore disc samplers.....	5
1.3.3 Mutual comparison of two co-deployed DPS devices.....	6
1.4 Bioanalysis .....	8

Fig. S1 Comparison of levels of PAHs and 4-nonylphenol per sampling surface of SR and ED samplers from 8 mobile and 2 stationary DPS samples. ....	7
Fig. S2 Comparison of PRC fractions retained in LDPE sheets in two co-deployed DPS devices (TOX PS and CHEM PS) in parallel in 8 Danube stretches .....	8
Fig. S3 Map of the sampling cruise with the stretches and stationary exposure site (red dot); blue dots indicate active LVSPE sampling sites from Neale et al. (2015a).....	10
Fig. S4 Overview of detected chemicals described in ToxCast database to be in/-active in eliciting ARE-mediated oxidative stress and anti-androgenic effects. The endpoints, where BEQ <sub>chem</sub> levels were calculated mainly with ToxCast database-derived REPs. ....	11
Fig. S5 Percentage of the bioanalytical equivalents (BEQ <sub>bio</sub> ) explained by the detected chemicals (BEQ <sub>chem</sub> ).....	12
Fig. S6 p53-mediated bioanalytical equivalents of mitomycin calculated from detected chemicals levels (BEQ <sub>chem</sub> ) in samples from a dynamic passive sampling of the Danube river; SR silicone rubber samples, ED Empore discs samples; S1, 2 – stationary sampling site; 1-8 sampled river stretches.....	13
Tab. S1 River stretches and stationary exposure localities sampled with passive samplers .....	13
Tab. S2 Concentrations [pg/L] of chemicals detected in SR sampler extracts; PAHs - polycyclic aromatic hydrocarbons; BDE - brominate diphenyl ethers .....	14
Tab. S3 Concentrations [pg/L] of chemicals detected in ED sampler extracts.....	18
Tab. S4 Sum molar concentrations of groups of chemicals detected in samples from dynamic passive sampling .....	26
Tab. S5 Relative effect potencies (REPs) of detected compounds.....	27

## 1 Passive sampling

The DPS devices were deployed on the frontal deck of the expedition ship. For sampling, the device was immersed in a flow-through system that consisted of a 600 L stainless steel tank. The river water in the tank was exchanged at a rate of approx. 3 m<sup>3</sup> h<sup>-1</sup> by a high-performance pump. The water intake to the chamber was provided by a vertical steel pipe positioned in front of the ship. The water sampling depth was approx. 0.5 m below the water level. Two sets of DPS samplers were employed in stationary sampling site Čunovo (Slovakia) located at the interface of stretches 1 and 2.

The silicone rubber (SR) sampler consisted of a single sheet of Altesil® silicone rubber with dimensions 14×28 cm and 0.5 mm thickness. The mass of a sampler was approx. 23 g and the surface area exposed to water 392 cm<sup>2</sup> (one side of the sheet). Prior to use all SR samplers were Soxhlet-extracted in ethyl acetate for 72 h to remove non-polymerized residues. SR samplers for chemical analyses were spiked prior to exposure with Performance Reference Compounds (PRCs) that were partially released during exposure. The employed PRC mixture contained perdeuterated biphenyl and 13 polychlorinated biphenyl (PCB) congeners that do not occur in technical mixtures (PCB 1, PCB 2, PCB 3, PCB 10, PCB 14, PCB 21, PCB 30, PCB 50, PCB 55, PCB 78, PCB 104, PCB 145, PCB 204). The residual concentration of PRCs was compared with the initial amount of PRCs analyzed in samplers that have not been exposed.

The ED sampler consisted of 10 solid phase extraction disks Empore® SDB-RPS with 47 mm diameter. The mass of a sampler was approx. 3.2 g and the surface area exposed to water was 173 cm<sup>2</sup>. Before exposure, Empore discs were washed and conditioned by subsequent immersing in 1) 100 ml acetone; 2) 100 ml isopropanol; 3) 100 ml methanol, 4) 2x 100 ml ultrapure water and kept immersed in ultrapure water until exposure. For this sampler type, PRCs were not used. ED sampling rate was determined indirectly from SR sampling rate and levels of PAHs and 4-nonylphenol that were detected both in SR and ED samples as described in detail in chapter S1.3 and Vrana et al. (submitted).

The sampling was performed only during cruising of the ship or when the ship was anchored outside harbors in areas without visible pollution point sources e.g. wastewater discharges, industrial areas next to the river or sites with visible oil film on the surface of the water. The sampling device was always switched off before entering harbors and resumed upon leaving the harbors. The samplers were mounted into the DPS device directly before sampling and recovered immediately after finishing the sampling. The recovered samplers were placed back into their respective storage containers, stored in a refrigerator at 4°C on board of the ship and transported to the processing laboratory once per week, where they were stored in a freezer at -20°C.

## **1.1 Sample processing**

### **1.1.1 Silicone rubber (SR) sheets**

Before extraction, SR samplers for chemical analyses were spiked with recovery internal standards (RIS; d<sub>8</sub>-naphthalene, d<sub>10</sub>-phenanthrene, d<sub>12</sub>-perylene, PCB 4, PCB 29, PCB 185, a mixture of <sup>13</sup>C BDEs, C<sub>13</sub> caffeine, d<sub>13</sub>-alachlor, d<sub>6</sub>-diuron, d<sub>10</sub>-simazine, 4-n-nonylphenol). The SR samplers were extracted for 8 hours in methanol using Soxhlet extraction. The volume of the extract was reduced using Kuderna-Danish (K-D) apparatus and under nitrogen flow to 2 ml. Extracts of samplers intended for bioanalysis were processed in the same way but without spiking any standards. After processing, they were split to aliquots for analysis by the different bioassays. 20 % aliquot of the sample for chemical analysis was used for instrumental analysis by LC/MS methods. The remaining 80 % aliquot of samples for chemical analysis was azeotropically transferred to hexane using K-D apparatus. Aliquots of the extract were divided into vials for different types of GC/MS analysis. The extract aliquots for analysis of PAHs were further cleaned-up by a silica gel column using diethyl ether/acetone elution. The extract aliquots for analysis of organochlorine compounds (OCs), polybrominated diphenyl ethers (PBDEs) and PRCs were purified by a clean-up using activated silica gel modified with sulfuric acid. Following clean-up, addition of internal standards and volume reduction using a K-D apparatus, samples were analyzed using a GC-MS/MS method for indicator PCBs, PBDEs, organochlorinated pesticides (OCPs) and PRCs.

### **1.1.2 Empore disks**

Unlike ED samplers for bioanalysis, the samplers for chemical analysis were spiked with recovery internal standards (C<sub>13</sub>-caffeine, C<sub>13</sub>-triclosan, m<sub>8</sub>-PFOA, n<sub>8</sub>-PFOS, C<sub>13</sub>-alachlor, d<sub>6</sub>-diuron, d<sub>10</sub>-simazine, deuterated EE2, n-nonylphenol). All samplers were then freeze-dried for 24 hours in the original containers that were used for sample storage and transport. The disks were extracted three times by overnight (12 h) slow shaking at room temperature with 70 ml acetone each. The combined extracts were reduced by vacuum rotary evaporation to 10 mL. After removal of particles by filtration through

a layer of anhydrous Na<sub>2</sub>SO<sub>4</sub> the extract was further reduced in volume to approx. 1 mL. The acetone extract was transferred to methanol by addition of methanol (20 mL) and subsequent evaporation and a nitrogen flow to further reduce in volume to 2 mL. The extract was split to aliquots for different types of analysis.

### 1.1.3 Low-density polyethylene (LDPE) sheets

LDPE samplers from both parallel DPS sampling devices were extracted twice by soaking overnight with *n*-pentane (100 mL). Recovery standards (for PAHs, PCBs and OCPs) were added to the extraction jar during the first extraction. Recovery standards were d<sub>8</sub>-naphthalene; d<sub>10</sub>-biphenyl; d<sub>8</sub>-acenaphthene; d<sub>10</sub>-dibenzothiophene, d<sub>10</sub>-pyrene, d<sub>12</sub>-benz[a]anthracene, d<sub>12</sub>-perylene, PCB 30, PCB 53, and PCB 204. The sample volume was reduced to 2 mL by a gentle stream of nitrogen at room temperature. Extracts were first split into two equal fractions by volume. One fraction was stored as a reserve and the other received a general clean-up using gel permeation chromatography (GPC). This post-GPC sample was again split into two equal fractions by volume; the first of these was reduced in volume using nitrogen and analyzed for PAHs; the second received treatment with 2×1 mL concentrated sulfuric acid, was reduced in volume and analyzed for PCBs and OCPs. Details of the procedure are described in Allan et al. (2013).

## 1.2 Chemical analysis

SR sampler extracts were analyzed using GC-MS/MS (GC 7890 / MS-MS Triple Quadrupole 7000B (Agilent), equipped with HT8 SGE Analytical Science column for PCB 28, PCB 52, PCB 101, PCB 118, PCB 153, PCB 138, PCB 180, and OCPs: α-hexachlorocyclohexane (HCH), β-HCH, γ-HCH, δ-HCH, p,p'-DDE, p,p'-DDD, o,p'-DDT and p,p'-DDT, hexachlorobenzene (HCB), and pentachlorobenzene (PeCB). Polycyclic aromatic hydrocarbons were analyzed using GC 7890 / MS5975 (Agilent) equipped with J&W Scientific fused silica DB-5MS column. Detection was performed in single ion monitoring mode, the temperature of the ionic source was 320°C and quadrupole temperature 150°C. One μL sample was injected in splitless mode at 280°C. Helium (purity 5.5) was used as carrier gas at a flow of 1.5 mL min<sup>-1</sup>. The GC instrument was operated with an initial oven temperature of 80°C (1 min hold), then ramped at 15°C min<sup>-1</sup> to 180°C and at 5°C min<sup>-1</sup> to 310°C which was held for 20 min.

PBDEs were analyzed by GC equipped with 15m × 0.25 mm × 0.10 μm RTX-1614 column (Restek, USA). HRMS (AutoSpec Premier) was operated in EI+ mode at the resolution of >10 000.

SR and ED sampler extracts were analyzed for alkylphenols using dansyl chloride derivatization and LC-MS/MS detection according to Pernica et al. (2015). 4-nonylphenol served as an internal standard. Chromatographic separation was performed using LC Agilent 1200 Infinity Series, equipped with chromatographic column ACE 5 C18, 150 × 4.6 mm i.d., 5 μm particle size (ACE, Scotland, UK). Water containing 7 mmol L<sup>-1</sup> formic acid (A) and methanol (B) was used as a mobile phase. The isocratic elution of 10 % (A) and 90 % (B) was used at a flow rate of 0.5 mL min<sup>-1</sup>. The Agilent 6410 Triple Quadrupole (Agilent Technologies, Santa Clara, CA, USA) was used for MS/MS analysis. The instrument was operated in the ESI-positive MRM mode. Two MS/MS transitions were used for analyses.

Chromatographic analyses of alkyl phosphates and polycyclic musks were performed using GC 6890 (Agilent, USA) coupled to MSD 5975 mass spectrometer (Agilent, USA) operated in EI+ mode. Compounds were separated on the column HP-5MS (30m x 0.25mm x 0.25μm film) in selected ion mode (SIM). Helium was used as mobile phase at 1.2 mL min<sup>-1</sup> at constant pressure. One or two μL of extract were injected in pulsed split-less mode at 280 °C. GC temperature program started at 70 °C

(hold for 2 min), ramped 25 °C min<sup>-1</sup> to 150 °C (hold for 0 min), ramped 3 °C min<sup>-1</sup> to 200 °C (hold for 0 min), ramped 8 °C min<sup>-1</sup> to 280 °C (hold for 10 min).

## 1.3 Estimation of sampling rates

### 1.3.1 Silicone rubber samplers


As described in detail in Vrana et al. (submitted), concentrations of chemicals dissolved in water were calculated from amounts of analytes accumulated in SR samplers. Amounts of analytes absorbed by the samplers follow a first-order rate law to equilibrium. Aqueous concentrations were calculated from the mass absorbed by the samplers, the in situ sampling rate ( $R_s$ ) of the compounds and their sampler-water partition coefficients (Smedes et al., 2009) as described in Smedes and Booij (2012). Sampling rates were estimated from dissipation of PRCs from SR samplers during exposure using nonlinear least squares method by Booij and Smedes (2010), considering the fraction of individual PRCs that remained in the SR after the exposure as a continuous function of their sampler-water partition coefficient and their molecular mass, with adjustable parameter  $B$ .  $R_s$  for a compound accumulated under water-boundary layer control was calculated as a function of its molecular mass  $R_s = B \times M^{0.47}$  (Rusina et al., 2010).

### 1.3.2 Empore disc samplers

For ED samplers, calibration is based on levels of 4-nonylphenol and 10 PAHs that were detected in both ED and SR samplers throughout all sampled river stretches. Surface specific sampling rates appeared to be well correlated and their values were close to equal. The  $R_s$  for ED sampler was calculated from  $R_s$  of SR and their respective sampling areas using equation S1.

$$\frac{R_s(ED)}{A(ED)} = 0.83 \frac{R_s(SR)}{A(SR)} \quad \text{Equation S1}$$

Thus, the overall ED sampling rate was calculated from SR sampling rate by comparing the levels of PAHs and 4-nonylphenol in SR and ED samplers as the median of sampling rates of the individual chemicals used for the calibration throughout the river stretches (Fig. S; for more details see Vrana et al., submitted).



**Fig. S1 Comparison of levels of PAHs and 4-nonylphenol per sampling surface of SR and ED samplers from 8 mobile and 2 stationary DPS samples.**


### 1.3.3 Mutual comparison of two co-deployed DPS devices

The two co-deployed DPS devices may exhibit some differences if their pumps differed in volume flow rate. Their mutual comparison was done using a LDPE stripe mounted into each device during each exposure. Sampling performance of the LDPE samplers was assessed by monitoring release of 6 deuterated PAH PRCs during exposure. Since the exposure time  $t$  and mass  $m_x$  of LDPE stripes deployed in both DPS devices was the same ( $0.71 \pm 0.02$  g), from Equation S2 the ratio of sampling rates of a PRC in both devices is related to the ratio of its retained fractions after exposure:

$$\frac{R_{s,TOX}}{R_{s,CHEM}} = \frac{\ln(f_{PRC,TOX})}{\ln(f_{PRC,CHEM})} \quad \text{Equation S2}$$

where the subscripts 'TOX' and 'CHEM' denote the two co-deployed DPS devices containing samplers intended for toxicological and chemical analysis, respectively. The comparison is relevant only for PRCs for which  $f_{PRC}$  was quantifiable and lower than 80 % (Fig. S2).

Those PRCs included  $d_{10}$ -FLT and  $d_{12}$ -CHR.  $d_{10}$ -ACE,  $d_{10}$ -FLU, and  $d_{10}$ -PHE were released from the samplers to concentrations below their limit of quantification, whereas more than 80 % of  $d_{12}$ -BeP remained in the samplers. For  $d_{10}$ -FLT and  $d_{12}$ -CHR the calculated  $R_{s,TOX}/R_{s,CHEM}$  ratio in the eight sampling stretches was  $1.00 \pm 0.11$  and  $0.93 \pm 0.14$ , respectively.



**Fig. S2 Comparison of PRC fractions retained in LDPE sheets in two co-deployed DPS devices (TOX PS and CHEM PS) in parallel in 8 Danube stretches**



## 1.4 Bioanalysis

### AhR-mediated response (CAFLUX)

This type of toxicity was assessed using the rat hepatoma cell line H4G1.1c2 stably transfected by EGFP under control of dioxin response element. The assay was performed as described in Nagy et al. (2002). Cells were seeded to black clear-bottom 96 well plate in 100  $\mu$ L of DMEM supplemented with 10% fetal bovine serum (FBS) in density  $2 \times 10^5$  cells/mL. After 24h incubation at 37°C, dilution series of extracts were added in 100  $\mu$ L of the cultivation medium with DMSO as a solvent (final concentration 0.5% v/v). After 24 h incubation, the medium was replaced and fluorescence measured. 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD) served as the positive reference compound. The data were expressed as EC<sub>20</sub> using a log-logistic dose-response model. Cytotoxic concentrations of extracts were excluded from the calculation based on viability data assessed with neutral red uptake assay (Freyberger and Schmuck, 2005).

### PXR-mediated response (HG5LN-hPXR)

The HG5LN-hPXR assay based on human HeLa cell line served for the assessment of the activation of the human pregnane X receptor (PXR) and was performed according to Creusot et al. (2010) with minor modifications. Briefly, HG5LN-hPXR cells in DMEM without phenol red supplemented with 5% fetal calf serum (FCS) were seeded in a white 96-well plate in 100  $\mu$ L of cell suspension at a density of  $6 \times 10^5$  cells/mL and incubated for 24 h at 37°C. The sampler extracts diluted in phenol red-free DMEM supplemented with 3% dextran-coated charcoal-treated fetal calf serum (DCC-FCS) were added and the exposed cells were incubated for a further 16 h. Before luminescence measurement, exposure medium was replaced with 50  $\mu$ L of medium supplemented with 0.3 mM luciferin. SR 12813 (Tetraethyl 2-(3,5-di-tert-butyl-4-hydroxyphenyl)ethenyl-1,1-bisphosphonate) was used as the positive reference compound. The data was expressed as EC<sub>20</sub> using a log-logistic dose-response model.

### (Anti-)androgenicity (MDA-kb2)

Human mammary carcinoma-derived cell line MDA-kb2 was used for the assessment of androgen receptor-mediated responses (androgenicity and anti-androgenicity). The cells were grown in Leibovitz's L-15 medium supplemented with 10 % FBS at 37 °C and normal air with high relative humidity. The experiment was performed in 96-well plates in L-15 medium supplemented with 10% dextran-charcoal-treated FBS. Androgen dihydrotestosterone (DHT) was used as a positive reference compound. For the assessment of androgenic effect, the cells were exposed to samples alone. To assess the anti-androgenic effect, cells were exposed to extracts in combination with the physiological ligand of the AR DHT (0.1 nM). The anti-androgenic effect was quantified using flutamide as model anti-androgen. The androgenic and anti-androgenic potential was expressed as EC<sub>20</sub> and IC<sub>20</sub>, respectively, using a log-logistic dose-response model. The viability of the cells in the assay was evaluated using neutral red uptake assay (Freyberger and Schmuck, 2005) and cytotoxic concentrations of the samples were excluded from data evaluation.

### Estrogenicity (MELN)

The MELN assay, based on human cell line MCF-7 transfected with the luciferase gene under control of estrogenic response element, was used for the assessment of estrogenicity according to the previously described procedure (Kinani et al., 2010). Briefly, 100  $\mu$ L of MELN cells suspension with a density of  $8 \times 10^4$  cells/mL was seeded to a white 96-well plate in phenol red-free DMEM supplemented

with 3% DCC-FCS. After plating at 37°C for 24 h, the cells were exposed to the sampler extracts and incubated for a further 16 h. Before luminescence measurement, exposure medium was replaced with 50 µL of medium supplemented with 0.3 mM luciferin. 17β-estradiol was used as the positive reference compound. The data were expressed as EC<sub>20</sub> using a log-logistic dose-response model. Cell viability was assessed using the MTT assay.

#### ARE-mediated oxidative stress response (AREc32)

The MCF7 human breast carcinoma cell line-derived AREc32 assay was applied to assess the activation of the Nrf2 pathway (Wang et al., 2006), with the assay conducted according to Escher et al. (2012). The sampler extracts were blown to dryness, then re-suspended in DMEM with GlutaMAX™ with 10% FBS and serially diluted tert-Butylhydroquinone (tBHQ) was used as the positive reference compound. Cells were exposed in 96-well plates for 24 hours and then luminescence was detected using Luciferase Assay System (Promega E1500) according to the Promega protocol. Using a linear concentration-effect curve, the data was expressed as EC<sub>IR1.5</sub> in units of REF, while the effect on cell viability detected with the MTS assay (Mosmann, 1983) expressed as EC<sub>10</sub> was determined from a log-logistic concentration-effect curve. EC<sub>IR1.5</sub> was calculated using equation S3.

$$EC_{IR1.5} = \frac{0.5}{slope} \quad \text{Equation S1}$$

#### p53-mediated response (p53RE-bla)

The CellSensor p53RE-bla HCT-116 assay (Life Technology, Australia) based on human colorectal carcinoma cell line HCT-116 was used to assess activation of p53 and was conducted according to Neale et al. (2015b). Briefly, the extracts were blown to dryness, then re-suspended in Opti-MEM with 0.5% dialyzed FBS and serially diluted. Eight microliters of the serially diluted extract were added to 32 µL of cells seeded at a density of 9.4×10<sup>5</sup> cells/mL in the black clear bottom 384-well plate. After incubating for 40 h at 37°C, 8 µL of FRET reagent with resazurin was added to each well and fluorescence was measured after incubating for 2.5 h at room temperature. Mitomycin was used as the positive reference compound with EC<sub>IR1.5</sub> derived from a linear concentration-effect curve and cell viability EC<sub>10</sub> derived from a log-logistic concentration-effect curve.

#### NF-κB-mediated response (NF-κB-bla)

The NF-κB-bla assay (Life Technology, Australia) based on the human monocyte THP-1 cell line, was used to assess the activation of NF-κB, which is indicative of inflammation response. The assay was performed as in Jin et al. (2015). The extracts were blown to dryness, then re-suspended in RPMI 1640 media with 10% dialyzed FBS and serially diluted. Eight microliters of the serially diluted extract were added to 32 µL of cells seeded at a density of 6.3×10<sup>5</sup> cells/mL in the black clear bottom 384-well plate. After incubating for 24 h at 37°C, 8 µL of FRET reagent with resazurin was added to each well and fluorescence was measured after incubating for 2 h at room temperature. Tumor Necrosis Factor-alpha (TNFα) was used as the positive reference compound. EC<sub>IR1.5</sub> values were derived from the linear concentration-effect curve, while the cell viability EC<sub>10</sub> was derived from the log-logistic concentration-effect curve.



**Fig. S1 Map of the sampling cruise with the stretches and stationary exposure site (red dot); blue dots indicate active LVSP sampling sites from Neale et al. (2015a).**

Analyzed chemicals: 267					
Detected chemicals not in ToxCast: 91					
Detected chemicals in ToxCast: 117					
<b>Inactive:</b>	<b>AhR:</b>	92	<b>Active:</b>	<b>AhR:</b>	22 (19 %)
	<b>ER:</b>	76		<b>ER:</b>	38 (33 %)
	<b>aAR:</b>	100		<b>aAR:</b>	16 (14 %)
	<b>ARE:</b>	73		<b>ARE:</b>	39 (35 %)
	<b>p53:</b>	102		<b>p53:</b>	12 (11 %)
<b>Inactive:</b>	<b>AhR:</b>	7478	<b>Active:</b>	<b>AhR:</b>	828 (10%)
	<b>ER:</b>	6948		<b>ER:</b>	358 (16 %)
	<b>aAR:</b>	7432		<b>aAR:</b>	874 (11 %)
	<b>ARE:</b>	5826		<b>ARE:</b>	696 (23 %)
	<b>p53:</b>	7461		<b>p53:</b>	845 (10 %)
ToxCast: 9076					

**Fig. S2 Overview of detected chemicals described in ToxCast database to be in/-active in eliciting the indicated effects.**





Fig. S3 Percentage of the bioanalytical equivalents (BEQ<sub>bio</sub>) explained by the detected chemicals (BEQ<sub>chem</sub>) in samples from a dynamic passive sampling of the Danube river; SR silicone rubber samples, ED Empore discs samples; S1, 2 – stationary sampling site; 1-8 sampled

river stretches; for ARE in ED extracts, 18 main identified drivers of the effect are given (total 42 quantified active chemicals are provided in Tab. S4)



**Fig. S4** p53-mediated mitomycin bioanalytical equivalents calculated from detected chemicals levels ( $BEQ_{chem}$ ) in samples from a dynamic passive sampling of the Danube river; SR silicone rubber samples, ED Empore discs samples; S1, 2 – stationary sampling site; 1-8 sampled river stretches

**Tab. S1** River stretches and stationary exposure localities sampled with passive samplers

Stretch/ sample number	Stretch start and end/stationary exposure site	River km	Sampling time span	Mean water temperature [°C]	Exposure time [d]	Parameter AB <sup>b</sup>	Water volume sampled by SR [L] <sup>a</sup>	Water volume sampled by ED [L] <sup>b</sup>
S1	Stationary exposure 1	1,852	19.8.-23.8.	21.3	4	904	245	90
S2	Stationary exposure 2	1,852	23.8.-28.8.	21.3	5	772.6	264	97
1	Passau–Bratislava	2,203-1,852	17.8.-22.8.	21.3	2	1208.8	236	62
2	Bratislava–Budapest	1,852-1,632	22.8.-26.8.	22	1.2	1055	98	31
3	Budapest–Vukovar	1,648-1,297	26.8.-2.9.	21.9	1.7	1179	151	51
4	Vukovar–Belgrade	1,297-1,154	2.9.-6.9.	22.8	1.6	1223.8	117	49
5	Belgrade–Turnu- Severin	1,154-930	6.9.-10.9.	22.1	2	1041.2	112	51
6	Turnu-Severin–Ruse	930-495	11.9.-17.9.	21.9	2	945.1	122	47
7	Ruse–Braila	495-170	17.9.-21.9.	19.2	1.4	821.2	90	29
8	Braila–Tulcea	170-71	21.9.-26.9.	18.7	1.3	791.2	73	26

<sup>a</sup> Volume of water extracted by the SR sampler during exposure; volume is calculated from PRCs levels for a model compound with a molecular mass of 300. <sup>b</sup> Volume of water extracted by ED sampler during sampling; volume estimate based on a comparison of levels of 10 PAHs in SR and samplers as described in 1.3.2.

<sup>b</sup> Exposure specific parameter for calculation of  $R_s$  in equations 1; for further details see Vrana et al.(submitted)

**Tab. S2 Concentrations [pg/L] of chemicals detected in SR sampler extracts; PAHs - polycyclic aromatic hydrocarbons; BDE - brominate diphenyl ethers**

Analyte [pg/L]	CAS Number	Stationary exposure		River stretch								
		S1	S2	1	2	3	4	5	6	7	8	
PAHs	Naphthalene	91-20-3	3,418	4,267	3,260	7,088	8,306	5,157	2,667	2,805	4,968	6,298
	Biphenyl	92-52-4	485	1,264	423	504	4,440	1,236	688	549	616	740
	Acenaphthylene	208-96-8	1,559	1,792	2,712	1,239	2,045	3,069	1,231	315	877	1,603
	Acenaphthene	83-32-9	770	1,173	838	619	11,283	1,942	835	286	279	639
	Fluorene	86-73-7	954	1,349	1,200	1,195	8,728	4,834	1,468	914	930	1,468
	Phenanthrene	85-01-8	1,917	2,800	3,086	2,441	5,527	14,755	2,717	2,178	1,975	3,502
	Anthracene	120-12-7	185	206	346	307	571	388	222	144	158	267
	Fluoranthene	206-44-0	3,527	4,440	2,672	3,516	4,659	2,464	2,654	2,371	1,463	2,386
	Pyrene	129-00-0	1,767	2,399	1,609	2,151	3,292	2,192	1,847	1,244	1,470	2,231
	Retene	483-65-8	996	1,155	617	599	358	243	258	119	137	187
	Benzo( <i>b</i> )fluorene	243-17-4	223	251	192	181	267	154	170	97	105	163
	Benzonaphthothiophene	205-43-6	82	130	132	71	89	68	61	35	64	83
	Benzo( <i>ghi</i> )fluoranthene	203-12-3	144	176	112	117	123	77	105	93	89	119
	Cyclopenta( <i>cd</i> )pyrene	27208-37-3	146	188	135	151	233	90	106	69	81	130
	Benzo( <i>a</i> )anthracene	56-55-3	146	188	<6	<6	234	91	106	69	81	131
	Triphenylene	217-59-4	360	442	403	403	371	295	402	384	392	425
	Chrysene	218-01-9	266	375	283	238	307	140	187	136	149	202
	Benzo( <i>b</i> )fluoranthene	205-99-2	158	129	87	109	106	41	79	100	51	63
	Benzo( <i>j</i> )fluoranthene	205-82-3	64	79	17	33	41	15	24	39	<15	<15
	Benzo( <i>k</i> )fluoranthene	207-08-9	34	27	<23	<23	<23	<23	<23	<23	<23	<23
	Benzo( <i>e</i> )pyrene	192-97-2	120	146	92	103	88	44	79	55	79	65
	Benzo( <i>a</i> )pyrene	50-32-8	<46	<46	<46	<46	<46	<46	70	<46	<46	<46
	Perylene	198-55-0	82	<66	<66	<66	<66	<66	<66	<66	<66	<66
	Indeno( <i>123cd</i> )pyrene	193-39-5	<21	<21	<21	<21	<21	<21	<21	<21	<21	<21
	Dibenzo( <i>ah</i> )anthracene	53-70-3	<13	<13	<13	<13	<13	<13	<13	<13	<13	<13

	Dibenzo( <i>ac</i> )anthracene	215-58-7	<18	<18	<18	<18	<18	<18	<18	<18	<18	<18
	Benzo( <i>ghi</i> )perylene	191-24-2	21	25	<17	<17	<17	<17	<17	<17	<17	<17
	Anthanthrene	191-26-4	<49	<49	<49	<49	<49	<49	<49	<49	<49	<49
	Coronene	191-07-1	<62	<62	<62	<62	<62	<62	<62	<62	<62	<62
	ΣPAHs 16 US EPA		14,721	19,169	16,092	18,902	45,058	35,073	14,083	10,562	12,402	18,790
	ΣPAHs		17,342	23,000	18,215	21,063	51,067	37,296	15,975	12,002	13,965	20,704
OCs	PCB 28	7012-37-5	49	56	39	91	64	44	190	80	173	141
	PCB 52	35693-99-3	82	94	69	121	59	53	121	16	102	98
	PCB 101	37680-73-2	21	23	16	20	11	22	17	3.3	6.7	24
	PCB 118	31508-00-6	4.2	4.5	3.0	3.9	2.3	2.7	3.5	4.0	2.5	2.9
	PCB 153	35065-27-1	31	33	22	28	16	25	19	30	12	14
	PCB 138	35065-28-2	22	24	16	21	12	19	14	19	8.5	10
	PCB 180	35065-29-3	7.8	8.2	5.5	6.9	3.3	5.0	4.0	5.1	2.6	4.6
	Σ PCB		217	244	171	291	167	172	369	158	307	295
	PeCB	608-93-5	16	20	14	22	96	84	56	23	31	29
	HCB	118-74-1	78	90	62	81	75	41	54	48	97	97
	a-HCH	319-84-6	5.2	6.0	3.7	2.2	3.7	1.6	2.2	7.4	5.0	12.2
	b-HCH	319-85-7	3.6	3.2	8.9	10	9.1	12	14	101	146	259
	d-HCH	58-89-9	2.7	2.4	7.1	9.2	7.6	10	12	84	132	235
	o,p'-DDE	3424-82-6	1.8	1.0	0.9	2.5	2.6	2.8	3.7	3.3	6.6	24.2
	p,p'-DDE	72-55-9	34	33	18	58	54	54	72	71	165	512
	o,p'-DDD	53-19-0	16	22	13	25	49	51	73	54	88	334
	p,p'-DDD	72-54-8	22	39	21	30	60	49	66	69	125	313
	o,p'-DDT	789-02-6	2.3	1.7	1.9	1.4	5.2	2.8	2.2	1.2	2.9	17.9
	p,p'-DDT	50-29-3	3.8	3.7	3.3	1.3	4.6	2.7	2.4	2.0	5.5	21.2
	Σ DDT		83	103	61	119	181	165	220	202	396	1,240
Σ OCPs		188	223	156	244	372	313	358	465	808	1,872	
PBDEs	BDE 28	41318-75-6	0.8	0.9	0.5	0.7	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4
	BDE 47	5436-43-1	0.8	0.7	6.8	<1.3	<1.3	<1.3	<1.3	<1.3	<1.3	<1.3
	BDE 66	189084-61-5	<0.03	0.04	0.3	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1



	BDE 100	189084-64-8	0.07	0.12	0.7	0.8	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
	BDE 99	60348-60-9	0.12	0.31	2.0	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
	BDE 85	182346-21-0	0.04	0.03	0.4	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
	BDE 154	207122-15-4	0.22	0.42	1.2	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6
	BDE 153	68631-49-2	<0.09	0.21	0.5	1.0	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3
	BDE 183	207122-16-5	1.84	3.30	10	<4.3	<4.3	<4.3	<4.3	<4.3	<4.3	<4.3
	Σ WFD BDEs		1.99	2.61	11.6	2.5	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Σ BDEs		3.86	5.98	22.2	2.5	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Alkylphenols	Bisphenol A	80-05-7	674	644	338	309	370	494	446	<0.2	251	310
	4-t-octylphenol	140-66-9	93	107	184	145	198	362	138	72	137	85
	4-nonylphenol	104-40-5	2,165	2,775	5,226	5,816	7,391	9,156	7,928	3,453	6,890	5,841
	Σ Alkylphenols		2,932	3,526	5,749	6,270	7,959	10,012	8,511	3,525	7,278	6,236
Musks	Cashmeran	33704-61-9	1,526	1,991	658	977	663	434	<341	<341	<341	<341
	Ceolestolide	13171-00-1	40	62	<34	<34	<34	54	<34	<34	<34	<34
	Phantolide	15323-35-0	<31	<31	<31	<31	<31	<31	<31	<31	<31	<31
	Traseolide	68140-48-7	87	129	68	70	75	81	74	39	69	<21
	Galaxolide	1222-05-5	17,021	25,063	10,104	10,848	16,722	15,831	22,312	9,183	12,096	8,864
	Tonalide	1506-02-1	1,473	2,648	1,069	976	1,521	2,003	3,008	837	1,307	868
	Σ Musks		18,582	27,839	11,240	11,894	18,317	17,915	25,395	10,060	13,471	9,732
NBFRs	TBECH	3322-93-8	0.941	1.488	1.275	0.756	0.807	0.604	1.243	0.323	0.225	0.230
	p-TBX	23488-38-2	0.018	0.012	0.030	0.039	0.056	0.015	<0.007	<0.007	<0.012	0.056
	BATE	3728-89-5	0.655	0.272	0.418	0.242	0.054	0.083	0.163	0.035	0.054	0.100
	TBCO	3194-57-8	<0.021	<0.019	<0.03	<0.061	<0.037	<0.039	<0.037	<0.04	<0.065	<0.072
	PBBZ	608-90-2	0.060	0.030	0.325	0.083	0.097	0.065	0.056	0.066	0.078	0.156
	TBCT	39569-21-6	0.006	<0.005	0.017	<0.016	<0.01	<0.01	<0.01	<0.011	<0.017	0.110
	PBT	87-83-2	0.090	0.071	0.127	0.136	1.299	0.364	0.184	0.170	0.381	0.565
	PBEB	85-22-3	0.04	<0.0016	<0.0024	<0.0049	0.1	<0.0031	<0.0029	<0.0032	<0.0052	0.2
	DPTE	35109-60-5	3.73	8.61	9.4	2.0	<0.17	<0.18	<0.17	<0.18	<0.3	<0.33
	HBB	87-82-1	0.14	0.13	0.19	<0.028	0.25	0.30	<0.017	<0.018	0.52	0.67
	PBBA	59947-55-1	<0.037	<0.034	<0.053	<0.107	<0.064	<0.067	<0.064	<0.069	<0.113	<0.125

	HCDBCO	51936-55-1	<0.023	<0.021	<0.033	<0.067	<0.041	<0.042	<0.04	<0.044	<0.071	<0.079
	BTBPE	37853-59-1	0.26	0.12	0.28	<0.03	<0.018	<0.019	<0.018	<0.019	<0.032	<0.035
	s-DP	13560-89-9	1.08	1.49	9.0	15.2	10.8	6.3	13.0	17.9	2.4	5.3
	a-DP	13560-89-9	0.47	0.80	2.5	6.3	2.0	1.6	2.2	3.2	1.8	4.4
	Σ NBFRRs		7.49	13.02	23.5	24.7	15.4	9.3	16.8	21.6	5.4	11.8
Alkyl phosphates	Triisobutyl phosphate	126-71-6	57,464	81,680	38,251	59,746	52,243	43,502	57,037	94,150	97,702	101,181
	Tributyl phosphate	126-73-8	5,067	8,584	3,289	15,329	4,591	4,048	6,574	8,745	6,950	9,694
	Tris(2-Chloroethyl)phosphate	115-96-8	<396	<360	<420	<418	<390	<397	<390	<433	<380	<392
	Tris(1-Chloro-2-propyl)phosphate	13674-84-5	56,811	56,211	31,360	35,714	37,372	22,414	31,220	37,939	28,631	32,109
	Dibutyl phenyl phosphate	2528-36-1	254	311	326	151	141	94	146	108	114	123
	Butyl diphenyl phosphate	2752-95-6	<27	110	92	<77	<46	<48	<46	<50	<81	<90
	Tris(1,3-Dichloro-2-propyl)phosphate	13674-87-8	7,209	13,399	6,752	3,738	5,814	2,700	2,803	3,585	4,325	3,729
	Tris(2-butoxyethyl)phosphate	78-51-3	14,211	23,996	17,628	6,554	10,276	5,738	6,299	2,447	1,678	1,604
	Tris(2-ethylhexyl)phosphate	78-42-2	<113	<105	<164	<329	<199	<207	<199	<214	<349	<386
	Triphenyl phosphate	115-86-6	811	2,208	913	344	567	427	275	181	255	202
	2-Ethylhexyldiphenyl phosphate	1241-94-7	45	142	108	52	82	60	65	<8	<13	<15
	Tri-o-tolyl phosphate	78-30-8	17	<4	20	57	14	31	15	67	27	37
	Tri-m-tolyl phosphate	228-312-4	34	83	63	47	29	20	12	11	<9	17
	Tri-p-tolyl phosphate	1038-95-5	<4	<4	15	32	12	11	<7	<7	<12	<13
	Σ Alkylphosphates		141,923	186,725	98,815	121,763	111,142	79,046	104,446	147,234	139,681	148,697

**Tab. S3 Concentrations [µg/L] of chemicals detected in ED sampler extracts**

	Analyte	CAS Number	Stationary exposure		River stretch							
			S1	S2	1	2	3	4	5	6	7	8
Pharmaceuticals	Ampicillin	69-52-3	89	< 60	< 100	< 190	136	172	216	< 120	238	< 260
ATBs	Azithromycin	104491-80-7	< 20	< 20	< 30	< 60	< 40	< 40	< 40	< 40	< 70	< 70
	Ciprofloxacin	85721-33-1	21	152	26	46	53	92	88	143	128	< 30
	Clarithromycin	116836-41-0	< 160	< 170	< 240	< 520	< 390	< 310	< 310	< 400	< 590	< 570
	Clindamycine	18323-44-9	737	1,249	776	1,855	1,375	983	1,492	1,414	1,139	914
	Difloxacin	91296-86-5	< 10	< 10	< 20	< 40	< 20	< 20	< 30	< 20	< 40	< 50
	Enoxacin	206873-63-4	103	24	42	75	41	66	< 30	116	69	107
	Enrofloxacin	112732-17-9	< 10	< 10	< 20	< 40	< 30	< 30	< 30	< 30	< 50	< 50
	Erythromycin	114-07-8	< 150	< 150	< 230	< 490	< 370	< 290	< 290	< 380	< 550	< 530
	Flumequine	143984-63-8	22.33	8.26	< 10	< 40	< 20	< 20	< 20	< 30	< 50	64.8
	Levofloxacin	100986-85-4	< 10	< 10	< 20	< 40	< 20	< 20	< 30	< 20	< 40	< 50
	Lomefloxacin	98079-51-7	< 10	< 10	< 20	< 40	< 30	< 30	< 30	< 30	< 50	< 50
	Norfloxacin	68077-27-0	< 10	< 10	< 10	< 30	< 20	< 20	27	< 20	< 30	< 30
	Oxolinic acid	14698-29-4	44	61	18	114	49	20	51	127	148	103
	Oxytetracycline	79-57-2	< 130	< 130	< 190	< 360	< 240	< 250	< 260	< 230	< 450	< 460
	Penicillin_V	132-98-9	< 20	< 10	< 20	< 50	< 30	< 30	< 30	< 40	< 70	< 60
	Roxithromycin	80214-83-1	< 90	< 90	< 130	< 240	< 140	< 140	< 150	< 170	< 280	< 280
	Sulfadiazine	116-44-9	402	754	485	749	1,061	512	844	< 360	828	< 760
	Sulfadimethoxine	1037-50-9	33	20	< 10	36	26	25	55	55	< 50	80
	Sulfamerazine	127-79-7	279	320	< 360	< 680	963	< 430	844	< 440	< 760	< 910
	Sulfamethazine	1981-58-4	458	651	840	3,060	3,909	2,090	2,611	2,553	18,326	12,078
	Sulfamethizole	144-82-1	301	217	533	< 420	< 240	328	393	717	< 480	1,448
	Sulfamethoxazole	129378-89-8	922	539	1,297	2,015	1,190	1,131	1,637	1,386	2,454	2,542
	Sulfamethoxypridazine	80-35-3	580	< 200	517	684	727	< 390	1,748	464	< 690	2,324
	Sulfamoxole	729-99-7	458	372	< 290	1,725	884	410	805	443	2,071	1,067

	Sulfaphenazole	526-08-9	15	26	13	52	26	51	24	< 30	< 50	50
	Sulfapyridine	144-83-2	8,059	7,796	6,465	12,304	12,572	11,453	10,780	5,127	7,593	8,154
	Sulfaquinoxaline	59-40-5	23	30	32	49	65	< 20	47	< 20	159	< 40
	Sulfasalazine	599-79-1	123	13	60	182	51	59	35	112	41	331
	Sulfathiazole	158269-46-6	< 250	< 200	< 320	< 620	< 350	< 390	< 390	< 400	< 690	< 840
	Trimethoprim	738-70-5	< 250	< 200	372	977	< 350	< 390	< 390	< 400	< 690	< 840
Cardiovascular	Amiodarone	1951-25-3	< 150	< 190	< 240	< 490	< 390	< 330	< 310	< 420	< 660	< 720
	Atenolol	29122-68-7	< 180	217	< 240	846	< 260	< 290	353	< 300	< 520	< 610
	Bisoprolol	66722-44-9	357	341	< 230	< 420	< 260	< 270	< 270	< 270	< 480	< 570
	Cilazapril	88768-40-5	< 10	< 10	< 10	< 20	< 10	< 10	< 10	< 20	< 30	< 30
	Diltiazem	144604-00-2	246	258	259	< 460	432	287	353	< 360	725	610
	Dipyridamole	58-32-2	301	465	< 370	< 780	< 570	< 470	609	738	< 860	< 840
	Disopyramide	3737-09-5	< 10	< 10	< 20	< 30	< 20	< 20	< 20	< 20	< 40	< 40
	Eprosartan	133040-01-4	51	145	115	260	139	84	55	63	207	324
	Flecainide	1403764-72-6	< 10	< 10	< 10	< 30	< 20	< 20	< 20	< 20	< 30	< 30
	Irbesartan	138402-11-6	10,002	10,656	9,907	14,583	16,442	8,216	7,952	4,241	6,695	4,839
	Metoprolol	13484-40-7	1,161	558	291	1,172	1,355	1,106	942	< 270	897	3,200
	Sotalol	27948-47-6	< 180	< 140	< 240	< 460	< 280	< 310	< 290	< 300	< 520	< 610
	Telmisartan	144701-48-4	1,708	2,241	1,018	2,669	2,986	1,578	1,335	2,152	1,967	1,943
	Valsartan	137862-53-4	190	85	158	163	295	96	130	68	273	110
	Verapamil	152-11-4	167	< 120	< 190	< 420	< 290	< 250	< 240	< 320	< 450	< 420
Psychoactive	Alprazolam	28981-97-7	647	743	323	1,660	1,709	1,045	1,178	2,110	2,830	1,753
	Amitriptyline	337376-15-5	< 130	176	< 210	< 460	334	< 270	< 270	< 340	< 480	< 500
	Bupropion	31677-93-7	659	475	372	< 420	334	< 270	< 260	< 270	< 480	< 570
	Carbamazepine	298-46-4	13,211	11,300	11,017	18,622	16,482	10,950	16,263	17,561	23,028	20,456
	Citalopram	128196-01-0	< 10	18	13	39	26	< 20	< 20	34	48	< 30
	Clonazepam	106955-87-7	15	11	53	117	22	47	47	76	66	110
	Codeine	70982-46-6	357	237	404	911	< 260	881	275	< 270	725	< 570
	Donepezil	120011-70-3	17	< 10	< 10	26	22	< 10	< 10	25	< 30	< 30
	Fluoxetine	100568-03-4	< 50	< 50	< 70	< 120	< 80	< 80	< 80	< 90	< 130	< 150

	Flupentixol	2709-56-0	27	27	37	78	114	< 30	51	91	197	91
	Fluphenazine	5002-47-1	268	114	66	133	< 60	82	334	165	190	278
	Haloperidol	337376-15-5	10	< 10	11	< 30	< 20	< 20	< 20	< 20	< 40	< 40
	Chlorprothixene	113-59-7	56	58	226	< 130	165	88	161	97	< 150	236
	Levomepromazine	60-99-1	< 100	< 100	< 150	< 320	< 240	195	< 190	< 250	< 350	< 340
	Maprotiline	10262-69-8	< 90	134	< 110	299	275	< 140	< 140	274	266	331
	Meclozine	1104-22-9	< 20	< 20	36	< 50	65	< 40	37	219	166	152
	Memantine	19982-08-2	13	8	< 0	< 10	12	< 10	12	< 10	< 20	< 20
	Mianserin	21535-47-7	< 130	< 130	259	< 420	< 310	< 250	< 260	< 340	< 480	< 460
	Mirtazapine	61337-67-5	< 10	< 10	< 20	< 30	< 20	< 20	< 20	< 20	< 40	< 40
	Nefazodone	337376-15-5	< 70	< 60	< 90	< 170	< 110	< 110	< 110	< 120	< 180	< 200
	Oxazepam	35295-88-6	9,310	7,641	4,477	6,543	6,306	4,016	5,144	3,966	5,142	5,144
	Paroxetine	110429-35-1	32	8	< 10	55	29	20	59	89	114	95
	Perphenazine	58-39-9	268	165	323	< 170	< 100	225	161	< 120	< 200	< 200
	Risperidone	106266-06-2	< 20	< 20	< 20	< 40	< 30	< 30	< 30	< 30	< 50	< 50
	Ropinirole	337376-15-5	< 160	< 120	< 210	< 390	< 240	< 250	< 260	< 250	< 450	< 530
	Sertaline	79617-96-2	290	77	91	192	171	328	120	< 110	656	< 180
	Tetracycline	60-54-8	< 190	< 190	< 260	< 490	< 310	< 350	< 330	< 320	< 590	< 610
	Tramadol	123154-38-1	179	176	115	215	161	107	118	103	124	103
	Trihexyphenidyl	144-11-6	< 80	< 70	< 100	< 190	< 120	< 120	< 130	< 140	< 200	< 230
	Venlafaxine	93413-44-6	80	53	37	65	65	35	41	44	45	< 30
Antihistamins	Azelastine	58581-89-8	< 70	< 60	< 80	< 160	< 100	< 100	< 110	< 110	< 170	< 190
	Clemastine	14976-57-9	< 130	< 120	< 180	< 360	< 200	< 200	< 220	< 250	< 410	< 420
	Cyproheptadine	129-03-3	123	< 110	< 180	< 360	< 280	348	242	< 270	583	< 380
	Desloratadine	100643-71-8	< 10	< 10	< 10	< 30	< 20	< 20	< 20	< 20	< 30	< 30
	Diphenhydramine	147-24-0	9	< 0	< 0	< 10	< 10	< 10	< 10	< 10	< 20	< 20
	Hydroxyzine	10246-75-0	< 70	91	116	< 170	112	< 110	< 120	< 120	< 180	< 210
	Loperamide	34552-83-5	< 120	< 120	< 180	< 390	< 290	< 230	< 240	< 300	< 450	< 420
	Orphenadrine	337376-15-5	123	< 100	158	< 330	< 240	< 190	< 200	< 250	< 350	< 350
Antifungals	Clotrimazole	117829-71-7	3,014	3,304	3,249	5,859	7,190	5,368	9,130	7,595	9,146	5,486

	Econazole	1069-66-5	47	30	44	62	59	< 30	< 30	< 30	93	213
	Fluconazole	123631-92-5	324	310	242	423	413	389	452	401	725	648
	Itraconazole	84625-61-6	< 70	< 90	< 110	< 220	< 180	< 160	< 150	< 200	< 310	< 340
	Ketoconazole	142128-59-4	625	1,136	485	1,823	373	2,623	746	2,131	< 480	2,858
	Miconazole	22832-87-7	50	68	57	124	47	59	71	196	466	91
	Sulconazole	61318-90-9	39	51	45	< 50	65	37	161	143	152	< 80
	Terbutaline	23031-25-6	< 170	< 130	< 230	< 420	< 260	< 270	< 270	< 270	< 480	< 570
Antidiabetics	Bromocriptine	22260-51-1	6.70	4.13	< 0	42.3	11.8	34.8	19.6	21.1	24.2	26.7
	Glibenclamide	10238-21-8	25	< 20	32	< 50	86	< 30	< 30	< 40	72	< 70
	Glimepiride	29094-61-9	< 30	< 30	< 40	< 90	< 70	< 60	< 60	< 80	< 120	< 130
	Repaglinide	135062-02-1	< 110	< 100	< 160	< 290	< 170	< 180	< 180	< 200	< 340	< 340
Statins	Atorvastatin	110862-48-1	56	< 40	< 60	< 110	< 70	< 70	< 70	84	< 120	< 130
	Bezafibrate	41859-67-0	37	102	39	49	53	156	310	95	211	217
	Fenofibrate	49562-28-9	99	207	226	273	1,179	820	864	186	380	< 220
	Rosuvastatin	1094100-06-7	97	9	21	78	28	55	45	53	< 20	290
Cancer treatment	Flutamide	13311-84-7	< 90	< 110	< 150	< 290	< 240	< 200	< 190	< 250	< 410	< 420
	Fulvestrant	129453-61-8	< 40	< 50	< 60	< 130	< 100	< 90	< 80	< 110	< 180	< 190
	Tamoxifen	10540-29-1	< 10	< 20	< 20	< 50	< 40	< 30	< 30	< 40	< 70	< 70
NSAID	Diclofenac	15307-86-5	< 30	< 40	< 50	< 110	< 90	< 70	< 70	< 90	< 150	< 160
Diuretic	Furosemide	106391-48-4	< 50	< 30	< 50	< 140	122	< 100	< 90	< 110	< 200	< 200
Other	Alfuzosin	337376-15-5	< 160	< 120	< 210	< 420	< 240	< 270	< 260	< 270	< 450	< 530
	Naloxone	465-65-6	290	248	< 230	< 460	< 260	533	< 270	295	< 520	< 610
	Ranitidine	66357-35-5	< 160	< 120	< 210	< 390	< 240	< 250	< 260	< 250	< 450	< 530
	Σ Pharmaceuticals		57,256	54,175	45,799	81,306	80,803	57,203	69,470	55,973	89,406	78,869
CUPs	Acetochlor	34256-82-1	11	45	< 3.2	< 6.5	57	42	138	498	592	965
	Alachlor	15972-60-8	21	45	< 16.3	< 66.1	< 19.8	< 20.7	29	< 42.6	101	56
	Atrazine	1912-24-9	3,025	4,339	7,349	4,703	6,995	3,400	4,817	4,767	18,295	8,482
	Azinphos methyl	86-50-0	< 2.2	< 2.1	< 3.2	< 6.5	< 3.9	< 4.1	< 3.9	< 4.2	< 6.9	< 7.6
	Carbaryl	63-25-2	< 1.1	< 1	< 1.6	< 3.3	< 2	< 2	< 2	< 2.1	< 3.5	< 3.8
	Carbendazim	10605-21-7	11	15	20	12	25	17	25	22	82	40

Chlorpyrifos	2921-88-2	179	257	289	117	351	188	415	329	1,482	608
Chlorsulfuron	64902-72-3	< 2.2	< 2.1	< 3.2	< 6.5	< 3.9	< 4.1	< 3.9	< 4.2	14	< 7.6
Chlortoluron	15545-48-9	1,815	1,717	3,463	2,118	2,439	603	687	673	1,992	780
Clopyralid	1702-17-6	< 22.2	< 20.8	< 32.5	< 66.1	< 39.5	< 41.2	< 39.7	< 42.6	< 69.7	< 76.2
Diazinon	333-41-5	141	165	116	144	245	299	677	608	1,415	1,363
Dimethachlor	50563-36-5	22	242	36	44	130	73	120	20	116	91
Dimethoate	60-51-5	24	33	77	32	34	18	37	24	92	43
Disulfoton	298-04-4	< 1.1	< 1	< 1.6	< 3.3	< 2	< 2	< 2	< 2.1	< 3.5	< 3.8
Diuron	330-54-1	1,228	1,825	2,409	1,238	1,770	804	1,266	1,079	3,533	1,504
Fenitrothion	122-14-5	< 66.6	< 62.2	< 97.6	< 198.6	< 118.6	< 123.8	< 119	< 127.9	< 208.8	< 228.2
Fenoxaprop ethyl	66441-23-4	< 1.1	< 0.4	< 0.6	< 1.3	< 2	< 0.8	< 2	< 2.1	3	< 3.8
Fenpropimorph	67564-91-4	10	13	15	< 1.3	5	3	2	< 0.8	< 3.5	< 1.5
Florasulam	145701-23-1	5.13	5.99	9.1	7.2	9.8	2.0	3.5	5.9	20.0	4.6
Fluroxypyr	69377-81-7	< 5.6	< 5.2	< 8.1	< 16.6	< 9.8	238	428	< 10.8	< 17.3	< 19.1
Fonofos	944-22-9	< 2.2	< 2.1	< 3.2	< 6.5	< 3.9	< 4.1	< 3.9	< 4.2	< 6.9	< 7.6
Isoproturon	34123-59-6	632	816	1,416	783	977	436	723	569	1,856	728
Malathion	121-75-5	< 5.6	14	23	18	16	47	214	31	89	35
Metamitron	41394-05-2	< 5.6	< 5.2	40	< 16.6	75	11	10	21	61	< 19.1
Metazachlor	67129-08-2	149	483	374	204	418	213	327	134	487	1,026
Metolachlor	51218-45-2	1,238	1,889	1,988	1,470	1,882	1,218	1,397	1,530	3,210	2,129
Metribuzin	21087-64-9	142	218	< 3.2	26	205	193	81	156	224	146
Parathion methyl	298-00-0	< 44.4	2,563	2,361	256	2,483	1,592	3,797	4,756	5,479	7,795
Pendimethalin	40487-42-1	95	172	162	113	200	88	95	69	97	53
Phosmet	732-11-6	< 2.2	< 5.2	17.1	8.5	16.9	6.1	5.9	12.7	11.7	< 19.1
Pirimicarb	23103-98-2	31	41	74	46	72	36	32	20	45	11
Prochloraz	67747-09-5	78	141	124	118	130	52	128	111	197	126
Propiconazole	60207-90-1	3,745	6,719	5,250	3,208	5,462	2,793	3,192	3,484	6,508	4,284
Pyrazon	216-920-2	67	87	175	195	296	120	191	208	831	412
Simazine	122-34-9	451	651	968	488	739	309	498	575	2,015	1,362
Tebuconazole	107534-96-3	3,114	4,490	3,331	2,864	3,911	2,458	3,508	4,169	7,473	5,174

	Temephos	3383-96-8	3.80	1.65	< 1	< 2	3.9	< 1.2	3.1	1.3	2.8	3.0
	Terbufos	13071-79-9	272	270	313	386	1,125	478	2,296	2,676	3,519	1,532
	Terbuthylazin	5915-41-3	1,029	1,481	2,708	1,885	2,409	1,320	1,873	1,818	7,241	3,198
	Tribenuron-methyl	101200-48-0	3.13	8.47	41.7	8.5	29.1	8.2	17.7	7.2	56.6	11.4
	Σ CUPs		17,543	28,746	33,152	20,492	32,509	17,067	27,033	28,373	67,140	41,961
Alkylphenols	Bisphenol A	80-05-7	5,426	5,771	6,821	6,207	7,862	17,471	27,537	12,921	14,894	14,449
	4-t-octylphenol	140-66-9	83	101	191	334	178	485	143	98	169	94
	4-nonylphenol	104-40-5	2,121	1,918	5,214	5,943	7,169	8,355	6,833	7,963	7,336	5,274
	Σ Alkylphenols		7,630	7,790	12,225	12,484	15,209	26,311	34,513	20,982	22,399	19,817
Steroids	Estriol	50-27-1	<8.93	<8.26	<25.86	<52.08	47	161	234	46	38	66
	17α-estradiol	57-91-0	<17.86	<16.52	<25.86	<52.08	<31.43	<32.78	<31.42	<33.76	<27.61	<30.48
	17β-estradiol	50-28-2	<17.86	<16.52	<25.86	<52.08	43	43	97	<33.76	57	56
	17α-ethinyl-estradiol	57-63-6	<25.86	<25.86	<25.86	<25.86	<25.86	<25.86	<25.86	<25.86	<25.86	<25.86
	Estrone	53-16-7	176	198	201	351	492	422	820	352	675	694
	Σ Steroids		176	198	201	351	582	626	1,151	398	770	816
PAHs	Naphthalene	91-20-3	129	426	1,152	3,531	<3	875	790	1,963	1,725	2,646
	Biphenyl	92-52-4	92	156	204	299	981	172	195	148	112	209
	Acenaphthylene	208-96-8	59	79	164	115	661	92	94	59	47	103
	Acenaphthene	83-32-9	72	129	175	140	2,445	195	125	72	30	155
	Fluorene	86-73-7	224	308	379	403	3,363	1,395	424	259	236	585
	Phenanthrene	85-01-8	1,121	1,507	2,601	2,048	5,136	10,235	2,098	1,948	1,458	3,403
	Anthracene	120-12-7	79	114	375	277	788	294	161	108	105	294
	Fluoranthene	206-44-0	1,972	2,134	1,967	2,504	3,633	1,542	1,832	1,515	1,152	1,953
	Pyrene	129-00-0	1,005	1,125	1,162	1,430	2,712	1,311	1,258	874	1,005	1,630
	Retene	483-65-8	482	439	328	272	605	98	128	127	179	123
	Benzo(b)fluorene	243-17-4	104	114	115	127	<12	119	116	80	51	99
	Benzonaphthothiophene	205-43-6	63	71	101	51	<7	58	64	40	51	65
	Benzo(ghi)fluoranthene	203-12-3	95	97	82	105	<5	75	89	85	81	91
	Cyclopenta(cd)pyrene	27208-37-3	<3	<2	<4	<7	<4	<5	<4	<5	<8	<9
	Benzo(a)anthracene	56-55-3	99	114	114	95	281	86	101	61	58	118



	Triphenylene	217-59-4	264	273	380	347	795	304	358	373	366	435
	Chrysene	218-01-9	189	211	231	166	1,012	278	169	143	120	142
	Benzo( <i>b</i> )fluoranthene	205-99-2	67	47	<12	<24	<14	<15	<14	<15	<25	<28
	Benzo( <i>j</i> )fluoranthene	205-82-3	36	36	<12	<24	<14	<15	<14	<16	<25	<28
	Benzo( <i>k</i> )fluoranthene	207-08-9	22	20	<13	<26	<16	<16	<16	<17	<27	<30
	Benzo( <i>e</i> )pyrene	192-97-2	64	62	64	74	<12	63	64	54	<21	<24
	Benzo( <i>a</i> )pyrene	50-32-8	7	11	<14	<28	<17	<18	<17	<18	<29	<33
	Perylene	198-55-0	18	19	<13	<26	<16	<17	<16	52	117	194
	Indeno(123 <i>cd</i> )pyrene	193-39-5	<8	<7	<11	<22	<13	<14	<13	<14	<23	<26
	Dibenzo( <i>ah</i> )anthracene	53-70-3	<9	<8	<13	<26	<16	<16	<16	<17	<27	<30
	Dibenzo( <i>ac</i> )anthracene	215-58-7	<9	<8	<13	<26	<16	<16	<16	<17	<27	<30
	Benzo( <i>ghi</i> )perylene	191-24-2	8	7	<7	<15	<9	<9	<9	<9	<15	<17
	Anthanthrene	191-26-4	<12	<11	<18	<36	<22	<23	<22	<23	<38	<42
	Coronene	191-07-1	<14	<13	<21	<42	<26	<27	<26	<27	<45	<49
	Σ PAHs		6,271	7,498	9,594	11,984	22,411	17,190	8,065	7,962	6,893	12,245
Sterols	Brassicasterol	474-67-9	5,465	3,631	1,253	1,927	3,042	2,293	2,479	3,925	1,519	1,166
	Campesterol	474-62-4	5,882	5,829	3,172	5,128	<196.44	3,988	5,480	4,972	7,294	6,127
	Desmosterol	313-04-2	977	1,135	1,042	<325.51	964	302	618	603	1,160	741
	Ergosterol	57-87-4	<66.98	324	<96.97	<195.31	<117.86	<122.93	<117.81	<126.58	<207.08	<228.6
	Fucoesterol	17605-67-3	2,044	1,204	1,422	<325.51	1,177	1,120	1,034	1,301	1,260	1,655
	Cholestanol	80-97-7	2,114	2,284	<161.62	<325.51	2,281	<204.89	<196.35	<210.97	<345.13	2,915
	Cholesterol	57-88-5	20,035	23,144	10,522	15,737	18,385	11,067	22,500	17,751	19,607	14,364
	Sitosterol	83-46-5	16,120	<20.65	5,990	10,776	9,077	6,421	6,775	8,936	7,298	12,106
	Spinasterol	481-18-5	9,786	7,531	2,196	7,967	7,497	3,159	2,577	6,274	5,788	<381.01
	Stigmastanol	19466-47-8	805	905	<161.62	<325.51	<196.44	889	<196.35	627	<345.13	<381.01
	Stigmasterol	83-48-7	8,782	6,791	4,286	4,429	7,572	5,514	5,995	1,821	5,072	11,735
	Σ Sterols		72,010	52,776	29,885	45,964	49,994	34,753	47,458	46,209	48,996	50,809
Flavonoids	Apigenin	520-36-5	18.98	18.07	12.9	14.6	24.6	34.8	20.6	21.1	19.0	19.1
	Biochanin A	491-80-5	30.70	35.11	22.6	9.8	11.8	11.3	3.9	<0.04	3.5	3.8
	Coumestrol	479-13-0	<0.22	<0.21	<0.32	<0.65	<0.39	<0.41	<0.39	<0.42	<0.69	<0.76

Daidzein	486-66-8	73	108	93	76	88	488	595	162	255	404
Equol	94105-90-5	207	250	318	150	138	1,332	821	268	2,571	9,449
Formononetin	485-72-3	268	302	328	270	177	163	231	141	178	389
Genistein	446-72-0	33	53	47	21	268	585	159	28	47	65
Naringenin	266-769-1	166	188	132	94	98	302	160	72	138	<0.08
$\Sigma$ Flavonoids		798	955	954	636	806	2,916	1,990	693	3,211	10,329

---

**Tab. S4 Sum molar concentrations of groups of chemicals detected in samples from dynamic passive sampling**

**1-8 samples from mobile sampling of river stretches; SR silicone rubber sampler, ED Empore™ disc sampler; No. number of analyzed chemicals in the category; <LOD concentrations below the limit of detection; for individual chemical identification see SI.**

Sampler	Chemicals (pmol L <sup>-1</sup> )	No.	S1	S2	1	2	3	4	5	6	7	8
ED	ATBs	31	48.34	46.17	43.51	89.63	87.56	66.47	81.53	46.24	122.21	108.15
	Cardiovascular	15	31.57	33.97	26.78	45.09	48.20	24.97	26.34	15.84	23.62	22.23
	Psychoactive	30	98.12	82.94	69.93	114.01	103.54	70.08	94.72	98.63	132.72	114.32
	Antihistamins	8	0.92	0.24	0.90	<LOD	0.30	1.21	0.84	<LOD	2.03	<LOD
	Antifungals	8	11.32	13.10	11.49	22.26	23.33	22.01	29.93	28.18	30.63	24.18
	Antidiabetics	4	0.06	0.01	0.07	0.06	0.19	0.05	0.03	0.03	0.18	0.04
	Statins	4	0.57	0.86	0.76	0.97	3.44	2.76	3.30	0.98	1.63	0.89
	Other pharm. <sup>1</sup>	8	0.89	0.76	<LOD	<LOD	0.37	1.63	<LOD	0.90	<LOD	<LOD
	CUPs	40	67.55	107.47	132.15	81.52	126.65	65.13	102.24	106.26	266.67	160.25
	Alkylphenols	3	0.65	0.73	0.74	1.30	2.14	2.28	4.20	1.46	2.84	3.00
	PAHs	29	33.80	34.47	54.47	55.78	67.84	116.80	152.33	93.21	99.35	87.69
	Steroids	5	31.78	39.30	53.14	71.50	123.13	96.11	44.11	46.47	39.98	71.16
	Phytosterols	11	178.93	132.94	74.58	117.35	124.81	86.46	119.31	115.55	122.70	126.43
	Flavonoids	8	3.05	3.66	3.69	2.44	3.06	11.47	7.85	2.72	13.04	42.36
SR	PAHs	29	100.66	133.97	108.06	130.08	315.96	223.61	93.50	70.97	87.78	128.33
	PCBs	7	0.72	0.80	0.56	0.99	0.58	0.57	1.32	0.54	1.11	1.05
	OCPs	12	0.63	0.75	0.53	0.82	1.28	1.07	1.20	1.55	2.68	6.06
	BDEs	9	0.01	0.02	0.06	0.01	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
	NBFRs	15	0.01	0.02	0.04	0.04	0.03	0.02	0.03	0.03	0.01	0.02
	Musks	6	144.03	215.80	46.69	50.77	74.10	138.94	<LOD	<LOD	<LOD	<LOD
	Alkylphenols	3	13.23	15.93	26.09	28.45	36.12	45.47	38.60	16.02	33.03	28.28
	Alkylphosphates	14	464.26	610.68	316.42	418.14	369.38	269.61	358.08	517.79	495.86	528.22

<sup>1</sup> furosemide, alfuzosin, naloxone, ranitidine

**Tab. S5 Relative effect potencies (REPs) of detected compounds**

REPs that were used for calculation of chemical data-based bioanalytical equivalents (BEQ<sub>chem</sub>) were obtained either from experiments from the current study or from literature. Literature was preferentially searched for REP values specific for the bioassays employed in the current study. Where no such REPs were available, we used data from analogical bioassays as indicated. When there was more than one relevant literature source available, geometrical mean of REP value was calculated.

Assay	Activation of AhR			Activation of PXR			Activation of ER			Inhibition of AR			Oxidative stress response			Activation of p53		
No. chemicals	23			27			26			22			11			12		
Reference compound	2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (TCDD)			SR 12813			17β-Estradiol			Flutamide			tert-Butylhydroquinone (tBHQ)			Mitomycin		
Chemical	REP	based on	Chemical	REP	based on	Chemical	REP	based on	Chemical	REP	based on	Chemical	REP	based on	Chemical	REP	based on	
4-Nonylphenol	1.0×10 <sup>-8</sup>	EC <sub>10</sub> <sup>1,4, a</sup>	17β-Estradiol	3.0×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	17β-Estradiol	1.0×10 <sup>+0</sup>	EC <sub>25</sub> <sup>3,10, i</sup>	4-Nonylphenol	0.02	IC <sub>20</sub> <sup>3,12, i</sup>	4-Nonylphenol	2.9×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, a</sup>	17β-Estradiol	9.8×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,15, v</sup>	
Benzo(a)anthracene	2.6×10 <sup>-5</sup>	EC <sub>25</sub> <sup>1,5, b, c, d</sup>	4-Nonylphenol	3.0×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	2,4'-DDE	6.3×10 <sup>-6</sup>	EC <sub>50</sub> <sup>3,10, j</sup>	4-t-Octylphenol	0.34	IC <sub>20</sub> <sup>3,12, i</sup>	Atenolol	1.5×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Apigenin	3.2×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(a)pyrene	1.5×10 <sup>-4</sup>	EC <sub>20</sub> <sup>1,6, e</sup>	4-t-Octylphenol	5.8×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	2,4'-DDT	1.7×10 <sup>-5</sup>	EC <sub>25</sub> <sup>3,10, i</sup>	Acetochlor	0.62	IC <sub>20</sub> <sup>3,12, r</sup>	Atorvastatin	3.5×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Benzo(a)pyrene	6.6×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(b)fluoranthene	4.6×10 <sup>-4</sup>	EC <sub>25</sub> <sup>1,5, b, c, d</sup>	Bezafibrate	6.2×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,13, o</sup>	4,4'-DDD	6.8×10 <sup>-7</sup>	EC <sub>20</sub> <sup>3,10, k</sup>	Alachlor	0.83	IC <sub>20</sub> <sup>3,12, r</sup>	Atrazine	1.0×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Benzo(b)fluoranthene	2.8×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(e)pyrene	6.6×10 <sup>-7</sup>	EC <sub>25</sub> <sup>1,5, b, c</sup>	Bisphenol A	5.8×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	4,4'-DDE	7.0×10 <sup>-7</sup>	EC <sub>50</sub> <sup>3,10, j</sup>	Benzo(a)pyrene	0.16	IC <sub>20</sub> <sup>3,12, a</sup>	Benzo(a)pyrene	4.8×10 <sup>+0</sup>	EC <sub>IR1.5</sub> <sup>3,14, a</sup>	Benzo(e)pyrene	1.7×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(ghi)perylene	2.6×10 <sup>-6</sup>	EC <sub>25</sub> <sup>1,5, b</sup>	Carbamazepine	4.6×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,13, o</sup>	4,4'-DDT	1.3×10 <sup>-6</sup>	EC <sub>50</sub> <sup>3,10, l</sup>	Bisphenol A	1.20	IC <sub>20</sub> <sup>3,12, a</sup>	Benzo(b)fluoranthene	8.4×10 <sup>+0</sup>	EC <sub>IR1.5</sub> <sup>3,14, a</sup>	Benzo(k)fluoranthene	4.0×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(j)fluoranthene	5.5×10 <sup>-4</sup>	EC <sub>25</sub> <sup>1,5, b, c</sup>	Chlorpyrifos	3.1×10 <sup>-3</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	4-Nonylphenol	3.3×10 <sup>-6</sup>	EC <sub>20</sub> <sup>3,10, k</sup>	Bromocriptine mesylate	1.68	IC <sub>20</sub> <sup>3,12, r</sup>	Bisphenol A	1.5×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, a</sup>	Biochanin A	5.6×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Benzo(k)fluoranthene	3.1×10 <sup>-3</sup>	EC <sub>25</sub> <sup>1,5, b, c, d</sup>	Clotrimazole	4.3×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	4-t-Octylphenol	1.1×10 <sup>-4</sup>	EC <sub>25</sub> <sup>3,10, i</sup>	Chloropyrifos	0.01	IC <sub>20</sub> <sup>3,12, a</sup>	Citalopram	3.0×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Bisphenol A	7.1×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Bisphenol A	1.4×10 <sup>-7</sup>	EC <sub>10</sub> <sup>1,6, e</sup>	Diazinon	1.1×10 <sup>-2</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	anthracene	7.9×10 <sup>-7</sup>	EC <sub>25</sub> <sup>3,10, n</sup>	Clotrimazole	2.23	IC <sub>20</sub> <sup>3,12, r</sup>	Genistein	2.7×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, a</sup>	Carbendazim	7.3×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, o</sup>	
Carbendazim	1.2×10 <sup>-8</sup>	EC <sub>10</sub> <sup>1,6, e</sup>	Diuron	4.3×10 <sup>-4</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	Benzo(a)pyrene	4.7×10 <sup>-6</sup>	EC <sub>10</sub> <sup>3,10, a</sup>	Diazinon	0.02	IC <sub>20</sub> <sup>3,12, e</sup>	Metoprolol	3.0×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Chlorprothixene	1.2×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Chlorpyrifos	1.1×10 <sup>-7</sup>	EC <sub>10</sub> <sup>1,4, a</sup>	Estrone	4.1×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	fluoranthene	3.4×10 <sup>-6</sup>	EC <sub>10</sub> <sup>3,10, a</sup>	Fenofibrate	0.59	IC <sub>20</sub> <sup>3,12, r</sup>	Propiconazole	2.7×10 <sup>-2</sup>	EC <sub>IR1.5</sub> <sup>3,14, u</sup>	Genistein	1.2×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,15, o</sup>	
Chrysene	6.5×10 <sup>-5</sup>	EC <sub>25</sub> <sup>1,5, b, c</sup>	Fenofibrate	2.5×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	Biochanin A	8.8×10 <sup>-5</sup>	EC <sub>50</sub> <sup>3,11, m</sup>	Ketoconazole	0.98	IC <sub>20</sub> <sup>3,12, r</sup>				Prochloraz	7.2×10 <sup>-4</sup>	EC <sub>IR1.5</sub> <sup>3,15, r</sup>	
Cyclopenta(cd)pyrene	6.5×10 <sup>-7</sup>	EC <sub>25</sub> <sup>1,5, c</sup>	Genistein	1.2×10 <sup>-2</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	Bisphenol A	4.4×10 <sup>-5</sup>	EC <sub>20</sub> <sup>3,10, k</sup>	Maprotiline	0.51	IC <sub>20</sub> <sup>3,12, r</sup>							
Daidzein	1.1×10 <sup>-6</sup>	LOEC <sup>2,8, f</sup>	Isoproturon	1.6×10 <sup>-3</sup>	EC <sub>IR1.5</sub> <sup>3,13, o</sup>	Chlorpyrifos	2.8×10 <sup>-7</sup>	EC <sub>10</sub> <sup>3,10, a</sup>	Memantine	0.58	IC <sub>20</sub> <sup>3,12, r</sup>							
Diazinon	1.0×10 <sup>-8</sup>	EC <sub>20</sub> <sup>1,6, a</sup>	Metolachlor	5.9×10 <sup>-2</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	Chrysene	2.2×10 <sup>-6</sup>	EC <sub>25</sub> <sup>3,10, q</sup>	Metolachlor	0.02	IC <sub>20</sub> <sup>3,12, e</sup>							
Diuron	1.8×10 <sup>-6</sup>	EC <sub>20</sub> <sup>2,9, h</sup>	o,p'-DDT	1.7×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	Daidzein	6.4×10 <sup>-5</sup>	EC <sub>50</sub> <sup>3,10, o</sup>	o,p'-DDD	0.31	IC <sub>20</sub> <sup>3,12, r</sup>							
Equol	1.0×10 <sup>-6</sup>	LOEC <sup>2,8, f</sup>	p,p'-DDD	4.1×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	Diazinon	2.9×10 <sup>-7</sup>	EC <sub>10</sub> <sup>3,10, a</sup>	Prochloraz	0.43	IC <sub>20</sub> <sup>3,12, r</sup>							
Fluoranthene	9.3×10 <sup>-7</sup>	EC <sub>25</sub> <sup>1,5, c</sup>	p,p'-DDE	3.0×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	hydrochloride	6.3×10 <sup>-5</sup>	EC <sub>10</sub> <sup>3,11, p</sup>	Propiconazole	0.07	IC <sub>20</sub> <sup>3,12, a</sup>							
Genistein	7.9×10 <sup>-8</sup>	EC <sub>10</sub> <sup>1,6, a</sup>	p,p'-DDT	2.9×10 <sup>-2</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	Equol	2.8×10 <sup>-4</sup>	EC <sub>50</sub> <sup>3,10, i</sup>	Triphenyl phosphate	0.03	IC <sub>20</sub> <sup>3,12, a</sup>							
Prochloraz	2.3×10 <sup>-6</sup>	EC <sub>20</sub> <sup>2,9, g</sup>	PCB 101	3.4×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	Estriol	1.7×10 <sup>-1</sup>	EC <sub>25</sub> <sup>3,10, i</sup>	Temephos	0.41	IC <sub>20</sub> <sup>3,12, r</sup>							

Propiconazole	1.9×10 <sup>-5</sup>	EC <sub>50</sub> <sup>2,8, h</sup>	PCB 118	7.0×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, t</sup>	Estrone	1.1×10 <sup>-1</sup>	EC <sub>IR1.5</sub> <sup>3,10, a</sup>	TDCPP <sup>w</sup> Verapamil hydrochloride	0.33	IC <sub>20</sub> <sup>3,12, r</sup>
Pyrene	4.1×10 <sup>-6</sup>	EC <sub>25</sub> <sup>1,5, b,c</sup>	PCB 138	5.3×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	Formononetin	2.3×10 <sup>-4</sup>	EC <sub>50</sub> <sup>3,11, m</sup>		0.27	IC <sub>20</sub> <sup>3,12, r</sup>
Terbutylazine	1.3×10 <sup>-5</sup>	EC <sub>50</sub> <sup>2,8, h</sup>	PCB 153	6.0×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	Genistein	2.0×10 <sup>-4</sup>	EC <sub>10</sub> <sup>3,10, a</sup>			
			PCB 180	6.0×10 <sup>-3</sup>	EC <sub>20</sub> <sup>3,13, s</sup>	Naringenin	2.2×10 <sup>-5</sup>	EC <sub>50</sub> <sup>3,11, m</sup>			
			Propiconazole	5.4×10 <sup>-3</sup>	EC <sub>10</sub> <sup>3,13, a</sup>	Terbutylazine	1.6×10 <sup>-7</sup>	EC <sub>10</sub> <sup>3,10, a</sup>			
			Terbutylazine Triphenyl phosphate	4.1×10 <sup>-3</sup>	EC <sub>10</sub> <sup>3,13, s</sup>	Triphenyl phosphate	1.4×10 <sup>-6</sup>	EC <sub>10</sub> <sup>3,10, a</sup>			
			1.7×10 <sup>-2</sup>	EC <sub>10</sub> <sup>3,13, a</sup>							

<sup>1</sup> rat-based cell line	<sup>5</sup> H4EII- <i>luc</i>	<sup>9</sup> Hepa1c1c7	<sup>13</sup> HG5LN-hPXR	<sup>a</sup> (Neale et al., 2017)	<sup>e</sup> (current study)	<sup>i</sup> (Kinani et al., 2010)	<sup>m</sup> (Procházková et al., 2017)	<sup>q</sup> (Lam et al., 2017)	<sup>u</sup> (Escher et al., 2013)
<sup>2</sup> mouse-based cell line	<sup>6</sup> H4G1.1c2	<sup>10</sup> MELN	<sup>14</sup> AREc32	<sup>b</sup> (Larsson et al., 2012)	<sup>f</sup> (Denison et al., 1998)	<sup>j</sup> (Pillon et al., 2005)	<sup>n</sup> (Machala et al., 2001)	<sup>r</sup> (US EPA ToxCast)	<sup>v</sup> (König et al., 2016)
<sup>3</sup> human-based cell line	<sup>7</sup> H4L1.1c4	<sup>11</sup> HeLa9903	<sup>15</sup> p53RE-bla	<sup>c</sup> (M Machala et al., 2001)	<sup>g</sup> (Takeuchi et al., 2008)	<sup>k</sup> (Creusot et al., 2013)	<sup>o</sup> (Neale et al., 2015a)	<sup>s</sup> (Creusot et al., 2010)	<sup>w</sup> Tris(1,3-Dichloro-2-propyl)phosphate
<sup>4</sup> DR-CALUX	<sup>8</sup> Hepa1.12cR	<sup>12</sup> MDA-kb2		<sup>d</sup> (Behnisch et al., 2003)	<sup>h</sup> (Ghisari et al., 2015)	<sup>l</sup> (Leusch et al., 2010)	<sup>p</sup> (Ceger et al., 2015)	<sup>t</sup> (Creusot, 2011)	<sup>x</sup> Tris(2-butoxyethyl)phosphate

## References

- Allan, I.J., Harman, C., Ranneklev, S.B., Thomas, K. V., Grung, M., 2013. Passive sampling for target and nontarget analyses of moderately polar and nonpolar substances in water. *Environ. Toxicol. Chem.* 32, 1718–1726.
- Behnisch, P.A., Hosoe, K., Sakai, S., 2003. Brominated dioxin-like compounds: in vitro assessment in comparison to classical dioxin-like compounds and other polyaromatic compounds. *Environ. Int.* 29, 861–877.
- Booij, K., Smedes, F., 2010. An improved method for estimating in situ sampling rates of nonpolar passive samplers. *Environ. Sci. Technol.* 44, 6789–6794.
- Ceger, P., Allen, D., Huang, R., Xia, M., Casey, W., 2015. Performance of the BG1Luc ER TA method in a qHTS format. *ALTEX* 32, 287–296.
- Creusot, N., 2011. Contribution de l'approche effect directed analysis à l'identification de perturbateurs endocriniens dans les milieux aquatiques.
- Creusot, N., Budzinski, H., Balaguer, P., Kinani, S., Porcher, J.-M., Aït-Aïssa, S., 2013. Effect-directed analysis of endocrine-disrupting compounds in multi-contaminated sediment: identification of novel ligands of estrogen and pregnane X receptors. *Anal. Bioanal. Chem.* 405, 2553–2566.
- Creusot, N., Kinani, S., Balaguer, P., Tapie, N., LeMenach, K., Maillot-Maréchal, E., Porcher, J.-M., Budzinski, H., Aït-Aïssa, S., 2010. Evaluation of an hPXR reporter gene assay for the detection of aquatic emerging pollutants: screening of chemicals and application to water samples. *Anal. Bioanal. Chem.* 396, 569–583.
- Denison, M.S., Phelan, D., Winter, G.M., Ziccardi, M.H., 1998. Carbaryl, a carbamate insecticide, is a ligand for the hepatic Ah (dioxin) receptor. *Toxicol. Appl. Pharmacol.* 152, 406–14.
- Escher, B.I., Dutt, M., Maylin, E., Tang, J.Y.M., Toze, S., Wolf, C.R., Lang, M., 2012. Water quality assessment using the AREc32 reporter gene assay indicative of the oxidative stress response pathway. *J. Environ. Monit.* 14, 2877.
- Escher, B.I., van Daele, C., Dutt, M., Tang, J.Y.M., Altenburger, R., 2013. Most oxidative stress response in water samples comes from unknown chemicals: the need for effect-based water quality trigger values. *Environ. Sci. Technol.* 47, 7002–11.
- Freyberger, A., Schmuck, G., 2005. Screening for estrogenicity and anti-estrogenicity: a critical evaluation of an MVLN cell-based transactivation assay. *Toxicol. Lett.* 155, 1–13.
- Ghisari, M., Long, M., Tabbo, A., Bonefeld-Jørgensen, E.C., 2015. Effects of currently used pesticides and their mixtures on the function of thyroid hormone and aryl hydrocarbon receptor in cell culture. *Toxicol. Appl. Pharmacol.* 284, 292–303.
- Jin, L., Gaus, C., Escher, B.I., 2015. Adaptive Stress Response Pathways Induced by Environmental Mixtures of Bioaccumulative Chemicals in Dugongs. *Environ. Sci. Technol.* 49, 6963–6973.
- Kinani, S., Bouchonnet, S.S., Creusot, N., Bourcier, S., Balaguer, P., Porcher, J.-M., Aït-Aïssa, S., Ait-Aïssa, S., 2010. Bioanalytical characterisation of multiple endocrine- and dioxin-like activities in sediments from reference and impacted small rivers. *Environ. Pollut.* 158, 74–83.
- König, M., Escher, B.I., Neale, P.A., Krauss, M., Hilscherová, K., Novák, J., Teodorović, I., Schulze, T., Seidensticker, S., Kamal Hashmi, M.A., Ahlheim, J., Brack, W., 2016. Impact of untreated wastewater on a major European river evaluated with a combination of in vitro bioassays and chemical analysis. *Environ. Pollut.*
- Lam, M.M., Engwall, M., Denison, M.S., Larsson, M., 2017. Methylated polycyclic aromatic hydrocarbons and/or their metabolites are important contributors to the overall estrogenic activity of polycyclic aromatic hydrocarbon-contaminated soils. *Environ. Toxicol. Chem.*
- Larsson, M., Orbe, D., Engwall, M., 2012. Exposure time-dependent effects on the relative potencies and additivity of PAHs in the Ah receptor-based H4IIE-luc bioassay. *Environ. Toxicol. Chem.* 31, 1149–1157.
- Leusch, F.D.L., de Jager, C., Levi, Y., Lim, R., Puijker, L., Sacher, F., Tremblay, L. a, Wilson, V.S., Chapman, H.F., 2010.

- Comparison of five in vitro bioassays to measure estrogenic activity in environmental waters. *Environ. Sci. Technol.* 44, 3853–60.
- Machala, M., Ciganek, M., Bláha, L., Minksová, K., Vondráček, J., 2001. Aryl hydrocarbon receptor-mediated and estrogenic activities of oxygenated polycyclic aromatic hydrocarbons and azaarenes originally identified in extracts of river sediments. *Environ. Toxicol. Chem.* 20, 2736–2743.
- Machala, M., Vondracek, J., Blaha, L., Ciganek, M., Neca, J. V, Vondráček, J., Bláha, L., 2001. Aryl hydrocarbon receptor-mediated activity of mutagenic polycyclic aromatic hydrocarbons determined using in vitro reporter gene assay. *Mutat. Res. Toxicol. Environ. Mutagen.* 497, 49–62.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65, 55–63.
- Nagy, S.R., Sanborn, J.R., Hammock, B.D., Denison, M.S., 2002. Development of a Green Fluorescent Protein-Based Cell Bioassay for the Rapid and Inexpensive Detection and Characterization of Ah Receptor Agonists. *Toxicol. Sci.* 65, 200–210.
- Neale, P.A., Ait-Aissa, S., Brack, W., Creusot, N., Denison, M.S., Deutschmann, B., Hilscherova, K., Hollert, H., Krauss, M., Novák, J., Schulze, T., Seiler, T.B., Serra, H., Shao, Y., Escher, B.I., 2015a. Linking in vitro effects and detected organic micropollutants in surface water using mixture toxicity modeling. *Environ. Sci. Technol.* 49, 14614–14624.
- Neale, P.A., Altenburger, R., Ait-Aïssa, S., Brion, F., Busch, W., de Aragão Umbuzeiro, G., Denison, M.S., Du Pasquier, D., Hilscherová, K., Hollert, H., Morales, D.A., Novák, J., Schlichting, R., Seiler, T.-B., Serra, H., Shao, Y., Tindall, A.J., Tollefsen, K.E., Williams, T.D., Escher, B.I., 2017. Development of a bioanalytical test battery for water quality monitoring: Fingerprinting identified micropollutants and their contribution to effects in surface water. *Water Res.* 123, 734–750.
- Neale, P.A., Stalter, D., Tang, J.Y.M., Escher, B.I., 2015b. Bioanalytical evidence that chemicals in tattoo ink can induce adaptive stress responses. *J. Hazard. Mater.* 296, 192–200.
- Pernica, M., Poloucká, P., Seifertová, M., Šimek, Z., 2015. Determination of alkylphenols in water samples using liquid chromatography–tandem mass spectrometry after pre-column derivatization with dansyl chloride. *J. Chromatogr. A* 1417, 49–56.
- Pillon, A., Boussioux, A.-M.M., Escande, A., Ait-Aïssa, S., Gomez, E., Fenet, H.H., Ruff, M., Moras, D., Vignon, F., Duchesne, M.-J.J., Casellas, C., Nicolas, J.-C.C., Balaguer, P., 2005. Binding of estrogenic compounds to recombinant estrogen receptor- $\alpha$ : Application to environmental analysis. *Environ. Health Perspect.* 113, 278–284.
- Procházková, T., Sychrová, E., Javůrková, B., Večerková, J., Kohoutek, J., Lepšová-Skácelová, O., Bláha, L., Hilscherová, K., 2017. Phytoestrogens and sterols in waters with cyanobacterial blooms - Analytical methods and estrogenic potencies. *Chemosphere* 170, 104–112.
- Rusina, T.P., Smedes, F., Koblížková, M., Klanová, J., 2010. Calibration of Silicone Rubber Passive Samplers: Experimental and Modeled Relations between Sampling Rate and Compound Properties. *Environ. Sci. Technol.* 44, 362–367.
- Smedes, F., Booij, K., 2012. Guidelines for passive sampling of hydrophobic contaminants in water using silicone rubber samplers. ICES Techniques in Marine Environmental Sciences No. 52., ICES TECHNIQUES IN MARINE ENVIRONMENTAL SCIENCES. International Council for the Exploration of the Sea, Denmark.
- Smedes, F., Geertsma, R.W., Van Der Zande, T., Booij, K., 2009. Polymer-water partition coefficients of hydrophobic compounds for passive sampling: Application of cosolvent models for validation. *Environ. Sci. Technol.* 43, 7047–7054.
- Takeuchi, S., Iida, M., Yabushita, H., Matsuda, T., Kojima, H., 2008. In vitro screening for aryl hydrocarbon receptor agonistic activity in 200 pesticides using a highly sensitive reporter cell line, DR-EcoScreen cells, and in vivo mouse liver cytochrome P450-1A induction by propanil, diuron and linuron. *Chemosphere* 74, 155–65.

- Vrana, B., Smedes, F., Allan, I., Rusina, T., Okonski, K., Hilscherová, K., Novák, J., Tarábek, P., Slobodník, J., n.d. Dynamic mobile passive sampling of trace organic compounds: evaluation of sampler performance in the Danube river. *Sci. Total Environ* - submitted.
- Wang, X.J., Hayes, J.D., Wolf, C.R., 2006. Generation of a Stable Antioxidant Response Element-Driven Reporter Gene Cell Line and Its Use to Show Redox-Dependent Activation of Nrf2 by Cancer Chemotherapeutic Agents. *Cancer Res.* 66, 10983–10994.